



## **BiotechLec: an interactive guide of commercial lectins for glycobiology and biomedical research applications**

Boris Schnider, Francisco Lorenzo Escudero, Anne Imberty, Frédérique Lisacek

### **► To cite this version:**

Boris Schnider, Francisco Lorenzo Escudero, Anne Imberty, Frédérique Lisacek. BiotechLec: an interactive guide of commercial lectins for glycobiology and biomedical research applications. *Glycobiology*, 2023, 33, pp.684-686. <10.1093/glycob/cwad034>. <hal-04100786>

**HAL Id: hal-04100786**

**<https://hal.science/hal-04100786v1>**

Submitted on 18 May 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

# BiotechLec: an interactive guide of commercial lectins for glycobiology and biomedical research applications

Boris Schneider<sup>1,2</sup>, Francisco Lorenzo Escudero<sup>1</sup>, Anne Imberty<sup>3\*</sup> and Frédérique Lisacek<sup>1,2,4\*</sup>

1. Proteome Informatics Group, SIB Swiss Institute of Bioinformatics, CH-1211 Geneva, Switzerland
2. Computer Science Department, University of Geneva, CH-1227 Geneva, Switzerland
3. University Grenoble Alpes, CNRS, CERMAV, 38000 Grenoble, France
4. Section of Biology, University of Geneva, CH-1205 Geneva, Switzerland

\* Authors for correspondence: Anne Imberty ([anne.imberty@cermav.cnrs.fr](mailto:anne.imberty@cermav.cnrs.fr)) & Frédérique Lisacek ([frederique.lisacek@sib.swiss](mailto:frederique.lisacek@sib.swiss))

## Keywords

Lectins, biotechnology, database, carbohydrate-binding, glycosylation

## Abstract

For decades lectins have been used as probes in glycobiology and this usage has gradually spread to other domains of Life Science. Nowadays, researchers investigate glycan recognition with lectins in diverse biotechnology and clinical applications addressing key questions regarding binding specificity. The latter is documented in scattered and heterogeneous sources and this situation begs for a centralized and easy-access reference. To address this need, an on-line solution called BiotechLec (<https://www.unilectin.eu/biotechlec>) is proposed in a new section of UniLectin, a platform dedicated to lectin molecular knowledge.

## Introduction

Because of their specificity for glycans, lectins are widely used as biotechnological tools (Cummings et al. 2022; Lis and Sharon 1986). Their application extends further than the domain of glycobiology since lectins are effective markers in cell typing and as such, frequently used in immunology, cancer research and clinical microbiology, to name only a few areas. One of the oldest usages of lectins is testing the agglutination and precipitation of glycoconjugates, cells, and membrane vesicle preparations, the first application of which is blood group typing by erythrocytes hemagglutination in 1953 (Morgan and Watkins

1953). Lectins provide further resources: they can be immobilized and utilized for the purification of glycoproteins, they are also classically employed to characterize cell-surface glycoconjugates through histochemistry or flow cytometry with cell sorting. For example, sorting residual human pluripotent stem cells from cell cultures is achieved with BC2L-CNt, a bacterial lectin (Haramoto et al. 2020). Lectins are handy in enzymology for assaying specific glycosyltransferases and glycosidases. In recent years, they have been immobilized on microarrays or other biosensors, to characterize the glycan moiety of glycoproteins, cell extracts, or whole cells (Ribeiro and Mahal 2013). They are also used for cell-type specific intracellular

delivery of cargo, for example by conjugation to nanodiamonds in brain cell imaging (Ghanimi Fard et al. 2022).

Lectins are often preferred to other glycan-binding proteins, such as antibodies offering higher specificity but usually costlier. Compared to carbohydrate-binding modules (CBMs), their multi-valency makes them effective reagents for affinity chromatography of glycoproteins and glycans and histology biomarkers. Traditionally, lectins made available in catalogues of (bio)chemicals have been purified from seeds and plant materials. However, an increasing number is now obtained from fungi and animals, as well as bacteria and viruses. While most of the commercial lectins are purified from natural sources, they can also be produced recombinantly, which offers advantages of purity and reproducibility and opens other engineering possibilities.

Such a wide and diverse range of applications together with a growing collection of solutions may become a source confusion for scientists addressing specific carbohydrate-binding related issues. The purpose of the present note is therefore to describe a means of supporting researchers in selecting the most suitable lectin(s) to solve the corresponding carbohydrate-binding questions.

## What is BiotechLec?

In 2018, we launched the UniLectin portal ([unilectin.eu](https://www.unilectin.eu)) with the ultimate aim of bringing the broadest coverage of lectin molecular knowledge. UniLectin is dedicated to the classification, curation and prediction of lectins (Imberty et al. 2021). The platform is composed of several modules, starting with Unilectin3D, a curated database with more than 2400 3D-structures, which sets the basis for classification (Bonnardel et al. 2019b). The other modules provide information on predicted lectomes (i) from all available translated genomes – LectomeXplore (Bonnardel et al. 2021) –, (ii) from specific origin – MycoLec (Lebreton et al. 2021) – and (iii) reflecting structural families – PropLec and TrefLec (Bonnardel et al. 2019a; Notova et al.

2022). Here, we introduce the BiotechLec module (<https://www.unilectin.eu/biotechlec>) that centralizes information about lectins that are commercially available and classically used for research. A recent review summarizing the activity of 57 of such lectins (Bojar et al. 2022) motivated the design of BiotechLec as an interactive complement. A larger screen of on-line catalogues (Vector Laboratories, Glycodiag, Elicityl and Wako companies) led to identify 77 lectins that were first organized in a tabular format, reflecting the classification adopted across the whole UniLectin platform and their main glycan specificity.


While a large majority is purified from plants (66%), the table shows that fungal (13%) and bacterial (10%) origins are also represented. Animal lectins are less commonly used and they originate mostly from invertebrates (snail, slug and lobster) or fish (eel). The consequence is that many of the listed lectins belong to a legume lectin class. Conversely, some classes are not represented such as I-type lectins, mainly found in mammals so far. It should be noted that some human lectins are available commercially, but they are not included in the database since they are not classically used for purification or diagnostic purposes.

Glycan specificity is represented by the monosaccharides that are used for eluting the lectin, as listed in catalog or literature, even though the specificities may involve more complex glycan sequences. Monosaccharides are depicted in the widely adopted Symbol Nomenclature for Glycans (SNFG) (Neelamegham et al. 2022). Galactose and GalNAc are overrepresented in the specificity, while sialic acids (NeuAc and NeuGc) have only a few binding representatives. The double Gal/GlcNAc specificity from two mushroom lectins (XCL and ABA) is explained by the occurrence of two distinct binding sites with different specificities on each monomer in the fungal fruit-body lectin class (Carrizo et al. 2005).

## How to explore BiotechLec

A three-colour code was adopted to characterize each item, symbolized by the lectin abbreviated name, in the table. A green box corresponds to a curated UniLectin entry that includes 3D-structural information while a blue box points to entries with amino acid sequence information and prediction based on similarity (the quality of this prediction is evaluated with a score calculated by the LectomeXplore engine (Bonnardel et al. 2021)). Three orange boxes are lectins lacking sequence information and therefore not linked to any UniLectin entry (all UniLectin entries are associated with an amino acid sequence and a link to a protein database). Each coloured box of the BiotechLec dataset is clickable. Information is provided in a small pop-up table. In the first column, the biological origin of the lectin is described through the species name (with link to the NCBI taxonomy browser) and the common name. Information on the protein includes the UniProt name and accession number ([www.uniprot.org](http://www.uniprot.org)), amino acid sequence length and reference to the InterPro database of protein families ([www.ebi.ac.uk/interpro](http://www.ebi.ac.uk/interpro)). For the small number of lectins lacking a protein sequence (not published or deposited) a UniProt reference is proposed nonetheless. It is chosen as the most likely one, considering the information available and not dismissing the possibility of isolectins. Corrections and additions are easily made and feedback on entries is welcome. The second column of the pop-up table contains information

from (i) the Protein Data Bank ([www.rcsb.org](http://www.rcsb.org)), when the 3D structure is available, (ii) the Unilectin portal and associated classification, and (iii) websites dedicated to the analysis of lectin specificity. The Unilectin entry directs either to the UniLectin3D module of curated lectins for which structural data are available, or to the LectomeXplore module of predicted lectins displaying sequence alignment with a class representative. When several crystal structures are available, the one with the most representative ligand, i.e. a natural ligand with documented high affinity, has been selected, but all other structures are available from UniLectin3D through the Unilectin link. It should be noted that when no crystal structure is available, a model generated by AlphaFold, is available in the corresponding UniProt entry. This is of interest for example for the tomato lectin (LEA) that is widely utilized for labeling vasculature but could never be crystallized and that appears to be modelled with occurrence of five hevein domains. Finally, glycan array data are available, either as raw data Consortium for Functional Genomics (<http://www.functionalglycomics.org>) or as analyzed results from CarboGrove ([carbogrove.org](http://carbogrove.org)) (Klamer et al. 2022). For both websites, one representative entry was selected to provide a clear view of oligosaccharide specificity, with concentrations low enough to avoid cross-reactivity and high enough for strong signal/noise ratio.



**Figure 1 :** Rows of the table are lectin fold classes defined in UniLectin. Columns are defined by monosaccharides best characterising the lectin specificity and pictured with the SNFG notation (<https://www.ncbi.nlm.nih.gov/glycans/snfg.html>). A table cell is filled when the fold of a commercial lectin matches a given glycan specificity. The distribution is uneven. For example, 11 lectins adopting the L-type legume fold bind to GalNAc. The colour code is green for curated and blue for predicted information. Orange is used in the few cases when no sequence data is available. Each item can be clicked on to prompt a summary table with various links to further protein details and glycan-binding information.

## Conclusion

BiotechLec was created to guide experimentalists in their choice for the right lectin depending on the aim of the assays that may be constrained by size, structure, multivalency and/or fine specificity. It should also help rationalizing results of assays built using commercial lectins. It is not intended as a full fledged database but a set of shortcuts to key information regarding lectins commonly used as probes. Other databases provide information on structure, sequence and specificity of an overlapping set of lectins, such as the Lectin Frontier Database containing approximately 400 entries (<https://acgg.asia/lfdb2/index>) and GlyCosmos Lectins (<https://glycosmos.org/lectins>) that combines several sources including UniProt and UniLectin into roughly 2300 entries. These

resources are general purpose in contrast with BioTechLec which a user-friendly interface to quickly access information on commercial lectins. This guide was also designed to easily account for regular updates reflecting, as mentioned above, researchers' comments and feedback on testing the listed lectins or newly published information. It can equally as easily accommodate the addition of new entries.

## Acknowledgements

We thank Silvano Alda for prototyping the interactive table. The development of resources of the Glyco@ExPASy initiative that includes

UniLectin is supported by the Swiss Federal Government through the State Secretariat for Education, Research and Innovation (SERI). Expasy is maintained by the Swiss Institute of Bioinformatics and hosted at the Vital-IT Competency Center. UniLectin is also supported by ANR PIA Glyco@Alps (ANR-15-IDEX-0002) and Labex Arcane/CBH-EUR-GS (ANR-17-EURE-0003).

## References

- Bojar D, Meche L, Meng G, Eng W, Smith DF, Cummings RD, Mahal LK. 2022. A useful guide to lectin binding: Machine-learning directed annotation of 57 unique lectin specificities. *ACS Chem Biol*, 17:2993-3012, 10.1021/acscchembio.1c00689
- Bonnardel F, Kumar A, Wimmerova M, Lahmann M, Perez S, Varrot A, Lisacek F, Imberty A. 2019a. Architecture and evolution of blade assembly in  $\beta$ -propeller lectins. *Structure*, 27:764–775, 10.1016/j.str.2019.02.002
- Bonnardel F, Mariethoz J, Perez S, Imberty A, Lisacek F. 2021. LectomeXplore, an update of UniLectin for the discovery of carbohydrate-binding proteins based on a new lectin classification. *Nucleic Acids Res*, 49:D1548-D1554, 10.1093/nar/gkaa1019
- Bonnardel F, Mariethoz J, Salentin S, Robin X, Schroeder M, Pérez S, Lisacek F, Imberty A. 2019b. UniLectin3D, a database of carbohydrate binding proteins with curated information on 3D structures and interacting ligands. *Nucleic Acids Res*, 47:D1236–D1244, 10.1093/nar/gky832
- Carrizo ME, Capaldi S, Perduca M, Irazoqui FJ, Nores GA, Monaco HL. 2005. The antineoplastic lectin of the common edible mushroom (*Agaricus bisporus*) has two binding sites, each specific for a different configuration at a single epimeric hydroxyl. *J Biol Chem*, 280:10614-10623, 10.1074/jbc.M411989200
- Cummings RD, Etzler M, Hahn MG, Darvill A, Godula K, Woods RJ, Mahal LK. 2022. Glycan-Recognizing Probes as Tools. In: Varki A, Cummings RD, Esko JD, Stanley P, Hart GW, Aebi M, Mohnen D, Kinoshita T, Packer NH, Prestegard JH, et al. editors. *Essentials of Glycobiology*. Cold Spring Harbor (NY). p. 645-662.
- Ghanimi Fard M, Khabir Z, Reineck P, Cordina NM, Abe H, Ohshima T, Dalal S, Gibson BC, Packer NH, Parker LM. 2022. Targeting cell surface glycans with lectin-coated fluorescent nanodiamonds. *Nanoscale Adv*, 4:1551-1564, 10.1039/d2na00036a
- Haramoto Y, Onuma Y, Mawaribuchi S, Nakajima Y, Aiki Y, Higuchi K, Shimizu M, Tateno H, Hirabayashi J, Ito Y. 2020. A technique for removing tumourigenic pluripotent stem cells using rBC2LCN lectin. *Regen Ther*, 14:306-314, 10.1016/j.reth.2020.03.017
- Imberty A, Bonnardel F, Lisacek F. 2021. UniLectin, a one-stop-shop to explore and study carbohydrate-binding proteins. *Curr Protoc*, 1:e305, 10.1002/cpz1.305
- Klamer ZL, Harris CM, Beirne JM, Kelly JE, Zhang J, Haab BB. 2022. CarboGrove: a resource of glycan-binding specificities through analyzed glycan-array datasets from all platforms. *Glycobiology*, 32:679-690, 10.1093/glycob/cwac022
- Lebreton A, Bonnardel F, Dai Y-C, Imberty A, Martin FM, Lisacek F. 2021. Comprehensive phylogenetic and bioinformatics survey of lectins in the fungal kingdom. *J Fungi*, 7:453, 10.3390/jof7060453
- Lis H, Sharon N. 1986. Lectins as molecules and as tools. *Annu Rev Biochem*, 55:35-67, 10.1146/annurev.bi.55.070186.000343

- Morgan WT, Watkins WM. 1953. The inhibition of the haemagglutinins in plant seeds by human blood group substances and simple sugars. *Br J Exp Pathol*, 34:94-103,
- Neelamegham S, Aoki-Kinoshita K, Bolton E, Frank M, Lisacek F, Lütteke h, O'Boyle N, Packer NH, Stanley P, Toukach P, *et al.* 2022. Updates to the Symbol Nomenclature for Glycans guidelines. *Glycobiology*, 29:620-624, 10.1093/glycob/cwz045
- Notova S, Bonnardel F, Rosato F, Siukstaite L, Schwaiger J, Bovin N, Varrot A, Römer W, Lisacek F, Imberty A. 2022. The choanoflagellate pore-forming lectin SaroL-1 punches holes in cancer cells by targeting tumor-related glycosphingolipid Gb3. *Comm Biol*, 5:594, 10.1038/s42003-022-03869-w
- Ribeiro JP, Mahal LK. 2013. Dot by dot: analyzing the glycome using lectin microarrays. *Curr Opin Chem Biol*, 17:827-831, 10.1016/j.cbpa.2013.06.009