



HAL
open science

Activation of Nod2 signalling upon Norovirus infection enhances antiviral immunity and susceptibility to colitis

Ghaffar Muharram, Marion Thépaut, Pierre-Emmanuel Lobert, Teddy Grandjean, Olivier Boulard, Myriam Delacre, Emmrich Wakeford, Richard Wheeler, Lionel Franz Poulin, Ivo Gomperts Boneca, et al.

► **To cite this version:**

Ghaffar Muharram, Marion Thépaut, Pierre-Emmanuel Lobert, Teddy Grandjean, Olivier Boulard, et al.. Activation of Nod2 signalling upon Norovirus infection enhances antiviral immunity and susceptibility to colitis. 2023. hal-04101382v2

HAL Id: hal-04101382

<https://hal.science/hal-04101382v2>

Preprint submitted on 13 Aug 2023 (v2), last revised 30 Oct 2023 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Activation of Nod2 signaling upon norovirus infection enhances antiviral immunity and**
2 **susceptibility to colitis**

3

4 **Authors:** Ghaffar Muharram^a, Marion Thépaut^a, Pierre-Emmanuel Lobert^c, Teddy Grandjean^a,
5 Olivier Boulard^b, Myriam Delacre^a, Emmrich Wakeford^a, Richard Wheeler^c, Lionel Franz
6 Poulin^b, Ivo Gomperts Boneca^c, Frank Lafont^a, Marie-Cécile Michallet^d, Didier Hober^c, Ken
7 Cadwell^{f,g,h}, Mathias Chamaillard^b.

8 **Affiliations :**

9 ^aUniv. Lille, CNRS, Inserm, CHU Lille, Institut Pasteur de Lille, U1019 - UMR 9017 - CIIL -
10 Centre d'Infection et d'Immunité de Lille, F-59000 Lille, France.

11

12 ^bLaboratory of Cell Physiology, INSERM U1003, University of Lille, Lille, France.

13 ^cUniv Lille, Faculté de Médecine, CHU Lille, Laboratoire de Virologie ULR3610, Lille,
14 France.

15

16 ^dTERI (Tumor Escape, Resistance and Immunity), Centre de recherche en cancérologie de
17 Lyon, Centre Léon Bérard, Université de Lyon, Université Claude Bernard Lyon 1, Inserm
18 1052, CNRS 5286, 69008 Lyon, France.

19

20 ^eInstitut Pasteur, Université Paris Cité, CNRS UMR6047, INSERM U1306, Unité de Biologie
21 et génétique de la paroi bactérienne, F-75015, Paris, France

22

23 ^fKimmel Center for Biology and Medicine at the Skirball Institute, New York University
24 Grossman School of Medicine, New York, NY 10016, USA

25 ^gDepartment of Microbiology, New York University Grossman School of Medicine, New
26 York, NY 10016, USA

27 ^hDivision of Gastroenterology and Hepatology, Department of Medicine, New York
28 University Langone Health, New York, NY 10016, USA

29

30 **Disclosure of interest:**

31 K.C. has received research support from Pfizer, Takeda, Pacific Biosciences, Genentech, and Abbvie;
32 consulted for or received honoraria from Vedanta, Genentech, and Abbvie; and holds U.S. patent
33 10,722,600 and provisional patent 62/935,035 and 63/157,225.

34 **Correspondence should be addressed to:**

35 Mathias Chamaillard, PhD, Phone number: +33359317427, Fax number: +33359317480, E-
36 mail: mathias.chamaillard@inserm.fr

37 Ghaffar Muharram, PhD, phone number: +33320871033, E-mail: [ghaffar.muhammad@pasteur-](mailto:ghaffar.muhammad@pasteur-lille.fr)
38 [lille.fr](mailto:ghaffar.muhammad@pasteur-lille.fr)

39

40 **Key words:** Antiviral immunity, Colitis, Macrophages NOD2, Norovirus, Signaling.

41

42 **Data availability statement:** the data that support the findings of this study are available from
43 the corresponding author, GM or MC, upon request on HAL

44 <https://hal.science/hal-04101382>

45

46

47

48

49 **Abstract.** Over 90% of epidemic nonbacterial gastroenteritis are caused by human Noroviruses
50 (NoV) which are persisting in a substantial subset of people allowing their spread worldwide.
51 It leads to a significant number of endemic cases and up to 70,000 children deaths in developing
52 countries. NoVs are primarily transmitted through the fecal-oral route. To date studies have
53 focused on the influence of the gut microbiota on enteric viral clearance by mucosal immunity.
54 In this study, the use of mouse norovirus S99 (MNoV_S99) and CR6 (MNoV_CR6), two
55 persistent strains, allowed us to provide evidence that the norovirus-induced exacerbation of
56 colitis severity relied on bacterial sensing by nucleotide-binding oligomerization domain 2
57 (Nod2). Consequently, *Nod2*-deficient mice showed reduced levels of gravity of Dextran
58 sodium sulphate (DSS)-induced colitis with both viral strains. And MNoV_CR6 viremia was
59 heightened in *Nod2*^{-/-} mice in comparison with animals hypomorphic for *Atg16ll1*, that are prone
60 to aggravated inflammation under DSS. Accordingly, the infection of macrophages derived
61 from WT mice promoted the phosphorylation of Signal Transducer and Activator of
62 Transcription1 (STAT1) and NOD2's expression levels. Higher secretion of Tumor Necrosis
63 Factor alpha (TNF α) following NOD2 activation and better viral clearance were measured in
64 these cells. By contrast, reduced levels of pSTAT1 and blunted downstream secretion of TNF α
65 were found in *Nod2*-deficient macrophages infected by MNoV_S99. Hence, our results uncover
66 a previously unidentified virus-host-bacterial interplay that may represent a novel therapeutic
67 target for treating noroviral origin gastroenteritis that may be linked with susceptibility to
68 several common illnesses such as Crohn's disease.

69
70
71
72
73
74
75
76
77
78

79 **Introduction.** The development of high throughput sequencing techniques progressively
80 helped us gain a better comprehension of the composition of the enteric virome¹. This is of
81 particular importance for unravelling how some enteric viruses can establish persistent
82 infections and subsequently influence our health^{2,3}. Amongst those, noroviruses (NoV) have
83 been studied for several years. These single stranded RNA (+) viruses, belonging to the family
84 of *Caliciviridae* are responsible for the majority of non-bacterial gastroenteritis worldwide.
85 Until recent progress in cultivation methods of noroviruses from genogroups I, II, IV, VIII and
86 IX⁴⁻⁶ that infect humans (HuNoVs), mechanistic details were mostly obtained from studies
87 using the mouse norovirus (MNoV)⁷. Indeed, MNoVs can be easily produced *in vitro* in
88 macrophages or dendritic cells⁸. Several MNoV strains have been described based on their
89 shedding time in the stool and classified as either persistent or acute (non-persistent) strains⁹.
90 Owing to the variety of circulating NoV strains, strain-dependent effects are expected to occur
91 and should be studied more in details. Even a single amino acid change in the NS1/2 viral
92 protein is sufficient to switch the acute MNoV_CW3 strain to a persistent state¹⁰, showing the
93 high degree of adaptability of NoV genomes and hence their versatility to counteract host
94 defence mechanisms. Type III interferon (IFN- λ) response has been shown to control the *in*
95 *vivo* propagation of the persistent MNoV_CR6 strain¹¹, while type I and II interferons mediate
96 innate immune responses against acute MNoV infections^{12,13}.
97 Multiple intracellular pattern recognition receptors (PRR) have been associated with indirect
98 noroviral detection, especially during NoV genome replication, when double stranded RNA
99 intermediates are generated¹⁴. As such, the Toll like receptor 3 (TLR3) that recognizes any
100 dsRNA in the endosomal compartment and the melanoma differentiation associated protein-5
101 (MDA-5), a member of the RIG-I like receptor (RLR) family that recognizes long dsRNA, have
102 been shown to participate in the detection of the MNoV_1 strain and its plaque isolate the
103 MNoV_CW3¹⁵. Amongst the Nod like receptor (NLR) family, it was shown that the loss of

104 *Nlrp6* impairs antiviral immunity to MNoV_1¹⁶. Besides its ability to sense muropeptides from
105 Gram positive and negative bacteria¹⁷ the nucleotide-binding oligomerization domain 2
106 (NOD2) was also shown to directly bind with the genome (ssRNA-) of the respiratory syncytial
107 virus (RSV) to induce antiviral responses¹⁸. The recruitment of MAVS by NOD2 was needed
108 to activate the interferon-regulatory factor 3 (IRF3) dependent signaling and promote IFN- β
109 production. In the same line, NOD2-dependent signaling was shown to disrupt intestinal
110 homeostasis and increase lethality in an *E. Coli* and MNoV_1 co-infection mouse model¹⁹.
111 Another molecular player that is essential to activate antiviral signaling is the cytosolic protein
112 signal transducer and activator of transcription 1 (STAT1). MNoV_1 infection was shown to
113 be lethal in STAT1-deficient mice¹². Furthermore, STAT1 dependent interferon responses were
114 shown to limit MNoV_1 replication and dissemination *in vivo*²⁰.
115 Interestingly, Cadwell and colleagues have demonstrated how infection with the MNoV
116 persistent strain CR6 rendered *Atg16l1^{HM}* mice more susceptible to DSS-induced colitis²¹.
117 Interestingly, the interaction of NOD2 with ATG16L1 was shown to regulate the autophagy
118 clearance of pathogenic bacteria²². Furthermore, it has been established that loss-of-function
119 point mutations of either *NOD2* or *ATG16L1* predispose humans to Crohn's disease^{23,24}.
120 Insights into the mechanisms of how NOD2 signaling is regulated upon noroviral infection and
121 whether it may subsequently modulate colitis susceptibility remained to be assessed.
122 In this study, we have investigated how NOD2 influences the pathogenesis of the two persistent
123 murine norovirus strains MNoV_S99 (*Berlin*)²⁵ and MNoV_CR6 *in vivo* in DSS-induced colitis
124 models. At the cellular level, the *Nod2*-dependent pro-inflammatory signaling pathways
125 activation were further examined in response to MNoV_S99 and MNoV_CR6 infection in
126 myeloid cells to determine how this in turn subsequently affect viral production.
127
128

129 **Results.**

130 **MNoV_S99 exacerbates the severity of DSS-induced inflammation**

131 Since MNoV_S99 is a natural strain commonly found in animal facilities that is able to establish
132 persistent infections²⁵, we assessed the implication of MNoV_S99 in the exacerbation of colitis.
133 Hence, we examined MNoV_S99's impact in a widely used chemically-induced colitis model
134 that is induced upon administration of Dextran Sodium Sulfate (DSS). Specifically, C57BL/6J
135 mice were first infected with a single dose of 5×10^7 TCID₅₀/mL of MNoV_S99 or mock-treated
136 with PBS (200µL). One week later, drinking water was replaced with either a 3% DSS or 5%
137 DSS solution for a period of seven consecutive days (Fig. 1A). Body weight-loss was monitored
138 on a daily basis together with signs of diarrhea and rectal bleeding. With both concentrations
139 of DSS, the pre-treatment with MNoV_S99 had induced a more significant body weight loss
140 (purple and orange curves Fig. 1B and C). After 7 days of DSS treatment, mice were euthanized
141 and the disease severity was evaluated. Histological analysis of colon sections revealed an
142 increased infiltration of inflammatory cells and more pronounced epithelial damages in infected
143 mice treated with DSS when compared to similarly challenged animals that were not infected
144 (Fig. 1D). Accordingly, increased muscle thickness was measured in colon sections from mice
145 treated with the virus in presence of both 3 and 5% DSS in comparison with Mock-treated
146 control mice (Fig. 1E). Similarly, histological scoring of the hematoxylin and eosin (H&E)
147 stained colonic sections expectedly showed significantly more severity in MNoV_S99-infected
148 mice treated with either 3% or 5% DSS than DSS alone (Fig. 1F). Upon mice sacrifice, resected
149 colons were measured. A significantly reduced length was obtained in mice infected with
150 MNoV_S99 and treated with 5% DSS in comparison with non-infected control animals (PBS)
151 (Fig. 1G). The increased level of colitis was also evidenced by a significantly augmented
152 secretion of the pro-inflammatory cytokine interleukin-6 (IL-6) in the supernatants of over-
153 night *in vitro* cultured colon explants from MNoV_S99 infected + 5% DSS treated mice versus

154 those from DSS only mice (Fig. 1H). Taken together, these data clearly show that MNoV_S99
155 infection **is able to lower the resistance of mice to chemically induced colitis.**

156

157 **MNoV_S99 associated inflammation is abrogated in *Nod2*^{-/-} mice**

158 In cells from the myeloid lineage, NOD2 was shown to participate in the sensing of *Citrobacter*
159 *rodentium* and of commensal bacteria that are translocating in response to DSS-induced
160 epithelial injury²⁶. To determine whether NOD2 is also implicated in the immunopathology of
161 MNoV in our preclinical model of colitis, we compared the severity of DSS-induced colitis in
162 the presence **or absence** of MNoV_S99 between WT and *Nod2*-deficient mice. In contrast to
163 what was observed in WT mice (Fig. 2A), the sensitizing effect of the virus on the daily weight-
164 loss was absent upon MNoV_S99 infection of *Nod2*^{-/-} mice (Fig. 2B). These data suggest that
165 the pro-inflammatory effect of MNoV_S99 observed in WT mice **may depend on the degree of**
166 **bacterial sensing through NOD2.** Indeed, DSS induces an abrasion of the epithelial barrier
167 allowing commensal bacteria to cross the mucosa and unleash a **NOD2-dependent** pro-
168 inflammatory response. This was further confirmed after mice were autopsied. Colon tissue
169 sections staining showed decreased levels of colonic inflammation from MNoV_S99 infected
170 *Nod2*^{-/-} vs WT mice (Fig. 2C). In agreement, muscle thickness quantification from these colon
171 sections showed significantly increased thickness in MNoV_S99-infected WT mice under 3%
172 DSS treatment vs 3% DSS alone (Fig. 2D). The variations of colon length measurements that
173 were induced upon MNoV_S99 infection in WT mice were absent **by contrast** in *Nod2*^{-/-} mice
174 (Fig. S1A). Similarly, differences in histological scores **and mucosal wall thickness** measured
175 from MNoV_S99-infected WT mice vs DSS alone were also **blunted** in *Nod2*^{-/-} infected mice
176 (Fig. S1B). And a clear trend was observed regarding the increased levels of secretion of
177 Tumour necrosis factor α (TNF α) in the supernatants of *in vitro* cultured colon explants from
178 WT mice infected with MNoV_S99 and treated with 3% DSS vs those with DSS only. This

179 effect was absent in the supernatants from *Nod2*^{-/-} mice infected and treated similarly (Fig.
180 S1C).

181 Overall, these data **demonstrate that** the excessive inflammatory response measured in the
182 presence of MNoV_S99 **depends on NOD2-dependent bacterial sensing** in our DSS-induced
183 colitis mouse model.

184

185 **Mouse noroviruses infections triggers NOD2-dependent pro-inflammatory signaling**

186 Mouse noroviruses are known to have a tropism for the myeloid cell lineages, namely
187 macrophages and dendritic cells. Recently it was shown that Tuft cells belonging to the gut
188 epithelial cellular lineages, that possess the CD300lf MNoV receptor can also be targeted by
189 the MNoV_CR6 strain²⁷.

190 Making use of the GFP-Nod2 reporter mice line, we first examined in which cellular lineages
191 Nod2 is expressed **within the intestinal mucosa**. We failed to observe **any** GFP-Nod2 expression
192 in epithelial cells. **By contrast, GFP** signal was detectable in the bulk of the cells below the
193 lamina propria where **intestinal mononuclear phagocytes** are **primarily** located (Fig. 3A).
194 Hence, **the NOD2-dependent response to MNoV_S99 infection was further** studied in
195 macrophages, monocytes and dendritic cells in our *in vitro* cellular models.

196 Interferon production being one of the most common response to the intracellular presence of
197 RNA genomes of foreign origin, we **next** examined if the STAT1 signaling pathway that lays
198 upstream of ATG1611-mediated interferon production was activated in response to murine
199 norovirus infection. To this end, bone marrow derived macrophages (BMDM) from WT mice
200 were infected with several **persistent and non-persistent** MNoV strains at a multiplicity of
201 infection (Moi) of 1, for 1h and pSTAT1 levels of expression from cell lysates were measured
202 by western blot (Fig. 3B). In contrast to the non-persistent strain MNoV_CW3 that caused a
203 striking increase in pSTAT1 levels, **infections with** the persistent strains S99 and CR6 induced

204 a much milder STAT1 signaling pathway activation. The CW1 strain was associated with the
205 lowest level of STAT1 activation.

206 Since MNoV_S99 is a natural strain commonly found in animal facilities that is able to establish
207 persistent infections, we used it to further analyze whether the response to viral infections may
208 rely on NOD2-associated signaling. Using the macrophage cell-line Raw264.7, we confirmed
209 that MNoV_S99 is able to induce phosphorylation of STAT1 (Tyrosine 701) at different times
210 at Moi 1 (Fig. S2). pSTAT1 was detectable as early as 15 minutes post-infection and its
211 expression further increased at 30 min and 1h, while only a slight increase of STAT1 level was
212 detectable at later time-points. In addition to the STAT1 activation through the Janus kinases
213 JAK1 and Tyk2, we also examined the status of NF κ B activation. For this purpose, pI κ B α levels
214 were quantified from infected cell lysates. Despite basal low-level activation in mock-infected
215 cells, pI κ B α intensity increased slightly at 30 min and 1h post-infection even if it failed to reach
216 significance. When checking AKT or the MAPK-ERK signaling axes in the same time frame,
217 MNoV_S99 failed to induce any activation of these signaling pathways (Fig. S2). Next, we
218 analyzed the response of BMDM derived from WT or *Nod2*^{-/-} mice that were infected with
219 increasing doses of MNoV_S99 for 30 or 60 minutes (Fig. 3C). STAT1 activation increased
220 over the course of the acute infection in cells derived from WT mice. In contrast, the levels of
221 pSTAT1 were barely detectable in *Nod2*^{-/-} cells at Moi 1. Similarly, with prolonged times of
222 infection (24h, Moi 0.1) we measured significantly lowered pStat1 levels in MNoV_S99-
223 infected *Nod2*^{-/-} cells in comparison with WT cells (Fig. 3D). However, with a higher dose (Moi
224 5) this lowered phosphorylation of STAT1 in response to MNoV_S99 infection progressively
225 vanished (Fig. 3C), suggesting NOD2-independent alternative mechanisms of action operate
226 when the viral load is too important.

227 To examine whether the Mitochondrial antiviral signaling (MAVS) adaptor protein is required
228 for sensing MNoV_S99, BMDM derived from Mavs-deficient mice were generated.

229 Interestingly, the higher pStat1 levels measured in WT cells (Moi 0.1, 1h) were significantly
230 decreased in **BMDM** from *Mavs*^{-/-} cells (Fig. 3E).

231 Altogether, these data show a mild but significant activation of **STAT1** following MNoV_S99
232 infection in our *in vitro* cellular models. These signaling responses can be at least partially
233 abrogated in the absence of MAVS and NOD2.

234

235 **Lower levels of inflammation in *Nod2*^{-/-} mice benefit noroviral propagation**

236 MNoV_CR6 infection in DSS-induced colitis was previously shown to promote inflammation
237 in a functional *Atg16l1* dependent manner²¹. Using this viral strain **that promote STAT1**
238 **phosphorylation in similar degree than MNoV_S99 (Fig. 3B)**, we checked how loss of *Nod2*
239 influenced resolution of inflammation in response to 3% DSS treatment **in mice. As shown with**
240 **the MNoV_S99 strain, loss of NOD2 improved the muscle thickening of the colon wall in mice**
241 **infected with MNoV_CR6 (Fig. 4A). However, a CR6 strain-specific effect can be observed**
242 **with these experiments since significantly increased inflammation was only measured in**
243 ***Atg16l1* hypomorph mice but not in WT mice as opposed to MNoV_S99.**

244 Next, we examined the impact of the pro-inflammatory signaling activation on viral
245 propagation in these mice. In the absence of DSS-induced inflammation no significant
246 differences could be found when comparing the viral load in the stool of WT, *Atg16l1*^{HM} or
247 *Nod2*^{-/-} mice (Fig. 4B). **By contrast, the measured viral load was significantly increased in ileum**
248 **and colon of DSS-treated *Nod2*^{-/-} mice in comparison with tissues of *Atg16l1*^{HM} mice (Fig. 4C).**
249 We have also analyzed the viral replication in BMDM derived from WT or *Nod2*^{-/-} mice
250 infected *in vitro* with MNoV_S99 (Moi 0.1 for 24h). In that case a slight but significant increase
251 in the number of viral genomes was measured from *Nod2*^{-/-} derived cultures (Fig. 4D).

252 **Hence, the lower levels of inflammation found in *Nod2*^{-/-} mice seems to favor a higher**
253 **propagation of these two persistent strains.**

254 **MNoV_S99 infection promotes bacterial sensing by NOD2**

255 To further explore the mechanisms behind the excessive inflammatory response associated with
256 noroviral infections, we examined **whether noroviral infection may enhance** *Nod2* expression
257 levels by RT-qPCR. **As what was previously observed with MNoV-1¹⁵, the transcription level**
258 **of NOD2 was significantly enhanced** in response to MNoV_S99 infection in **either** Raw264.7
259 cells or BMDM derived from WT mice (Fig. 5A and 5B).

260 This increased *Nod2* expression following norovirus infection could explain **at least partially**
261 the viral-induced pathogenesis that we observed in our colitis models. **Indeed, the greater**
262 **expression of NOD2 should spike up when suddenly intestinal mononuclear phagocytes**
263 interact with intestinal commensal bacteria following DSS-induced epithelial abrasion. **We next**
264 **evaluated whether this heightened expression of NOD2 in response to MNoV_S99 infection**
265 **may enhance bacterial sensing. While MNoV_S99 infection did induce Stat1 (but not I κ B α)**
266 **activation** in monocytes isolated from WT mice (Fig. 5C and 5D, MNoV_S99+, MDP-
267 condition), MDP alone triggered potent I κ B α activation without activating Stat1 (Fig. 5C,
268 MNoV_S99-, MDP+ condition). Interestingly, MNoV_S99 infection followed by secondary
269 MDP treatment was associated with significant **phosphorylation of both Stat1 and I κ B α** (Fig.
270 5D, MNoV_S99+, MDP+ vs mock treated cells). In the same line of investigation, we noticed
271 a significantly augmented pro-inflammatory cytokine TNF α response to MDP in dendritic cells
272 that were first primed with noroviral RNA (Fig. 5E). **Next, the secretion of TNF α was measured**
273 **in the supernatants from WT and *Nod2*^{-/-} BMDM cultures** infected with MNoV_S99 followed
274 by MDP treatment. TNF α levels were increased with increasing doses of virus in WT cells,
275 while they remained undetectable (Moi 0.1) or much lower (Moi 1) in *Nod2*^{-/-} cells (Fig. 5F).
276 **Equally of importance, viral titers measured in supernatants showed significantly decreased**
277 **yields when infected cells were treated with MDP** (Fig. 5G).

278 Overall, these data show that the increased *Nod2* transcription levels following noroviral
279 infections may at least partially promote an inflammatory response to bacterial MDP, leading
280 ultimately to a limitation in the viral propagation. Hence the lowered inflammation in *Nod2*^{-/-}
281 mice is associated with increased dissemination of MNoV_S99 and MNoV_CR6 strains,
282 suggesting that bacterial sensing through NOD2 may regulate their persistence.

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303 Discussion

304 Intestinal homeostasis is tightly regulated by the interplay between the microorganisms found
305 in the gut microbiota, intestinal epithelial cells and mononuclear phagocytes. The breach in gut
306 flora homeostasis termed dysbiosis is becoming central in the onset of chronic inflammatory
307 bowel disease, such as Crohn's disease^{28,29}. With the development of molecular biology
308 techniques, alterations in the virome relevant to disease pathogenesis are thoroughly under
309 investigation². As such, the role played by noroviruses was studied in several gastrointestinal
310 inflammatory models. An endemically circulating mouse norovirus strain was shown to
311 promote intestinal barrier dysfunctions and inflammatory lesions caused by loss of tolerance to
312 commensal bacteria in *III0*^{-/-} mice³⁰. The secondary challenge with a pathogenic *E. coli* strain
313 following MNoV-1's infection can even cause lethal lesions due to the induction of an
314 unresolved proinflammatory cytokine secretion in a *Nod1/2* dependent manner¹⁹. The
315 hypothesis of an environmental-driver in genetically susceptible hosts that could be at the origin
316 of Crohn's like disease was also explored with the MNoV_CR6 strain in the *Atg16l1*^{HM} mice
317 presenting defective autophagy in a DSS colitis model²¹.

318 NOD2 is very well known for its pathogen associated molecular pattern recognition activity
319 against bacteria¹⁷, and recently it appears it can also participate in detection of certain viruses¹⁸.
320 Using pro-inflammatory signaling activation downstream of NOD2 as a read-out, we have
321 analyzed if NOD2 can also participate in MNoV detection and elimination. All our *in vitro*
322 experiments with monocytic lineage derived primary cells (e.g. BMDM, BMDC and BM) and
323 cell line (e.g. Raw264.7) infected with different MNoV strains showed activation of the STAT1
324 signaling pathway implicated in the pro-inflammatory IFN and TNF α responses. By contrast,
325 these responses were abrogated in *Nod2*^{-/-} and *Mavs*^{-/-} mice derived BMDM.

326 Amongst enteric viruses commonly found in the microbiota, noroviruses have been proposed
327 to be good candidates in the hypothesis of an environmental-driver in genetically susceptible

328 hosts at the origin of Crohn's disease onset. MNoV_CR6 strain was shown to induce colitis in
329 *Atg16l1^{HM}* mice presenting defective autophagy in a DSS colitis model²¹.

330 Several PRR are implicated in viral sensing. MDA-5 and TLR3 have been shown to participate
331 in detection of either MNoV_1 or CW3¹⁵. These are both strains which do not persist *in vivo*,
332 contrary to the S99 (*Berlin*)²⁵ and CR6 that were used in our study.

333 Our *in vivo* data show significantly exacerbated inflammation when applying either 3% or 5%
334 DSS colitis protocol to WT or *Atg16l1^{HM}* mice. The excessive MNoV_S99 or MNoV_CR6
335 associated inflammatory response was abrogated in *Nod2^{-/-}* mice or in BMDM derived from
336 these mice.

337 Hence, NOD2 dependent signaling adds a layer of regulation in **the viral sensing and**
338 **propagation**. While in our norovirus + DSS model knocking-down NOD2 prevents gut
339 inflammation, it has the opposite effect with influenza A virus (IAV) infection in the lung of
340 *Nod2^{-/-}* or *Ripk2^{-/-}* mice³¹. Indeed, the NOD2-RIPK2 signaling axis has been shown to
341 downmodulate the NLRP3 dependent mitophagy activation and the associated inflammatory
342 response in the lung tissue of infected mice.

343 Noroviruses propagation or prolonged persistence is tightly regulated by the interplay between
344 several signaling pathways. MNoV_1 replication was shown to be inhibited in macrophages
345 following LPS mediated NF- κ B activation that in turn triggers IFN- β dependent JAK-STAT
346 antiviral activity³². Interestingly, in our cellular models, in addition to the phosphorylation of
347 STAT1 that we observed when cells were infected with MNoV_S99, **the heightened *Nod2***
348 **expression levels preceded a greater ability of cells to respond to bacterial MDP**. Consequently,
349 the subsequent treatment with MDP induced an increased Nf κ B signaling pathway via I κ B α
350 activation. The sequential activation of these two pro-inflammatory signaling pathways resulted
351 in the inhibition of the MNoV_S99 propagation on one hand and the augmented production of
352 the pro-inflammatory cytokine TNF α **in response to MDP** on the other hand. Hence, the

353 aggravated inflammatory pathology we observed in our DSS colitis model can be explained by
354 this NOD2-dependent signaling loop in MNoV_S99 or MNoV_CR6 infected mice.

355 In conclusion, our study shows a **NOD2-dependent bacterial sensing that prevents**
356 **dissemination of the persistent MNoV_S99 and MNoV_CR6 strains.**

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378 **Methods:**

379 Unless otherwise indicated, all media and reagents were purchased from ThermoFischer
380 Scientific.

381

382 **Colitis mouse model and viral infection**

383 All animal experiments were approved by the local ethical committee n° CEEA-
384 2016030717519903. Age and gender related *Nod2*^{-/-} and wild type C57BL/6J mice (CDTA,
385 Orléans, France) were housed 4-6 per cage, under strict pathogens-free environment. Mice were
386 gavaged with 5×10^7 TCID₅₀/mL of MNoV_S99 diluted in 200µl PBS or mock treated with PBS
387 only. 7 days post-infection, drinking water was substituted with a 3-5% dextran sodium sulphate
388 (DSS) (35,000-40,000 MW; TdB Consultancy) solution replaced every 2 days for a total
389 duration of 7 days. 3% DSS colitis experiments with MNoV_CR6 were conducted as previously
390 described in²¹.

391 **Cells and virus**

392 Mouse leukemic monocyte macrophage cell line RAW264.7 were purchased from ECCAC
393 (Sigma-Aldrich) and maintained in Dulbecco's modified Eagle's medium (DMEM),
394 supplemented with 10% (v/v) foetal bovine serum, 1% penicillin/streptomycin and 1% sodium
395 pyruvate.

396 Bone marrow derived macrophages (BMDM) were isolated from femurs of wild-type, *Nod2*^{-/-}
397 or *Mavs*^{-/-} mice of C57BL/6J background³³. Using a 26G ½" needle, bone marrow cells were
398 flushed out of the bones with Iscove's modified Dulbecco's medium (IMDM), supplemented
399 with 10% foetal bovine serum, 1% penicillin/streptomycin, 1% non-essential amino acid, 1%
400 sodium pyruvate, 1% glutamine, and 20% (v/v) conditioned-media from L929 cells. Red blood
401 cells were lysed using a 160mM NH₄CL and 170mM Tris solution for 5 minutes at RT. 3 to

402 6×10^6 viable cells were plated in non-cell-culture-treated petri dishes and grown for 5-7 days
403 in the above mentioned fully supplemented IMDM medium.

404 Bone marrow derived dendritic cells were isolated similarly as BMDM cells. Then, 2×10^6
405 viable cells were plated in non-cell-culture-treated petri dishes and grown for 7 days in RPMI
406 1640, supplemented with 10% foetal bovine serum, 1% penicillin/streptomycin, 1% L-
407 glutamine, 1% HEPES and 20% (v/v) conditioned-media from J558 cell line producing murine
408 granulocyte monocyte colony stimulating factor (GM-CSF).

409 Monocytes were isolated from bone marrow cells by using the mouse Monocyte Isolation Kit
410 (Miltenyi Biotec) and a QuadroMACS separator.

411 The MNoV-S99 strain (GenBank accession no. DQ911368) was provided by Prof. P. Maris
412 from ANSES Fougères Laboratory (France) and was propagated in RAW 264.7 cells. MNoV-
413 CR6 concentrated stocks were prepared as described in³⁴. Briefly, supernatant from 293T cells
414 transfected with a plasmid containing the viral genome was applied to RAW264.7 cells to
415 amplify virus production, and virions were concentrated by ultracentrifugation and
416 resuspension in endotoxin-free PBS. Concentration of stocks was determined by plaque assay.

417 For in vitro infection experiments, $3-5 \times 10^5$ viable cells were seeded in 6 wells plates. The next
418 day, just before viral infection, cells were counted from 1 well to determine precisely the
419 multiplicity of infection (Moi). In the remaining wells medium was removed and replaced with
420 variable quantities of MNoV_S99 from a common viral stock in a final volume of 1 mL fresh
421 medium per well. Mock treated cells were incubated in parallel with matching quantities of
422 culture media from non-infected cells that was generated at the same time as viral stocks.

423 At various time-points after infection, supernatant was removed and cells were washed once
424 with PBS, before lysis in RIPA buffer for WB analysis or RLT buffer for total RNA extraction
425 (RNAeasy kit, Quiagen) for RT-qPCR experiments.

426

427 **Immunohistochemistry, ELISA, Western Blot, RT-qPCR and Titration**

428 For immunohistochemistry analysis, formalin-fixed colon samples were embedded in paraffin.
429 5µm sections were stained with H&E. Slides were imaged with an AxioPlan 2 (Zeiss)
430 microscope and blindly scored for inflammation severity by two investigators as previously
431 described³⁵. Briefly, tissue lesions were scored from 0 to 3, so was the infiltration of
432 inflammatory cells in the lamina propria. A combined score was calculated, ranging from 0 (no
433 changes) to 6 (wide-spread cellular infiltrations and extensive tissue damage).

434 The muscularis propria thickness was measured from light microscopy images of H&E stained
435 colon sections using Image J (NCBI). 5 measurements from different locations were performed
436 for each animal and compared with colon sections from mock treated animals.

437 For ELISA analysis, briefly 1cm colon samples were recovered from mice, luminal content was
438 flushed with PBS, samples were cut opened and resuspended over-night in 250µl DMEM
439 supplemented with 1% penicillin/streptomycin in 24 wells plate at 37°C. The following day,
440 supernatants were collected and stored at -80°C before cytokine measurement. TNFα and
441 Interleukin-6 proteins levels were measured specifically using pre-coated 96 wells ELISA
442 plates (R&D System) following manufacturers indications.

443 For TNFα ELISA from BMDC, 5×10^5 cells were plated in 24 wells plate and treated with
444 MNoV_S99's ssRNA (extracted with RNeasy Quiagen kit) with Lipofectamine 2000¹⁸ ± MDP
445 or MDP-DD (Invivogen) over-night and supernatants were measured for TNFα levels.

446 For TNFα ELISA from BMDM, 2×10^5 cells were plated in 96 wells plate infected with
447 MNoV_S99 (Moi 0.1 or 1) for 6h prior to MDP treatment (10ug/mL) over-night and
448 supernatants were measured for TNFα levels.

449 For western blot analysis, total proteins from cell lysates were quantified by BCA protein assay
450 kit (Pierce), equalized amounts of proteins (10-20µg) were resolved by 4-15% gradient SDS-
451 PAGE gels and transferred to nitrocellulose membranes. Membranes were blocked in 0,05%

452 PBS-Tween, 5% skimmed milk solution, before incubation subsequently with primary and
453 secondary antibodies diluted (see Table 1) in blocking solution. Following secondary antibodies
454 incubations, signals of interest were detected using a chemiluminescence reader
455 (ImageQuant™ Las 4000, GE Healthcare Life Sciences), images were processed and band ROI
456 were quantified with Image J software (NCBI). Mean band intensities \pm SD of protein of
457 interest were normalized to β -ACTIN. Relative values to control mock-infected conditions were
458 compared between different experiments.

459 For *Nod2* quantification from MNoV_S99 infected BMDM or RAW264.7 cells, total RNA
460 extraction was performed using the RNeasy Mini Kit (Qiagen). One μ g of RNA was reversed
461 transcribed in the presence of 2.5mM of oligo-dT using the Reverse Transcriptor kit (Roche)
462 following the manufacturer's instructions. 75ng of cDNAs were used for the qPCR analyses
463 using Q5 High-Fidelity 2X Master Mix (New England BioLabs). *Nod2* and *Actb* were
464 specifically amplified using the Fwd_5'GGAAACAACATTGGCAGCAT3' and
465 Rev_5'TCTTGAGTCCTTCTGCGAGA3' Fwd_5'TTCTTTGCAGCTCCTTCGTT3' and
466 Rev_5'ATGGAGGGGAATACAGCCC3' respectively. *Actb* was used as an internal reference
467 gene in order to normalize the transcript levels. Relative mRNA levels ($2^{-\Delta\Delta Ct}$) were
468 determined by comparing (a) the PCR cycle thresholds (Ct) for *Nod2* and *Actb* (ΔCt) and (b)
469 ΔCt values for treated and control groups ($\Delta\Delta Ct$).

470 MNoV_S99 load was determined by measuring MNoV genome copy numbers by RT-qPCR
471 analysis as previously described³⁶. Briefly, total RNA was extracted from BMDM cell cultures
472 using the RNeasy Mini Kit (Qiagen). 5 μ L of RNA were used per reaction of RT-qPCR. Viral
473 genome levels were determined using the Takyon™Dry No Rox One-Step RT Probe
474 Mastermix (Eurogentec). Viral genomes were specifically amplified using the
475 Fwd_5'GTGCGCAACACAGAGAAAACG3' and Rev_5'CGGGCTGAGCTTCCTGC3'
476 primers that target MNoV ORF1 region combined with the TaqMan probe 5'[6-FAM]-

477 **CTAGTGTCTCCTTTGGAGCACCTA-[BHQ1]3'** using the Stratagene Mx3005P (Agilent
478 Technologies) **with 1 RT cycle and 40 cDNA amplification cycles**. The quantification of viral
479 genome copies was obtained using a standard curve (**ranging from 2×10^1 to 2×10^6 genome**
480 **copies**) with serial dilutions of the plasmid pMNoV1 corresponding to the full genome of
481 MNoV_1 (Dr Christiane Wobus, University of Michigan, USA). Viral load for MNoV_CR6
482 strain was determined by measuring genome copy numbers by RT-qPCR analysis as previously
483 described in²¹.

484 The net production of MNoV_S99 was determined using the TCID₅₀ method as described in³⁵.
485 Briefly, Raw264.7 cells were infected at Moi 1 with MNoV_S99 for 2h, before adding MDP
486 (100ng/mL) or not in the supernatant for 24h. Titers (n = 3) were obtained from each
487 supernatant. The net production is calculated by normalizing the amount of viruses produced
488 with the amount of viruses used for the infection.

489

490 ***NOD2*^{GFP} immunohistochemistry imaging**

491 *NOD2*^{GFP} mice³⁷ in the C57BL/6 background were maintained at the Institut Pasteur Central
492 Animal facility. We used heterozygous *NOD2*^{GFP} mice since the presence of one functional
493 copy of *NOD2* ensures that physiological responses dependent on functional NOD2 are not
494 impaired. Heterozygous *NOD2*^{GFP} female mice (8-12 weeks of age) were sacrificed and several
495 centimeters of the distal ileum were removed, opened longitudinally and washed in PBS. The
496 tissue was spread flat over a 4% w/v LMP agarose (Merck) pad and fixed with 4% v/v
497 paraformaldehyde (VWR). The ileal tissue was then embedded in a 4% LMP agarose block and
498 cut into 200µm-thick sections using a Microm HM 650 V Vibration microtome.

499 For staining, tissues were blocked and **permeabilized overnight in solution containing 0.4%**
500 **Triton X-100 (Merck)** and marked with an anti-GFP (rabbit polyclonal antibody A-11122,
501 Invitrogen) primary antibody and secondary Alexa Fluor 488 goat anti-rabbit antibody

502 (Invitrogen). Tissues sections were counter stained with DAPI (1µg/mL, BD Biosciences) and
503 Phalloidin- iFluor 647 (1:200 dilution, Abcam) and mounted using ProLong Gold Antifade
504 reagent. Confocal acquisitions were performed using a Leica HyD SP5 confocal microscope.
505 Image analysis was performed using Icy version 2.4.2.0³⁸.

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

525

526

527 **References:**

- 528 1. Cadwell, K. Expanding the role of the virome: commensalism in the gut. *J. Virol.* **89**,
529 1951–1953 (2015).
- 530 2. Norman, J. M. *et al.* Disease-specific alterations in the enteric virome in inflammatory
531 bowel disease. *Cell* **160**, 447–460 (2015).
- 532 3. Pfeiffer, J. K. & Virgin, H. W. Viral immunity. Transkingdom control of viral
533 infection and immunity in the mammalian intestine. *Science* **351**, aad5872 (2016).
- 534 4. Ettayebi, K. *et al.* Replication of human noroviruses in stem cell-derived human
535 enteroids. *Science* **353**, 1387–1393 (2016).
- 536 5. Estes, M. K. *et al.* Human Norovirus Cultivation in Nontransformed Stem Cell-
537 Derived Human Intestinal Enteroid Cultures: Success and Challenges. *Viruses* **11**, E638
538 (2019).
- 539 6. Ludwig-Begall, L. F., Mauroy, A. & Thiry, E. Noroviruses-The State of the Art,
540 Nearly Fifty Years after Their Initial Discovery. *Viruses* **13**, 1541 (2021).
- 541 7. Newman, K. L. & Leon, J. S. Norovirus immunology: Of mice and mechanisms. *Eur.*
542 *J. Immunol.* **45**, 2742–2757 (2015).
- 543 8. Wobus, C. E. *et al.* Replication of Norovirus in cell culture reveals a tropism for
544 dendritic cells and macrophages. *PLoS Biol.* **2**, e432 (2004).
- 545 9. Thackray, L. B. *et al.* Murine noroviruses comprising a single genogroup exhibit
546 biological diversity despite limited sequence divergence. *J. Virol.* **81**, 10460–10473 (2007).
- 547 10. Nice, T. J., Strong, D. W., McCune, B. T., Pohl, C. S. & Virgin, H. W. A single-
548 amino-acid change in murine norovirus NS1/2 is sufficient for colonic tropism and
549 persistence. *J. Virol.* **87**, 327–334 (2013).
- 550 11. Nice, T. J. *et al.* Interferon- λ cures persistent murine norovirus infection in the
551 absence of adaptive immunity. *Science* **347**, 269–273 (2015).
- 552 12. Karst, S. M., Wobus, C. E., Lay, M., Davidson, J. & Virgin, H. W. STAT1-dependent
553 innate immunity to a Norwalk-like virus. *Science* **299**, 1575–1578 (2003).
- 554 13. Hwang, S. *et al.* Nondegradative role of Atg5-Atg12/ Atg16L1 autophagy protein
555 complex in antiviral activity of interferon gamma. *Cell Host Microbe* **11**, 397–409 (2012).
- 556 14. Thorne, L. G. & Goodfellow, I. G. Norovirus gene expression and replication. *J. Gen.*
557 *Virol.* **95**, 278–291 (2014).
- 558 15. McCartney, S. A. *et al.* MDA-5 recognition of a murine norovirus. *PLoS Pathog.* **4**,
559 e1000108 (2008).
- 560 16. Wang, P. *et al.* Nlrp6 regulates intestinal antiviral innate immunity. *Science* **350**, 826–
561 830 (2015).
- 562 17. Chamailard, M., Girardin, S. E., Viala, J. & Philpott, D. J. Nods, Nalps and Naip:
563 intracellular regulators of bacterial-induced inflammation. *Cell. Microbiol.* **5**, 581–592
564 (2003).
- 565 18. Sabbah, A. *et al.* Activation of innate immune antiviral responses by Nod2. *Nat.*
566 *Immunol.* **10**, 1073–1080 (2009).
- 567 19. Kim, Y.-G. *et al.* Viral infection augments Nod1/2 signaling to potentiate lethality
568 associated with secondary bacterial infections. *Cell Host Microbe* **9**, 496–507 (2011).
- 569 20. Mumphrey, S. M. *et al.* Murine norovirus 1 infection is associated with
570 histopathological changes in immunocompetent hosts, but clinical disease is prevented by
571 STAT1-dependent interferon responses. *J. Virol.* **81**, 3251–3263 (2007).
- 572 21. Cadwell, K. *et al.* Virus-plus-susceptibility gene interaction determines Crohn’s
573 disease gene Atg16L1 phenotypes in intestine. *Cell* **141**, 1135–1145 (2010).
- 574 22. Travassos, L. H. *et al.* Nod1 and Nod2 direct autophagy by recruiting ATG16L1 to the
575 plasma membrane at the site of bacterial entry. *Nat. Immunol.* **11**, 55–62 (2010).

- 576 23. Hugot, J. P. *et al.* Association of NOD2 leucine-rich repeat variants with susceptibility
577 to Crohn's disease. *Nature* **411**, 599–603 (2001).
- 578 24. Hampe, J. *et al.* A genome-wide association scan of nonsynonymous SNPs identifies a
579 susceptibility variant for Crohn disease in ATG16L1. *Nat. Genet.* **39**, 207–211 (2007).
- 580 25. Niendorf, S., Klemm, U., Mas Marques, A., Bock, C.-T. & Höhne, M. Infection with
581 the Persistent Murine Norovirus Strain MNV-S99 Suppresses IFN-Beta Release and
582 Activation of Stat1 In Vitro. *PLoS One* **11**, e0156898 (2016).
- 583 26. Hrdý, J. *et al.* *Lactobacillus reuteri* 5454 and *Bifidobacterium animalis* ssp. *lactis* 5764
584 improve colitis while differentially impacting dendritic cells maturation and antimicrobial
585 responses. *Sci. Rep.* **10**, 5345 (2020).
- 586 27. Wilen, C. B. *et al.* Tropism for tuft cells determines immune promotion of norovirus
587 pathogenesis. *Science* **360**, 204–208 (2018).
- 588 28. Hold, G. L. *et al.* Role of the gut microbiota in inflammatory bowel disease
589 pathogenesis: what have we learnt in the past 10 years? *World J. Gastroenterol.* **20**, 1192–
590 1210 (2014).
- 591 29. Kim, D. H. & Cheon, J. H. Pathogenesis of Inflammatory Bowel Disease and Recent
592 Advances in Biologic Therapies. *Immune Netw.* **17**, 25–40 (2017).
- 593 30. Basic, M. *et al.* Norovirus triggered microbiota-driven mucosal inflammation in
594 interleukin 10-deficient mice. *Inflamm. Bowel Dis.* **20**, 431–443 (2014).
- 595 31. Lupfer, C. *et al.* Receptor interacting protein kinase 2-mediated mitophagy regulates
596 inflammasome activation during virus infection. *Nat. Immunol.* **14**, 480–488 (2013).
- 597 32. Yu, P., Li, Y., Wang, Y., Peppelenbosch, M. P. & Pan, Q. Lipopolysaccharide restricts
598 murine norovirus infection in macrophages mainly through NF- κ B and JAK-STAT signaling
599 pathway. *Virology* **546**, 109–121 (2020).
- 600 33. Michallet, M.-C. *et al.* TRADD protein is an essential component of the RIG-like
601 helicase antiviral pathway. *Immunity* **28**, 651–661 (2008).
- 602 34. Kernbauer, E., Ding, Y. & Cadwell, K. An enteric virus can replace the beneficial
603 function of commensal bacteria. *Nature* **516**, 94–98 (2014).
- 604 35. Hwang, S. *et al.* Murine norovirus: propagation, quantification, and genetic
605 manipulation. *Curr. Protoc. Microbiol.* **33**, 15K.2.1-61 (2014).
- 606 36. Wirtz, S. *et al.* Chemically induced mouse models of acute and chronic intestinal
607 inflammation. *Nat. Protoc.* **12**, 1295–1309 (2017).
- 608 37. Barreau, F. *et al.* CARD15/NOD2 is required for Peyer's patches homeostasis in mice.
609 *PLoS One* **2**, e523 (2007).
- 610 38. de Chaumont, F. *et al.* Icy: an open bioimage informatics platform for extended
611 reproducible research. *Nat. Methods* **9**, 690–696 (2012).
- 612
- 613
- 614
- 615
- 616
- 617
- 618

619 **Acknowledgments:**

620 For the *NOD^{GFP}* tissue imaging, we thank the UTechS PBI, a member of the France–
621 BioImaging infrastructure network supported by the French National Research Agency (ANR-
622 10–INSB–04, Investissement d’Avenir program) for microscope usage and assistance. We
623 thank the members of the Institut Pasteur Central Animal Facility for their assistance with
624 animal studies. I.G.B. laboratory was supported by Investissement d’Avenir program,
625 Laboratoire d’Excellence "Integrative Biology of Emerging Infectious Diseases" (ANR-10-
626 LABX-62-IBEID), by the Investissement d’Avenir program (RHU Torino Lumière ANR-16-
627 RHUS-0008), by the French National Research Agency (ANR-16-CE15-0021) and by R&D
628 grants from Danone and MEIJL.

629 M.C. was supported by grants from the Fondation pour la Recherche Médicale
630 (DEQ20130326475) and the European Union's European Regional Development Fund. We
631 thank T. Durand and MP. Fourmaux for excellent technical assistance.

632

633 **Authors’ contribution.** G.M. and M.T. performed the majority of the experimental procedures.
634 T.G. contributed to the preclinical studies in mice. M.D., T.G. and O.B. performed RT-qPCR
635 and ELISA analysis. M.D. contributed to histological analysis of colon sections. E.W.
636 performed viral genome quantification by RT-qPCR. R.W. and I.G.B. performed histological
637 analysis of tissue sections from NOD2- β -Gal reporter mice. All authors contributed to
638 interpretation of raw data and critically reviewed and/or modified the manuscript. G.M. and
639 M.C. conceived, designed and wrote the paper.

640

641 **Figures legends:**

642 **Figure 1: MNoV_S99 aggravates inflammation in DSS-induced colitis**

643 **A)**- In vivo experimental design, C57BL/6J mice were orally gavaged with PBS (200 μ L) or MNoV_S99
644 (5×10^7 TCID₅₀/mL), 7 days later, all the mice were given DSS in drinking water for a week. Mice were
645 weighed daily for weight loss comparison. **B)**- 3% DSS-induced colitis \pm MNoV_S99 (n = 6). **C)**- 5%
646 DSS-induced colitis \pm MNoV_S99 (n = 6). In B) and C), statistical differences were determined by 2
647 ways ANOVA test, * p < 0.05, ** p < 0.01, *** p < 0.001. **D)**- Representative images from H&E
648 staining of colon sections from mock-treated (Control) or 3% DSS \pm MNoV_S99 treated mice. Insets
649 showing higher magnifications from each image are presented next to each image. **E)**- Muscle wall
650 thickness was determined with Image J from colon section images. Percentages of increase are plotted
651 after baseline subtraction relatively to the values obtained from Control mice (n = 6). **F)**- Histological
652 intestinal epithelial inflammation scores comparison between Control, 3% or 5% DSS \pm MNoV_S99
653 treated mice (n = 6). **G)**- Colon length comparison between Control and 5% DSS \pm MNoV_S99 treated
654 mice **H)**- IL6 secretion levels from colon explants from mock or 5% DSS \pm MNoV_S99 treated mice (n
655 = 6). Statistical differences were determined with Student's t-Test in E, F, G and H.

656

657 **Figure 2: MNoV_S99 associated inflammation is NOD2 dependent**

658 Weight loss comparison in **A)**- 3% DSS-induced colitis \pm MNoV_S99 in WT mice (n = 6) and **B)**- in
659 *Nod2*^{-/-} mice (n = 6), statistical differences were determined by 2 ways ANOVA test, * p < 0,05, ** p <
660 0,01, *** p < 0,001. **C)**- Representative images of H&E staining of colon sections from mice in A) and
661 B). Insets showing higher magnifications are presented below each image. **D)**- Percentages of increase
662 of muscle wall thickness from colon section images are plotted after baseline subtraction relatively to
663 the values obtained from WT mice treated only with DSS (n = 6), statistical differences were determined
664 with Student's t-Test.

665

666 **Figure 3: MNoV_S99 associated pro-inflammatory signaling is NOD2 and MAVS-dependent in**
667 **myeloid lineage cells**

668 **A)**- Representative image showing an **intestinal villus** from a *Nod2*^{GFP} mouse. NOD2-GFP is shown in
669 green, Actin is stained with Phalloidin in grey and nuclei were stained with DAPI in blue; scale bare
670 represents 50 μ m. **B)**- Representative western blot showing STAT1 signaling pathway activation after
671 1h infection (Moi 1) with the mentioned MNoV strains in BMDM. Below are shown quantification of
672 pSTAT1 bands intensities normalized to β -actin and relative to mock-infected cells (n = 3, mean \pm
673 SEM), statistical differences were determined by 1 way ANOVA test * p < 0,05, ** p < 0,005. **C)**-
674 STAT1 activation in response to MNoV_S99 (Moi 1 or 5) in BMDM from WT vs *Nod2*^{-/-} infected for
675 30 or 60 minutes. The lower panel shows quantification of pSTAT1, bands intensities normalized to β -
676 actin and relative to mock-infected cells (n = 3, mean \pm SEM). Statistical differences were determined
677 by 2 ways ANOVA test * p < 0,05. **D)**- STAT1 activation in response to MNoV_S99 (Moi 0.1) in WT
678 or *Nod2*^{-/-} BMDM infected for 24h. Relative quantification of pSTAT1 (n = 3, mean \pm SEM) is shown
679 in the lower panel, statistical differences were determined by 2 ways ANOVA test *** p < 0,0005. **E)**-
680 STAT1 activation in response to MNoV_S99 (Moi 0.1) in BMDM from WT vs *Mavs*^{-/-} infected for 60
681 minutes. Relative quantification of pSTAT1 (n = 4, mean \pm SEM) are shown below. The lower panel
682 shows the mock vs Moi 0.1 infected cells quantification of pSTAT1. Statistical differences were
683 determined by 2 ways ANOVA test * p < 0,05.

684

685 **Figure 4: NOD2-dependent noroviral load**

686 **A)**- WT, *Nod2*^{-/-} and *Atg16l1*^{HM} mice (n = 5 each) were treated with 3% DSS and infected with
687 MNoV_CR6 (3×10^7 PFU). Inflammation was measured as percentage of increase in muscle thickness
688 at the anal-rectal junction in colonic section images plotted after baseline subtraction relatively to the
689 values obtained from WT mice. **B)**- The viral load measured in stools from WT, *Nod2*^{-/-} and *Atg16l1*^{HM}
690 mice at day 3, 5 and 7 post-infection with MNoV_CR6 (3×10^7 PFU) (n = 8). **C)**- The viral load measured
691 from indicated tissue samples from MNoV_CR6 infected (3×10^7 PFU) WT, *Nod2*^{-/-} and *Atg16l1*^{HM} mice
692 that were treated for 1 week with D55 3% (n = 5). **D)**- MNoV_S99 genome quantification from WT and
693 *Nod2*^{-/-} BMDM cells infected with MNoV_S99 (Moi 0.1 for 24h) (n = 3). All the statistical differences
694 were measured with Student's t-Test.

695 **Figure 5: NOD2-dependent pro-inflammatory signaling associated with MNoV_S99 infection and**
 696 **bacterial MDP**

697 Quantification of *Nod2* mRNA levels measured by RT-qPCR in **A)**- in Raw264.7 cells infected with
 698 MNoV_S99 (Moi 0.1 or 1, for 24h) and in **B)**- in BMDM from WT mice (Moi 0.1, for 24h) normalized
 699 to *ActB* and relative to mock-infected cells (n = 3). **C)**- Representative western blot showing STAT1
 700 and IκBα signaling pathways modulation in cell lysates from monocytes that were either infected with
 701 MNoV_S99 (Moi 5, 2h), treated with MDP (10ng/mL, 2h), or a combination of both. **D)**- Quantification
 702 and comparison of pSTAT1 and pIκBα signals between MNoV_S99 infected alone cells or in
 703 combination with MDP (10ng/mL, 2h), (n = 3), with the band intensity being normalized to β-ACTIN
 704 and relative to mock-treated cells. **E)** TNFα production by BMDC subjected to MNoV_S99's ssRNA
 705 (10μg/mL) alone, or supplemented with either MDP (1μg/mL) or MDP-DD (1μg/mL) overnight. **F)**
 706 TNFα production by BMDM cells infected with MNoV_S99 (Moi 0.1 or 1) for 6h, prior to being treated
 707 with MDP (10μg/mL) overnight. **G)** Net production of MNoV_S99 in Raw264.7 cells infected with
 708 MNoV_S99 (Moi 1 for 2h before MDP treatment (100ng/mL) for 24h). Statistical differences were
 709 analyzed with Student's t Test.

710
 711

712 **Supplementary Figures legends:**

713

714 **Supplementary Figure 1: Analysis of MNoV_S99 associated NOD2 dependent inflammation**
 715 **measurements from mice showed in Figure 2**

716 **A)** Colon length comparison between WT and *Nod2*^{-/-} from 3% DSS ± MNoV_S99 treated mice,
 717 statistical differences were determined with Student's t-Test, * p <0,02. **B)**- Histological scores
 718 comparison between WT and *Nod2*^{-/-} ± MNoV_S99 treated mice, statistical differences were determined
 719 with Student's t-Test, * p <0,05. **C)**- TNFα production of explants (ON) from WT and *Nod2*^{-/-} 3% DSS
 720 ± MNoV_S99 treated mice was measured by ELISA.

721

722 **Supplementary Figure 2: MNoV_S99 associated pro-inflammatory signaling pathways activation**

723 Representative western blot showing STAT1, IκBα, AKT and ERK signaling pathways' modulation in
 724 cell lysates from Raw264.7 that were either infected with MNoV_S99 (Moi 1) for various times or mock
 725 treated for 60 min. β-ACTIN was used as loading control.

726

727 **Tables:**

Table1: Antibodies used for Western Blotting		
Primary antibodies	Dilution	Supplier
anti-phosphoSTAT1 (Tyr 701) (58D6) (85kDa)	1/1000	Cell Signaling Technology
anti-STAT1(D1K9Y)(85kDa)	1/1000	
anti-phosphoIκBα (Ser32/36) (5A5) (40kDa)	1/1000	
anti-IκBα (44D4) (39kDa)	1/1000	
anti-phosphoAKT (Ser473) (D9E) (60kDa)	1/1000	
anti-AKT (60kDa)	1/1000	
anti-phospho44/42 MAPK (Erk1/2) (Trh202/Tyr204) (197G2) (42, 44 kDa)	1/1000	
anti-p44/42 (Erk1/2) (42, 44 kDa)	1/1000	
anti-β-ACTIN (45kDa)	1/500	
Secondary antibodies	Dilution	Supplier
anti-Rabbit_HRP (#711-035-152)	1/5000	Jackson
anti-Mouse_HRP (#115-005-003)	1/5000	Laboratories

728