



Quasi-rational solutions of the Hirota equation depending on multi-parameters and rogue waves

Pierre Gaillard

► To cite this version:

| Pierre Gaillard. Quasi-rational solutions of the Hirota equation depending on multi-parameters and rogue waves. 2023. hal-04101514

HAL Id: hal-04101514

<https://hal.science/hal-04101514v1>

Preprint submitted on 20 May 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Quasi-rational solutions of the Hirota equation depending on multi-parameters and rogue waves

Pierre Gaillard,

Université de Bourgogne Franche Comté,
Institut de mathématiques de Bourgogne,
9 avenue Alain Savary BP 47870
21078 Dijon Cedex, France :
E-mail : Pierre.Gaillard@u-bourgogne.fr

Abstract

Quasi-rational solutions to the Hirota equation are given. We construct explicit expressions of these solutions for the first orders.

As a byproduct, we get quasi-rational solutions to the focusing NLS equation and also rational solutions to the mKdV equation.

We study the patterns of these configurations in the (x, t) plane.

Key Words : Hirota equation, quasi-rational solutions.

PACS numbers :

33Q55, 37K10, 47.10A-, 47.35.Fg, 47.54.Bd

1 Introduction

We consider the Hirota (H) equation in the following normalization

$$iu_t + \alpha(u_{xx} + 2|u|^2u) - i\beta(u_{xxx} + 6|u|^2u_x) = 0, \quad (1)$$

with as usual the subscript meaning the partial derivatives and α, β real numbers.

Hirota proposed this equation in 1973 in a slightly different formulation

$$iu_t + \rho_1 u_{xx} + \delta_1 |u|^2 u + i\sigma_1 u_{xxx} + 3i\alpha_1 |u|^2 u_x = 0,$$

with $\alpha_1, \rho_1, \sigma_1, \delta_1$ real numbers satisfying $\alpha_1\rho_1 = \sigma_1\delta_1$ and constructed a kind of soliton in [1].

This last equation can be rewritten as (1) by choosing $\alpha_1 = -2\beta$, $\delta_1 = 2\alpha$, $\rho_1 = \alpha$, $\sigma_1 = -\beta$.

Maccari introduced in 1998 a two-dimensional extension of this equation. This equation is used to identify many kinds of nonlinear phenomena in the fields physics, in particular in optical fibers. It describes the evolution of the slowly varying amplitude of a nonlinear train in weakly nonlinear systems. Hirota equation is an integrable equation which has a number of physical applications, such as the propagation of optical pulses [3] or in plasma physics, [4].

A lot of methods to solve this equation have been presented such as general projective Riccati equation method [5], Darboux transformation [6, 7] or the trace method [8].

Here, we construct quasi-rational solutions for the first orders.

2 Quasi-rational solutions of order 1 to the Hirota equation

Theorem 2.1 *The function $v(x, t)$ defined by*

$$v(x, t) = \left(1 - 4 \frac{1 + 4i\alpha t}{1 + (2x + 12\beta t)^2 + 16\alpha^2 t^2} \right) e^{2iat} \quad (2)$$

is a solution to the Hirota equation (1)

$$iu_t + \alpha(u_{xx} + 2|u|^2 u) - i\beta(u_{xxx} + 6|u|^2 u_x) = 0.$$

Proof: It is sufficient to replace the expression of the solution given by (2) and check that (1) is verified.

We give another type of solution with one real parameter a_1 .

Theorem 2.2 *The function $v(x, t)$ defined by*

$$v(x, t) = \left(1 - 4 \frac{1 + i(4\alpha t - 24a_1)}{1 + (2x + 12\beta t)^2 + (4\alpha t - 24a_1)^2} \right) e^{i(2\alpha t - 6a_1)} \quad (3)$$

is a solution to the Hirota equation (1)

$$iu_t + \alpha(u_{xx} + 2|u|^2 u) - i\beta(u_{xxx} + 6|u|^2 u_x) = 0,$$

with a_1 arbitrary real parameter.

Proof: It is sufficient to replace the expression of the solution given by (3) and check that (1) is verified.

Remark 2.1 *It is possible to add other real parameters but it does not change drastically the structure of the solution as in the case of one parameter a_1 . So, we do not mention these parameters in the following.*




Figure 1. Solution of order 1 to (1); to the left $a_1 = 0$; in the center $a_1 = 0.2$; to the right $a_1 = 10$; with $\alpha = \beta = 1$.

Remark 2.2 If we choose, $\alpha = 1$, $\beta = 0$ and the parameters equal to 0, we recover the classical Peregrine breather solution to the NLS equation

$$iu_t + u_{xx} + 2|u|^2u = 0, \quad (4)$$

$$v(x, t) = \left(1 - 4 \frac{1 + 4it}{1 + 4x^2 + 16t^2} \right) e^{2it}. \quad (5)$$

This solution was first found by Peregrine [9].

It was recovered by the present author by using degeneracy of Riemann theta functions in [18].

Remark 2.3 If we choose, $\alpha = 0$, $\beta = 1$ and the parameters equal to 0, we recover a non singular rational solution to the mKdV equation

$$u_t = u_{xxx} + 6|u|^2u_x \quad (6)$$

$$v(x, t) = \left(1 - \frac{4}{1 + (2x + 12t)^2} \right). \quad (7)$$

This solution is different from those constructed by the author in [58].

3 Quasi-rational solutions of order 2 to the Hirota equation

Theorem 3.1 The function $v(x, t)$ defined by

$$v(x, t) = \left(1 - 12 \frac{n(x, t)}{d(x, t)} \right) e^{2i\alpha t} \quad (8)$$

with

$$n(x, t) = (2x + 12\beta t)^4 + 6(16\alpha^2 t^2 + 1)(2x + 12\beta t)^2 + 192\beta t(2x + 12\beta t) + 1280\alpha^4 t^4 + 288\alpha^2 t^2 - 3 + i(4\alpha t(2x + 12\beta t)^4 + 2(64\alpha^3 t^3 - 12\alpha t)(2x + 12\beta t)^2 + 768\alpha t^2 \beta(2x + 12\beta t) + 1024\alpha^5 t^5 + 128\alpha^3 t^3 - 60\alpha t)$$

and

$$d(x, t) = ((2x + 12\beta t)^2 + 16\alpha^2 t^2 + 1)^3 - 192\beta t(2x + 12\beta t)^3 - 24(16\alpha^2 t^2 - 1)(2x + 12\beta t)^2 + 576(16\alpha^2 t^2 + 1)\beta t(2x + 12\beta t) + 6144\alpha^4 t^4 + 8 + 1536\alpha^2 t^2 + 9216\beta^2 t^2$$

is a solution to the Hirota equation (1).

Proof: Replacing the expression of the solution given by (8), we check that the relation (1) is verified.

We can give also the solution depending on 2 real parameters.

The function $v(x, t)$ defined by

Theorem 3.2

$$v(x, t) = \left(1 - 12 \frac{n(x, t)}{d(x, t)}\right) e^{i(2\alpha t - 6a_1)} \quad (9)$$

with

$$n(x, t) = (2x + 12\beta t + 60b_1)^4 + 6((4\alpha t - 24a_1)^2 + 1)(2x + 12\beta t + 60b_1)^2 - 24(-8\beta t - 80b_1)(2x + 12\beta t + 60b_1) + 5(4\alpha t - 24a_1)^4 + 18(4\alpha t - 24a_1)^2 - 384a_1(4\alpha t - 24a_1) - 3 + i((4\alpha t - 24a_1)(2x + 12\beta t + 60b_1)^4 + 2((4\alpha t - 24a_1)^3 - 12\alpha t + 168a_1)(2x + 12\beta t + 60b_1)^2 - 24(4\alpha t - 24a_1)(-8\beta t - 80b_1)(2x + 12\beta t + 60b_1) + (4\alpha t - 24a_1)^5 + 2(4\alpha t - 24a_1)^3 - 192a_1(4\alpha t - 24a_1)^2 - 60\alpha t + 552a_1)$$

and

$$d(x, t) = ((2x + 12\beta t + 60b_1)^2 + (4\alpha t - 24a_1)^2 + 1)^3 + 24(-8\beta t - 80b_1)(2x + 12\beta t + 60b_1)^3 - 24((4\alpha t - 24a_1)^2 - 48a_1(4\alpha t - 24a_1) - 1)(2x + 12\beta t + 60b_1)^2 - 72((4\alpha t - 24a_1)^2 + 1)(-8\beta t - 80b_1)(2x + 12\beta t + 60b_1) + 24(4\alpha t - 24a_1)^4 - 384a_1(4\alpha t - 24a_1)^3 + 96(4\alpha t - 24a_1)^2 - 3456a_1(4\alpha t - 24a_1) + 36864a_1^2 + 144(-8\beta t - 80b_1)^2 + 8$$

is a solution to the Hirota equation (1).

Proof: Replacing the expression of the solution given by (9), we check that the relation (1) is verified.




Figure 2. Solution of order 1 to (1); to the left $a_1 = 0, b_1 = 0$; in the center $a_1 = 1, b_1 = 0$; to the right $a_1 = 10, b_1 = 0$; with $\alpha = \beta = 1$.

We observe a very fast evolution of the structure of the solution according to the parameter a_1 toward three peaks.




Figure 3. Solution of order 1 to (1); to the left $a_1 = 0, b_1 = 0.01$; in the center $a_1 = 0, b_1 = 0.05$; to the right $a_1 = 0, b_1 = 1$; with $\alpha = \beta = 8$.

The evolution previously observed is the same as in the case of the parameter a_1 .

Remark 3.1 If we choose, $\alpha = 1, \beta = 0$ and the parameters equal to 0, we recover solutions to the NLS equation

$$iu_t + u_{xx} + 2|u|^2u = 0, \quad (10)$$

$$u_k(x, t) = (1 - 12 \frac{n(x, t)}{d(x, t)}) e^{2it} \quad (11)$$

with

$$n(x, t) = 16x^4 + 24(16t^2 + 1)x^2 - 3 + 1280t^4 + 288t^2 + i(64tx^4 + 8(64t^3 -$$

$$12t)x^2 + 1024t^5 + 128t^3 - 60t)$$

and

$$d(x, t) = (4x^2 + 16t^2 + 1)^3 + 8 - 96(16t^2 - 1)x^2 + 6144t^4 + 1536t^2.$$

We recover the solutions given by the author in [18].

Remark 3.2 If we choose, $\alpha = 0$, $\beta = 1$ and the parameters equal to 0, we recover rational solutions to the mKdV equation

$$u_t = u_{xxx} + 6|u|^2u_x \quad (12)$$

$$u(x, t) = (1 - 12 \frac{n(x, t)}{d(x, t)}) \quad (13)$$

with

$$n(x, t) = (2x + 12t)^4 + 6(2x + 12t)^2 + 192t(2x + 12t) - 3$$

$$\text{and } v(x, t) = ((2x + 12t)^2 + 1)^3 - 192t(2x + 12t)^3 + 24(2x + 12t)^2 + 576t(2x + 12t) + 8 + 9216t^2.$$

These solutions are different from those presented in [64].

4 Quasi-rational solutions of order 3 to the Hirota equation

Theorem 4.1 The function $v(x, t)$ defined by

$$v(x, t) = \left(1 - 24 \frac{n(x, t)}{d(x, t)}\right) e^{2i\alpha t} \quad (14)$$

with

$$\begin{aligned} n(x, t) = & (2x + 12\beta t)^{10} + 15(1 + 16\alpha^2 t^2)(2x + 12\beta t)^8 + (210 - 960\alpha^2 t^2 + \\ & 12800\alpha^4 t^4)(2x + 12\beta t)^6 + (92160\alpha^2 t^3 \beta + 5760\beta t)(2x + 12\beta t)^5 + (7200\alpha^2 t^2 - \\ & 38400\alpha^4 t^4 + 286720\alpha^6 t^6 - 450 + 345600\beta^2 t^2)(2x + 12\beta t)^4 + (4915200\alpha^4 t^5 \beta - \\ & 57600\beta t + 1843200\alpha^2 t^3 \beta)(2x + 12\beta t)^3 + (1728000\alpha^4 t^4 + 1720320\alpha^6 t^6 + 2949120t^8 \alpha^8 - \\ & 43200\alpha^2 t^2(5 + 256\beta^2 t^2) - 675 - 691200\beta^2 t^2)(2x + 12\beta t)^2 + (55050240\alpha^6 t^7 \beta + \\ & 1382400\alpha^2 t^3 \beta - 86400\beta t - 22118400\beta^3 t^3 - 7372800\alpha^4 t^5 \beta)(2x + 12\beta t) + 8970240\alpha^6 t^6 + \\ & 32440320t^8 \alpha^8 + 11534336t^{10} \alpha^{10} + 675 + 115200\alpha^4 t^4(-17 + 1792\beta^2 t^2) + 10800\alpha^2 t^2(-3 + \\ & 1024\beta^2 t^2) + 1728000\beta^2 t^2 + i((-60\alpha t + 320\alpha^3 t^3)(2x + 12\beta t)^8 + (-600\alpha t - \\ & 8960\alpha^3 t^3 + 10240\alpha^5 t^5)(2x + 12\beta t)^6 + (122880\alpha^3 t^4 \beta - 23040\alpha t^2 \beta)(2x + 12\beta t)^5 + \\ & (-28800\alpha^3 t^3 - 215040\alpha^5 t^5 + 163840\alpha^7 t^7 + 1800\alpha t(-3 + 768\beta^2 t^2))(2x + 12\beta t)^4 - \\ & 14254080\alpha^7 t^7 + (3932160\alpha^5 t^6 \beta - 230400\alpha t^2 \beta + 2457600\alpha^3 t^4 \beta)(2x + 12\beta t)^3 + \\ & (1751040\alpha^5 t^5 - 983040\alpha^7 t^7 + 1310720t^9 \alpha^9 - 57600\alpha^3 t^3(7 + 256\beta^2 t^2) + 2700\alpha t(7 + \\ & 1024\beta^2 t^2))(2x + 12\beta t)^2 + 6553600t^9 \alpha^9 + (31457280\alpha^7 t^8 \beta + 43200\alpha t(-24\beta t - \\ & 2048\beta^3 t^3) - 5529600\alpha^3 t^4 \beta - 41287680\alpha^5 t^6 \beta)(2x + 12\beta t) + 4194304t^{11} \alpha^{11} + \\ & 4\alpha t(2x + 12\beta t)^{10} + 92160\alpha^5 t^5(-107 + 1792\beta^2 t^2) - 14400\alpha^3 t^3(11 + 5120\beta^2 t^2) - \\ & 2700\alpha t(-7 + 3584\beta^2 t^2)) \end{aligned}$$

and

$$\begin{aligned}
d(x, t) = & 2024 + (-1920 \alpha^2 t^2 + 120)(2x + 12\beta t)^8 + (7680 \alpha^2 t^2 - 61440 \alpha^4 t^4 + 2320 + \\
& 138240 \beta^2 t^2)(2x + 12\beta t)^6 + (1474560 \alpha^4 t^5 \beta - 552960 \alpha^2 t^3 \beta + 51840 \beta t)(2x + \\
& 12\beta t)^5 + (-368640 \alpha^4 t^4 + 3360 + 3840 \alpha^2 t^2(56 + 8640 \beta^2 t^2) + 2073600 \beta^2 t^2)(2x + \\
& 12\beta t)^4 + (31457280 \alpha^6 t^7 \beta + 16588800 \alpha^2 t^3 \beta + 88473600 \alpha^4 t^5 \beta - 345600 \beta t + \\
& 44236800 \beta^3 t^3)(2x + 12\beta t)^3 + (55050240 \alpha^6 t^6 + 15728640 t^8 \alpha^8 - 61440 \alpha^4 t^4(-326 + \\
& 2880 \beta^2 t^2) + 7680 \alpha^2 t^2(-76 + 8640 \beta^2 t^2) + 12144 - 6220800 \beta^2 t^2)(2x + 12\beta t)^2 + \\
& (188743680 \alpha^8 t^9 \beta + 47185920 \alpha^6 t^7 \beta - 66355200 \alpha^4 t^5 \beta + 1036800 \alpha^2 t^2(40 \beta t - \\
& 2048 \beta^3 t^3) + 648000 \beta t - 132710400 \beta^3 t^3)(2x + 12\beta t) + 22809600 \beta^2 t^2 + 243793920 t^8 \alpha^8 + \\
& 125829120 t^{10} \alpha^{10} + 2123366400 \beta^4 t^4 - 960 \beta t(2x + 12\beta t)^9 + 327680 \alpha^6 t^6(191 + \\
& 4032 \beta^2 t^2) + 61440 \alpha^4 t^4(599 + 8640 \beta^2 t^2) + 384 \alpha^2 t^2(3881 + 777600 \beta^2 t^2) + (1 + \\
& (2x + 12\beta t)^2 + 16 \alpha^2 t^2)^6
\end{aligned}$$

is a solution to the Hirota equation (1).

Proof: We have to check that the relation (1) is verified when we replace the expression of the solution given by (14).

We can give also the solutions to the Hirota depending on 4 real parameters. Because of the length of the expression, we only give it in the appendix.

We give patterns of the modulus of the solutions in function of the parameters.




Figure 4. Solution of order 3 to (1); to the left $a_1 = 0, b_1 = 0, a_2 = 0, b_2 = 0$; in the center $a_1 = 1, b_1 = 0, a_2 = 0, b_2 = 0$; to the right $a_1 = 2, b_1 = 0, a_2 = 10, b_2 = 0$; with $\alpha = \beta = 1$.




Figure 5. Solution of order 3 to (1); to the left $a_1 = 0, b_1 = 0, a_2 = 0.1, b_2 = 0$; in the center $a_1 = 0, b_1 = 10, a_2 = 1, b_2 = 0$; to the right $a_1 = 10, b_1 = 0, a_2 = 8, b_2 = 0$; with $\alpha = \beta = 1$.




Figure 6. Solution of order 3 to (1); to the left $a_1 = 0, b_1 = 0.1, a_2 = 0, b_2 = 0$; in the center $a_1 = 0, b_1 = 1, a_2 = 0, b_2 = 0$; to the right $a_1 = 2, b_1 = 10, a_2 = 0, b_2 = 0$; with $\alpha = \beta = 1$.




Figure 7. Solution of order 3 to (1); to the left $a_1 = 0, b_1 = 0, a_2 = 0, b_2 = 0.01$; in the center $a_1 = 0, b_1 = 10, a_2 = 0, b_2 = 0.1$; to the right $a_1 = 10, b_1 = 0, a_2 = 0, b_2 = 1$; with $\alpha = \beta = 1$.

In the evolution of the structure of the solutions according to the different parameters, we note the appearance of triangles with six peaks with more at least speed according to the parameters.

Remark 4.1 If we choose, $\alpha = 1, \beta = 0$ and the parameters equal to 0, we recover solutions to the NLS equation

$$iu_t + u_{xx} + 2|u|^2u = 0, \quad (15)$$

$$u_k(x, t) = (1 - 24 \frac{n(x, t)}{d(x, t)})e^{2i\alpha t} \quad (16)$$

with

$$n(x, t) = 16x^4 + 24((4t - 24000)^2 + 1)x^2 + 9215999997 + 5(4t - 24000)^4 + 18(4t - 24000)^2 - 1536000t + i(16(4t - 24000)x^4 + 8((4t - 24000)^3 - 12t + 168000)x^2 + (4t - 24000)^5 + 2(4t - 24000)^3 - 192000(4t - 24000)^2 - 60t + 552000)$$

and

$$d(x, t) = (4x^2 + (4t - 24000)^2 + 1)^3 + 119808000008 - 96((4t - 24000)^2 - 192000t + 115199999)x^2 + 24(4t - 24000)^4 - 384000(4t - 24000)^3 + 96(4t - 24000)^2 - 13824000t.$$

We recover the solutions constructed in [24].

Remark 4.2 If we choose, $\alpha = 0, \beta = 1$ and the parameters equal to 0, we recover rational solutions to the mKdV equation

$$u_t = u_{xxx} + 6|u|^2u_x \quad (17)$$

$$u_k(x, t) = (1 - 24 \frac{n(x, t)}{d(x, t)})e^{2i\alpha t} \quad (18)$$

with

$$n(x, t) = (2x + 12t)^4 + 6(2x + 12t)^2 + 192t(2x + 12t) - 3$$

$$\text{and } d(x, t) = ((2x + 12t)^2 + 1)^3 - 192t(2x + 12t)^3 + 24(2x + 12t)^2 + 576t(2x + 12t) + 8 + 9216t^2.$$

These solutions are different from those given in [58, 64].

5 Conclusion

Quasi-rational solutions to the Hirota equation have been given for the first orders. As a byproduct, we have recovered quasi-rational solutions to the NLS equation and rational solutions to the mKdV equation.

We can mention some other recent works about this equation. For example,

in [10], by using the inverse scattering transform an explicit soliton solution formula for the Hirota equation has been constructed. This formula allows to get, as a particular case, the N-soliton solution, the breather solution and, most relevantly, a new class of solutions called multi-pole soliton solutions.

By using the Hirota direct method, lump-soliton to the Hirota equation were given in [11].

In [12], the mixed localized wave solutions of the Hirota equation are constructed through the modified Darboux transformation. One of them is the mixed 1-breather and 1-rogue wave solution and the other two are the mixed 1-breather and 2-rogue wave solution, and the mixed 2-breather and 1-rogue wave solution. These mixed localized wave solutions are presented graphically by choosing proper parameters and we discuss their dynamic behavior and control briefly.

References

- [1] R. Hirota, Exact envelope-soliton solutions of a nonlinear wave equation, *J. Math. Phys.*, V. 14, 805809, 1973.
- [2] A. Maccari, , A generalized Hirota equation in 2+1 dimensions, *J. Math. Phys.*, V. 39, 65476551, 1998
- [3] S.M. Hoseini, T.R. Marchant, Soliton perturbation theory for a higher order Hirota equation, *Math. Comput. Simul.*, V. 80, 770778, 2008
- [4] R.F. Zhang, B.L. Guo, Global attractor for Hirota equation, *Appl. Math. J. Chin. Univ.*, V. 23, N. 1, 5764, 2008
- [5] Q. Wang, Y. Chen, B. Li, H. Zhang, New exact travelling solutions to Hirota equation and (1+1)-dimensional dispersive long wave equation,, *Commun. Theor. Phys.*, V. 41, 821828, 2004
- [6] L. Li, Z. Wu, L. Wang, J. He, High-order rogue waves for the Hirota equation, *Ann. Phys.*, V. 334, 198211, 2013
- [7] A. Ankiewicz, J.M. Soto-Crespo, N. Akhmediev, Rogue waves and rational solutions of the Hirota equation, *Phys. Rev. E*, V. 81, 046602-18, 2010
- [8] J.J. Shu, Exact n-envelope-soliton solutions of the Hirota equation, *Opt. Appl.*, V. 33, 539546, 2003
- [9] D. Peregrine, Water waves, nonlinear Schrodinger equations and their solutions, *J. Aust. Math. Soc. B*, V. 25, 1643, 1983
- [10] F. Demontis, G. Ortenzi, C. Van der Mee, Exact solutions of the Hirota equation and vortex filaments motion, *Phys. D*, V. 313, 6180, 2015

- [11] Y. Zhou, S. Manukure, W.X. Ma, Lump and lump-soliton solutions to the HirotaSatsumaIto equation, Comm. In Nonlin. Sci And Num. Sim., V. 68, 56-62, 2019
- [12] D.Y. Liu, H.M. Yu, Mixed localized wave solutions of the Hirota equation, Appl. Math. Let., V. 118, 107154-1-7, 2021
- [13] P. Gaillard, V.B. Matveev, Wronskian addition formula and its applications, Max-Planck-Institut für Mathematik, MPI 02-31, V. **161**, 2002
- [14] P. Gaillard, A new family of deformations of Darboux-Pöschl-Teller potentials, Lett. Math. Phys., V. **68**, 77-90, 2004
- [15] P. Gaillard, V.B. Matveev, New formulas for the eigenfunctions of the two-particle Calogero-Moser system, Lett. Math. Phys., V. **89**, 1-12, 2009
- [16] P. Gaillard, V.B. Matveev, Wronskian and Casorai determinant representations for Darboux-Pöschl-Teller potentials and their difference extensions, J. Phys A : Math. Theor., V. **42**, 404409-1-16, 2009
- [17] P. Dubard, P. Gaillard, C. Klein, V. B. Matveev, On multi-rogue wave solutions of the NLS equation and positon solutions of the KdV equation, Eur. Phys. J. Spe. Top., V. **185**, 247-258, 2010
- [18] P. Gaillard, Families of quasi-rational solutions of the NLS equation and multi-rogue waves, J. Phys. A : Meth. Theor., V. **44**, 435204-1-15, 2011
- [19] P. Gaillard, Wronskian representation of solutions of the NLS equation and higher Peregrine breathers, J. Math. Sciences : Adv. Appl., V. **13**, N. 2, 71-153, 2012
- [20] P. Gaillard, Degenerate determinant representation of solution of the NLS equation, higher Peregrine breathers and multi-rogue waves, J. Math. Phys., V. **54**, 013504-1-32, 2013
- [21] P. Gaillard, Wronskian representation of solutions of NLS equation and seventh order rogue waves, J. Mod. Phys., V. **4**, N. 4, 246-266, 2013
- [22] P. Gaillard, V.B. Matveev, Wronskian addition formula and Darboux-Pöschl-Teller potentials, J. Math., V. **2013**, ID 645752, 1-10, 2013
- [23] P. Gaillard, Two-parameters determinant representation of seventh order rogue waves solutions of the NLS equation, J. Theor. Appl. Phys., V. **7**, N. **45**, 1-6, 2013
- [24] P. Gaillard, Deformations of third order Peregrine breather solutions of the NLS equation with four parameters, Phys. Rev. E, V. **88**, 042903-1-9, 2013

- [25] P. Gaillard, Two parameters deformations of ninth Peregrine breather solution of the NLS equation and multi rogue waves, *J. Math.*, V. **2013**, 1-111, 2013
- [26] P. Gaillard, Six-parameters deformations of fourth order Peregrine breather solutions of the NLS equation, *J. Math. Phys.*, V. **54**, 073519-1-22, 2013
- [27] P. Gaillard, The fifth order Peregrine breather and its eight-parameters deformations solutions of the NLS equation, *Commun. Theor. Phys.*, V. **61**, 365-369, 2014
- [28] P. Gaillard, Ten parameters deformations of the sixth order Peregrine breather solutions of the NLS equation, *Phys. Scripta*, V. **89**, 015004-1-7, 2014
- [29] P. Gaillard, Higher order Peregrine breathers, their deformations and multi-rogue waves, *J. Of Phys. : Conf. Ser.*, V. **482**, 012016-1-7, 2014
- [30] P. Gaillard, M. Gastineau, Eighteen parameter deformations of the Peregrine breather of order ten solutions of the NLS equation, *Int. J. Mod. Phys. C*, V. **26**, N. 2, 1550016-1-14, 2014
- [31] P. Gaillard, Two parameters wronskian representation of solutions of nonlinear Schrödinger equation, eight Peregrine breather and multi-rogue waves, *J. Math. Phys.*, V. **5**, 093506-1-12, 2014
- [32] P. Gaillard, Hierarchy of solutions to the NLS equation and multi-rogue waves, *J. Phys. : Conf. Ser.*, V. **574**, 012031-1-5, 2015
- [33] P. Gaillard, Tenth Peregrine breather solution of the NLS, *Ann. Phys.*, V. **355**, 293-298, 2015
- [34] P. Gaillard, M. Gastineau, The Peregrine breather of order nine and its deformations with sixteen parameters solutions of the NLS equation *Phys. Lett. A*, V. **379**, 1309-1313, 2015
- [35] P. Gaillard, Other 2N-2 parameters solutions to the NLS equation and 2N+1 highest amplitude of the modulus of the N-th order AP breather, *J. Phys. A: Math. Theor.*, V. **48**, 145203-1-23, 2015
- [36] P. Gaillard, Multi-parametric deformations of the Peregrine breather of order N solutions to the NLS equation and multi-rogue waves, *Adv. Res.*, V. **4**, N. **5**, 346-364, 2015
- [37] P. Gaillard, Higher order Peregrine breathers solutions to the NLS equation, *Jour. Phys. : Conf. Ser.*, V. **633**, 012106-1-6, 2016
- [38] P. Gaillard, M. Gastineau Patterns of deformations of Peregrine breather of order 3 and 4, solutions to the NLS equation with multi-parameters, *Journal of Theoretical and Applied Physics*, V. **10**, 1-7, 2016

- [39] P. Gaillard, M. Gastineau Twenty parameters families of solutions to the NLS equation and the eleventh Peregrine breather, *Commun. Theor. Phys.*, V. **65**, N. **2**, 136-144, 2016
- [40] P. Gaillard, Rational solutions to the KPI equation and multi rogue waves, *Annals Of Physics*, V. **367**, 1-5, 2016
- [41] P. Gaillard, M. Gastineau Twenty two parameters deformations of the twelfth Peregrine breather solutions to the NLS equation, *Adv. Res.*, V. **10**, 83-89, 2016
- [42] P. Gaillard, Towards a classification of the quasi rational solutions to the NLS equation, *Theor. And Math. Phys.*, V. **189**, N. **1**, 1440-1449, 2016
- [43] P. Gaillard, Fredholm and Wronskian representations of solutions to the KPI equation and multi-rogue waves, *Jour. of Math. Phys.*, V. **57**, 063505-1-13, doi: 10.1063/1.4953383, 2016
- [44] P. Gaillard, M. Gastineau Families of deformations of the thirteenth Peregrine breather solutions to the NLS equation depending on twenty four parameters, *Jour. Of Bas. And Appl. Res. Int.*, V. **21**, N. 3, 130-139, 2017
- [45] P. Gaillard, From Fredholm and Wronskian representations to rational solutions to the KPI equation depending on $2N^2$ parameters, *Int. Jour. of Appl. Sci. And Math.*, V. **4**, N. 3, 60-70, 2017
- [46] P. Gaillard, Families of rational solutions of order 5 to the KPI equation depending on 8 parameters, *New Hor. in Math. Phys.*, V. 1, N. 1, 26-31, 2017
- [47] P. Gaillard, 6-th order rational solutions to the KPI Equation depending on 10 parameters, *Jour. Of Bas. And Appl. Res. Int.*, V. **21**, N. 2, 92-98, 2017
- [48] P. Gaillard, N -Order rational solutions to the Johnson equation depending on $2N - 2$ parameters, *Int. Jour. of Adv. Res. in Phys. Sci.*, V. **4**, N. 9, 19-37, 2017
- [49] P. Gaillard, Families of rational solutions to the KPI equation of order 7 depending on 12 parameters, *Int. Jour. of Adv. Res. in Phys. Sci.*, V. **4**, N. 11, 24-30, 2017
- [50] P. Gaillard, Rational solutions to the Johnson equation and rogue waves, *Int. Jour. of Inn. In Sci. and Math.*, V. **6**, N. 1, 14-19, 2018
- [51] P. Gaillard, Multiparametric families of solutions of the KPI equation, the structure of their rational representations and multi-rogue waves, *Theo. And Mat. Phys.*, V. **196**, N. 2, 1174-1199, 2018

- [52] P. Gaillard, The Johnson equation, Fredholm and wronskian representations of solutions and the case of order three, *Adv. In Math. Phys.*, V. **2018**, 1-18, 2018
- [53] P. Gaillard, Families of Solutions of Order 5 to the Johnson Equation Depending on 8 Parameters, *NHIMP*, V. **2**, N. 4, 53-61, 2018
- [54] P. Gaillard, Multiparametric families of solutions to the Johnson equation, *J. Phys. : Conf. Series*, V. **1141**, 012102-1-10, 2018
- [55] P. Gaillard, Rational solutions to the Boussinesq equation, *Fund. Jour. Of Math. And Appl.*, V. , 109-112, 2019
- [56] P. Gaillard, Differential relations for the solutions to the NLS equation and their different representations, *Comm. In Adv. Math. Sci.*, V. **2**, N. 4, 1-4, 2019
- [57] P. Gaillard, Multi-parametric families of solutions of order N to the Boussinesq and KP equations and the degenerate rational case, *UJMA*, V. **3**, N. 2, 44-52, 2020
- [58] P. Gaillard, The mKdV equation and multi-parameters rational solutions, *Wave Motion*, V. **100**, 102667-1-9, 2021
- [59] P. Gaillard, Degenerate Riemann theta functions, Fredholm and wronskian representations of the solutions to the KdV equation and the degenerate rational case, *Jour. Of Geom. And Phys.*, V. **161**, 104059-1-12, 2021
- [60] P. Gaillard, 2N parameter solutions to the Burgers equation, *Jour. Of Appl. Nonlin. Dyn.*, V. **1**, N. **1**, 1-6, 2021
- [61] P. Gaillard, Multi-Parametric rational Solutions to the KdV equation, *Asi. Jour. Of Res. and Rev. In Phys.*, V. **3**, N. 3, 14-21, 2021
- [62] P. Gaillard, Multiparametric solutions to the Gardner equation and the degenerate rational case, *Jour. Of Appl. Ana. And Appl.*, V. **11**, N. 4, 2102-2113, 2021
- [63] P. Gaillard, Other families of rational solutions to the KPI equation, *Asi. Res. Jour Of Math.*, V. **17**, N. 6, 27-34, 2021
- [64] P. Gaillard, Rational solutions to the mKdV equation associated to particular polynomials, *Wave Motion*, V. **107**, 102824-1-11, 2021
- [65] P. Gaillard, Multiparametric Rational Solutions of Order N to the KPI Equation and the Explicit Case of Order 3, *Arch. Of cur. Res. Int.*, V. **21**, N . 6, 58-71, 2021
- [66] P. Gaillard, Rational solutions to the KPI equation from particular polynomials, *Wave Motion*, V. **108**, 102828-1-9, 2022

- [67] P. Gaillard, Rogue Waves of the Lakshmanan Porsezian Daniel Equation Depending on Multi-parameters, As. Jour. Of Adv. Res. And Rep., V 16, N. 3, 32-40, 2022
- [68] P. Gaillard, Quasi-rational and rational solutions to the defocusing NLS equation, Jour. Of Math. Sci. And Mod. Res. And Rep., V 5, N. 1, 24-34, 2022
- [69] P. Gaillard, Rational solutions to the KdV equation in terms of particular polynomials, Jour. Of Nonlin. Mod. and Ana., V. 4, N. 4, 615-626, 2022
- [70] P. Gaillard, Rational solutions to the Painlevé II equation from particular polynomials, Int. J. Of Appl. Nonlin. Sci., V. 3, N. 3, 189-196, 2022
- [71] P. Gaillard, Multi-lump solutions to the KPI equation with a zero degree of derivation, Jour. Of Theor. And Appl. Phys., V. 16, N. 4, 1-8, 2022
- [72] P. Gaillard, Families of solutions to the KPI equation given by an extended Darboux transformation, Part. Diff. Equ. And Appl., V. 3, N. 80, 1-14, 2022

Appendix

Solution of order 3 to the Hirota equation depending on 4 real parameters :
The function $v(x, t)$ defined by

$$v(x, t) = \left(1 - 24 \frac{n(x, t)}{d(x, t)} \right) e^{2 i \alpha t} \quad (19)$$

with

$$\begin{aligned} n(x, t) = & 675 + 15(1 + (4\alpha t - 24a_1 + 120a_2)^2)(2x + 12\beta t + 60b_1 - 280b_2)^8 + \\ & (210 - 60(4\alpha t - 24a_1 + 120a_2)^2 + 50(4\alpha t - 24a_1 + 120a_2)^4 + 480(4\alpha t - 24a_1 + 120a_2)(16a_1 - 160a_2))(2x + 12\beta t + 60b_1 - 280b_2)^6 + (-720(4\alpha t - 24a_1 + 120a_2)^2(-8\beta t - 80b_1 + 560b_2) + 5760\beta t - 11520b_1 + 564480b_2)(2x + 12\beta t + 60b_1 - 280b_2)^5 + (450(4\alpha t - 24a_1 + 120a_2)^2 - 150(4\alpha t - 24a_1 + 120a_2)^4 + 70(4\alpha t - 24a_1 + 120a_2)^6 + 1200(4\alpha t - 24a_1 + 120a_2)^3(16a_1 - 160a_2) - 450 + 5400(-8\beta t - 80b_1 + 560b_2)^2 - 1800(16a_1 - 160a_2)^2 + 3600(4\alpha t - 24a_1 + 120a_2)(16a_1 - 224a_2))(2x + 12\beta t + 60b_1 - 280b_2)^4 + (-2400(4\alpha t - 24a_1 + 120a_2)^4(-8\beta t - 80b_1 + 560b_2) + 28800(4\alpha t - 24a_1 + 120a_2)(-8\beta t - 80b_1 + 560b_2)(16a_1 - 160a_2) - 57600\beta t - 806400b_1 + 7257600b_2 - 7200(4\alpha t - 24a_1 + 120a_2)^2(-16\beta t - 128b_1 + 672b_2))(2x + 12\beta t + 60b_1 - 280b_2)^3 + (6750(4\alpha t - 24a_1 + 120a_2)^4 + 420(4\alpha t - 24a_1 + 120a_2)^6 + 45(4\alpha t - 24a_1 + 120a_2)^8 - 2700(4\alpha t - 24a_1 + 120a_2)^2(5+4(-8\beta t - 80b_1 + 560b_2)^2 - 12(16a_1 - 160a_2)^2) - 675 - 10800(-8\beta t - 80b_1 + 560b_2)^2 - 10800(16a_1 - 160a_2)^2 + 21600(4\alpha t - 24a_1 + 120a_2)(32a_1 - 384a_2) - 7200(4\alpha t - 24a_1 + 120a_2)^3(32a_1 - 128a_2))(2x + 12\beta t + 60b_1 - 280b_2)^2 + (-1680(4\alpha t - 24a_1 + 120a_2)^6(-8\beta t - 80b_1 + 560b_2) - 28800(4\alpha t - 24a_1 + 120a_2)^3(-8\beta t - 80b_1 + 560b_2)(16a_1 - 160a_2) - 10800(4\alpha t - 24a_1 + 120a_2)^2(-8\beta t - 272b_1 + 3248b_2) - 86400\beta t - 1209600b_1 + 10886400b_2 + 43200(-8\beta t - 80b_1 + 560b_2)^3 + 43200(-8\beta t - 80b_1 + 560b_2)(16a_1 - 160a_2)^2 + \end{aligned}$$

$$\begin{aligned}
& 3600(4\alpha t - 24a_1 + 120a_2)^4(-8\beta t + 80b_1 - 1680b_2) - 86400(4\alpha t - 24a_1 + 120a_2)((-8\beta t - 80b_1 + 560b_2)(16a_1 - 160a_2) + (16a_1 - 160a_2)(32b_1 - 448b_2) - 64(-8\beta t - 80b_1 + 560b_2)a_2)(2x + 12\beta t + 60b_1 - 280b_2) - 720(4\alpha t - 24a_1 + 120a_2)^7(16a_1 - 160a_2) + 450(4\alpha t - 24a_1 + 120a_2)^4(-17 + 28(-8\beta t - 80b_1 + 560b_2)^2 + 12(16a_1 - 160a_2)^2) - 3600(4\alpha t - 24a_1 + 120a_2)^3(48a_1 - 1376a_2) - 720(4\alpha t - 24a_1 + 120a_2)^5(272a_1 - 3168a_2) + 10800(4\alpha t - 24a_1 + 120a_2)(-16a_1 + 224a_2 + 4(-8\beta t - 80b_1 + 560b_2)^2(16a_1 - 160a_2) + 4(16a_1 - 160a_2)^3) + 675(4\alpha t - 24a_1 + 120a_2)^2(-3 + 16(-8\beta t - 80b_1 + 560b_2)^2 + 16(16a_1 - 160a_2)^2 - 128(-8\beta t - 80b_1 + 560b_2)(32b_1 - 448b_2) - 8192(16a_1 - 160a_2)a_2) - 86400(-8\beta t - 80b_1 + 560b_2)(32b_1 - 448b_2) - 11059200(16a_1 - 160a_2)a_2 + (2x + 12\beta t + 60b_1 - 280b_2)^{10} + 2190(4\alpha t - 24a_1 + 120a_2)^6 + 27000(-8\beta t - 80b_1 + 560b_2)^2 + 91800(16a_1 - 160a_2)^2 + 495(4\alpha t - 24a_1 + 120a_2)^8 + 11(4\alpha t - 24a_1 + 120a_2)^{10} + 86400(32b_1 - 448b_2)^2 + 353894400a_2^2 + i(-21600(-8\beta t - 80b_1 + 560b_2)^2(16a_1 - 160a_2) + (4\alpha t - 24a_1 + 120a_2)(2x + 12\beta t + 60b_1 - 280b_2)^{10} + (-60\alpha t + 840a_1 - 6600a_2 + 5(4\alpha t - 24a_1 + 120a_2)^3)(2x + 12\beta t + 60b_1 - 280b_2)^8 + (-600\alpha t - 240a_1 + 58800a_2 - 140(4\alpha t - 24a_1 + 120a_2)^3 + 10(4\alpha t - 24a_1 + 120a_2)^5 + 240(4\alpha t - 24a_1 + 120a_2)^2(16a_1 - 160a_2))(2x + 12\beta t + 60b_1 - 280b_2)^6 + (-240(4\alpha t - 24a_1 + 120a_2)^3(-8\beta t - 80b_1 + 560b_2) - 1440(-8\beta t - 80b_1 + 560b_2)(16a_1 - 160a_2) + 720(4\alpha t - 24a_1 + 120a_2)(-8\beta t - 176b_1 + 1904b_2))(2x + 12\beta t + 60b_1 - 280b_2)^5 + (-450(4\alpha t - 24a_1 + 120a_2)^3 - 210(4\alpha t - 24a_1 + 120a_2)^5 + 10(4\alpha t - 24a_1 + 120a_2)^7 + 300(4\alpha t - 24a_1 + 120a_2)^4(16a_1 - 160a_2) + 450(4\alpha t - 24a_1 + 120a_2)(-3 + 12(-8\beta t - 80b_1 + 560b_2)^2 - 4(16a_1 - 160a_2)^2) - 14400a_1 + 259200a_2 + 1800(4\alpha t - 24a_1 + 120a_2)^2(16a_1 - 224a_2))(2x + 12\beta t + 60b_1 - 280b_2)^4 + (-480(4\alpha t - 24a_1 + 120a_2)^5(-8\beta t - 80b_1 + 560b_2) + 14400(4\alpha t - 24a_1 + 120a_2)^2(-8\beta t - 80b_1 + 560b_2)(16a_1 - 160a_2) + 7200(4\alpha t - 24a_1 + 120a_2)(-8\beta t - 48b_1 + 112b_2) - 2400(4\alpha t - 24a_1 + 120a_2)^3(-16\beta t - 128b_1 + 672b_2) - 14400(-8\beta t - 80b_1 + 560b_2)(16a_1 - 160a_2) - 14400(16a_1 - 160a_2)(32b_1 - 448b_2) + 921600(-8\beta t - 80b_1 + 560b_2)a_2)(2x + 12\beta t + 60b_1 - 280b_2)^3 + (1710(4\alpha t - 24a_1 + 120a_2)^5 - 60(4\alpha t - 24a_1 + 120a_2)^7 + 5(4\alpha t - 24a_1 + 120a_2)^9 - 900(4\alpha t - 24a_1 + 120a_2)^3(7 + 4(-8\beta t - 80b_1 + 560b_2)^2 - 12(16a_1 - 160a_2)^2) + 675(4\alpha t - 24a_1 + 120a_2)(7 + 16(-8\beta t - 80b_1 + 560b_2)^2 + 16(16a_1 - 160a_2)^2) - 345600a_1 + 4492800a_2 - 21600(-8\beta t - 80b_1 + 560b_2)^2(16a_1 - 160a_2) - 21600(16a_1 - 160a_2)^3 + 691200(4\alpha t - 24a_1 + 120a_2)^2a_2 - 1800(4\alpha t - 24a_1 + 120a_2)^4(64a_1 - 448a_2))(2x + 12\beta t + 60b_1 - 280b_2)^2 + (-240(4\alpha t - 24a_1 + 120a_2)^7(-8\beta t - 80b_1 + 560b_2) - 7200(4\alpha t - 24a_1 + 120a_2)^4(-8\beta t - 80b_1 + 560b_2)(16a_1 - 160a_2) + 10800(4\alpha t - 24a_1 + 120a_2)(-24\beta t - 400b_1 + 3920b_2 + 4(-8\beta t - 80b_1 + 560b_2)^3 + 4(-8\beta t - 80b_1 + 560b_2)(16a_1 - 160a_2)^2) + 3600(4\alpha t - 24a_1 + 120a_2)^3(-24\beta t - 176b_1 + 784b_2) + 720(4\alpha t - 24a_1 + 120a_2)^5(-56\beta t - 400b_1 + 1680b_2) + 21600(-8\beta t - 80b_1 + 560b_2)(16a_1 - 160a_2) + 43200(16a_1 - 160a_2)(32b_1 - 448b_2) - 2764800(-8\beta t - 80b_1 + 560b_2)a_2 - 43200(4\alpha t - 24a_1 + 120a_2)^2((-8\beta t - 80b_1 + 560b_2)(16a_1 - 160a_2) + (16a_1 - 160a_2)(32b_1 - 448b_2) - 64(-8\beta t - 80b_1 + 560b_2)a_2))(2x + 12\beta t + 60b_1 - 280b_2) - 90(4\alpha t - 24a_1 + 120a_2)^8(16a_1 - 160a_2) + 90(4\alpha t - 24a_1 + 120a_2)^5(-107 + 28(-8\beta t - 80b_1 + 560b_2)^2 + 12(16a_1 - 160a_2)^2) + 5400(4\alpha t - 24a_1 + 120a_2)^2(176a_1 - 2464a_2 + 4(-8\beta t - 80b_1 + 560b_2)^2(16a_1 - 160a_2) + 4(16a_1 - 160a_2)^3) -
\end{aligned}$$

$$\begin{aligned}
& 120(4\alpha t - 24a_1 + 120a_2)^6(80a_1 - 1248a_2) + 900(4\alpha t - 24a_1 + 120a_2)^4(464a_1 - \\
& 4000a_2) - 225(4\alpha t - 24a_1 + 120a_2)^3(11 + 80(-8\beta t - 80b_1 + 560b_2)^2 + 80(16a_1 - \\
& 160a_2)^2 + 128(-8\beta t - 80b_1 + 560b_2)(32b_1 - 448b_2) + 8192(16a_1 - 160a_2)a_2) - \\
& 675(4\alpha t - 24a_1 + 120a_2)(-7 + 56(-8\beta t - 80b_1 + 560b_2)^2 + 88(16a_1 - 160a_2)^2 - \\
& 128(-8\beta t - 80b_1 + 560b_2)(32b_1 - 448b_2) - 128(32b_1 - 448b_2)^2 - 524288a_2^2) + \\
& 5529600(-8\beta t - 80b_1 + 560b_2)^2a_2 - 5529600(16a_1 - 160a_2)^2a_2 - 172800(-8\beta t - \\
& 80b_1 + 560b_2)(16a_1 - 160a_2)(32b_1 - 448b_2) + 64800(16a_1 - 160a_2)^3 - 870(4\alpha t - \\
& 24a_1 + 120a_2)^7 + 25(4\alpha t - 24a_1 + 120a_2)^9 + (4\alpha t - 24a_1 + 120a_2)^{11} - 151200a_1 + \\
& 1857600a_2)
\end{aligned}$$

and

$$\begin{aligned}
d(x, t) = & 2024 - 777600(-8\beta t - 80b_1 + 560b_2)(32b_1 - 448b_2) - 82944000(16a_1 - \\
& 160a_2)a_2 + (-120(4\alpha t - 24a_1 + 120a_2)^2 + 360(4\alpha t - 24a_1 + 120a_2)(16a_1 - \\
& 160a_2) + 120)(2x + 12\beta t + 60b_1 - 280b_2)^8 + (480(4\alpha t - 24a_1 + 120a_2)^2 - \\
& 240(4\alpha t - 24a_1 + 120a_2)^4 + 960(4\alpha t - 24a_1 + 120a_2)^3(16a_1 - 160a_2) + 2320 + \\
& 2160(-8\beta t - 80b_1 + 560b_2)^2 + 5040(16a_1 - 160a_2)^2 - 1440(4\alpha t - 24a_1 + \\
& 120a_2)(64a_1 - 960a_2))(2x + 12\beta t + 60b_1 - 280b_2)^6 + (-720(4\alpha t - 24a_1 + \\
& 120a_2)^4(-8\beta t - 80b_1 + 560b_2) - 17280(4\alpha t - 24a_1 + 120a_2)(-8\beta t - 80b_1 + \\
& 560b_2)(16a_1 - 160a_2) + 4320(4\alpha t - 24a_1 + 120a_2)^2(-8\beta t - 176b_1 + 1904b_2) + \\
& 51840\beta t + 103680b_1 + 2177280b_2)(2x + 12\beta t + 60b_1 - 280b_2)^5 + (-1440(4\alpha t - \\
& 24a_1 + 120a_2)^4 + 720(4\alpha t - 24a_1 + 120a_2)^5(16a_1 - 160a_2) + 240(4\alpha t - 24a_1 + \\
& 120a_2)^2(56 + 135(-8\beta t - 80b_1 + 560b_2)^2 - 45(16a_1 - 160a_2)^2) + 32400(4\alpha t - \\
& 24a_1 + 120a_2)(16a_1 - 288a_2) + 7200(4\alpha t - 24a_1 + 120a_2)^3(48a_1 - 544a_2) + \\
& 3360 + 32400(-8\beta t - 80b_1 + 560b_2)^2 - 54000(16a_1 - 160a_2)^2 + 86400(-8\beta t - \\
& 80b_1 + 560b_2)(32b_1 - 448b_2) + 5529600(16a_1 - 160a_2)a_2)(2x + 12\beta t + 60b_1 - \\
& 280b_2)^4 + (-960(4\alpha t - 24a_1 + 120a_2)^6(-8\beta t - 80b_1 + 560b_2) + 57600(4\alpha t - \\
& 24a_1 + 120a_2)^3(-8\beta t - 80b_1 + 560b_2)(16a_1 - 160a_2) - 43200(4\alpha t - 24a_1 + \\
& 120a_2)^2(-24\beta t - 272b_1 + 2128b_2) - 7200(4\alpha t - 24a_1 + 120a_2)^4(-48\beta t - \\
& 448b_1 + 2912b_2) - 345600\beta t - 5529600b_1 + 53222400b_2 - 86400(-8\beta t - 80b_1 + \\
& 560b_2)^3 - 86400(-8\beta t - 80b_1 + 560b_2)(16a_1 - 160a_2)^2 + 172800(4\alpha t - 24a_1 + \\
& 120a_2)((-8\beta t - 80b_1 + 560b_2)(16a_1 - 160a_2) - (16a_1 - 160a_2)(32b_1 - 448b_2) + \\
& 64(-8\beta t - 80b_1 + 560b_2)a_2))(2x + 12\beta t + 60b_1 - 280b_2)^3 + (13440(4\alpha t - \\
& 24a_1 + 120a_2)^6 + 240(4\alpha t - 24a_1 + 120a_2)^8 - 240(4\alpha t - 24a_1 + 120a_2)^4(-326 + \\
& 45(-8\beta t - 80b_1 + 560b_2)^2 - 135(16a_1 - 160a_2)^2) + 480(4\alpha t - 24a_1 + 120a_2)^2(-76 + \\
& 135(-8\beta t - 80b_1 + 560b_2)^2 + 1215(16a_1 - 160a_2)^2) - 129600(4\alpha t - 24a_1 + \\
& 120a_2)^3(32a_1 - 256a_2) - 12960(4\alpha t - 24a_1 + 120a_2)^5(32a_1 - 256a_2) - 64800(4\alpha t - \\
& 24a_1 + 120a_2)(-96a_1 + 1280a_2 + 4(-8\beta t - 80b_1 + 560b_2)^2(16a_1 - 160a_2) + \\
& 4(16a_1 - 160a_2)^3) + 12144 - 97200(-8\beta t - 80b_1 + 560b_2)^2 + 32400(16a_1 - \\
& 160a_2)^2 + 518400(32b_1 - 448b_2)^2 - 33177600(16a_1 - 160a_2)a_2 + 2123366400a_2^2)(2x + \\
& 12\beta t + 60b_1 - 280b_2)^2 + (-360(4\alpha t - 24a_1 + 120a_2)^8(-8\beta t - 80b_1 + 560b_2) - \\
& 17280(4\alpha t - 24a_1 + 120a_2)^5(-8\beta t - 80b_1 + 560b_2)(16a_1 - 160a_2) - 1440(4\alpha t - \\
& 24a_1 + 120a_2)^6(-8\beta t - 240b_1 + 2800b_2) + 32400(4\alpha t - 24a_1 + 120a_2)^4(-8\beta t + \\
& 112b_1 - 2128b_2) + 64800(4\alpha t - 24a_1 + 120a_2)^2(40\beta t + 752b_1 - 7728b_2 + \\
& 4(-8\beta t - 80b_1 + 560b_2)^3 + 4(-8\beta t - 80b_1 + 560b_2)(16a_1 - 160a_2)^2) - 777600(4\alpha t - \\
& 24a_1 + 120a_2)((-8\beta t - 80b_1 + 560b_2)(16a_1 - 160a_2) + 2(16a_1 - 160a_2)(32b_1 - \\
& 448b_2) - 128(-8\beta t - 80b_1 + 560b_2)a_2) - 172800(4\alpha t - 24a_1 + 120a_2)^3(3(-8\beta t - \\
& 80b_1 + 560b_2)(16a_1 - 160a_2) + (16a_1 - 160a_2)(32b_1 - 448b_2) - 64(-8\beta t -
\end{aligned}$$

$$\begin{aligned}
& 80b_1 + 560b_2)a_2) + 648000\beta t + 8553600b_1 - 74390400b_2 + 259200(-8\beta t - 80b_1 + 560b_2)^3 + 1296000(-8\beta t - 80b_1 + 560b_2)(16a_1 - 160a_2)^2 - 1036800(-8\beta t - 80b_1 + 560b_2)^2(32b_1 - 448b_2) + 1036800(16a_1 - 160a_2)^2(32b_1 - 448b_2) - 132710400(-8\beta t - 80b_1 + 560b_2)(16a_1 - 160a_2)a_2)(2x + 12\beta t + 60b_1 - 280b_2) + 120(-8\beta t - 80b_1 + 560b_2)(2x + 12\beta t + 60b_1 - 280b_2)^9 + 1440(32b_1 - 448b_2)(2x + 12\beta t + 60b_1 - 280b_2)^7 - 120(4\alpha t - 24a_1 + 120a_2)^9(16a_1 - 160a_2) + 80(4\alpha t - 24a_1 + 120a_2)^6(191 + 63(-8\beta t - 80b_1 + 560b_2)^2 + 27(16a_1 - 160a_2)^2) - 2160(4\alpha t - 24a_1 + 120a_2)^5(240a_1 - 4576a_2) + 21600(4\alpha t - 24a_1 + 120a_2)^3(-368a_1 + 3488a_2 + 4(-8\beta t - 80b_1 + 560b_2)^2(16a_1 - 160a_2) + 4(16a_1 - 160a_2)^3) - 1440(4\alpha t - 24a_1 + 120a_2)^7(80a_1 - 864a_2) + 240(4\alpha t - 24a_1 + 120a_2)^4(599 + 135(-8\beta t - 80b_1 + 560b_2)^2 - 225(16a_1 - 160a_2)^2 - 360(-8\beta t - 80b_1 + 560b_2)(32b_1 - 448b_2) - 23040(16a_1 - 160a_2)a_2) - 16200(4\alpha t - 24a_1 + 120a_2)(496a_1 - 6240a_2 + 80(-8\beta t - 80b_1 + 560b_2)^2(16a_1 - 160a_2) + 16(16a_1 - 160a_2)^3 + 128(-8\beta t - 80b_1 + 560b_2)(16a_1 - 160a_2)(32b_1 - 448b_2) - 4096(-8\beta t - 80b_1 + 560b_2)^2a_2 + 4096(16a_1 - 160a_2)^2a_2) + 24(4\alpha t - 24a_1 + 120a_2)^2(3881 + 12150(-8\beta t - 80b_1 + 560b_2)^2 + 28350(16a_1 - 160a_2)^2 + 21600(-8\beta t - 80b_1 + 560b_2)(32b_1 - 448b_2) + 21600(32b_1 - 448b_2)^2 + 88473600a_2^2) + 1036800(-8\beta t - 80b_1 + 560b_2)^2(16a_1 - 160a_2)^2 + 356400(-8\beta t - 80b_1 + 560b_2)^2 + 874800(16a_1 - 160a_2)^2 + 3720(4\alpha t - 24a_1 + 120a_2)^8 + 120(4\alpha t - 24a_1 + 120a_2)^{10} + 518400(32b_1 - 448b_2)^2 + 2123366400a_2^2 + (1 + (2x + 12\beta t + 60b_1 - 280b_2)^2 + (4\alpha t - 24a_1 + 120a_2)^2)^6 + 518400(-8\beta t - 80b_1 + 560b_2)^4 + 518400(16a_1 - 160a_2)^4
\end{aligned}$$

is a solution to the Hirota equation (1).