



# Explicit Formulae for Some Polynomial Continued Fractions

David Naccache, Ofer Yifrach-Stav

## ► To cite this version:

David Naccache, Ofer Yifrach-Stav. Explicit Formulae for Some Polynomial Continued Fractions. 2023. hal-04105623v1

HAL Id: hal-04105623

<https://hal.science/hal-04105623v1>

Preprint submitted on 24 May 2023 (v1), last revised 18 Apr 2024 (v15)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

Public Domain

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Explicit Formulae for Some Polynomial Continued Fractions

David Naccache<sup>1</sup> and Ofer Yifrach-Stav<sup>1</sup>

DIÉNS, ÉNS, CNRS, PSL University, Paris, France  
45 rue d'Ulm, 75230, Paris CEDEX 05, France  
[ofer.friedman@ens.fr](mailto:ofer.friedman@ens.fr), [david.naccache@ens.fr](mailto:david.naccache@ens.fr)

**Abstract.** In a previous article we gave a collection of continued fractions involving Catalan's constant. In this note we provide the general formula governing those continued fractions.

We also show that a single formula involving  $\log 2$  discovered by the Ramanujan machine is part of a large family of which we provide examples.

## 1 Introduction

In a previous article [2] we gave a collection of continued fractions involving Catalan's constant. In this note we provide the general formula governing those continued fractions.

We also show that a single formula involving  $\log 2$  discovered by the Ramanujan Machine Project [1, 3, 4] is part of a large family of which we provide examples.

## 2 Notations

We denote by  $n!!$  the semifactorial of, i.e. the product of all the integers from 1 up to  $n$  having the same parity as  $n$ :

$$n!! = \prod_{k=0}^{\lceil \frac{n}{2} \rceil - 1} (n - 2k) = n(n-2)(n-4)\cdots$$

Because in all the following we will only apply semifactorials to odd numbers, this can simplified as:

$$n!! = \prod_{k=1}^{\frac{n+1}{2}} (2k-1) = n(n-2)(n-4)\cdots 3 \cdot 1$$

We denote by Catalan's constant by  $G = 0.91596559\ldots$  and let  $C_n$  be the  $n$ -th Catalan number:

$$C_n = \frac{1}{n+1} \binom{2n}{n} = \frac{(2n)!}{(n+1)! n!} = \prod_{k=2}^n \frac{n+k}{k} \quad \text{for } n \geq 0$$

The first Catalan numbers are:

$$1, 1, 2, 5, 14, 42, 132, 429, 1430, 4862, 16796, 58786, \dots$$

### 3 The Explicit Formulae

Consider:

$$Q_{c,\kappa} = 2\kappa + 1 + \sum_{n=1}^{\infty} \left( \frac{-2n^2(n+2\kappa)(n+c)}{3n^2 + (3+4\kappa)n + 2\kappa + 1} \right)$$

Define the functions:

$$\begin{aligned} \Delta_{c,\kappa}(\alpha, \beta) &= \begin{cases} \alpha + \beta c & \text{if } c < 2 \\ -2c(2c-1)(2(c-\kappa)-1)^2 \Delta_{c-2,\kappa} & \text{if } c \geq 2 \\ + (8c^2 + (2-8\kappa)c - 2\kappa + 1) \Delta_{c-1,\kappa} \end{cases} \\ \Gamma_{c,\kappa}(\alpha, \beta) &= (2c-1)!! G + \Delta_{c-1,\kappa}(\alpha, \beta) \cdot \prod_{i=0}^{\kappa} (2(c-i)-1) \\ \delta_{\kappa} &= \frac{4^{\kappa-1}}{(2\kappa-1)C_{\kappa-1}} \text{ and } \rho_{\kappa} = \frac{\delta_{\kappa}(-1)^k(1-2k)}{(2k)!(2k-3)!!} \\ \alpha_{\kappa} &= \rho_{\kappa} \Delta_{k-1,1}(1, -2) \text{ and } \beta_{\kappa} = -\rho_{\kappa} (2k-3)^2 \Delta_{k-1,2}(1, 12) - \alpha_{\kappa} \\ \text{Then } \forall \kappa, c \in \mathbb{N}^2, \quad Q_{c,\kappa} &= \frac{\delta_{\kappa}(2c)!}{\Gamma_{c,\kappa}(\alpha_{\kappa}, \beta_{\kappa})} \end{aligned}$$

$Q_{c,\kappa}$  is hence an explicitly computable fraction of the form:

$$Q_{c,\kappa} = \frac{a_0}{a_1 + a_2 G} \text{ where } a_0, a_1, a_2 \in \mathbb{Z}$$

```

1 QNum[c_, k_] :=
2 2 k + 1 +
3 ContinuedFractionK[-2 n^2 (n + 2 k) (n + c), 3 n^2 + (3 + 4 k) n + 2 k +
4 1, {n, 1, 2000}];
5 Del[a_, b_, c_, k_] := If[c < 2, a + b c,
6 -2 c (2 c - 1) (2 (c - k) - 1)^2 Del[a, b, c - 2,
7 k] + (8 c^2 + (2 - 8 k) c - 2 k + 1) Del[a, b, c - 1, k]];
8
9 Gam[a_, b_, c_, k_] := ((2 c - 1)!!)^2 Catalan +
10 Product[(2 (c - j) - 1), {j, 0, k - 1}] * Del[a, b, c - 1, k];
11
12 d[k_] := 4^(k - 1)/(2 k - 1)/CatalanNumber[k - 1];
13
14 Rho[k_] := d[k] (-1)^(k) (1 - 2 k)/((2 k)! (2 k - 3)!!);


```

```

15
16 Alp[k_] := Rho[k] Del[1, -2, k - 1, 1];
17 Bet[k_] := - Rho[k] (2 k - 3)^2 Del[1, 12, k - 1, 2] - Alp[k];
18
19 QFor[c_, k_] := d[k] (2 c)!/Gam[Alp[k], Bet[k], c, k];
20
21 Union[Flatten[Table[N[QFor[c, k], 200] == N[QNum[c, k], 200], {c, 1, 20}, {
22 {k, 1, 20}]]]

```

---



**Fig. 1.** Functional dependency between the functions computing  $Q_{c,k}$

Note that, as described, the complexity of  $\Delta_{c,k}$  is exponential in  $c$ , however, using classical Fibonacci memoization, this complexity can be reduced to  $O(c)$  thereby resulting in a very efficient algorithm for computing  $Q_{c,k}$ .

#### 4 Low Degree Continued Fractions Involving log 2

Continued fractions involving  $\log 2$  can be also obtained with lower degree polynomials, e.g.:

$$\frac{2}{L(\frac{1}{2}, 1, c-1)} = \frac{2}{\sum_{n=0}^{\infty} \frac{e^{\pi i n}}{(n+c-1)}} = \frac{1}{2^{c-2} \log(2) - \sum_{j=1}^{c-2} \frac{2^{c-j-2}}{j}} = c + \sum_{n=1}^{\infty} \left( \frac{-2n^2}{3n+c} \right)$$

---

```

1 Q[c_] := c + ContinuedFractionK[-2 n^2, 3 n + c, {n, 1, 2000}];
2 For[c = 2, c <= 25,
3   r = N[Q[c], 80];
4   v = FindIntegerNullVector[{1, r, N[Log[2] r, 80]}];
5   Print[{b, c,
6     Simplify[
7       2/LerchPhi[1/2, 1, -1 + c] == -v[[1]]/(v[[2]] + Log[2] v[[3]]) ==
8       1/(Log[2] 2^(c - 2) - Sum[2^(c - j - 2)/j, {j, c - 2}]]}]};
9 c++];
```

---

We get a similar behavior for:

$$R_c = c + \sum_{n=1}^{\infty} \mathcal{K}\left(\frac{-2n^2 - 2n}{3n + c}\right)$$

where  $2^{c-4} \cdot (c-3) \cdot a_0 = a_2$  and for which we provide numerical examples in Table 5.

---

```

1 R[c_] := c + ContinuedFractionK[-2 n^2 - 2 n, 3 n + c, {n, 1, 2000}];
2 For[c = 3, c <= 20,
3   r = N[R[c], 200];
4   v = FindIntegerNullVector[{1, r, N[Log[2] r, 200]}];
5   Print[{c,
6     2^(c - 4)*(c - 3)*v[[1]] ==
7     v[[3]], -v[[1]]/(v[[2]] + Log[2] v[[3]])}];
8 c++];
```

---

## 5 Further Generalizations

$Q_{c,k} = Q_{1,c,k}$  is a particular case  $j = 1$  of a more general continued fraction:

$$Q_{j,c,k} = j(2-j+2\kappa) + \sum_{n=1}^{\infty} \mathcal{K}\left(\frac{-2n(c+n)(j+n-1)(1-j+2\kappa+n)}{j(2-j+2\kappa)+(3+4\kappa)n+3n^2}\right)$$

When  $Q_{j,c,k}$  is not a rational,  $Q_{j,c,k}$  converges to:

$$Q_{j,c,k} = \begin{cases} \frac{a_0}{a_1 + a_2 G} & \text{if } j \text{ is odd} \\ \frac{a_0}{a_1 + a_2 \log 2} & \text{if } j \text{ is even} \end{cases}$$

In the examples given in the appendix we adopt the notation:

$$\frac{a_0}{a_1 + a_2 \cdot \text{constant}} = T(0) + \sum_{n=1}^{\infty} \mathcal{K}\left(\frac{P(n)}{T(n)}\right)$$

$a_0$	$a_1$	$a_2$	$P(n)/(-2n)$	$T(n) - 3n^2$
-7351344	-32375839	46558512	$n(3+n)(17+n)$	$72 + 43n$
-2450448	-1768477	2450448	$n(1+n)(17+n)$	$36 + 39n$
-1081080	-16147379	23279256	$n(5+n)(15+n)$	$96 + 43n$
-793800	-232217	322560	$(4+n)(9+n)^2$	$100 + 39n$
-504504	-16140515	23279256	$n(7+n)(13+n)$	$112 + 43n$
-436590	7989199	-11531520	$(3+n)(9+n)(11+n)$	$120 + 43n$
-180180	-1001393	1441440	$(2+n)(5+n)(13+n)$	$84 + 39n$
-174636	-8069449	11639628	$n(9+n)(11+n)$	$120 + 43n$
-72072	-850133	1225224	$n(5+n)(13+n)$	$84 + 39n$
-72072	-251099	360360	$n(3+n)(13+n)$	$56 + 35n$
-72072	-52279	72072	$n(1+n)(15+n)$	$32 + 35n$
-45045	124048	-180180	$(1+n)(3+n)(13+n)$	$56 + 35n$
-28028	-999391	1441440	$(2+n)(7+n)(13+n)$	$112 + 43n$
-27720	-20417	27720	$n(1+n)(11+n)$	$24 + 27n$
-22050	27649	-40320	$(3+n)(7+n)(9+n)$	$80 + 35n$
-19305	424423	-612612	$(1+n)(5+n)(15+n)$	$96 + 43n$
-17640	-250007	360360	$n(7+n)(9+n)$	$80 + 35n$
-14700	-153907	221760	$(2+n)(7+n)(9+n)$	$80 + 35n$
-10395	124741	-180180	$(1+n)(5+n)(11+n)$	$72 + 35n$
-9450	76691	-110880	$(1+n)(5+n)(9+n)$	$60 + 31n$
-7350	-62563	90090	$n(7+n)^2$	$64 + 31n$
-7007	424566	-612612	$(1+n)(7+n)(13+n)$	$112 + 43n$
-6300	-14087	20160	$(2+n)(5+n)(9+n)$	$60 + 31n$
-6006	99839	-144144	$(1+n)(5+n)(13+n)$	$84 + 39n$
-4900	-21043	30240	$(2+n)(7+n)^2$	$64 + 31n$
-2520	-1879	2520	$n(1+n)(9+n)$	$20 + 23n$
-2450	28781	-41580	$(1+n)(7+n)^2$	$64 + 31n$
-1225	367	-560	$(3+n)(7+n)^2$	$64 + 31n$
-525	-2413	3465	$n(5+n)(7+n)$	$48 + 27n$
-450	1151	-1680	$(1+n)(5+n)^2$	$36 + 23n$
-350	1739	-2520	$(1+n)(5+n)(7+n)$	$48 + 27n$
-180	-299	420	$n(3+n)(5+n)$	$24 + 19n$
-105	142	-210	$(1+n)(3+n)(7+n)$	$32 + 23n$
-18	7	-12	$(1+n)(3+n)^2$	$16 + 15n$
-6	-5	6	$n(1+n)(3+n)$	$8 + 11n$

**Table 1.** Examples of convergence to  $\frac{a_0}{a_1+a_2 \log 2}$

$a_0$	$a_1$	$a_2$	$P(n)/(-2n)$	$T(n) - 3n^2$
1	1	-1	$n(1+n)^2$	4 + 7n
9	11	-15	$n(3+n)^2$	16 + 15n
50	147	-210	$n(5+n)^2$	36 + 23n
60	47	-60	$n(1+n)(5+n)$	12 + 15n
90	-79	120	$(1+n)(3+n)(5+n)$	24 + 19n
420	319	-420	$n(1+n)(7+n)$	16 + 19n
420	887	-1260	$n(3+n)(7+n)$	32 + 23n
900	361	-480	$(2+n)(5+n)^2$	36 + 23n
1890	-3443	5040	$(1+n)(3+n)(9+n)$	40 + 27n
2100	2377	-3360	$(2+n)(5+n)(7+n)$	48 + 27n
5544	50035	-72072	$n(5+n)(11+n)$	72 + 35n
6468	-249713	360360	$(1+n)(7+n)(11+n)$	96 + 39n
7560	19409	-27720	$n(3+n)(9+n)$	40 + 27n
8316	-19031	27720	$(1+n)(3+n)(11+n)$	48 + 31n
15444	-49705	72072	$(1+n)(3+n)(15+n)$	64 + 39n
22050	-499279	720720	$(1+n)(7+n)(9+n)$	80 + 35n
24255	-76586	110880	$(3+n)(7+n)(11+n)$	96 + 39n
25740	200107	-288288	$(2+n)(5+n)(15+n)$	96 + 43n
37800	250427	-360360	$n(5+n)(9+n)$	60 + 31n
38808	849671	-1225224	$n(7+n)(11+n)$	96 + 39n
39690	-1997851	2882880	$(1+n)(9+n)^2$	100 + 39n
41580	154327	-221760	$(2+n)(5+n)(11+n)$	72 + 35n
58212	3997025	-5765760	$(2+n)(9+n)(11+n)$	120 + 43n
79380	2123957	-3063060	$n(9+n)^2$	100 + 39n
83160	251561	-360360	$n(3+n)(11+n)$	48 + 31n
87318	-8491859	12252240	$(1+n)(9+n)(11+n)$	120 + 43n
97020	1999321	-2882880	$(2+n)(7+n)(11+n)$	96 + 39n
132300	3997907	-5765760	$(2+n)(9+n)^2$	100 + 39n
198450	-1227581	1774080	$(3+n)(9+n)^2$	100 + 39n
216216	852707	-1225224	$n(3+n)(15+n)$	64 + 39n
360360	263111	-360360	$n(1+n)(13+n)$	28 + 31n
630630	-3990557	5765760	$(3+n)(7+n)(13+n)$	112 + 43n
918918	-3383801	4900896	$(1+n)(3+n)(17+n)$	72 + 43n
1746360	2475007	-3548160	$(4+n)(9+n)(11+n)$	120 + 43n
46558512	33464927	-46558512	$n(1+n)(19+n)$	40 + 43n

**Table 2.** Examples of convergence to  $\frac{a_0}{a_1+a_2 \log 2}$ . The entry in blue is the one reported by the Ramanujan Project.

$a_0$	$a_1$	$a_2$	$P(n)/(-2n)$	$T(n) - 3n^2$
-80281600	-10675439	9459450	$(2+n)(3+n)(14+n)$	$45 + 35n$
-15728640	392683	-727650	$(4+n)^2(12+n)$	$65 + 35n$
-13107200	-263867	-28350	$(4+n)(5+n)(10+n)$	$55 + 31n$
-10485760	-93699	-103950	$(4+n)(5+n)(12+n)$	$65 + 35n$
-7372800	-884203	727650	$(2+n)(3+n)(12+n)$	$39 + 31n$
-6291456	149419	-257250	$(4+n)(8+n)^2$	$81 + 35n$
-5242880	-86807	9450	$(5+n)(6+n)(10+n)$	$77 + 35n$
-3932160	-116317	3150	$(4+n)(5+n)(8+n)$	$45 + 27n$
-3276800	-158859	22050	$(2+n)(5+n)(10+n)$	$33 + 27n$
-2621440	-48609	-1050	$(5+n)(6+n)(8+n)$	$63 + 31n$
-2359296	-168445	103950	$(2+n)(4+n)(12+n)$	$39 + 31n$
-491520	50593	-66150	$(3+n)(4+n)(10+n)$	$55 + 31n$
-327680	-21271	9450	$(2+n)(4+n)(10+n)$	$33 + 27n$
-230400	-1909	-22050	$(2+n)^2(5+n)$	$9 + 11n$
-196608	-184547	198450	$(3+n)(6+n)(10+n)$	$77 + 35n$
-163840	-4981	-22050	$n(5+n)(8+n)$	$9 + 19n$
-131072	-2951	-630	$(4+n)^2(8+n)$	$45 + 27n$
-122880	-13079	9450	$(2+n)(3+n)(10+n)$	$33 + 27n$
-61440	-2467	-3150	$(2+n)(4+n)(5+n)$	$15 + 15n$
-61440	791	-9450	$n(5+n)(6+n)$	$7 + 15n$
-51200	2839	-9450	$n(4+n)(5+n)$	$5 + 11n$
-49152	-1919	90	$(4+n)^2(6+n)$	$35 + 23n$
-36864	-2693	-450	$(2+n)(4+n)(6+n)$	$21 + 19n$
-18432	-419	-3150	$n(4+n)(6+n)$	$7 + 15n$
-18432	-419	-3150	$n(3+n)(8+n)$	$9 + 19n$
-8192	-487	-54	$(4+n)^3$	$25 + 19n$
-3072	121	-630	$n(4+n)^2$	$5 + 11n$
-2048	-129	-90	$(2+n)(4+n)^2$	$15 + 15n$
-2048	-43	-30	$(3+n)(4+n)(6+n)$	$35 + 23n$
-768	-77	-18	$(2+n)(3+n)(4+n)$	$15 + 15n$
-288	31	-90	$n(2+n)(3+n)$	$3 + 7n$

**Table 3.** Examples of convergence to  $\frac{a_0}{a_1+a_2G}$

$a_0$	$a_1$	$a_2$	$P(n)/(-2n)$	$T(n)-3n^2$
192	13	18	$(2+n)^2(3+n)$	$9+11n$
384	1	90	$n(3+n)(4+n)$	$5+11n$
2304	389	450	$n(3+n)(6+n)$	$7+15n$
3072	179	-18	$(3+n)(4+n)^2$	$25+19n$
4608	133	450	$(2+n)^2(4+n)$	$9+11n$
4608	383	-90	$(2+n)(3+n)(6+n)$	$21+19n$
11520	-1373	3150	$n(2+n)(4+n)$	$3+7n$
12288	973	-750	$(3+n)(6+n)^2$	$49+27n$
12288	1145	-630	$(2+n)(3+n)(8+n)$	$27+23n$
16384	-543	1050	$(3+n)(4+n)(8+n)$	$45+27n$
81920	3983	1350	$(4+n)^2(5+n)$	$25+19n$
89600	-10891	22050	$n(2+n)(5+n)$	$3+7n$
98304	2263	150	$(4+n)(6+n)^2$	$49+27n$
98304	35389	-36750	$(3+n)(6+n)(8+n)$	$63+31n$
122880	6563	3150	$(2+n)(5+n)(6+n)$	$21+19n$
147456	21365	22050	$n(4+n)(8+n)$	$9+19n$
163840	6789	450	$(4+n)(5+n)(6+n)$	$35+23n$
262144	710401	-771750	$(3+n)(8+n)^2$	$81+35n$
294912	18013	-3150	$(2+n)(4+n)(8+n)$	$27+23n$
524288	27787	-22050	$(4+n)(6+n)(10+n)$	$77+35n$
786432	19099	-5250	$(4+n)(6+n)(8+n)$	$63+31n$
983040	25979	-450	$(5+n)(6+n)^2$	$49+27n$
1310720	1723	28350	$(4+n)^2(10+n)$	$55+31n$
2949120	168821	22050	$(2+n)(5+n)(8+n)$	$27+23n$
9830400	-1833409	2182950	$(3+n)(4+n)(12+n)$	$65+35n$
12582912	184025	-7350	$(5+n)(8+n)^2$	$81+35n$
39321600	1965547	-727650	$(2+n)(5+n)(12+n)$	$39+31n$
165150720	12969199	-9459450	$(2+n)(4+n)(14+n)$	$45+35n$
330301440	17687791	-9459450	$(2+n)(5+n)(14+n)$	$45+35n$

Table 4. Examples of convergence to  $\frac{a_0}{a_1+a_2 G}$

$c$	$a_0$	$a_1$	$a_2$
3	2	1	0
4	1	1	-1
5	-1	-3	4
6	-2	-17	24
7	-3	-67	96
8	12	667	-960
9	5	666	-960
10	30	9319	-13440
11	-105	-74537	107520
12	-280	-447187	645120
13	126	447173	-645120
14	63	491884	-709632
15	231	3935051	-5677056
16	2772	102311095	-147603456
17	-1287	-102310996	147603456
18	-6006	-1023109531	1476034560
19	45045	16369749493	-23616552960
20	720720	556571437717	-802962800640

Table 5. Examples of convergence to  $\frac{a_0}{a_1+a_2 \log 2}$  for  $R_c$

## References

1. N. B. David, G. Nimri, U. Mendlovic, Y. Manor, and I. Kaminer. On the Connection Between Irrationality Measures and Polynomial Continued Fractions, 2021.
2. D. Naccache and O. Yifrach-Stav. On catalan constant continued fractions. In S. El Hajji, S. Mesnager, and E. M. Souidi, editors, *Codes, Cryptology and Information Security*, pages 43–54, Cham, 2023. Springer Nature Switzerland.
3. G. Raayoni, S. Gottlieb, Y. Manor, G. Pisha, Y. Harris, U. Mendlovic, D. Haviv, Y. Hadad, and I. Kaminer. Generating Conjectures on Fundamental Constants with the Ramanujan Machine. *Nature*, 590(7844):67–73, Feb 2021.
4. G. Raayoni, G. Pisha, Y. Manor, U. Mendlovic, D. Haviv, Y. Hadad, and I. Kaminer. The Ramanujan Machine: Automatically Generated Conjectures on Fundamental Constants. *CoRR*, abs/1907.00205, 2019.