

# Automorphisms of projective structures Maycol Falla Luza, Frank Loray

### ▶ To cite this version:

Maycol Falla Luza, Frank Loray. Automorphisms of projective structures. 2023. hal-04106621

## HAL Id: hal-04106621 https://hal.science/hal-04106621

Preprint submitted on 25 May 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

#### AUTOMORPHISMS OF PROJECTIVE STRUCTURES

#### MAYCOL FALLA LUZA<sup>1</sup>, FRANK LORAY<sup>2</sup>

ABSTRACT. We study the problem of classifying local projective structures in dimension two having non trivial Lie symmetries. In particular we obtain a classification of flat projective structures having positive dimensional Lie algebra of projective vector fields.

#### 1. INTRODUCTION

Let II be a projective structure defined on some neighborhood U of  $0 \in \mathbb{C}^2$  by

(1) y'' = f(x, y, y')

with  $f(x, y, z) = A(x, y) + B(x, y)z + C(x, y)z^{2} + D(x, y)z^{3}$ 

The projectivized tangent bundle M = P(TU) is naturally a contact manifold and each solution on U lifts uniquely as a Legendrian curve on M defining a foliation  $\mathcal{G}$  on M. If U is a small euclidean ball, the space of  $\mathcal{G}$ -leaves is a complex surface  $U^*$  which contains a rational curve  $C_0$  of self-intersection +1 given by the image of  $\mathbb{P}(T_0U)$  (equivalently,  $C_0$  corresponds to the solutions of 1 passing through  $0 \in U$ ). We call the pair  $(U^*, C_0)$ the dual neighborhood of (U, 0). For a survey of such duality, we refer to [6].

A local diffeomorphism  $\Psi$  in U (fixing 0 or not) is an automorphism of the projective structure II, if it sends geodesics to geodesics of (1). As a consequence,  $\Psi$  acts also on the dual neighborhood  $(U^*, C_0)$ , inducing a diffeomorphism  $\check{\Psi}$  from the neighborhood of  $C_0$  onto the neighborhood of (itself or another) (+1)-rational curve C inside  $U^*$ . Conversely, such a diffeomorphism  $\check{\Psi}$  in  $U^*$ , between neighborhoods of (+1)-rational curves, induces an automorphism  $\Psi$  of the projective structure II as above. Lie showed that the pseudo-group of automorphisms of the projective structure forms a Lie pseudogroup denoted by Aut(II). Vector fields whose local flow belong to this pseudo-group are called infinitesimal symmetries and form a Lie algebra denoted by  $\mathfrak{aut}(\Pi)$ . Elements of  $\mathfrak{aut}(\Pi)$  obviously correspond to germs of holomorphic vector fields on the dual (germ of) neighborhood  $(U^*, C_0)$ , and we denote by  $\mathfrak{aut}(U^*, C_0)$  the corresponding Lie algebra. Clearly, we have

$$\mathfrak{aut}(\Pi) \simeq \mathfrak{aut}(U^*, C_0).$$

In [8], Lie gives a classification of the possible infinitesimal symmetry algebras for projective structures, showing that they must be isomorphic to one of the following algebras

(2) 
$$\{0\}, \mathbb{C}, \operatorname{aff}(\mathbb{C}), \operatorname{sl}_2(\mathbb{C}) \text{ or } \operatorname{sl}_3(\mathbb{C})$$

where  $\operatorname{aff}(\mathbb{C})$  is the non commutative 2-dimensional Lie algebra corresponding to the affine group acting on the line: it is spanned by X and Y satisfying [X, Y] = X.

In their paper [2], R. Bryant, G. Manno and V. Matveev classified two-dimensional local metrics (U, g) whose underlying projective structure  $(U, \Pi_g)$  is such that dim  $\mathfrak{aut}(\Pi_g) =$ 2. In [1] a local obstruction to the existence of a Levi-Civita connection within a given projective structure on a surface is given. More recently, in [10] the case dim  $\mathfrak{aut}(\Pi_g) = 1$ is solved. These problems were settled by Lie himself. As a biproduct, they provide in [2, section 2.3] a list of almost unique normal forms for generic local projective structures

Date: May 25, 2023.

Key words and phrases. Automorphism, Projective Structure, Rational Curves.

The authors thank Brazilian-French Network in Mathematics and CAPES/COFECUB Project Ma 932/19 "Feuilletages holomorphes et intéractions avec la géométrie". Falla Luza acknowledges support from CNPq (Grant number 402936/2021-3). Loray is supported by CNRS and Centre Henri Lebesgue, program ANR-11-LABX-0020-0.

 $(U,\Pi)$  with dim  $\mathfrak{aut}(\Pi) = 2$ . In section 3, we give a precise statement of this, completed with other possible dimensions dim  $\mathfrak{aut}(\Pi) = 1, 2, 3, 8$ . We mention here the recent paper [4] where the authors establish an explicit correspondence between two-dimensional projective structures admitting a projective vector field and a class of solutions to the  $SU(\infty)$  Toda equation. There, our preliminary version of the current work [5, section 6] is used by the authors.

Then we focus on flat projective structures. These are structures whose solutions are given by a pencil of (transverse) foliations or, equivalently whose dual surface  $(U^*, C_0)$ admits a semi-local fibration transverse to  $C_0$ . A non-linearizable projective structure admits at most two flat structures, see [6, Theorem 5.1] or [7, Theorem A]. In other words, if  $(U, \Pi)$  admits at least three flat structures then there is a local biholomorphism sending solutions of  $\Pi$  in solutions of the linear structure y'' = 0. We use this result in order to prove our main result, Theorem 4.1 where we classify flat projective structures admiting positive dimensional Lie algebra of symmetries, see Section 4.

**Organization of the paper.** In Section 2 we present some preliminar properties about projective structures with one vector field of symmetries and webs with non trivial automorphism Lie algebra. In Section 3 we establish a normal form of projective structures having non trivial Lie symmetries, see Theorem 3.1. Finally in Section 4 we present our main result, Theorem 4.1 classifying flat projective structures having Lie symmetries. A preliminary version of this work was posted in [5, section 6].

#### 2. Preliminaries

Let us start with some considerations in the case the projective structure  $(U,\Pi)$  is invariant by one (regular) vector field, say  $\partial_{y}$ .

**Lemma 2.1.** Let  $X = \partial_y$  be a non trivial symmetry of a projective structure  $(U, \Pi)$  and let  $\check{X}$  be the dual vector field on  $(U^*, C_0)$ . Then the differential equation for the projective structure takes the form

(3) 
$$y'' = A(x) + B(x)(y') + C(x)(y')^2 + D(x)(y')^3$$

and we have the following possibilities:

- $D(0) \neq 0$  and  $\check{X}$  is regular on  $(U^*, C_0)$ , with exactly one tangency with  $C_0$ ;
- D(0) = 0 but  $D \not\equiv 0$  and  $\check{X}$  has an isolated singular point on  $(U^*, C_0)$ ;
- $D \equiv 0$  and  $\check{X}$  has a curve  $\Gamma$  of singular points on  $(U^*, C_0)$ ; moreover,  $\Gamma$  is transversal to  $C_0$  and the saturated foliation  $\mathcal{F}_{\check{X}}$  defines a fibration transversal to  $C_0$ .

*Proof.* The normal form (3) follows from a straithforward computation. Clearly, X has a singular point at  $p \in U^*$  if, and only if, the corresponding geodesic in U is a trajectory of X (i.e. is X-invariant). Therefore, up to shrinking the neighborhoods U and  $U^*$ , we have 3 possibilities:

- $D \neq 0$  on U and X is regular on  $(U^*, C_0)$ ;
- D vanishes exactly along x = 0 which is therefore geodesic, and X has an isolated singular point at the corresponding point on  $(U^*, C_0)$ ;
- $D \equiv 0$ , the foliation dx = 0 defined by X is geodesic, and X has a curve  $\Gamma$  of singular points on  $(U^*, C_0)$ .

In the first case, the restriction  $\check{X}|_{C_0}$  cannot be identically tangent to  $TC_0 \simeq \mathcal{O}_{\mathbb{P}^1}(2)$ , otherwise it would have a singular point; it thus defines a non trivial section of  $NC_0 \simeq \mathcal{O}_{\mathbb{P}^1}(1)$  which must have a single zero, meaning that  $\check{X}$  has a single tangency with  $C_0$ . The second case we are not interested, since moving to a nearby point of U, we can assume that we are in the first case.

In the third case, each trajectory of X is geodesic, so is the induced foliation dx = 0. By duality, this foliation defines a cross section to  $C_0$  consisting of singular points of  $\check{X}$ . The saturated foliation  $\mathcal{F}_{\check{X}}$  is locally defined by  $\check{X}$  oustide  $\Gamma$ , and by the vector field  $\frac{1}{f}\check{X}$  near  $\Gamma$ , where f is a reduced equation of  $\Gamma$ . But  $\frac{1}{f}\check{X}$  induces a non zero section of  $NC_0$  (near  $\Gamma$ ) since it must be less vanishing;  $\frac{1}{f}X$  is therefore transversal to  $C_0$ , and so is  $\mathcal{F}_{\check{X}}$ .

**Lemma 2.2** (Cartan [3, p.78-83]). Let  $\mathcal{W} = \mathcal{F}_0 \boxtimes \mathcal{F}_1 \boxtimes \mathcal{F}_\infty$  be a regular 3-web on  $(\mathbb{C}, 0)$ , and let  $\mathfrak{aut}(\mathcal{W})$  be the Lie algebra of vector fields whose flow preserve  $\mathcal{W}$ . If  $\mathfrak{aut}(\mathcal{W}) \neq 0$ then we are in one of the two cases, up to change of coordinates:

- $\mathfrak{aut}(\mathcal{W}) = \mathbb{C}\partial_y$  and  $\mathcal{W} = dy \boxtimes (dy dx) \boxtimes (dy + f(x)(dy dx))$ , with f analytic, not of the form  $f(x) = ae^{bx}$ ,  $a, b \in \mathbb{C}$ ; •  $aut(\mathcal{W}) = \mathbb{C}\langle \partial_x, \partial_y, x\partial_x + y\partial_y \rangle$  and  $\mathcal{W} = dy \boxtimes (dy - dx) \boxtimes dx$ .

Lemma 2.3. Under assumptions and notations of Lemma 2.1, assume that we are in the last case  $D \equiv 0$ . If the singular set  $\Gamma$  of  $\check{X}$  is a fiber of the fibration defined by  $\mathcal{F}_{\check{X}}$  on  $(U^*, C_0)$ , then  $(U, \Pi)$  is linearizable.

*Proof.* Take 3 different fibers of  $\mathcal{F}_{\check{X}}$  different from  $\Gamma$ , they define a 3-web  $\mathcal{W} = \mathcal{F}_0 \boxtimes$  $\mathcal{F}_1 \boxtimes \mathcal{F}_\infty$  which is invariant by X. By Cartan's Lemma 2.2, we can assume  $X = \partial_y$  and  $\mathcal{W} = dy \boxtimes (dy - dx) \boxtimes (dy + f(x)(dy - dx))$  so that the flat structure of  $\Pi$  is defined by the pencil dy + zf(x)(dy - dx) = 0. But the foliation dx = 0 defined by X is, by assumption, belonging to the pencil, which means that  $\Pi$  is also defined by the hexagonal 3-web  $dy \boxtimes (dy - dx) \boxtimes dx$ , thus linearizable.  $\square$ 

#### 3. Classification of projective structures with Lie symmetries

The problem of this section is to classify those local projective structures  $(U, \Pi)$  having non trivial Lie symmetry, i.e. such that dim  $\mathfrak{aut}(\Pi) > 0$ , up to change of coordinates. However, in this full generality, the problem is out of reach; indeed, it includes for instance the problem of classification of germs of holomorphic vector fields (with arbitrary complicated singular points), which is still challenging. Instead of this, and in the spirit of Lie's work, we produce a list of possible normal forms up to change of coordinates for such a  $(U,\Pi)$  at a generic point  $p \in U$ , i.e. outside a closed analytic subset consisting of singular features. For instance, a non trivial vector field is regular at a generic point and can be rectified to  $\partial_y$ ; we only consider this constant vector field in the case dim  $\mathfrak{aut}(\Pi) =$ 1. The following resumes some results of [2, section 2.3].

**Theorem 3.1.** Let  $(U,\Pi)$  be a projective structure with  $\mathfrak{aut}(\Pi) \neq \{0\}$ . Then, at the neighborhood of a generic point  $p \in U$ , the pair  $(\Pi, \mathfrak{aut}(\Pi))$  can be reduced by local change of coordinate to one of the following normal forms:

- (i)  $\mathfrak{aut}(\Pi) = \mathbb{C} \cdot \partial_y$  and (A, B, C, D) =(i.a) (A(x), B(x), 0, 1) with  $A, B \in \mathbb{C}\{x\}$ ; (i.b)  $(A(x), 0, e^x, 0)$  with  $A \in \mathbb{C}\{x\}$ ; (ii)  $\operatorname{\mathfrak{aut}}(\Pi) = \mathbb{C}\langle \partial_y, \partial_x + y \partial_y \rangle$  and (A, B, C, D) =(ii.a)  $(\alpha e^x, \beta, 0, e^{-2x})$  with  $\alpha, \beta \in \mathbb{C}$ ,  $(\alpha, \beta) \neq (0, 2), (0, \frac{1}{2});$ (ii.b)  $(\alpha e^x, 0, e^{-x}, 0)$  with  $\alpha \in \mathbb{C}$ ; (iii)  $\operatorname{aut}(\Pi) = \mathbb{C}\langle \partial_y, \partial_x + y\partial_y, y\partial_x + \frac{y^2}{2}\partial_y \rangle$  and  $(A, B, C, D) = (0, \frac{1}{2}, 0, e^{-2x});$ (iv)  $\operatorname{aut}(\Pi) = \operatorname{sl}_3(\mathbb{C})$  and (A, B, C, D) = (0, 0, 0, 0).

These normal forms are unique, except for the case (i.a), which is unique up to the  $\mathbb{C}^*$ -action:

$$(A(x), B(x), 0, 1) \xrightarrow{\lambda \in \mathbb{C}^*} (\lambda^3 A(\lambda^2 x), \lambda^2 B(\lambda^2 x), 0, 1).$$

**Remark 3.2.** The normal forms for  $\mathfrak{aut}(\Pi)$  in the statement correspond to the list of transitive local actions of Lie algebras listed in (2), except that  $sl_2(\mathbb{C})$  has also exotic representations generated by

$$X = \partial_y, \quad Y = \partial_x + y\partial_y \quad \text{and} \quad Z = (y + c_1 e^x)\partial_x + (\frac{y^2}{2} + c_2 e^{2x})\partial_y, \quad c_1, c_2 \in \mathbb{C}.$$

Only the standard one occurs as symmetry algebra of a projective structure.

Remark 3.3. Case (iii) corresponds to the special structure

$$\Pi_0 : y'' = (xy' - y)^3$$

at the neighborhood of any point  $p \neq (0,0)$  and is invariant under the standard action of  $sl_2(\mathbb{C})$ 

$$\mathfrak{aut}(\Pi_0) = \mathbb{C} \langle x \partial_y, \frac{1}{2} (-x \partial_x + y \partial_y), -\frac{1}{2} y \partial_x \rangle$$

(see [11, Theorem 3]). However, at p = (0,0), the Lie algebra is singular, which is excluded from the list of Theorem 3.1. Case (iv) corresponds to the linearizable case y'' = 0.

Proof of Theorem 3.1. We essentially follow [2, section 2.3]. Let us start with the case  $\mathfrak{aut}(\Pi) = \mathbb{C}\langle X \rangle$ . At a generic point  $p \in U$ , the vector field X is regular and we can choose local coordinates such that  $X = \partial_y$ . One easily deduce that the equation (1) for the projective structure, being X-invariant, takes the form y'' = f(x, y, y') with

(4) 
$$f(x,y,z) = A(x) + B(x)z + C(x)z^{2} + D(x)z^{3}.$$

The normalizing coordinates for X are unique up to a change of the form

$$\Phi: (x, y) \mapsto (\psi(x), y + \phi(x)), \quad \psi(0) = \phi(0) = 0, \ \psi'(0) \neq 0.$$

The projective structure  $\Phi^*\Pi$  is defined by

$$f(x, y, z) = \widehat{A}(x) + \widehat{B}(x)z + \widehat{C}(x)(z)^2 + \widehat{D}(x)(z)^3$$

where (we decompose for simplicity)

(5) when 
$$\Phi = (\psi(x), y)$$
, then 
$$\begin{cases} A = A \circ \psi \cdot (\psi')^2 \\ \widehat{B} = B \circ \psi \cdot \psi' + \frac{\psi''}{\psi'} \\ \widehat{C} = C \circ \psi \\ \widehat{D} = \frac{D \circ \psi}{\psi'} \end{cases}$$

(6) when 
$$\Phi = (x, y + \phi(x))$$
, then 
$$\begin{cases} \widehat{A} = A + B\phi' + C(\phi')^2 + D(\phi')^3 - \phi'' \\ \widehat{B} = B + 2C\phi' + 3D(\phi')^2 \\ \widehat{C} = C + 3D\phi' \\ \widehat{D} = D \end{cases}$$

If  $D \neq 0$ , then we can assume at a generic point that  $D \neq 0$ . We can normalize  $\hat{D} = 1$  by setting  $\psi^{-1}(x) := \int_0^x \frac{d\zeta}{D(\zeta)}$  in the first change, and then normalize  $\hat{C} = 0$  by setting  $\phi'(x) = -C/3$  (which does not change D = 1): (A(x), B(x), 0, 1).

Since we are interested in the Lie algebra, more than a given vector field, then we can also change  $\Phi(x, y) = (x, ay)$  with  $a \neq 0$  and get the form

(7) when 
$$\Phi = (x, ay)$$
, then  $(\widehat{A}, \widehat{B}, \widehat{C}, \widehat{D}) = (a^{-1}A, B, aC, a^2D)$ .

Finally,  $\Phi = (a^2 x, ay)$ , a combination of (5) and (7), yields the new normal form

$$(a^{3}A(a^{2}x), a^{2}B(a^{2}x), 0, 1)$$

whence the  $\mathbb{C}^*$ -action of the statement.

Suppose now  $D \equiv 0$ . If C would be constant then, by (8),  $L_1 = L_2 = 0$  and  $\Pi$  is linearizable. Passing to a generic point, we can assume  $C'(0) \neq 0$  and use changes (5) and (7) to normalize  $C = e^x$ . Finally by using (6), we arrive in the unique desired normal form  $(A(x), 0, e^x, 0)$ . In this case the equation is never linearizable, since by (8) we get  $(L_1, L_2) = (0, 2e^{2x})$ .

Now we study the case  $\mathfrak{aut}(\Pi) = \mathbb{C}\langle X, Y \rangle$ , with [X, Y] = X. By [2, Lemma 1], we know that, at a generic point, we can find coordinates where  $X = \partial_y, Y = \partial_x + y\partial_y$ . The invariance of the projective structure by both the flows of X and Y yields

$$(A, B, C, D) = (\alpha e^x, \beta, \gamma e^{-x}, \delta e^{-2x})$$

were  $\alpha$ ,  $\beta$ ,  $\gamma$  and  $\delta$  are constants. The normalizing coordinates for the Lie algebra are unique up to a change

$$\Phi = (x, ay + be^x)$$

with  $a, b \in \mathbb{C}$ ,  $a \neq 0$ . If  $\delta \neq 0$ , we obtain a unique normal form  $(\alpha e^x, \beta, 0, e^{-2x})$ . By [2, Lemma 4], the cases  $(\alpha, \beta) = (0, 2)$  and  $(0, \frac{1}{2})$  have more symmetries: they respectively correspond to the  $sl_3(\mathbb{C})$  and  $sl_2(\mathbb{C})$  cases.

When  $\delta = 0$ , we shall have  $\gamma \neq 0$  (otherwise  $\Pi$  would be linearizable), and we can normalize  $(\widehat{A}, \widehat{B}, \widehat{C}, \widehat{D}) = (\alpha e^x, 0, e^{-x}, 0)$ , with  $\alpha \in \mathbb{C}$ ; this normal form is unique. The case  $\mathfrak{aut}(\Pi) = \mathrm{sl}_2(\mathbb{C})$  follows directly from [2, Lemma 4].

In Theorem 3.1, normal forms (i) contain some models with larger symmetry Lie algebra, and we end the section by determining them.

In the case  $\mathfrak{aut}(\Pi) = \mathrm{sl}_3(\mathbb{C})$ . First of all, we have from [9] that the projective structure is linearizable when Liouville invariants  $L_1$  and  $L_2$  given by

(8) 
$$\begin{cases} L_1 = 2B_{xy} - C_{xx} - 3A_{yy} - 6AD_x - 3A_xD + 3(AC)_y + BC_x - 2BB_y, \\ L_2 = 2C_{xy} - B_{yy} - 3D_{xx} + 6A_yD + 3AD_y - 3(BD)_x - B_yC + 2CC_x. \end{cases}$$

are identically zero, and we get:

 $(L_1, L_2) = (-3A'(x), -3B'(x))$  for model (i.a), and

$$(L_1, L_2) = (-e^{-x}, -2e^{-2x})$$
 for model (i.b).

Linearizability only occur in model (i.a) when A and B are simultaneously constant.

In the case  $\mathfrak{aut}(\Pi) = \operatorname{aff}(\mathbb{C})$ . There must exists a vector field  $v \in \mathfrak{aut}(\Pi)$  such that

$$[\partial_y, v] = \partial_y \quad \text{or} \quad c \cdot v, \quad c \in \mathbb{C}.$$

This implies that v takes the respective form

$$v = \alpha(x)\partial_x + (y + \beta(x))\partial_y$$
 or  $e^{cy}(\alpha(x)\partial_x + \beta(x)\partial_y).$ 

We can furthermore assume  $\alpha(0) \neq 0$  so that the local action is transitive together with  $\partial_y$ ; moreover,  $c \neq 0$ , otherwise the two vector fields commute, which is excluded in the non linearizable case. Let us firstly discuss the case of normal form (i.a). In the case where  $\partial_y$  is stabilized by  $\mathfrak{aut}(\Pi)$ , by using [2, formula (3)] (a PDE system for a vector field to be a symmetry of a projective structure) for  $v = \alpha \partial_x + (y + \beta) \partial_y$ , one easily deduce that

$$v = 2(x + \alpha_0)\partial_x + (y + \beta_0)\partial_y$$
 and  $(A, B, C, D) = \left(\frac{\gamma_0}{(x + \alpha_0)^{3/2}}, \frac{\delta_0}{(x + \alpha_0)}, 0, 1\right),$ 

with  $\alpha_0, \beta_0, \gamma_0, \delta_0 \in \mathbb{C}$ ,  $\alpha_0 \neq 0$ . The second case  $v = e^{cy}(\alpha(x)\partial_x + \beta(x)\partial_y)$  is less explicit. The invariance of the projective structure in normal form (i.a) allows us to express everything in terms of  $\alpha(x)$  and its derivatives:

$$\beta = \frac{\alpha' - c^2 \alpha}{2c} \quad \text{and} \quad (A, B, C, D) = \left(\frac{\alpha''' - c^4 \alpha'}{4c^3 \alpha}, -\frac{3\alpha'' + c^4 \alpha}{4c^2 \alpha}, 0, 1\right),$$

and finally yields the following differential equation for  $\alpha$ 

$$c^{4}(\alpha\alpha'' - (\alpha')^{2}) - 3c^{2}(\alpha\alpha''' - \alpha'\alpha'') + 2\alpha\alpha'''' + \alpha'\alpha''' - 3(\alpha'')^{2} = 0.$$

Once we know the 3-jet of  $\alpha$ , then we can deduce all the coefficients by mean of this differential equation. Mind that we can set  $\alpha(0) = 1$  so that we get a 4-parameter family of projective structures, taking into account the constant c, that can further be normalized to c = 1 by using the  $\mathbb{C}^*$ -action. Equivalently, the family of projective structures is given by the solutions of the system of differential equations

$$A' = \frac{27cA^2 + 9AB' - 3c(B+c^2)B' + c(4B+c^2)(B+c^2)^2}{6(B+c^2)}$$
$$B'' = -\frac{27cA(cA-B') - 12(B')^2 - 9c^2(B+c^2)B' + c^2(4B+c^2)(B+c^2)^2}{6(B+c^2)}$$

and we can recover  $\alpha$  and  $\beta$  by:

$$\frac{\alpha'}{\alpha} = -\frac{3cA+B'}{B+c^2}$$
 and  $\beta = \frac{\alpha'-c^2\alpha}{2c}$ .

For normal forms (i.b), the discussion is similar, easier though, and one find  $v = -\partial_x + (y+c)\partial_y$ ,  $c \in \mathbb{C}$ , with projective structure  $(\alpha_0 e^{-x}, 0, e^x, 0)$ .

In the case  $\mathfrak{aut}(\Pi) = \mathrm{sl}_2(\mathbb{C})$ . We just note that  $\partial_y$  must be contained in a 2-dimensional affine Lie subalgebra, and we are in a particular case of the previous one.

#### 4. Symmetries of flat projective structures

We say that a projective structure  $\Pi$  defined on U by 1 is *flat* (or *foliated*) if the geodesics are tangent to a pencil of foliations  $\{\mathcal{F}_z : \omega_z = \omega_0 + z\omega_\infty\}$ , where  $\omega_0$  and  $\omega_\infty$  are 1-forms on U satisfying  $\omega_0 \wedge \omega_\infty \neq 0$ .

Here, we classify those projective structures having simultaneously a flat structure and Lie symmetries. In other words, we describe which models in the list of Theorem 3.1 have a flat structure, and how many. As one can see on [6, Section 3], the flatness condition is equivalent to the existence of a semi-local fibration on the dual surface  $(U^*, C_0)$  transverse to  $C_0$ . In [6, Theorem 5.1] we show that a given projective structure, if not linearizable, has at most 2 flat structures, see also [7, Theorem A].

**Theorem 4.1.** Let  $(U, \Pi)$  be a flat projective structure with Lie symmetries:  $\mathfrak{aut}(\Pi) \neq \{0\}$ . Then, at the neighborhood of a generic point  $p \in U$ , the pair  $(\Pi, \mathfrak{aut}(\Pi))$  and pencil of geodesic foliations  $\mathcal{F}_z : \omega_0 + z\omega_\infty$  can be reduced by local change of coordinate to one of the following normal forms:

(i) 
$$\operatorname{aut}(\Pi) = \mathbb{C} \cdot \partial_y$$
,  $(A, B, C, D) =$   
(i.a.1)  $(0, 0, 1 + g', g)$  and  $\mathcal{F}_z : \underbrace{e^y(dx + g(x)dy)}_{\omega_0} + z \underbrace{dy}_{\omega_\infty}$ ;  
(i.a.2)  $(0, 0, g', 1)$  and  $\mathcal{F}_z : \underbrace{-(dx + (g(x) + y)dy)}_{\omega_0} + z \underbrace{dy}_{\omega_\infty}$ ;  
(i.b)  $(0, 0, g', 0)$  and  $\mathcal{F}_z : \underbrace{dx + g(x)dy}_{\omega_0} + z \underbrace{dy}_{\omega_\infty}$ ;  
(ii)  $\operatorname{aut}(\Pi) = \mathbb{C}\langle \partial_y, \partial_x + y \partial_y \rangle$  and  $(A, B, C, D) =$ 

(ii.a)  $(\alpha e^x, \beta, 0, e^{-2x})$  with  $\alpha, \beta \in \mathbb{C}$  belonging to the cubic nodal curve

$$\begin{cases} \mathbb{C} & \to & \Gamma = \{27\alpha^2 + 4\beta^3 - 12\beta^2 + 9\beta - 2 = 0\} \subset \mathbb{C}^2 \\ \gamma & \mapsto & (\gamma(2\gamma^2 - 1), 2 - 3\gamma^2) \end{cases}$$

and the corresponding flat structure is given by

$$\mathcal{F}_z:\underbrace{e^x(\gamma y + (2\gamma^2 - 1)e^x)dx - (y + 2\gamma e^x)dy}_{\omega_0} + z\underbrace{(dy - \gamma e^x dx)}_{\omega_\infty}.$$

Here, we exclude the cases  $\alpha = 0$  corresponding to (iii) and (iv) below. (ii.b.1)  $\left(\frac{1-\lambda^2}{4}e^x, 0, e^{-x}, 0\right)$  with  $\lambda \in \mathbb{C}^*$ , and

$$\mathcal{F}_{z}:\underbrace{e^{\lambda x}\left[dy-\left(\frac{1-\lambda}{2}\right)e^{x}dx\right]}_{\omega_{0}}+z\underbrace{\left[dy-\left(\frac{1+\lambda}{2}\right)e^{x}dx\right]}_{\omega_{\infty}}$$

(ii.b.2) 
$$(\underline{e^x}, 0, e^{-x}, 0)$$
 and  $\mathcal{F}_z : \underbrace{(1 - \frac{x}{2})e^x dx + x dy}_{\omega_0} + z \underbrace{(dy - \frac{1}{2}e^x dx)}_{\omega_{\infty}};$   
iii)  $\mathfrak{aut}(\Pi) = \mathbb{C}\langle \partial_y, \partial_x + y \partial_y, y \partial_x + \frac{y^2}{2} \partial_y \rangle$  and  $(A, B, C, D) = (0, \frac{1}{2}, 0, e^{-2x});$ 

(iv)  $aut(\Pi) = sl_3(\mathbb{C}) and (A, B, C, D) = (0, 0, 0, 0).$ 

Case (iii) corresponds to the case (ii.a) with  $\gamma = \pm \frac{1}{\sqrt{2}}$ ; the two values of  $\gamma$  provide the two flat structures for  $\Pi$  in this case. Case (iv) corresponds to the case (ii.a) with  $\gamma = 0$ ; in that case, all flat structures are described in [6, Example 3.3].

**Lemma 4.2.** Let  $(U,\Pi)$  be a projective structure with Lie symmetry  $X = \partial_y$  and flat structure  $\mathcal{F}_{\omega_z}$ , with  $\omega_z = \omega_0 + z\omega_\infty$ . If  $(U,\Pi)$  is not linearizable, then

- the flow  $\phi_X^t$  of X must preserve the flat structure,
- the flow  $\phi_X^t$  must preserve at least one foliation of the pencil, say  $\mathcal{F}_{\omega_{\infty}}$ ,
- no element of the pencil  $\mathcal{F}_{\omega_z}$  can coincide with the foliation  $\mathcal{F}_X : \{dx = 0\}$ , and after change of coordinate  $y \mapsto y + \phi(x)$ , we may furthermore assume  $\omega_{\infty} = dy$ .

In particular, at the neighborhood of a generic point  $p \in U$ , we may furthermore assume  $\omega_{\infty} = dy$  in convenient coordinates.

Proof. The vector field induces an action on geodesics, and therefore on the dual space  $(U^*, C_0)$ ; denote by  $\check{X}$  the infinitesimal generator. Let  $\mathcal{H}$  be the transverse fibration corresponding to the flat structure. If  $\mathcal{H}$  is not invariant by the flow  $\phi_{\check{X}}^t$ , then we obtain a 1-parameter family  $\mathcal{H}_t = (\phi_{\check{X}}^t)_* \mathcal{H}$  and deduce from [6, Theorem 5.1] that  $\Pi$  is linearizable, contradiction. Therefore,  $\check{X}$  preserves  $\mathcal{H}$  and acts on the space of leaves  $\simeq \mathbb{P}^1_z$ . In particular, it has a fixed point, corresponding to an X-invariant foliation in the pencil, say  $\mathcal{F}_{\omega_{\infty}}$ .

Assume for contradiction that the foliation  $\mathcal{F}_X$ , defined by dx = 0, coincides with one of the  $\mathcal{F}_{\omega_z}$ 's; since it is X-invariant, we can assume  $z = \infty$ . Therefore, we are in the third case of Lemma 2.1:  $\check{X}$  has a curve  $\Gamma \subset U^*$  of singular points transversal to  $C_0$ . Moreover,  $\Gamma$  is  $\mathcal{H}$ -invariant and  $\check{X}$  defines another transverse fibration  $\mathcal{F}_{\check{X}}$ . If  $\Gamma$  is invariant by  $\mathcal{F}_{\check{X}}$ , then Lemma 2.3 implies that  $(U, \Pi)$  is linearizable, contradiction. Therefore,  $\Gamma$  is not invariant by  $\mathcal{F}_{\check{X}}$ , and in particular, the fibrations  $\mathcal{F}_{\check{X}}$  and  $\mathcal{H}$  do not coincide. Consider the tangency set  $\Sigma := \operatorname{tang}(\mathcal{F}_{\check{X}}, \mathcal{H})$ . Since  $\mathcal{H}$  is  $\check{X}$ -invariant,  $\Sigma$  must be  $\check{X}$ -invariant. Clearly,  $\Sigma$  is not contained in the singular set  $\operatorname{sing}(\check{X}) = \Gamma$  and  $\Sigma$  is thus  $\mathcal{F}_{\check{X}}$ -invariant. Again, this means that  $\Sigma$  is a common fiber of  $\mathcal{F}_{\check{X}}$  and  $\mathcal{H}$ , and the proof of Lemma 2.3 implies that  $(U, \Pi)$  is linearizable (see also [6, Theorem 5.3]), contradiction.

Proof of Theorem 4.1. Let us start with the case  $\mathfrak{aut}(\Pi) = \mathbb{C} \cdot X$  with  $X = \partial_y$ , and assume  $(U, \Pi)$  not linearizable. Then, by Lemma 4.2, X preserves the pencil of foliations and acts on the parameter space  $z \in \mathbb{P}^1$  fixing  $z = \infty$ . More precisely, we can assume  $\omega_{\infty} = dy$  and that the action on the pencil is induced by one of the following vector fields

$$z\partial_z, \quad \partial_z \quad \text{or} \quad 0.$$

In the first case, we must have that  $(\phi_X^t)_*\omega_z$  is proportional to  $\omega_{e^tz}$  for any  $t, z \in \mathbb{C}$ ; since  $\omega_z = \omega_0 + z\omega_\infty$  and  $\omega_\infty$  is  $\phi_X^t$ -invariant, we deduce

$$(\phi_X^t)_*\omega_0 = e^{-t}\omega_0.$$

This implies that  $\omega_0 = e^y(f(x)dx + g(x)dy)$  for some functions  $f, g \in \mathbb{C}\{x\}, f(0) \neq 0$ . After taking  $\int f(x)dx$  as a new coordinate, we get the normal form

$$\omega_0 = e^y (dx + g(x)dy).$$

We easily derive the projective structure  $\Pi$  by derivating " $\omega_0/\omega_\infty$ ":

$$0 = \left(e^{y}\left(\frac{1}{y'}+g\right)\right)' = e^{y}y'\left(\frac{1}{y'}+g\right) + e^{y}\left(-\frac{y''}{(y')^{2}}+g'\right)$$
$$e^{y''} = \left(1+e^{y'}\right)\left(e^{y'}\right)^{2} + \left(e^{y}\right)\left(e^{y'}\right)^{3} \quad \text{i.e.} \quad \left(A,B,C,D\right) = \left(0,0,1+e^{y'}\right)$$

$$\Rightarrow y'' = (1+g')(y')^2 + (g)(y')^3$$
, i.e.  $(A, B, C, D) = (0, 0, 1+g', g)$ 

If the action is now induced by  $\partial_z$ , then we get

$$(\phi_X^t)_*\omega_0 = \omega_0 + t\omega_\infty$$

which gives  $\omega_0 = f(x)dx + (g(x) - y)dy$ , where again we can normalize  $f \equiv 1$  which gives the projective structure

$$\omega_0 = -(dx + (g(x) + y)dy \text{ and } (A, B, C, D) = (0, 0, g', 1).$$

Finally, when the action is trivial on the parameter space z, we get that  $\omega_0$  is also invariant, i.e. of the form f(x)dx + g(x)dy; we can again normalize  $f \equiv 1$  and get

$$\omega_0 = dx + g(x)dy$$
 and  $(A, B, C, D) = (0, 0, g', 0).$ 

Let us now consider the case  $\mathfrak{aut}(\Pi) = \mathbb{C}\langle X, Y \rangle$  with  $X = \partial_y$  and  $Y = \partial_x + y \partial_y$ , and still assume  $(U, \Pi)$  not linearizable. Like before, the Lie algebra preserves the pencil  $\mathcal{F}_z$ and induces an action on the parameter space of the form

$$(X,Y)|_z = (\partial_z, z\partial_z), \quad (0, \lambda z\partial_z), \quad (0, \partial_z), \quad \text{or} \quad (0,0),$$

with  $\lambda \in \mathbb{C}^*$ . We note that we cannot normalize  $\lambda = 1$  by homothecy since Y has to satisfy [X, Y] = X in the (x, y)-variables; different values of  $\lambda$  will correspond to different projective structures.

In any case for  $(X, Y)|_z$ ,  $\mathcal{F}_{\infty}$  is fixed, and this means that we can write

$$\omega_{\infty} = d(y - \gamma e^x)$$

for some  $\gamma \in \mathbb{C}$ . Indeed, the invariance by X means that the leaves of  $\mathcal{F}_{\infty}$  are  $\partial_y$ -translates of the leaf y = f(x) passing through the origin, i.e. we can choose  $\omega_{\infty} = d(y - f(x))$ ; then, the invariance by Y gives the special form  $f(x) = \gamma e^x$ . Here, we have used Lemma 4.2 to insure that, maybe passing to a generic point  $p \in U$ , we can assume that  $\mathcal{F}_{\infty}$  is not vertical at p.

If  $(X, Y)|_z = (\partial_z, z \partial_z)$ , then we can check that

$$\omega_0 = (\alpha e^{2x} + \gamma e^x y)dx + (\beta e^x - y)dy$$

for some constants  $\alpha, \beta \in \mathbb{C}$ . This normalization is unique up to a change of coordinate of the form  $\Phi = (x, ay + be^x)$  preserving the Lie algebra; this allow to reduce the corresponding projective structure  $\Pi$  into the normal form (ii.a) of Theorem 3.1, yielding after straightforward computation the formulae (ii.a) of Theorem 4.1.

If  $(X, Y)|_z = (0, \lambda z \partial_z)$ , then we find

$$\omega_0 = e^{\lambda x} \left( \alpha e^x dx + \beta e^x dy \right)$$

which gives after normalization

$$\omega_z = e^{\lambda x} \left[ dy - \left(\frac{1-\lambda}{2}\right) e^x dx \right] + z \left[ dy - \left(\frac{1+\lambda}{2}\right) e^x dx \right]$$
  
and  $(A, B, C, D) = \left(\frac{1-\lambda^2}{4} e^x, 0, e^{-x}, 0\right).$ 

If  $(X, Y)|_z = (0, \partial_z)$ , then we find

$$\omega_0 = (\alpha + \gamma x)e^x dx + (\beta - x)dy$$

which gives after normalization

$$\omega_z = (1 - \frac{x}{2})e^x dx + x dy + z (dy - \frac{1}{2}e^x) \quad \text{and} \quad (A, B, C, D) = \left(\frac{e^x}{4}, 0, e^{-x}, 0\right).$$

Finally, if  $(X, Y)|_z = (0, 0)$ , then we find

$$\omega_0 = \alpha e^x dx + \beta dy$$

which gives after normalization (A, B, C, D) = (0, 1, 0, 0), which is linearizable.

**Remark 4.3.** Projective structures of Theorem 4.1 (i) can be put in normal form as in Theorem 3.1. For instance, in the case (i.a.1), using change(5), one easily get the following form

$$\omega_z = e^{f(x)y}(dx + dy) + zdy$$
 and  $(A, B, C, D) = (0, f', 1 + f', 1)$ 

The final normalization (6) is not so explicit, but turning the other way round, we can easily check that a normalized projective structure (A, B, 0, 1) comes from such a flat structure iff it satisfies

$$A^{2} = -\frac{(4B^{2} + 5B - 3B' + 1)^{2}}{3(4B + 1)}$$

and in this case, f(x) (and the flat structure) is given by

$$f' = \frac{1}{2} + \frac{1}{2}\sqrt{-3(4B+1)}.$$

In a very similar way, the projective structure (A, B, 0, 1) comes from the case (i.a.2) iff

$$A^2 = -\frac{(4B^2 - 3B')^2}{108B}$$

and in this case, g(x) (and the flat structure) is given by

$$q' = \sqrt{-3B}.$$

Finally, one easily check by similar computations that any normal form (i.b) of Theorem 3.1, i.e.  $(A(x), 0, e^x, 0)$ , is also flat, i.e. comes from (i.b) of Theorem 4.1.

#### References

- R. BRYANT, M. DUNAJSKI, M. EASTWOOD Metrisability of two-dimensional projective structures. J. Differ. Geom. 83(3) 465 -500 (2009).
- [2] R. BRYANT, G. MANNO, V. MATVEEV, A solution of a problem of Sophus Lie: normal forms of two-dimensional metrics admitting two projective vector fields. Math. Ann. 340 (2008) 437-463.
- [3] É. CARTAN, Les sous-groupes des groupes continus de transformations. Ann. Scient. Éc. Norm. Sup. 25 (1908) 57-194.
- [4] M. DUNAJSKI, A. WATERHOUSE Einstein metrics, projective structures and the SU(∞) Toda equation Journal of Geometry and Physics, volume 147, January 2020, 103523.
- M. FALLA LUZA AND F. LORAY, Projective structures and neighborhoods of rational curves arXiv:1707.07868v3 (preliminary version of [6]).
- [6] M. FALLA LUZA AND F. LORAY, Projective Structures, Neighborhoods of Rational Curves and Painlevé equations. Moscow Mathematical Journal, volume 23, issue 1, 2023, pp. 59-95.
- [7] M. FALLA LUZA AND F. LORAY, On the number of fibrations transverse to a rational curve in complex surfaces. Comptes Rendus Mathematique, volume 354, Issue 5, May 2016, Pages 470-474.
- [8] S. LIE, Untersuchungen uber geodätische Curven. Math. Ann. 20 (1882) 357-454.
- [9] R. LIOUVILLE, Sur les invariants de certaines équations différentielles et sur leurs applications. Journal de l'École Polytechnique 59 (1889) 7-76.
- [10] V. MATVEEV Two-dimensional metrics admitting precisely one projective vector field. Math. Ann. 312 (2012) 865-909.
- [11] Y. R. ROMANOVSKII Computation of Local Symmetries of Second-Order Ordinary Differential Equations by the Carton Equivalence Method. Mathematical Notes, Vol. 60, No. I, 1996.
- 1 UFF, Universidad Federal Fluminense, Rua Prof. Marcos Waldemar de Freitas Reis, S/N Bloco H, 40 Andar Campus do Gragoatá, Niterói, RJ, Brazil, Brasil

2 UNIV RENNES, CNRS, IRMAR, UMR 6625, F-35000 RENNES, FRANCE

 $Email \ address: \ ^1 \ {\tt hfalla@id.uff.br}$ 

Email address: <sup>2</sup> frank.loray@univ-rennes1.fr