



HAL
open science

The genome sequence of the malaria mosquito, *Anopheles funestus*, Giles, 1900

Diego Ayala, Ousman Akone-Ella, Pierre Kengne, Harriet Johnson, Haynes Heaton, Joanna Collins, Ksenia Krasheninnikova, Sarah Pelan, Damon-Lee Pointon, Ying Sims, et al.

► To cite this version:

Diego Ayala, Ousman Akone-Ella, Pierre Kengne, Harriet Johnson, Haynes Heaton, et al.. The genome sequence of the malaria mosquito, *Anopheles funestus*, Giles, 1900. Wellcome Open Research, 2022, 7, pp.287. 10.12688/wellcomeopenres.18445.1 . hal-04107233

HAL Id: hal-04107233

<https://hal.science/hal-04107233v1>

Submitted on 26 May 2023










HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



DATA NOTE

REVISED **The genome sequence of the malaria mosquito, *Anopheles funestus*, Giles, 1900 [version 2; peer review: 2 approved]**

Diego Ayala ^{1,2}, Ousman Akone-Ella², Pierre Kengne ^{1,2}, Harriet Johnson³, Haynes Heaton⁴, Joanna Collins⁵, Ksenia Krasheninnikova⁵, Sarah Pelan ⁵, Damon-Lee Pointon ⁵, Ying Sims⁵, James Torrance ⁵, Alan Tracey⁵, Marcela Uliano-Silva⁵, Katharina von Wyszczetki⁵, Jonathan Wood ⁵, DNA Pipelines collective, Shane McCarthy ^{5,6}, Daniel Neafsey^{7,8}, Alex Makunin ⁵, Mara Lawniczak ⁵

¹MIVEGEC, IRD, Montpellier, 34394, France

²ESV-GAB, Centre Interdisciplinaire de Recherches Médicales de Franceville (CIRMF), Franceville, BP 769, Gabon

³Scientific Operations, Wellcome Sanger Institute, Hinxton, CB10 1SA, UK

⁴CSSE, Auburn University, Auburn, AL, 36849, USA

⁵Tree of Life, Wellcome Sanger Institute, Hinxton, CB10 1SA, UK

⁶Department of Genetics, University of Cambridge, Cambridge, CB2 3EH, UK

⁷Department of Immunology and Infectious Diseases, Harvard T.H. Chan School of Public Health, Boston, MA, 02115, USA

⁸Infectious Disease and Microbiome Program, Broad Institute, Cambridge, MA, 02142, USA

v2 First published: 25 Nov 2022, 7:287
<https://doi.org/10.12688/wellcomeopenres.18445.1>
 Latest published: 27 Mar 2023, 7:287
<https://doi.org/10.12688/wellcomeopenres.18445.2>

Abstract

We present a genome assembly from an individual female *Anopheles funestus* (the malaria mosquito; Arthropoda; Insecta; Diptera; Culicidae). The genome sequence is 251 megabases in span. The majority of the assembly is scaffolded into three chromosomal pseudomolecules with the X sex chromosome assembled. The complete mitochondrial genome was also assembled and is 15.4 kilobases in length.

Keywords



Anopheles funestus, African malaria mosquito, genome sequence, chromosomal inversions



This article is included in the [Tree of Life gateway](#).

Open Peer Review

Approval Status  

	1	2
version 2 (revision) 27 Mar 2023		
version 1 25 Nov 2022	 view	 view

1. **Igor V Sharakhov**, Virginia Polytechnic Institute and State University, Blacksburg, USA

2. **Ben Matthews** , University of British Columbia, Vancouver, Canada

Any reports and responses or comments on the

article can be found at the end of the article.

Corresponding author: Mara Lawniczak (mara@sanger.ac.uk)

Author roles: **Ayala D:** Methodology, Writing – Review & Editing; **Akone-Ella O:** Methodology; **Kengne P:** Methodology; **Johnson H:** Methodology; **Heaton H:** Investigation, Methodology; **Collins J:** Data Curation; **Krasheninnikova K:** Methodology; **Pelan S:** Data Curation; **Pointon DL:** Data Curation; **Sims Y:** Data Curation; **Torrance J:** Data Curation; **Tracey A:** Data Curation; **Uliano-Silva M:** Methodology; **von Wyszetzki K:** Methodology; **Wood J:** Data Curation; **McCarthy S:** Methodology; **Neafsey D:** Conceptualization, Funding Acquisition; **Makunin A:** Formal Analysis, Methodology, Project Administration, Visualization, Writing – Original Draft Preparation; **Lawniczak M:** Conceptualization, Funding Acquisition, Investigation, Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by a Bill & Melinda Gates Foundation Award (INV-009760) to MKNL. The Wellcome Sanger Institute is funded by Wellcome (206194, <https://doi.org/10.35802/206194>), which supports MKNL. Field collection was supported by an ANR grant (ANR-18-CE35-0002-01 –WILDING). SAM is supported from a Wellcome Grant (207492, <https://doi.org/10.35802/207492>). *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

Copyright: © 2023 Ayala D *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Ayala D, Akone-Ella O, Kengne P *et al.* **The genome sequence of the malaria mosquito, *Anopheles funestus*, Giles, 1900 [version 2; peer review: 2 approved]** Wellcome Open Research 2023, 7:287 <https://doi.org/10.12688/wellcomeopenres.18445.2>

First published: 25 Nov 2022, 7:287 <https://doi.org/10.12688/wellcomeopenres.18445.1>

REVISED Amendments from Version 1

The changes we made to the *An. Funestus* genome note are as follows

- added small note on need of additional validation for small variants relative to FUMOZ
- fixed assembly accessions
- added chromosome arm names to HiC plot (Figure 4)
- updated synteny plot (Figure 5) with chromosome arms, inversions & assembly gaps information
- clarified methods for DNA shearing, purification, quantification and fragment size estimation
- fixed minor typos and referencing issues

Any further responses from the reviewers can be found at the end of the article

Species taxonomy

Animalia; Arthropoda; Insecta; Diptera; Culicidae; Anophelinae; Anopheles; *Anopheles funestus*; Giles, 1900 (NCBI txid:62324).

Background

The mosquito *Anopheles funestus* is one of the major malaria vectors in Sub-Saharan Africa¹. Although it can have a sparse and patchy distribution, this mosquito species is present nearly everywhere across the continent from the savannahs of West-Africa, the rainforest of Central Africa, through the dry valleys of East Africa until the Red Island of Madagascar². *Anopheles funestus* breeds in natural and artificial, permanent or semi-permanent water bodies such swamps or rice fields. It is a member of a species group containing at least thirteen species, among which it is the most medically important species³. Its prominent role in the transmission of the malaria parasites is due to its close relation to humans, which provide shelters, breeding sites, and blood meals¹. Although this association makes it highly susceptible to vector campaigns such as indoor residual spraying (IRS) and insecticide treated nets (ITNs), this mosquito species has become resistant to multiple insecticides in many parts of Africa⁴. Therefore, any program aiming at eradicating malaria cannot ignore this species.

At the genetic level, *Anopheles funestus* has been historically neglected in comparison to the members of the *Anopheles gambiae* complex. Multiple studies using genetic markers, such as microsatellites, chromosomal inversions, or DNA sequences have revealed the extraordinary genetic and inversion polymorphism of this species. This genetic richness is likely to underlie its ecological plasticity⁵, its ability to overcome insecticide pressures⁶, and incipient speciation⁷. The first complete genome draft of this mosquito appeared in 2015, originating from a colony derived from wild individuals collected in Mozambique (Fumoz)⁸. Later, the quality of the reference genome for Fumoz was improved using long read sequencing from multiple individuals and Hi-C data⁹. Here, as part of the *Anopheles* Reference Genomes Project (PRJEB5169), we present a chromosomally complete genome sequence for *Anopheles funestus*, based on a single female specimen from La Lopé, Gabon.

Genome sequence report

The genome was sequenced from a single female *Anopheles funestus* collected from La Lopé, Gabon (-0.187, 11.611). A total of 56-fold coverage in Pacific Biosciences single-molecule long reads (N50 10.684 kb) and 68-fold coverage in 10X Genomics read clouds were generated. Primary assembly contigs were scaffolded with chromosome conformation Hi-C data from a female sibling. Manual assembly curation corrected four missing joins or misjoins, reducing the scaffold number by 0.6%.

The final assembly has a total length of 251 Mb in 330 sequence scaffolds with a scaffold N50 of 84.637 Mb (Table 1). 92.38% of the assembly sequence was assigned to three chromosomal-level scaffolds, representing two autosomes (numbered and oriented against the AfunF3 assembly

Table 1. Genome data for *Anopheles funestus*, idAnoFuneDA-416_04.

Project accession data	
Assembly identifier	idAnoFuneDA-416_04
Species	<i>Anopheles funestus</i>
Specimen	idAnoFuneDA-416_04
NCBI taxonomy ID	62324
BioProject	PRJEB53265
BioSample ID	ERS10527360
Isolate information	female, whole organism
Raw data accessions	
PacificBiosciences SEQUEL I	ERR9439501
10X Genomics Illumina	ERR9356795, ERR9356796, ERR9356797, ERR9356798
Hi-C Illumina	ERR9356794
Genome assembly	
Assembly accession	GCA_943734645
Accession of alternate haplotype	GCA_943734845
Span (Mb)	250,713
Number of contigs	349
Contig N50 length (Mb)	24.105
Number of scaffolds	330
Scaffold N50 length (Mb)	84.637
Longest scaffold (Mb)	102.883
BUSCO* genome score	97.6

* BUSCO scores based on the diptera_odb10 (3285) set using \$BUSCO_VERSION. C= complete [S= single copy, D=duplicated], F=fragmented, M=missing, n=number of orthologues in comparison. A full set of BUSCO scores is available at <https://blobtoolkit.genomehubs.org/view/Anopheles%20funestus/dataset/CALSEJ01/busco>.

(9; GCA_003951495.1)), and the X sex chromosome (Figure 1–Figure 4; Table 2). Synteny analysis against the AfunF3 assembly revealed multiple inversions and translocations (Figure 5), correspondence of four largest inversions to known polymorphic inversions in *Anopheles funestus* was revealed based on population genomics¹⁰ and cytogenetics¹¹ data (Table 3), smaller inversions and other variant types will require additional validation.

The assembly has a BUSCO 5.3.2¹² completeness of 97.6% using the diptera_odb10 reference set. While not fully phased, the assembly deposited is of one haplotype. Contigs corresponding to the second haplotype have also been deposited.

Methods

Sample acquisition and nucleic acid extraction

Anopheles funestus offspring were reared from a wild caught gravid female collected from La Lopé, Gabon (latitude -0.187, longitude 11.611) by Ousman Akone-ella. A single female idAnoFuneDA-416_04 was used for Pacific BioSciences and 10x genomics, its sibling female idAnoFuneDA-416_06 was used for Arima Hi-C.

For the high molecular weight (HMW) DNA extraction for *Anopheles* mosquitoes, one whole insect was disrupted by manual grinding with a blue plastic pestle in Qiagen MagAttract lysis buffer and then extracted using the Qiagen MagAttract HMW DNA extraction kit with two minor




Figure 1. Genome assembly of *Anopheles funestus*, idAnoFuneDA-416_04: metrics. The BlobToolKit Snailplot shows N50 metrics and BUSCO gene completeness. The main plot is divided into 1,000 size-ordered bins around the circumference with each bin representing 0.1% of the 250,713,484 bp assembly. The distribution of chromosome lengths is shown in dark grey with the plot radius scaled to the longest chromosome present in the assembly (102,883,511 bp, shown in red). Orange and pale-orange arcs show the N50 and N90 chromosome lengths (84,636,641 and 355,752 bp), respectively. The pale grey spiral shows the cumulative chromosome count on a log scale with white scale lines showing successive orders of magnitude. The blue and pale-blue area around the outside of the plot shows the distribution of GC, AT and N percentages in the same bins as the inner plot. A summary of complete, fragmented, duplicated and missing BUSCO genes in the diptera_odb10 set is shown in the top right. An interactive version of this figure is available at [https://blobtoolkit.genomehubs.org/view/Anopheles funestus/dataset/CALSEJ01/snail](https://blobtoolkit.genomehubs.org/view/Anopheles_funestus/dataset/CALSEJ01/snail).




Figure 2. Genome assembly of *Anopheles funestus*, idAnoFuneDA-416_04: GC coverage. BlobToolKit GC-coverage plot. An interactive version of this figure is available at [https://blobtoolkit.genomehubs.org/view/Anopheles funestus/dataset/CALSEJ01/blob#Filters](https://blobtoolkit.genomehubs.org/view/Anopheles_funestus/dataset/CALSEJ01/blob#Filters).

modifications. These modifications include using half volumes of the kit recommendations due to small sample size (*Anopheles* mosquitoes typically weigh 2–3 mg) and running two elutions of 100 μ l each to increase DNA yield. The quality of the DNA was evaluated using an Agilent FemtoPulse to ensure that most DNA molecules were larger than 30 kb, and preferably > 100 kb. Single mosquito extractions ranged in total estimated DNA yield from 192 ng to 800 ng, with an average yield of 500 ng. Low molecular weight DNA was removed from using an 0.8X AMPure XP purification. A small aliquot (<~5% of the total volume) of HMW DNA

was set aside for 10X Linked Read sequencing and the rest of the DNA was sheared to an average fragment size of 12–20 Kb using a Diagenode Megaruptor 3 at speeds ranging from 27 to 30. Sheared DNA was purified using AMPure PB beads with a 1.8X ratio of beads to sample to remove the shorter fragments and concentrate the DNA sample. The concentration and quality of the sheared and purified DNA was assessed using a Nanodrop spectrophotometer and Qubit Fluorometer with the Qubit dsDNA High Sensitivity Assay kit. Fragment size distribution was evaluated by running the sheared and cleaned sample on the FemtoPulse system once




Figure 3. Genome assembly of *Anopheles funestus*, idAnoFuneDA-416_04: cumulative sequence. BlobToolKit cumulative sequence plot. An interactive version of this figure is available at [https://blobtoolkit.genomehubs.org/view/Anopheles funestus/dataset/CALSEJ01/cumulative#Filters](https://blobtoolkit.genomehubs.org/view/Anopheles_funestus/dataset/CALSEJ01/cumulative#Filters).

more. The median DNA fragment size was 15 kb and the median yield of sheared DNA was 200 ng, with samples typically losing about 50% of the original estimated DNA quantity through the process of shearing and purification.

Sheared DNA was purified using AMPure PB beads with a 1.8X ratio of beads to sample to remove the shorter fragments and concentrate the DNA sample. The concentration of the sheared and purified DNA was assessed using a Nanodrop spectrophotometer and Qubit Fluorometer and Qubit dsDNA High Sensitivity Assay kit. Fragment size distribution was evaluated by running the sample on the FemtoPulse system once more on the sheared and cleaned sample.

For Hi-C samples, a separate sibling whole insect specimen idAnoFuneDA-416_06 was used as input material for the

Arima V2 Kit according to the manufacturer's instructions for animal tissue. This approach of using a sibling was taken in order to enable all material from a single specimen to contribute to the PacBio data generation given we were not always able to meet the minimum suggested guidance of starting with > 300 ng of HMW DNA from a specimen. Samples proceeded to the Illumina library prep stage even if they were suboptimal (too little tissue) going into the Arima reaction.

To assist with annotation, which will be made available through VEuPathDB Vectorbase in due course, RNA was extracted from separate whole unrelated insect specimens idAnoFuneDA-146_02, idAnoFuneDA-367_03, and idAnoFuneDA-367_04 using TRIzol, according to the manufacturer's instructions. RNA was then eluted in 50 µl RNase-free




Figure 4. Genome assembly of *Anopheles funestus*, idAnoFuneDA-416_04: Hi-C contact map. Hi-C contact map of the idAnoFuneDA-416_04 assembly, visualised in HiGlass. Chromosomes are arranged in size order from left to right and top to bottom. The interactive Hi-C map can be viewed at <https://genome-note-higlass.tol.sanger.ac.uk/l/?d=ajmC2VieTICjrBC-4LXsA>.

Table 2. Chromosomal pseudomolecules in the genome assembly of *Anopheles funestus*, idAnoFuneDA-416_04.

INSDC accession	Chromosome	Size (Mb)	Gaps
OX030923.1	2RL	102.884	3
OX030924.1	3RL	84.637	4
OX030925.1	X	22.264	2

water and its concentration assessed using a Nanodrop spectrophotometer and Qubit Fluorometer using the Qubit RNA Broad-Range (BR) Assay kit. Analysis of the integrity of the RNA was done using Agilent RNA 6000 Pico Kit and Eukaryotic Total RNA assay. Samples were not always ideally preserved for RNA, so qualities varied but all were sequenced anyway.

Sequencing

We prepared libraries as per the PacBio procedure and checklist for SMRTbell Libraries using Express TPK 2.0 with low DNA input. Every library was barcoded to support multiplexing. Final library yields ranged from 20 ng to 100 ng, representing only about 25% of the input sheared DNA. Libraries from two specimens were typically multiplexed on a single 8M SMRT Cell. Sequencing complexes were made using Sequencing Primer v4 and DNA Polymerase v2.0. Sequencing




Figure 5. Synteny between genome assemblies of *Anopheles funestus*, AfunF3 and idAnoFuneDA-416_04. Black dots represent locations of assembly gaps, blue vertical lines – approximate centromere locations. Four largest inversions identified as 2Rh, 3Ra, 3Rb, and 3La (Table 3).

Table 3. Known large-scale inversions between genome assemblies of *Anopheles funestus*, AfunF3 and idAnoFuneDA-416_04 identified by syri.
Coordinates given for AfunF3 (ref) and idAnoFuneDA-416_04 (q).

Inversion	Chromosome	Start in ref	End in ref	Start in q	End in q
2Rh	2RL	29671756	38731257	29517507	38117412
3Ra	3RL	1995204	11122881	2428547	12234590
3Rb	3RL	21099051	33133207	21361107	34095918
3La	3RL	66185724	86544782	57224763	76848507

was carried out on the Sequel II system with 24 hour run time and 2 hour pre-extension. A 10X Genomics Chromium read cloud sequencing library was also constructed according to the manufacturer's instructions (this product is no longer available). Only 0.5ng of DNA was used and only 25–50% of the gel emulsion was put forward for library prep due to the small genome size. For Hi-C data generation, following the Arima HiC 2 reaction, samples were processed through Library Preparation using a NEB Next Ultra II DNA Library Prep Kit and sequenced aiming for 100x depth. RNA libraries were created using the directional NEB Ultra II stranded kit. Sequencing was performed by the Scientific Operations core at the Wellcome Sanger Institute on Pacific Biosciences SEQUEL II (HiFi), Illumina NovaSeq 6000 (10X and Hi-C), or Illumina HiSeq 4000 (RNAseq).

Genome assembly

Assembly was carried out with Hifiasm¹³; haplotypic duplication was identified and removed with purge_dups¹⁴. One round of polishing was performed by aligning 10X Genomics read data to the assembly with longranger align, calling variants with freebayes¹⁵. The assembly was then scaffolded with Hi-C data¹⁶ using SALSA2¹⁷. The assembly was checked for contamination as described previously¹⁸. Manual curation was performed using gEVAL¹⁹, HiGlass²⁰ and Pretext (<https://github.com/wtsi-hpag/PretextView>). The mitochondrial genome was assembled using MitoHiFi²¹, which performs annotation using MitoFinder²². The genome was analysed and BUSCO scores generated within the BlobToolKit environment²³. Synteny analysis was performed with syri v1.6²⁴ and visualised with plotsr 0.5.3²⁵. Table 4 contains a list of all software tool versions used, where appropriate.

Ethics/compliance issues

The genetic resources accessed and utilised under this project were done so in accordance with the UK ABS legislation (Nagoya Protocol (Compliance) (Amendment) (EU Exit)

Table 4. Software tools used.

Software tool	Version	Source
hifiasm	0.14	13
purge_dups	1.2.3	14
SALSA2	2.2-4c80ac1	17
longranger align	2.2.2	https://support.10xgenomics.com/genome-exome/software/pipelines/latest/advanced/other-pipelines
freebayes	1.3.1	15
MitoHiFi	2	21
gEVAL	N/A	19
HiGlass	1.11.6	20
PretextView	0.1.x	https://github.com/wtsi-hpag/PretextView
BlobToolKit	2.6.2	23

Regulations 2018 (SI 2018/1393)) and the national ABS legislation within the country of origin, where applicable.

Data availability

European Nucleotide Archive: *Anopheles funestus* genome assembly, idAnoFuneDA-416_04. Accession number PRJEB53266; <https://identifiers.org/bioprotect/PRJEB53266>.

The genome sequence is released openly for reuse. The *Anopheles funestus* genome sequencing initiative is part of the Anopheles

Reference Genomes project. All raw sequence data and the assembly have been deposited in INSDC databases. Raw data and assembly accession identifiers are reported in [Table 1](#).

Author information

Members of the Sanger Scientific Operations: DNA Pipelines collective are listed here: <https://doi.org/10.5281/zenodo.4790456>.

References

- Dia I, Wamdaogo M, Ayal D: **Advances and Perspectives in the Study of the Malaria Mosquito *Anopheles funestus***. In: Manguin S, editor. *Anopheles mosquitoes - New insights into malaria vectors*. InTech; 2013. [Publisher Full Text](#)
- Sinka ME, Bangs MJ, Manguin S, et al.: **The dominant *Anopheles* vectors of human malaria in Africa, Europe and the Middle East: occurrence data, distribution maps and bionomic précis**. *Parasit Vectors*. 2010; **3**: 117. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Harbach RE: **The Phylogeny and Classification of *Anopheles***. In: Manguin S, editor. *Anopheles mosquitoes - New insights into malaria vectors*. InTech; 2013. [Publisher Full Text](#)
- Coetzee M, Koekemoer LL: **Molecular Systematics and Insecticide Resistance in the Major African Malaria Vector *Anopheles funestus***. *Annu Rev Entomol*. 2013; **58**: 393–412. [PubMed Abstract](#) | [Publisher Full Text](#)
- Ayala D, Acevedo P, Pombi M, et al.: **Chromosome inversions and ecological plasticity in the main African malaria mosquitoes**. *Evolution*. 2017; **71**(3): 686–701. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Wondji CS, Irving H, Morgan J, et al.: **Two duplicated P450 genes are associated with pyrethroid resistance in *Anopheles funestus*, a major malaria vector**. *Genome Res*. 2009; **19**(3): 452–459. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Michel AP, Guelbeogo WM, Grushko O, et al.: **Molecular differentiation between chromosomally defined incipient species of *Anopheles funestus***. *Insect Mol Biol*. 2005; **14**(4): 375–387. [PubMed Abstract](#) | [Publisher Full Text](#)
- Neafsey DE, Waterhouse RM, Abai MR, et al.: **Mosquito genomics. Highly evolvable malaria vectors: the genomes of 16 *Anopheles* mosquitoes**. *Science*. 2015; **347**(6217): 1258522. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ghurye J, Koren S, Small ST, et al.: **A chromosome-scale assembly of the major African malaria vector *Anopheles funestus***. *GigaScience*. 2019; **8**(6): giz2063. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lukindu M, Love RR, Guelbeogo MW, et al.: **High-Throughput Genotyping of Common Chromosomal Inversions in the Afrotropical Malaria Mosquito *Anopheles Funestus***. *Insects*. 2020; **11**(10): 693. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sharakhov I, Braginets O, Grushko O, et al.: **A microsatellite map of the African human malaria vector *Anopheles funestus***. *J Hered*. 2004; **95**(1): 29–34. [PubMed Abstract](#) | [Publisher Full Text](#)
- Simão FA, Waterhouse RM, Ioannidis P, et al.: **BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs**. *Bioinformatics*. 2015; **31**(19): 3210–3212. [PubMed Abstract](#) | [Publisher Full Text](#)
- Cheng H, Concepcion GT, Feng X, et al.: **Haplotype-resolved *de novo* assembly using phased assembly graphs with hifiasm**. *Nat Methods*. 2021; **18**(2): 170–175. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Guan D, McCarthy SA, Wood J, et al.: **Identifying and removing haplotypic duplication in primary genome assemblies**. *Bioinformatics*. 2020; **36**(9): 2896–2898. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Garrison E, Marth G: **Haplotype-based variant detection from short-read sequencing**. arXiv [q-bio.GN]. 2012. [Publisher Full Text](#)
- Rao SSP, Huntley MH, Durand NC, et al.: **A 3D Map of the Human Genome at Kilobase Resolution Reveals Principles of Chromatin Looping**. *Cell*. 2014; **159**(7): 1665–1680. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ghurye J, Rhie A, Walenz BP, et al.: **Integrating Hi-C links with assembly graphs for chromosome-scale assembly**. *PLoS Comput Biol*. 2019; **15**(8): e1007273. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Howe K, Chow W, Collins J, et al.: **Significantly improving the quality of genome assemblies through curation**. *GigaScience*. 2021; **10**(1): g1aa153. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Chow W, Brugger K, Caccamo M, et al.: **gEVAL - a web-based browser for evaluating genome assemblies**. *Bioinformatics*. 2016; **32**(16): 2508–2510. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kerpedjiev P, Abdennur N, Lekschas F, et al.: **HiGlass: web-based visual exploration and analysis of genome interaction maps**. *Genome Biol*. 2018; **19**(1): 125. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Uliano-Silva M, Nunes JGF, Krasheninnikova K, et al.: **marcelauliano/MitoHiFi: mitohifi_v2.0**. 2021. [Publisher Full Text](#)
- Allio R, Schomaker-Bastos A, Romiguier J, et al.: **MitoFinder: Efficient automated large-scale extraction of mitogenomic data in target enrichment phylogenomics**. *Mol Ecol Resour*. 2020; **20**(4): 892–905. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Challis R, Richards E, Rajan J, et al.: **BlobToolKit - Interactive quality assessment of genome assemblies**. *G3 (Bethesda)*. 2020; **10**(4): 1361–1374. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Goel M, Sun H, Jiao WB, et al.: **SyRI: finding genomic rearrangements and local sequence differences from whole-genome assemblies**. *Genome Biol*. 2019; **20**(1): 277. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Goel M, Schneeberger K: **plots: visualizing structural similarities and rearrangements between multiple genomes**. *Bioinformatics*. 2022; **38**(10): 2922–2926. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:  

Version 1

Reviewer Report 21 December 2022

<https://doi.org/10.21956/wellcomeopenres.20454.r53476>

© 2022 Matthews B. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Ben Matthews 

¹ Department of Zoology, University of British Columbia, Vancouver, BC, Canada

² Department of Zoology, University of British Columbia, Vancouver, BC, Canada

In their data note entitled: "The genome sequence of the malaria mosquito, *Anopheles funestus*, Giles, 1900," Ayala and colleagues clearly describe the generation of a PacBio long read + 10X genomics genome assembly from a single female offspring from a single wild female mosquito collected in La Lopé, Gabon, with additional scaffolding provided from a sibling via Hi-C. The resulting assembly provides a marked improvement over the previous reference genome, though reveals some discrepancies in the form of inversions and other structural rearrangements that will be important to dig into.

Rationale:

Anopheles funestus is a major malaria vector and thus any information about its biology is critically important for control efforts. A complete and contiguous reference genome is an important piece for a variety of research surrounding the biology of this deadly vector.

Protocols appropriate, and work technically sound:

The collection, HMW DNA extraction, DNA QC, library prep, sequencing, and assembly informatics are clearly described and appropriate. The resulting assembly is technically sound given current state-of-the-art assembly.

Sufficient details of methods and materials:

Generally very clear, with one exception. On page four of the methods, the following sentence is vague: "Following shearing, samples were cleaned using a SPRI clean up sometimes with a bead ratio intended to help remove fragments below 3 kb." In particular, the word sometimes here is confusing - for the assembly in question, was a SPRI clean up used to remove fragments below 3kb? If this approach is used sometimes, how does one decide when and when not to apply it?

What is the bead ratio? This information is required to make the methods complete.

Datasets clearly presented in useable and accessible format:

Table 1 and links provided within the paper provide complete access to the datasets and raw sequencing reads in standard formats at standard repositories.

Is the rationale for creating the dataset(s) clearly described?

Yes

Are the protocols appropriate and is the work technically sound?

Yes

Are sufficient details of methods and materials provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Mosquito genomics, behaviour, neuroscience

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 19 December 2022

<https://doi.org/10.21956/wellcomeopenres.20454.r53455>

© 2022 Sharakhov I. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Igor V Sharakhov

¹ Department of Entomology, Virginia Polytechnic Institute and State University, Blacksburg, USA

² Department of Entomology, Virginia Polytechnic Institute and State University, Blacksburg, USA

The article by Ayala and coauthors present a new genome assembly obtained from an individual female *Anopheles funestus*. The genome was sequenced using the PacBio procedure, polished by 10X Genomics Chromium reads, and scaffolded into three chromosomal pseudomolecules using Hi-C data obtained from a single sibling mosquito. The complete mitochondrial genome was also assembled. The rationale for creating the *Anopheles funestus* genome is clearly described. The previous genome assembly originated from a colony derived from wild individuals, while the new genome assembly is obtained from a single mosquito and is of higher quality. The protocols are appropriate and the work is technically sound. The manuscript provides sufficient details of

methods and materials to allow replication by others. The datasets are clearly presented in a useable and accessible format.

Minor comments:

- The Fumoz genome assembly was improved not just by using long read sequencing but also by Hi-C.
- In Table 1, Assembly accession and Accession of alternate haplotype numbers are mixed up here or on the NCBI website.
- "The median DNA fragment size was 15 kb and the median concentration of sheared DNA was 200 ng," change to "The median DNA fragment size was 15 kb and the median amount of sheared DNA was 200 ng,"
- Label chromosome arms in Figure 4.
- A reference is missing for the Pretext software.
- Figure 5 shows a number of small inversions, duplications and translocations, which could be misassemblies in one of two genomes. Please indicate methods of validation of genomic arrangements.

Is the rationale for creating the dataset(s) clearly described?

Yes

Are the protocols appropriate and is the work technically sound?

Yes

Are sufficient details of methods and materials provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Mosquito evolutionary genetics, cytogenetics, genomics

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.
