

Protocol for emergency dosimetry based on EPR spectrometry on nails and hairs

Lucille Remy, Céline Calas, Francois Trompier

▶ To cite this version:

Lucille Remy, Céline Calas, Francois Trompier. Protocol for emergency dosimetry based on EPR spectrometry on nails and hairs. 11th International Congress of the International Radiation Protection Association, May 2004, Madrid, Spain. hal-04108245

HAL Id: hal-04108245 https://hal.science/hal-04108245

Submitted on 16 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Protocol for emergency dosimetry based on EPR spectrometry on nails and hairs

L. REMY, C. CALAS, F. TROMPIER

External dosimetry department, Institute for Radiological protection and Nuclear Safety (IRSN) BP 17 92262 Fontenay-aux-Roses cedex, France Contact : francois.trompier@irsn.fr

CONTEXT

In the case of a radiological accident due to an external overexposure involving a limited number of persons from the public or workers without dosemeters, dosimetry by Electronic Paramagnetic Resonance (EPR) spectrometry on tooth enamel, among other techniques such as biological dosimetry, can be an efficient tool. However, for an accident involving a large number of victims, most of dosimetric techniques are limited by the necessary short delay of the answer or the invasive sampling. Indeed, in this specific case, it is very important in a first time and as quick as possible to sort population according to the severity of the exposure. Nevertheless, EPR dosimetry on materials easily sampled on the victims (hairs or nails) may be a pertinent tool. In this context, the objective of this work was to study the EPR properties of nails or hairs to propose a first protocol for fast triage of population.



peak to peak amplitude of native signal (before irradiation), F: fading correction factor depending on time elapsed between irradiation and measurement t, temperature T° and humidity H and A: dose coefficient

Dose estimation Protocol

The protocol can be divided in different stages. A first estimation can be done very quickly using averaged value of main characteristics (fading constant, radiation sensitivity, amplitude of background signal) previously determined on a large set of samples. At each next stage, the determination of the real values of the different constants for each sample allows to increase the accuracy of the dose estimation.



Conclusion

This study showed that it was possible, starting from EPR measurement of nails or hairs, to carry out a population triage according to the severity of the exposure

