



HAL
open science

Characterization of weighted Hardy spaces on which all composition operators are bounded

Pascal Lefèvre, Daniel Li, Hervé Queffélec, Luis Rodríguez-Piazza

► **To cite this version:**

Pascal Lefèvre, Daniel Li, Hervé Queffélec, Luis Rodríguez-Piazza. Characterization of weighted Hardy spaces on which all composition operators are bounded. 2023. hal-04108498v1

HAL Id: hal-04108498

<https://hal.science/hal-04108498v1>

Preprint submitted on 27 May 2023 (v1), last revised 23 Oct 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Characterization of weighted Hardy spaces on which all composition operators are bounded

*Pascal Lefèvre, Daniel Li,
Hervé Queffélec, Luis Rodríguez-Piazza*

January 25, 2023

Abstract. We give a complete characterization of the sequences $\beta = (\beta_n)$ of positive numbers for which all composition operators on $H^2(\beta)$ are bounded, where $H^2(\beta)$ is the space of analytic functions f on the unit disk \mathbb{D} such that $\sum_{n=0}^{\infty} |a_n|^2 \beta_n < +\infty$ if $f(z) = \sum_{n=0}^{\infty} a_n z^n$. We prove that all composition operators are bounded on $H^2(\beta)$ if and only if β is essentially decreasing and slowly oscillating. We also prove that every automorphism of the unit disk induces a bounded composition operator on $H^2(\beta)$ if and only if β is slowly oscillating. We give applications of our results.

MSC 2010. primary: 47B33 ; secondary: 30H10

Key-words. automorphism of the unit disk; composition operator; slowly oscillating sequence; weighted Hardy space

1 Introduction

Let $\beta = (\beta_n)_{n \geq 0}$ be a sequence of positive numbers such that

$$(1.1) \quad \liminf_{n \rightarrow \infty} \beta_n^{1/n} \geq 1.$$

The associated weighted Hardy space $H^2(\beta)$ is the Hilbertian space of analytic functions $f(z) = \sum_{n=0}^{\infty} a_n z^n$ such that

$$(1.2) \quad \|f\|^2 := \sum_{n=0}^{\infty} |a_n|^2 \beta_n < \infty.$$

Condition (1.1) is equivalent to the inclusion $H^2(\beta) \subseteq \mathcal{H}ol(\mathbb{D})$. Indeed, if (1.1) holds, we have $H^2(\beta) \subseteq \mathcal{H}ol(\mathbb{D})$ since $|a_n|^2 \beta_n$ is bounded and thanks to the Hadamard formula. Conversely, testing the inclusion $H^2(\beta) \subseteq \mathcal{H}ol(\mathbb{D})$ on the function $f(z) = \sum_{n=1}^{\infty} \frac{1}{n\sqrt{\beta(n)}} z^n \in H^2(\beta)$, we get (1.1) from the Hadamard formula.

Condition (1.1) will therefore be assumed throughout this paper, without repeating it.

When $\beta_n \equiv 1$, we recover the usual Hardy space H^2 ; the Bergman space corresponds to $\beta_n = 1/(n+1)$, and the Dirichlet space to $\beta_n = n+1$.

Recall that a symbol is a (non constant) analytic self-map $\varphi: \mathbb{D} \rightarrow \mathbb{D}$, and the associated composition operator $C_\varphi: H^2(\beta) \rightarrow \mathcal{H}ol(\mathbb{D})$ is defined as:

$$(1.3) \quad C_\varphi(f) = f \circ \varphi.$$

An important question in the theory is to decide when C_φ is bounded on $H^2(\beta)$, i.e. when $C_\varphi: H^2(\beta) \rightarrow H^2(\beta)$.

This question appears in the literature in several places. For instance, it is Problem 1 in the thesis of Nina Zorboska [22, Pb 1, p. 49]. This thesis contains many interesting results, in particular Proposition 3.1 and Proposition 4.2 of the present paper (actually we discovered the content of Zorboska's thesis once the present paper was almost finished). See also Question 36 raised by Deddens in [18, p. 122.c].

When $H^2(\beta)$ is the usual Hardy space H^2 (i.e. when $\beta_n \equiv 1$), it is well-known [17, pp. 13–17] that all symbols generate bounded composition operators. On the other hand, for the Dirichlet space, corresponding to $\beta_n = n+1$, not all composition operators are bounded since there exist symbols φ not belonging to the Dirichlet space (e.g. any infinite Blaschke product).

Note that, by definition of the norm of $H^2(\beta)$, all rotations R_θ , defined by $R_\theta(z) = e^{i\theta}z$, with $\theta \in \mathbb{R}$, induce bounded and surjective composition operators on $H^2(\beta)$ and send isometrically $H^2(\beta)$ into itself.

Our goal in this paper is characterizing the sequences β for which all composition operators act boundedly on the space $H^2(\beta)$, i.e. send $H^2(\beta)$ into itself.

In Shapiro's presentation for the Hardy space H^2 , the main point is the case $\varphi(0) = 0$ and a subordination principle for subharmonic functions (Littlewood's subordination principle). The case of automorphisms is claimed simple, using an integral representation for the norm and some change of variable. For general weights β , the situation is different, as we will see in this paper, and it turns out that the conditions on β for the boundedness of the composition operators C_φ on $H^2(\beta)$ are not the same depending on whether we consider the class of all symbols such that $\varphi(0) = 0$, or the class of symbols $\varphi = T_a$, where

$$(1.4) \quad T_a(z) = \frac{a+z}{1+\bar{a}z}$$

for $a \in \mathbb{D}$.

It is clear that when these two classes of composition operators are bounded, then all composition operators are bounded. Recall that every symbol φ can be written as the composition $\varphi = T_a \circ \psi$ where $\psi(0) = 0$ and $a = \varphi(0)$; and then $C_\varphi = C_\psi \circ C_{T_a}$.

In many occurrences, the weight β is defined as

$$(1.5) \quad \beta_n = \int_0^1 t^n d\sigma(t)$$

where σ is a positive measure on $(0, 1)$; more specifically the following definition is often used: let $G: (0, 1) \rightarrow \mathbb{R}_+$ be an integrable function and let H_G^2 be the space of analytic functions $f: \mathbb{D} \rightarrow \mathbb{C}$ such that:

$$(1.6) \quad \|f\|_{H_G^2}^2 := \int_{\mathbb{D}} |f(z)|^2 G(1 - |z|^2) dA(z) < \infty.$$

Such weighted Bergman type spaces are used, for instance, in [10], [11] and in [13]. We have $H_G^2 = H^2(\beta)$ with:

$$(1.7) \quad \beta_n = 2 \int_0^1 r^{2n+1} G(1 - r^2) dr = \int_0^1 t^n G(1 - t) dt,$$

and the sequence $\beta = (\beta_n)_n$ is *non-increasing* (actually, the above representation (1.5) is equivalent, by the Hausdorff moment theorem, to a high regularity of the sequence β , namely its *complete monotony*).

When the weight β is non-increasing (or more generally, essentially decreasing), all the symbols vanishing at the origin induce a bounded composition operator. This was proved by C. Cowen [4, Corollary, page 31], using Hadamard multiplication. We can also use Kacnel'son's theorem (see [2] or [12, Theorem 3.12]). Actually that follows from an older theorem of Goluzin [8] (see [5, Theorem 6.3]), which itself uses a self-refinement observed by Rogosinski of Littlewood's principle ([5, Theorem 6.2]).

For weights defined as in (1.5), we dispose of integral representations for the norm in $H^2(\beta)$, and, as in the Hardy space case, this integral representation rather easily allows us to decide when the boundedness of C_{T_a} on $H^2(\beta)$ occurs. This is not always the case, as shown by T. Kriete and B. MacCluer in [11]. They consider spaces of Bergman type $A_G^2 := H_G^2$, where $\tilde{G}(r) = G(1 - r^2)$, defined as the spaces of analytic functions in \mathbb{D} such that $\int_{\mathbb{D}} |f(z)|^2 \tilde{G}(|z|) dA < \infty$, for a positive non-increasing continuous function \tilde{G} on $[0, 1)$. They prove [11, Theorem 3] that for

$$\tilde{G}(r) = \exp\left(-B \frac{1}{(1-r)^\alpha}\right), \quad B > 0, \quad 0 < \alpha \leq 2,$$

and

$$\varphi(z) = z + t(1-z)^\gamma, \quad 1 < \gamma \leq 3, \quad 0 < t < 2^{1-\gamma},$$

then φ is a symbol and C_φ is bounded on A_G^2 if and only if $\gamma \geq \alpha + 1$.

Here

$$\beta_n = \int_0^1 t^n e^{-B/(1-\sqrt{t})^\alpha} dt \lesssim \exp(-cn^{\alpha/(\alpha+1)}).$$

Point out that β is non-increasing, so for every symbol φ fixing the origin, the composition operator C_φ is bounded. Nevertheless, choosing $\gamma < \alpha + 1$, there exist symbols inducing an unbounded composition operator, hence not all the C_{T_a} are bounded. Actually, for every $\alpha \in (0, 2]$, no C_{T_a} is bounded because β has no polynomial lower estimate (see Proposition 4.5 below).

Let us describe the contents of this paper. In Section 2, we introduce several notion of growth or regularity for a sequence β : essentially decreasing, polynomial decay and polynomial growth, slow oscillation, and give some connections between them. In Section 3, we consider the composition operators whose symbol vanishes at the origin. We show that in order for all these operators to be bounded, it is necessary that β be bounded above. We show that β is essentially decreasing if and only if all these operators are bounded and $\sup_{\varphi(0)=0} \|C_\varphi\| < +\infty$. In Theorem 3.3, we give a sufficient condition for having all the composition operators C_φ with $\varphi(0) = 0$ bounded, allowing us to give an example of a sequence β for which this happens though $\sup_{\varphi(0)=0} \|C_\varphi\| = +\infty$ (Theorem 3.7). In Section 4, we prove that all C_{T_a} are bounded on $H^2(\beta)$ if and only if β is slowly oscillating (Theorem 4.7 and Theorem 4.10). We hence have:

Theorem 1.1. *Let β be a sequence of positive numbers, and let*

$$T_a(z) = \frac{a + z}{1 + \bar{a}z}$$

for $a \in \mathbb{D}$. The following assertions are equivalent:

- 1) for some $a \in \mathbb{D} \setminus \{0\}$, the map T_a induces a bounded composition operator C_{T_a} on $H^2(\beta)$;
- 2) for all $a \in \mathbb{D}$, the maps T_a induce bounded composition operators C_{T_a} on $H^2(\beta)$;
- 3) β is slowly oscillating.

The deep implication is 2) \Rightarrow 3). Its proof requires some sharp estimates on the mean of Taylor coefficients of T_a for a belonging to a subinterval of $(0, 1)$. Once we found the equivalence of 1) and 2), we realized that it already appeared in Zorboska's thesis [22].

In Section 5, we show (Theorem 5.1) that if β is slowly oscillating, and moreover all composition operators are bounded on $H^2(\beta)$, then β is essentially decreasing. We thus obtain the following theorem.

Theorem 1.2. *Let β be a sequence of positive numbers. Then all composition operators on $H^2(\beta)$ are bounded if and only if β is essentially decreasing and slowly oscillating.*

For the notion of essentially decreasing and slowly oscillating sequences, see Definition 2.1 and Definition 2.2

We end the paper with some results about multipliers.

A first version of this paper, not including the complete characterization given here, was put on arXiv on 30 November 2020 (and a second version on 21 March 2022) under the title *Boundedness of composition operators on general weighted Hardy spaces of analytic functions*.

2 Definitions, notation, and preliminary results

The open unit disk of \mathbb{C} is denoted \mathbb{D} and we write \mathbb{T} its boundary $\partial\mathbb{D}$. We set $e_n(z) = z^n$, $n \geq 0$.

The weighted Hardy space $H^2(\beta)$ defined in the introduction is a Hilbert space with the canonical orthonormal basis

$$(2.1) \quad e_n^\beta(z) = \frac{1}{\sqrt{\beta_n}} z^n, \quad n \geq 0,$$

and the reproducing kernel K_w given for all $w \in \mathbb{D}$ by

$$(2.2) \quad K_w(z) = \sum_{n=0}^{\infty} e_n^\beta(z) \overline{e_n^\beta(w)} = \sum_{n=0}^{\infty} \frac{1}{\beta_n} \bar{w}^n z^n.$$

Note that H^2 is continuously embedded in $H^2(\beta)$ if and only if β is bounded above. In particular, this is the case when β is non-increasing. In this paper, we need a slightly more general notion.

Definition 2.1. *A sequence of positive numbers $\beta = (\beta_n)_{n \geq 0}$ is said to be essentially decreasing if, for some constant $C \geq 1$, we have, for all $m \geq n \geq 0$:*

$$(2.3) \quad \beta_m \leq C \beta_n.$$

Note that saying that β is essentially decreasing means that the shift operator on $H^2(\beta)$ is power bounded.

If β is essentially decreasing, and if we set:

$$\tilde{\beta}_n = \sup_{m \geq n} \beta_m,$$

the sequence $\tilde{\beta} = (\tilde{\beta}_n)$ is non-increasing and we have $\beta_n \leq \tilde{\beta}_n \leq C \beta_n$. In particular, $H^2(\beta) = H^2(\tilde{\beta})$ (with equivalent norms) and H^2 is continuously embedded in $H^2(\beta)$.

Definition 2.2. *We say that a sequence β is slowly oscillating if there are positive constants $c < 1 < C$ such that*

$$(2.4) \quad c \leq \frac{\beta_m}{\beta_n} \leq C \quad \text{when } n/2 \leq m \leq 2n.$$

We may remark that this is equivalent to the existence of some function $\rho: (0, \infty) \rightarrow (0, \infty)$ which is bounded above on each compact subset of $(0, \infty)$ and for which $\beta_m/\beta_n \leq \rho(m/n)$, equivalently

$$\frac{1}{\rho(n/m)} \leq \frac{\beta_m}{\beta_n} \leq \rho(m/n).$$

Definition 2.3. *The sequence of positive numbers $\beta = (\beta_n)$ is said to have polynomial decay if there are positive constants c and α such that, for all integers $n \geq 1$:*

$$(2.5) \quad \beta_n \geq c n^{-\alpha}.$$

That means that $H^2(\beta)$ is continuously embedded in the weighted Bergman space $\mathfrak{B}_{\alpha-1}^2$ of the analytic functions $f: \mathbb{D} \rightarrow \mathbb{C}$ such that

$$\|f\|_{\mathfrak{B}_{\alpha-1}^2}^2 := \alpha \int_{\mathbb{D}} |f(z)|^2 (1 - |z|^2)^{\alpha-1} dA(z) < \infty$$

since $\mathfrak{B}_{\alpha-1}^2 = H^2(\gamma)$ with $\gamma_n \approx n^{-\alpha}$.

Definition 2.4. *The sequence of positive numbers $\beta = (\beta_n)$ is said to have polynomial growth if there are positive constants C and γ such that, for all integers $n \geq 1$:*

$$(2.6) \quad \beta_n \leq C n^\gamma.$$

The following simple proposition links those notions.

Proposition 2.5.

- 1) *Every slowly oscillating sequence β has polynomial decay and growth.*
- 2) *There are sequences that are essentially decreasing, and with polynomial decay, but not slowly oscillating.*
- 3) *There are bounded sequences that are slowly oscillating, but not essentially decreasing.*

Proof. 1) This is clear, because, for some $c \in (0, 1)$, if $2^j \leq n < 2^{j+1}$, then

$$\beta_n \geq c \beta_{2^j} \geq c^{j+1} \beta_1 \geq c \beta_1 n^{-\alpha},$$

with $\alpha = \log(1/c)/\log 2$; and, for some $C > 1$,

$$\beta_n \leq C \beta_{2^j} \leq C^{j+1} \beta_1 \leq C \beta_1 n^\gamma,$$

with $\gamma = \log C/\log 2$.

- 2) Let $\delta > 0$. We set $\beta_0 = \beta_1 = 1$ and for $n \geq 2$:

$$\beta_n = \frac{1}{(k!)^\delta} \quad \text{when } k! < n \leq (k+1)!.$$

The sequence β is non-increasing.

For n and k as above, we have:

$$\beta_n = \frac{1}{(k!)^\delta} \geq \frac{1}{n^\delta};$$

hence β has arbitrarily slow polynomial decay. However we have, for $k \geq 2$:

$$\frac{\beta_{2(k!)}}{\beta_{k!}} = \frac{(k!)^{-\delta}}{[(k-1)!]^{-\delta}} = \frac{1}{k^\delta} \xrightarrow{k \rightarrow \infty} 0,$$

so β is not slowly oscillating.

3) We define β_n as follows. Let (a_k) be an increasing sequence of positive square integers such that $\lim_{k \rightarrow \infty} a_{k+1}/a_k = \infty$, for example $a_k = 4^{k^2}$, and let $b_k = \sqrt{a_k a_{k+1}}$; with our choice, this is an integer and we clearly have $a_k < b_k < a_{k+1}$. We set:

$$\beta_n = \begin{cases} a_k/n & \text{for } a_k \leq n < b_k \\ (a_k/b_k^2) n = (1/a_{k+1}) n & \text{for } b_k \leq n < a_{k+1}. \end{cases}$$

This sequence (β_n) is slowly oscillating by construction. Indeed, since the other cases are obvious, it suffices to check that for $a_k \leq n/2 < b_k \leq n < a_{k+1}$, the quotient β_m/β_n remains lower and upper bounded when $n/2 \leq m \leq n$ (it will then be automatically also satisfied when $n \leq m \leq 2n$). But for $n/2 \leq m < b_k$, we have

$$\frac{\beta_m}{\beta_n} = \frac{a_k/m}{n/a_{k+1}} = \frac{a_k a_{k+1}}{mn} = \frac{b_k^2}{mn},$$

which is $\leq 2b_k^2/n^2 \leq 2$ and $\geq b_k^2/n^2 \geq (n/2)^2/n^2 = 1/4$; and for $b_k \leq m$, we have

$$\frac{\beta_m}{\beta_n} = \frac{m/a_{k+1}}{n/a_{k+1}} = \frac{m}{n} \in \left[\frac{1}{2}, 1\right].$$

However, though (β_n) is bounded, since $\beta_n \leq 1$ for $a_k \leq n < b_k$ and, for $b_k \leq n < a_{k+1}$,

$$\beta_n \leq \beta_{a_{k+1}-1} = \frac{1}{a_{k+1}} (a_{k+1} - 1) \leq 1,$$

it is not essentially decreasing, since

$$\frac{\beta_{a_{k+1}-1}}{\beta_{b_k}} = \frac{1}{\sqrt{a_k a_{k+1}}} (a_{k+1} - 1) \sim \sqrt{\frac{a_{k+1}}{a_k}} \xrightarrow{k \rightarrow \infty} \infty. \quad \square$$

Now we are going to recall some well known facts about matrix representation of an operator T defined on a Hilbert space with an orthonormal basis $(e_n)_{n \geq 0}$ and how it is translated in our framework.

The entry $a_{m,n}$ (where $m, n \geq 0$) is defined by the m^{th} coordinate of $T(e_n)$:

$$a_{m,n} = e_m^*(T(e_n))$$

where $e_k^*(x)$ stands for the k^{th} coordinate of the vector x .

We shall use the notation $\widehat{f}(k)$ for the k^{th} Fourier coefficient of a function $f \in L^1(-\pi, \pi)$:

$$\widehat{f}(k) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) e^{-ikt} dt.$$

Let us point out that when the operator is the composition operator C_φ associated to the symbol φ , viewed on $H^2(\beta)$, its matrix representation in the basis $(e_n^\beta)_{n \geq 0}$ has an entry (m, n) which can be written as

$$(e_m^\beta)^*(C_\varphi(e_n^\beta)) = \frac{\sqrt{\beta_m}}{\sqrt{\beta_n}} e_m^*(\varphi^n) = \frac{\sqrt{\beta_m}}{\sqrt{\beta_n}} \widehat{\varphi^n}(m)$$

since the m^{th} Taylor coefficient of φ^n coincides with its m^{th} Fourier coefficient.

3 Boundedness of composition operators whose symbol vanishes at the origin

3.1 Necessary conditions

We begin with this simple observation (see [22, Prop. 3.1]).

Proposition 3.1. *If all composition operators with symbol vanishing at 0 are bounded on $H^2(\beta)$, then the sequence β is bounded above.*

Proof. Let $f \in H^\infty$. Write $f = A\varphi + f(0)$ where A is a constant and φ a symbol vanishing at 0. We have $\varphi = C_\varphi(z) \in H^2(\beta)$, by hypothesis. So that $f \in H^2(\beta)$ and $H^\infty \subseteq H^2(\beta)$. It follows (by the closed graph theorem, since the convergence in norm implies pointwise convergence) that there exists a constant M such that $\|f\|_{H^2(\beta)} \leq M \|f\|_\infty$ for all $f \in H^\infty$. Testing that with $f(z) = z^n$, we get $\beta_n \leq M^2$. \square

Let us point out that boundedness of β_n does not suffice. For example, let (β_n) be a sequence such that $\beta_{4k+2}/\beta_{2k+1} \xrightarrow{k \rightarrow \infty} \infty$ (for instance $\beta_{2k} = 1$ and $\beta_{2k+1} = 1/(k+1)$); if $\varphi(z) = z^2$, then $\|C_\varphi(z^{2n+1})\|^2 = \|z^{2(2n+1)}\|^2 = \beta_{2(2n+1)}$; since $\|z^{2n+1}\|^2 = \beta_{2n+1}$, the operator C_φ is not bounded on $H^2(\beta)$.

A partial characterization is given in the next proposition.

Proposition 3.2. *The following assertions are equivalent:*

1) *all symbols φ such that $\varphi(0) = 0$ induce bounded composition operators C_φ on $H^2(\beta)$ and*

$$(3.1) \quad \sup_{\varphi(0)=0} \|C_\varphi\| < \infty;$$

2) *β is an essentially decreasing sequence.*

Of course, by the uniform boundedness principle, (3.1) is equivalent to:

$$\sup_{\varphi(0)=0} \|f \circ \varphi\| < \infty \quad \text{for all } f \in H^2(\beta).$$

Let us point out an important fact: we shall see in Theorem 3.7 that there are weights β for which all composition operators C_φ with $\varphi(0) = 0$ are bounded, but $\sup_{\varphi(0)=0} \|C_\varphi\| = +\infty$.

Proof. 2) \Rightarrow 1) We may assume that β is non-increasing. Then the Goluzin-Rogosinski theorem ([5, Theorem 6.3]) gives the result; in fact, writing $f(z) = \sum_{n=0}^{\infty} c_n z^n$ and $(C_\varphi f)(z) = \sum_{n=0}^{\infty} d_n z^n$, it says that

$$\sum_{0 \leq k \leq n} |d_k|^2 \leq \sum_{0 \leq k \leq n} |c_k|^2 \quad \forall n \geq 0,$$

and hence, by Abel summation:

$$\|C_\varphi f\|^2 = \sum_{n=0}^{\infty} |d_n|^2 \beta_n \leq \sum_{n=0}^{\infty} |c_n|^2 \beta_n = \|f\|^2,$$

leading to C_φ bounded and $\|C_\varphi\| \leq 1$. This result was also proved by C. Cowen [4, Corollary of Theorem 7]. Alternatively, we can use a result of Kacnel'son ([9]; see also [2], [3, Corollary 2.2], or [12, Theorem 3.12]).

1) \Rightarrow 2) Set $M = \sup_{\varphi(0)=0} \|C_\varphi\|$. Let $m > n$, and take

$$\varphi(z) = \varphi_{m,n}(z) = z \left(\frac{1 + z^{m-n}}{2} \right)^{1/n}.$$

Then $\varphi(0) = 0$ and $[\varphi(z)]^n = \frac{z^n + z^m}{2}$; hence

$$\frac{1}{4} (\beta_n + \beta_m) = \|\varphi^n\|^2 = \|C_\varphi(e_n)\|^2 \leq \|C_\varphi\|^2 \|e_n\|^2 \leq M^2 \beta_n,$$

so β is essentially decreasing. □

Remark. Let us mention the following example. For $0 < r < 1$, let $\beta_n = \pi n r^{2n}$, for $n \geq 1$ and $\beta_0 = 1$. This sequence is eventually decreasing, so it is essentially decreasing. The quantity $\|f\|_{H^2(\beta)}^2 - |f(0)|^2$ is the area of the part of the Riemann surface on which $r\mathbb{D}$ is mapped by f . E. Reich [15], generalizing Goluzin's result [8] (see [5, Theorem 6.3]), proved that for all symbols φ such that $\varphi(0) = 0$, the composition operator C_φ is bounded on $H^2(\beta)$ and

$$\|C_\varphi\| \leq \sup_{n \geq 1} \sqrt{n} r^{n-1} \leq \frac{1}{\sqrt{2e}} \frac{1}{r \sqrt{\log(1/r)}}.$$

For $0 < r < 1/\sqrt{2}$, Goluzin's theorem asserts that $\|C_\varphi\| \leq 1$.

Note that this sequence β is not slowly oscillating since $\beta_{2n}/\beta_n = 2r^{2n}$. Hence, from Theorem 4.10 below, we get that no composition operator C_{T_a} is bounded on $H^2(\beta)$.

However, that the weight β is essentially decreasing is not necessary for the boundedness of all composition operators C_φ , with symbol φ vanishing at 0, as we will see later (Theorem 3.7).

3.2 Sufficient condition

Theorem 3.3. *Let $\beta = (\beta_n)_{n=0}^\infty$ be a sequence of positive numbers with polynomial decay and growth. Assume that β is weakly decreasing, i.e.:*

$$(3.2) \quad \text{For every } \delta > 0, \text{ there exists a positive constant } C = C(\delta) \text{ such that} \\ \beta_m \leq C \beta_n \quad \text{whenever } m > (1 + \delta)n.$$

Then, for all symbols $\varphi: \mathbb{D} \rightarrow \mathbb{D}$ vanishing at 0, the composition operator C_φ is bounded on $H^2(\beta)$.

Let us point out that (3.2) implies that β is bounded. For a counterexample with an exponential weight, see [22, Ex. 1, p. 14-15].

To prove Theorem 3.3, we need several lemmas.

Lemma 3.4. *Let $\varphi: \mathbb{D} \rightarrow \mathbb{D}$ be an analytic self-map such that $\varphi(0) = 0$ and $|\varphi'(0)| < 1$. Then there exists $\rho > 0$ such that, for every integers n and m ,*

$$|\widehat{\varphi^n}(m)| \leq \exp\left(-\frac{1}{2}[(1 + \rho)n - m]\right).$$

Proof. Since $\varphi(0) = 0$, we can write $\varphi(z) = z\varphi_1(z)$. Since $|\varphi'(0)| < 1$, we have $\varphi_1: \mathbb{D} \rightarrow \mathbb{D}$. Let $M(r) = \sup_{|z|=r} |\varphi_1(z)|$. Cauchy's inequalities say that $|\widehat{\varphi_1^n}(m)| \leq [M(r)]^n / r^m$. We have $M(r) < 1$, so there exists a positive number $\rho = \rho(r)$ such that $M(r) = r^\rho$. We get:

$$|\widehat{\varphi^n}(m)| = |\widehat{\varphi_1^n}(m - n)| \leq \frac{r^{\rho n}}{r^{m-n}} = r^{(1+\rho)n-m},$$

and the result follows, by taking $r = e^{-1/2}$. □

The next lemma is a variant of the following result of V. È. Kacnel'son ([9]; see also [2], [3, Corollary 2.2], or [12, Theorem 3.12]).

Theorem 3.5 (V. È. Kacnel'son). *Let H be a separable complex Hilbert space and $(e_i)_{i \geq 0}$ a fixed orthonormal basis of H . Let $M: H \rightarrow H$ be a bounded linear operator. We assume that the matrix of M with respect to this basis is lower-triangular: $\langle Me_j | e_i \rangle = 0$ for $i < j$.*

Let $(\gamma_j)_{j \geq 0}$ be a non-decreasing sequence of positive real numbers and Γ the (possibly unbounded) diagonal operator such that $\Gamma(e_j) = \gamma_j e_j$, $j \geq 0$. Then the operator $\Gamma^{-1}M\Gamma: H \rightarrow H$ is bounded and moreover:

$$\|\Gamma^{-1}M\Gamma\| \leq \|M\|.$$

This variant is used implicitly in [12, page 13].

Lemma 3.6. *Let $A: \ell_2 \rightarrow \ell_2$ be a bounded operator represented by the matrix $(a_{m,n})_{m,n}$, i.e. $a_{m,n} = \langle A e_n, e_m \rangle$, where $(e_n)_{n \geq 0}$ is the canonical basis of ℓ_2 .*

Let (d_n) be a sequence of positive numbers such that, for every m and n :

$$(3.3) \quad d_m < d_n \implies a_{m,n} = 0.$$

Then, D being the (possibly unbounded) diagonal operator with entries d_n , we have:

$$\|D^{-1}AD\| \leq \|A\|.$$

The proof is the same as that of Kacnel'son's theorem, but we reproduce it for the convenience of the reader. Actually we propose two different proofs.

Proof 1. Let \mathbb{C}_0 be the right-half plane $\mathbb{C}_0 = \{z \in \mathbb{C}; \Re z > 0\}$. We set $H_N = \text{span}\{e_n; n \leq N\}$ and

$$A_N = P_N A J_N,$$

where P_N is the orthogonal projection from ℓ_2 onto H_N and J_N the canonical injection from H_N into ℓ_2 . We consider, for $z \in \overline{\mathbb{C}_0}$:

$$A_N(z) = D^{-z} A_N D^z: H_N \rightarrow H_N,$$

where $D^z(e_n) = d_n^z e_n$.

If $(a_{m,n}(z))_{m,n}$ is the matrix of $A_N(z)$ on the basis $\{e_n; n \leq N\}$ of H_N , we clearly have:

$$a_{m,n}(z) = a_{m,n}(d_n/d_m)^z.$$

In particular, we have, thanks to (3.3):

$$a_{m,n}(z) = 0 \quad \text{if } d_m < d_n,$$

and

$$|a_{m,n}(z)| \leq \sup_{k,l} |a_{k,l}| := M, \quad \text{for all } z \in \overline{\mathbb{C}_0}.$$

Since $\|A_N(z)\|^2 \leq \|A_N(z)\|_{HS}^2 = \sum_{m,n \leq N} |a_{m,n}(z)|^2 \leq (N+1)^2 M^2$, we get:

$$\|A_N(z)\| \leq (N+1)M \quad \text{for all } z \in \overline{\mathbb{C}_0}.$$

Let us consider the function $u: \overline{\mathbb{C}_0} \rightarrow \overline{\mathbb{C}_0}$ defined by:

$$(3.4) \quad u_N(z) = \|A_N(z)\|.$$

This function u_N is continuous on $\overline{\mathbb{C}_0}$, bounded above by $(N+1)M$, and subharmonic in \mathbb{C}_0 . Moreover, thanks to (3.3), the maximum principle gives:

$$\sup_{\overline{\mathbb{C}_0}} u_N(z) = \sup_{\partial \mathbb{C}_0} u_N(z).$$

Since $\|D^z\| = \|D^{-z}\| = 1$ for $z \in \partial\mathbb{C}_0$, we have $\|A_N(z)\| \leq \|A_N\|$ for $z \in \partial\mathbb{C}_0$, and we get:

$$\sup_{\overline{\mathbb{C}_0}} u_N(z) \leq \|A_N\| \leq \|A\|.$$

In particular $u_N(1) \leq \|A\|$, and, letting N going to infinity, we obtain that $\|D^{-1}AD\| \leq \|A\|$.

Proof 2. Since d_n is positive, we can write $d_n = e^{-\rho_n}$ where $\rho_n \in \mathbb{R}$. Let $x = (x_n)_{n \geq 0}$ and $y = (y_n)_{n \geq 0} \in \ell^2$ with finite support, we are interested in controlling the sum

$$S = \sum_{m,n} a_{m,n} \frac{d_n}{d_m} x_n \overline{y_m}$$

which can also be written

$$S = \sum_{m,n} a_{m,n} e^{-|\rho_n - \rho_m|} x_n \overline{y_m}.$$

since the non trivial part of the sum runs over the pairs (m, n) such that $d_m \geq d_n$ i.e. $\rho_n \geq \rho_m$.

Now we introduce the function $f(t) = \frac{1}{\pi(1+t^2)}$ for $t \in \mathbb{R}$, which is positive and belongs to the unit ball of $L^1(\mathbb{R})$. Moreover, its Fourier transform satisfies, for every $x \in \mathbb{R}$,

$$\mathcal{F}(f)(-x) = \int_{\mathbb{R}} f(t) e^{ixt} dt = e^{-|x|}.$$

We get

$$S = \int_{\mathbb{R}} f(t) \left(\sum_{m,n} a_{m,n} x_n e^{i\rho_n t} \overline{y_m e^{i\rho_m t}} \right) dt = \int_{\mathbb{R}} f(t) \langle A(x(t)), y(t) \rangle_{\ell^2} dt$$

where

$$x(t) = (x_n e^{i\rho_n t})_{n \geq 0} \quad \text{and} \quad y(t) = (y_n e^{i\rho_n t})_{n \geq 0}.$$

We obtain

$$|S| \leq \int_{\mathbb{R}} f(t) \|A\| \|x(t)\| \|y(t)\| dt = \int_{\mathbb{R}} f(t) \|A\| \|x\| \|y\| dt = \|A\| \|x\| \|y\|$$

since $\|f\|_{L^1(\mathbb{R})} = 1$.

Since x and y are arbitrary, this proves $\|D^{-1}AD\| \leq \|A\|$. \square

Proof of Theorem 3.3. First, if $|\varphi'(0)| = 1$, we have $\varphi(z) = \alpha z$ for some α with $|\alpha| = 1$, and the result is trivial.

So, we assume that $|\varphi'(0)| < 1$. Then, by Lemma 3.4, there exists $\rho > 0$ such that, for all m, n :

$$|\widehat{\varphi}^n(m)| \leq \exp\left(-\frac{1}{2}[(1+\rho)n - m]\right).$$

Since $\varphi(0) = 0$, we also know that $\widehat{\varphi}^n(m) = 0$ if $m < n$.

Take $\delta = \rho/2$ and use property (3.2): there exists $M \geq 1$ such that:

$$\frac{\beta_m}{\beta_n} \leq M \quad \text{when } m \geq (1+\delta)n.$$

Define now a new sequence $\gamma = (\gamma_n)$ as:

$$\gamma_n = \max\left\{\beta_n, \sup_{m > (1+\delta)n} \beta_m\right\}.$$

We have:

- 1) $\beta_n \leq \gamma_n \leq M \beta_n$;
- 2) $\gamma_m \leq \gamma_n$ if $m \geq (1+\delta)n$.

Item 1) implies that $H^2(\gamma) = H^2(\beta)$, and we are reduced to prove that $C_\varphi: H^2(\gamma) \rightarrow H^2(\gamma)$ is bounded.

Let $A = (a_{m,n})_{m,n} = (\widehat{\varphi}^n(m))_{m,n}$. We have to prove that

$$B = (\gamma_m^{1/2} \gamma_n^{-1/2} a_{m,n})_{m,n}$$

represents a bounded operator on ℓ_2 .

Define the matrix

$$A_1 = (a_{m,n} \mathbb{1}_{\{(m,n); m \leq (1+\delta)n\}})_{m,n}$$

and set $A_2 = A - A_1$. Define analogously B_1 and $B_2 = B - B_1$.

Then A_1 is a Hilbert-Schmidt operator, because (recall that $a_{m,n} = 0$ if $m < n$), we have

$$\begin{aligned} \sum_{n=1}^{\infty} \sum_{m=1}^{(1+\delta)n} |a_{m,n}|^2 &\leq \sum_{n=1}^{\infty} \sum_{m=n}^{(1+\delta)n} \exp(-[(1+\rho)n - m]) \\ &\leq \sum_{n=1}^{\infty} (\delta n + 1) \exp(-\delta n) < \infty. \end{aligned}$$

Now, β is bounded and has polynomial decay, so, for some positive constants C_1, C_2 , and α , we have:

$$\begin{aligned} \sum_{n=1}^{\infty} \sum_{m=n}^{(1+\delta)n} \frac{\gamma_m}{\gamma_n} |a_{m,n}|^2 &\leq \sum_{n=1}^{\infty} \sum_{m=n}^{(1+\delta)n} \frac{C_1}{n^{-\alpha}} \exp(-\delta n) \\ &\leq \sum_{n=1}^{\infty} C_2 (1+\delta) n^{\alpha+1} \exp(-\delta n) < \infty, \end{aligned}$$

meaning that B_1 is a Hilbert-Schmidt operator.

Since A is bounded, it follows that $A_2 = A - A_1$ is bounded. Remark that, writing $A_2 = (\alpha_{m,n})_{m,n}$, we have, with $d_n = 1/\sqrt{\gamma_n}$:

$$d_m < d_n \implies \gamma_m > \gamma_n \implies m < (1 + \delta)n \implies \alpha_{m,n} = 0.$$

Hence we can apply Lemma 3.6 to the matrix A_2 , and it ensues that B_2 is bounded, and therefore that $B = B_1 + B_2$ is bounded as well, as wanted. \square

As a corollary of Theorem 3.3, we can provide the following example.

Theorem 3.7. *There exists a bounded sequence β , with polynomial decay, but which is not essentially decreasing, and for which every composition operator with symbol vanishing at 0 is bounded on $H^2(\beta)$.*

We hence have $\sup_{\varphi(0)=0} \|C_\varphi\| = +\infty$.

It should be noted that for this weight, the composition operators are not all bounded, as we will see in Proposition 4.11.

Proof. Define $\beta_n = 1$ for $n \leq 3!$, and, for $k \geq 3$:

$$\left\{ \begin{array}{l} \beta_n = \frac{1}{k!} \quad \text{for } k! < n \leq (k+1)! - 2 \text{ and for } n = (k+1)! \\ \beta_n = \frac{1}{(k+1)!} \quad \text{for } n = (k+1)! - 1. \end{array} \right.$$

Note that, for $m > n$, we have $\beta_m > \beta_n$ only if $n = (k+1)! - 1$ and $m = (k+1)! = n + 1$, for some $k \geq 3$.

However β is not essentially decreasing since, for every $k \geq 3$, we have $\beta_{n+1}/\beta_n = k + 1$ if $n = (k+1)! - 1$.

The sequence β has a polynomial decay because $\beta_n \geq 1/(2n)$ for all $n \geq 1$. In fact, for $k \geq 3$, we have $\beta_n \geq (k+1)/n \geq 1/n$ if $k! < n \leq (k+1)! - 2$ or if $n = (k+1)!$; and for $n = (k+1)! - 1$, we have $n\beta_n = [(k+1)! - 1]/(k+1)! \geq 1/2$. It has a polynomial growth since it is bounded above, by 1.

Now, it remains to check (3.2) in order to apply Theorem 3.3 and finish the proof of Theorem 3.7. Note first that we have $\beta_m/\beta_n \leq 1$ if $m \geq n + 2$. Next, for given $\delta > 0$, there exists an integer N such that $(1 + \delta)n \geq n + 2$ for every $n \geq N$, so $\beta_m/\beta_n \leq 1$ if $m \geq (1 + \delta)n$ and $n \geq N$. It suffices to take $C = \max_{1 \leq n \leq N} \beta_{n+1}/\beta_n$ to obtain (3.2). The last assertion follows from Proposition 3.2. \square

4 Boundedness of composition operators of symbol T_a

Recall that for $a \in \mathbb{D}$, we defined

$$(4.1) \quad T_a(z) = \frac{a+z}{1+\bar{a}z}, \quad z \in \mathbb{D}.$$

It is well-known that T_a is an automorphism of \mathbb{D} and that $T_a(0) = a$ and $T_a(-a) = 0$.

Though we do not really need this, we may remark that $(T_a)_{a \in (-1,1)}$ is a group and $(T_a)_{a \in (0,1)}$ is a semigroup. It suffices to see that $T_a \circ T_b = T_{a*b}$, with:

$$(4.2) \quad a * b = \frac{a+b}{1+ab}.$$

In this section, we are going to prove a necessary and sufficient condition in order that all composition operators C_{T_a} for $a \in \mathbb{D}$ are bounded on $H^2(\beta)$. Namely, we have the following theorem, the proof of which will occupy Section 4.2 and Section 4.3.

Theorem 4.1. *All composition operator C_{T_a} , with $a \in \mathbb{D}$ are bounded on $H^2(\beta)$ if and only if β is slowly oscillating.*

Before that, let us note the following fact (see also [22, Prop. 3.6]). Recall that if φ and ψ are two symbols, then $C_\varphi \circ C_\psi = C_{\psi \circ \varphi}$.

Proposition 4.2. *If C_{T_a} is bounded on $H^2(\beta)$ for some $a \in \mathbb{D} \setminus \{0\}$, then C_{T_b} is bounded on $H^2(\beta)$ for all $b \in \mathbb{D}$.*

Moreover the maps C_{T_b} are uniformly bounded on the compact subsets of \mathbb{D} .

We decompose the proof into lemmas. The first one was first proved in [22] (see also [6, Proposition 2.1]), and follows from the fact that if $b = \rho e^{i\theta}$ and R_θ is the rotation $R_\theta(z) = e^{i\theta}z$, which induces a unitary operator C_{R_θ} on $H^2(\beta)$, then $T_b = R_\theta \circ T_\rho \circ R_{-\theta}$ and $C_{T_b} = C_{R_{-\theta}} \circ C_{T_\rho} \circ C_{R_\theta}$.

Lemma 4.3. *The composition operator C_{T_b} is bounded if and only if $C_{T_{|b|}}$ is bounded, with equal norms.*

Lemma 4.4. *Let $r \in (0, 1)$ such that C_{T_r} is bounded. For any $b \in \mathbb{D}$ satisfying $|b| \leq \frac{2r}{1+r^2}$, C_{T_b} is bounded and we have $\|C_{T_b}\| \leq \|C_{T_r}\|^2$.*

Proof. Let S be the circle $C(0, r)$ and $u: S \rightarrow \mathbb{R}_+$ be the continuous function defined by

$$(4.3) \quad u(s) = \left| \frac{s+r}{1+\bar{s}r} \right|.$$

By connectedness, $u(S)$ contains the segment $[0, \frac{2r}{1+r^2}] = [u(-r), u(r)]$. Let now $b \in D(0, \frac{2r}{1+r^2})$. By the above, there exists $s \in S$ such that $|b| = u(s)$. That

means that $|T_b(0)| = |b| = |u(s)| = |T_s(r)| = |(T_s \circ T_r)(0)|$. Therefore, $T_b(0) = e^{i\alpha}(T_s \circ T_r)(0)$ for some $\alpha \in \mathbb{R}$, and hence, by Schwarz's lemma, there is some $\theta \in \mathbb{R}$ such that $T_b = R_\alpha \circ T_s \circ T_r \circ R_\theta$. We then have $C_{T_b} = C_{R_\theta} \circ C_{T_r} \circ C_{T_s} \circ C_{R_\alpha}$. Since C_{R_θ} and C_{R_α} are unitary, we get using Lemma 4.3 for C_{T_s} :

$$\|C_{T_b}\| = \|C_{T_r} \circ C_{T_s}\| \leq \|C_{T_r}\| \|C_{T_s}\| = \|C_{T_r}\|^2. \quad \square$$

Proof of Proposition 4.2. It suffices to use Lemma 4.3 and Lemma 4.4 and do an iteration, in noting that if $r_0 = |a| > 0$ and $r_{n+1} = \frac{2r_n}{1+r_n^2} = r_n * r_n$, then $(r_n)_{n \geq 0}$ increases to 1. \square

4.1 An elementary necessary condition

We begin by an elementary necessary condition. It is implied by Theorem 4.10, but its statement deserves to be pointed out. Moreover, its proof is simple and highlights the role of the reproducing kernel.

Proposition 4.5. *Let $a \in (0, 1)$ and assume that T_a induces a bounded composition operator on $H^2(\beta)$. Then β has polynomial decay.*

Proof. Since

$$\|K_x\|^2 = \sum_{n=0}^{\infty} \frac{x^{2n}}{\beta_n},$$

we have $\|K_x\| \leq \|K_y\|$ for $0 \leq x \leq y < 1$.

We define by induction a sequence $(u_n)_{n \geq 0}$ with:

$$u_0 = 0 \quad \text{and} \quad u_{n+1} = T_a(u_n).$$

Since $T_a(1) = 1$ (recall that $a \in (0, 1)$), we have:

$$1 - u_{n+1} = \int_{u_n}^1 T_a'(t) dt = \int_{u_n}^1 \frac{1 - a^2}{(1 + at)^2} dt;$$

hence

$$\frac{1 - a}{1 + a} (1 - u_n) \leq 1 - u_{n+1} \leq (1 - a^2)(1 - u_n).$$

Let $0 < x < 1$. We can find $N \geq 0$ such that $u_N \leq x < u_{N+1}$. Then:

$$1 - x \leq 1 - u_N \leq (1 - a^2)^N.$$

On the other hand, since $C_{T_a}^* K_z = K_{T_a(z)}$ for all $z \in \mathbb{D}$, we have:

$$\|K_x\| \leq \|K_{u_{N+1}}\| \leq \|C_{T_a}\| \|K_{u_N}\| \leq \|C_{T_a}\|^{N+1} \|K_{u_0}\| = \frac{1}{\sqrt{\beta_0}} \|C_{T_a}\|^{N+1}.$$

Let $s \geq 0$ such that $(1 - a^2)^{-s} = \|C_{T_a}\|$. We obtain:

$$(4.4) \quad \|K_x\| \leq \frac{1}{\sqrt{\beta_0}(1 - x)^s} \|C_{T_a}\|.$$

But

$$\|K_x\|^2 = \sum_{k=0}^{\infty} \frac{x^{2k}}{\beta_k};$$

so we get, for any $k \geq 2$:

$$\frac{x^{2k}}{\beta_k} \leq \frac{1}{\beta_0(1-x)^{2s}} \|C_{T_a}\|^2.$$

Taking $x = 1 - \frac{1}{k}$, we obtain $\beta_k \geq C k^{-2s}$. □

Remarks. 1) For example, when $\beta_n = \exp[-c(\log(n+1))^2]$, with $c > 0$, no T_a induces a bounded composition operator on $H^2(\beta)$, though all symbols φ with $\varphi(0) = 0$ are bounded, since β is decreasing, as we saw in Proposition 3.2.

2) For the Dirichlet space \mathcal{D}^2 , we have $\beta_n = n+1$, but all the maps T_a induce bounded composition operators on \mathcal{D}^2 (see [12, Remark before Theorem 3.12]). In this case β has polynomial growth though it is not bounded above.

3) However, even for decreasing sequences, a polynomial decay for β is not enough for some T_a to induce a bounded composition operator. Indeed, we saw in Proposition 2.5 an example of a decreasing sequence β with polynomial decay but not slowly oscillating, and we will see in Theorem 4.10 that this condition is needed for having some T_a inducing a bounded composition operator.

4) In [6], Eva Gallardo-Gutiérrez and Jonathan Partington give estimates for the norm of C_{T_a} , with $a \in (0, 1)$, when C_{T_a} is bounded on $H^2(\beta)$. More precisely, they proved that if β is bounded above and C_{T_a} is bounded, then

$$\|C_{T_a}\| \geq \left(\frac{1+a}{1-a}\right)^\sigma,$$

where $\sigma = \inf\{s \geq 0; (1-z)^{-s} \notin H^2(\beta)\}$, and

$$\|C_{T_a}\| \leq \left(\frac{1+a}{1-a}\right)^\tau,$$

where $\tau = \frac{1}{2} \sup \Re W(A)$, with A the infinitesimal generator of the continuous semigroup (S_t) defined as $S_t = C_{T_{\tanh t}}$, namely $(Af)(z) = f'(z)(1-z^2)$, and $W(A)$ its numerical range.

For $\beta_n = 1/(n+1)^\nu$ with $0 \leq \nu \leq 1$, the two bounds coincide, so they get $\|C_{T_a}\| = \left(\frac{1+a}{1-a}\right)^{(\nu+1)/2}$.

5) We saw in the proof of Proposition 4.5 that if C_{T_a} is bounded on $H^2(\beta)$ for some $a \in (0, 1)$, then the reproducing kernels K_w have, by (4.4), a *slow growth*:

$$(4.5) \quad \|K_w\| \leq \frac{C}{(1-|w|)^s}$$

for positive constants C and s . Actually, we have the following equivalence.

Proposition 4.6. *The sequence β has polynomial decay if and only if the reproducing kernels K_w of $H^2(\beta)$ have a slow growth.*

Proof. The sufficiency is easy and seen at the end of the proof of Proposition 4.5. For the necessity, we only have to see that:

$$\|K_w\|^2 = \frac{1}{\beta_0} + \sum_{n=1}^{\infty} \frac{|w|^{2n}}{\beta_n} \leq \frac{1}{\beta_0} + \delta^{-1} \sum_{n=1}^{\infty} n^\alpha |w|^{2n} \leq \frac{C}{(1 - |w|^2)^{\alpha+1}}. \quad \square$$

4.2 Sufficient condition

The following sufficient condition explains in particular why all composition operators C_{T_a} are bounded on the Dirichlet space.

Theorem 4.7. *If β is slowly oscillating, then all symbols that extend analytically in a neighborhood of \mathbb{D} induce a bounded composition operator on $H^2(\beta)$. In particular, all C_{T_a} for $a \in \mathbb{D}$ are bounded on $H^2(\beta)$.*

To prove Theorem 4.7, we begin by a very elementary fact.

Lemma 4.8. *Let $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ have an analytic extension to an open neighborhood Ω of \mathbb{D} . Then, there are a constant $b > 0$ and an integer $\lambda > 1$ such that*

$$|\widehat{\varphi^n}(m)| \leq \begin{cases} e^{-bn} & \text{if } n \geq \lambda m, \\ e^{-bm} & \text{if } m \geq \lambda n. \end{cases}$$

Proof. Let $R > 1$ such that $\overline{D(0, R)} \subseteq \Omega$. For $0 < r \leq R$, we set

$$M(r) = \sup_{|z|=r} |\varphi(z)|.$$

Take any $r \in (0, 1)$, for instance $r = e^{-1}$. We have $M(r) < 1$, so we can write $M(r) = e^{-\rho}$, for some positive ρ .

Cauchy's inequalities give:

$$|\widehat{\varphi^n}(m)| \leq \frac{[M(r)]^n}{r^m} = e^{m-\rho n}.$$

Choose $\lambda_1 = \max(2, 2/\rho)$ and $b_1 = \rho - \lambda_1^{-1}$ then $|\widehat{\varphi^n}(m)| \leq e^{-b_1 n}$ if $n \geq \lambda_1 m$.

For the second inequality, write $R =: e^\beta$, with $\beta > 0$. Let $\alpha > 0$ with $M(R) \leq e^\alpha$. Cauchy's inequalities again give:

$$|\widehat{\varphi^n}(m)| \leq \frac{[M(R)]^n}{R^m} \leq e^{\alpha n - \beta m}.$$

Choose $\lambda_2 = \max(2, 2\alpha/\beta)$ and $b_2 = \beta - \alpha\lambda_2^{-1}$ then $|\widehat{\varphi^n}(m)| \leq e^{-b_2 m}$ if $m \geq \lambda_2 n$. We get the conclusion taking $b = \min(b_1, b_2)$ and choosing an integer $\lambda \geq \max(\lambda_1, \lambda_2)$. \square

Lemma 4.9. *Let (β_n) be a slowly oscillating sequence of positive numbers. Let $A = (a_{m,n})_{m,n}$ be the matrix of a bounded operator on ℓ_2 . Assume that, for some integer $\lambda > 1$, and some constants c, b , we have:*

- 1) $|a_{m,n}| \leq ce^{-bn}$ when $n \geq \lambda m$;
- 2) $|a_{m,n}| \leq ce^{-bm}$ when $m \geq \lambda n$.

Then the matrix $\tilde{A} = \left(a_{m,n} \sqrt{\frac{\beta_m}{\beta_n}} \right)_{m,n}$ also defines a bounded operator on ℓ_2 .

Proof. In the sequel $\|\cdot\|$ stands for the ℓ^2 -norm.

Since β is slowly oscillating, it has polynomial decay and growth: for some $\alpha, \gamma > 0$ and $\delta \in (0, 1)$, we have $\delta(n+1)^{-\alpha} \leq \beta_n \leq \delta^{-1}(n+1)^\gamma$.

The matrix \tilde{A} is Hilbert-Schmidt far from the diagonal since

$$\begin{aligned} \sum_{n=1}^{\infty} \sum_{\lambda m < n} |a_{m,n}|^2 \frac{\beta_m}{\beta_n} &\lesssim \sum_{n=1}^{\infty} \sum_{\lambda m < n} (n+1)^{\alpha+\gamma} |a_{m,n}|^2 \\ &\lesssim \sum_{n=1}^{\infty} (n+1)^{\alpha+\gamma+1} e^{-2bn} < +\infty, \end{aligned}$$

and

$$\begin{aligned} \sum_{n=0}^{\infty} \sum_{m > \lambda n} |a_{m,n}|^2 \frac{\beta_m}{\beta_n} &\lesssim \sum_{n=0}^{\infty} \sum_{m > \lambda n} (n+1)^{\alpha+\gamma} |a_{m,n}|^2 \\ &\lesssim \sum_{n=0}^{\infty} (n+1)^{\alpha+\gamma} \left(\sum_{m > \lambda n} e^{-2bm} \right) < +\infty. \end{aligned}$$

Since β_m/β_n remains bounded from above and below around the diagonal, the matrix \tilde{A} behaves like A near the diagonal. More precisely, if I, J are blocks of integers such that $(m, n) \in I \times J$ implies that $n/\lambda^2 \leq m \leq \lambda^2 n$, then, with obvious notations, the slow oscillation of β gives, for some $C > 0$:

$$\begin{aligned} \left| \sum_{(m,n) \in I \times J} a_{m,n} x_n \overline{y_m} \sqrt{\frac{\beta_m}{\beta_n}} \right| &\leq \|A\| \left(\sum_{(m,n) \in I \times J} |x_n|^2 |y_m|^2 \frac{\beta_m}{\beta_n} \right)^{1/2} \\ &\leq C^{1/2} \|A\| \|P_J x\| \|P_I y\|. \end{aligned}$$

For $k = 0, 1, 2, \dots$, let $J_k = [\lambda^k, \lambda^{k+1}[$ and, for $k = 1, 2, \dots$, we define $I_k = [\lambda^{k-1}, \lambda^{k+2}[$. We also define $I_0 = [0, \lambda^2[$.

We define the matrix R , whose entries are

$$r_{m,n} = \begin{cases} \sqrt{\frac{\beta_m}{\beta_n}} a_{m,n} & \text{if } (m, n) \in \bigcup_{k=0}^{\infty} (I_k \times J_k) \\ 0 & \text{elsewhere.} \end{cases}$$

Let H_k be the subspace of the sequences $(x_n)_{n \geq 0}$ in ℓ_2 such that $x_n = 0$ for $n \notin I_k$, i.e. $H_k = \text{span}\{e_n; n \in I_k\}$, and $\tilde{H}_k = \text{span}\{e_n; n \in J_k\}$. Let P_k be (the matrix of) the orthogonal projection of ℓ_2 with range H_k and Q_k that with range \tilde{H}_k . Then $R_k = P_k A Q_k$ is the matrix with entries $a_{m,n}$ when $(m,n) \in I_k \times J_k$ and 0 elsewhere. By the above discussion, we have

$$|(R_k x | y)| \leq C^{1/2} \|A\| \|Q_k x\| \|P_k y\|.$$

Point out that, for every $y \in \ell^2$, we have $\sum \|P_k y\|^2 \leq 3\|y\|^2$ since each integer belongs to at most 3 intervals I_k .

In the same way, for every $x \in \ell^2$, we have $\sum \|Q_k x\|^2 \leq \|x\|^2$ since the subspaces \tilde{H}_k are orthogonal.

Summing up over k , we get the boundedness of $R = \sum_{k=0}^{\infty} R_k$.

Now let us check when the entries of R do not coincide with the entries of \tilde{A} . Actually it happens when (m,n) does not belong to the union of the $I_k \times J_k$. When $n \geq 1$, it means that n belongs to some J_p but $m \notin I_p$: either $m < \lambda^{p-1}$ or $m \geq \lambda^{p+2}$, hence either $m/n < \lambda^{-1}$ or $m/n > \lambda$. Therefore the non zero entries (m,n) of $\tilde{A} - R$ satisfy either $n > \lambda m$ or $m > \lambda n$.

That ends the proof since we have seen at the beginning that $\tilde{A} - R$ is Hilbert-Schmidt. \square

Remark. The proof shows that, instead of 1) and 2), it is enough to have:

$$\sum_{m < C_1 n} n^{\alpha+1} |a_{m,n}|^2 < \infty \quad \text{and} \quad \sum_{m > C_2 n} m^{\alpha} |a_{m,n}|^2 < \infty.$$

Moreover the proof also shows that when β is slowly oscillating, if we set $E = \{(m,n); C_1 n \leq m \leq C_2 n\}$, for some $C_1, C_2 > 0$, then the matrix $(\sqrt{\beta_m/\beta_n} \mathbf{1}_E(m,n))$ is a Schur multiplier over *all* the bounded matrices, while Kacnel'son's theorem (Theorem 3.5) says that, if $\gamma = (\gamma_n)$ is non-increasing, the matrix (γ_m/γ_n) is a Schur multiplier of all bounded *lower-triangular* matrices.

Proof of Theorem 4.7. Thanks to Lemma 4.8, the hypothesis of Lemma 4.9 are fulfilled by the matrix whose entries are $a_{m,n} = \tilde{\varphi}^n(m)$. It follows (with the notations of Lemma 4.9) that \tilde{A} is bounded on ℓ^2 , which means exactly that T_a is bounded on $H^2(\beta)$. \square

4.3 Necessary condition

The main theorem of this section is the following.

Theorem 4.10. *If the composition operator C_{T_a} is bounded on $H^2(\beta)$ for some $a \in \mathbb{D} \setminus \{0\}$, then β is slowly oscillating.*

Let us give right now a corollary of this result.

Proposition 4.11. *For the weight β constructed in the proof of Theorem 3.7, no automorphism T_a with $0 < a < 1$ can be bounded.*

Proof. Indeed, it is clear that β is not slowly oscillating, since

$$\beta_{(k+1)!-1}/\beta_{(k+1)!} = 1/(k+1) \xrightarrow[k \rightarrow \infty]{} 0. \quad \square$$

To prove Theorem 4.10, we need estimates on the Taylor coefficients of T_a^n . Actually the Taylor coefficients of T_a^n are the Fourier coefficients of $x \in \mathbb{R} \mapsto T_a^n(e^{ix})$ and we shall denote them with the same notation $\widehat{T_a^n}$. Sharp such estimates are given in the papers [19] and [20], and we thank R. Zarouf for interesting informations in this respect. Our method, using stationary phase and the van der Corput lemma, is a variant of that used in [19], [20], and goes back at least to [7]. However, we need minorations of $|\widehat{T_a^n}(m)|$ when m is close to n , and Szehr and Zarouf's estimates show that this quantity oscillates and, for individual a , can be too small for our purpose, so we cannot use them, and have to prove an estimate *in mean* for a in some subinterval of $(0, 1)$.

We begin with a standard fact, that we give with its proof for the convenience of the reader.

Lemma 4.12. *Let $a \in (0, 1)$ and let*

$$P_{-a}(x) = \frac{1 - a^2}{1 + 2a \cos x + a^2}$$

be the Poisson kernel at the point $-a$. Then, for all $x \in [-\pi, \pi]$:

$$(4.6) \quad T_a(e^{ix}) = \exp[iV_a(x)],$$

where

$$(4.7) \quad V_a(x) = \int_0^x P_{-a}(t) dt.$$

Proof. For $t \in [-\pi, \pi]$, write:

$$\psi(t) := \frac{e^{it} + a}{1 + a e^{it}} = \exp(i v(t)),$$

with v a real-valued, C^1 -function on $[-\pi, \pi]$ such that $v(0) = 0$. This is possible since $|\psi(e^{it})| = 1$ and $\psi(0) = 1$. Differentiating both sides with respect to t , we get:

$$i e^{it} \frac{1 - a^2}{(1 + a e^{it})^2} = i v'(t) \frac{e^{it} + a}{1 + a e^{it}}.$$

This implies

$$v'(t) = \frac{1 - a^2}{|1 + a e^{it}|^2} = P_{-a}(t),$$

and the result follows since $v(0) = 0 = V_a(0)$. \square

Let us note that, with V_a the function of Lemma 4.12, the Fourier formulas give, since $\widehat{T}_a^n(m)$ is real, or since $nV_a(x) - mx$ is odd:

$$(4.8) \quad 2\pi \widehat{T}_a^n(m) = \int_{-\pi}^{\pi} \exp(i[nV_a(x) - mx]) dx = 2\Re I_{m,n},$$

where

$$(4.9) \quad I_{m,n} = \int_0^{\pi} \exp i[nV_a(x) - mx] dx.$$

Now the main ingredient for proving Theorem 4.10 is the following.

Proposition 4.13. *Let $I := [\frac{1}{2}, \frac{2}{3}]$. There exist constants $\alpha > 1$, e.g. $\alpha = 5/4$, and $\delta \in (0, 1/2)$ such that, for n large enough ($n \geq n_0$), it holds*

$$(4.10) \quad \int_I |\widehat{T}_a^n(m)|^2 da \geq \frac{\delta}{n} \quad \text{for all } m \in [\alpha^{-1}n, \alpha n].$$

Proof. We will set once and for all

$$(4.11) \quad q = \frac{m}{n},$$

so that $\alpha^{-1} \leq q \leq \alpha$ where $\alpha = 5/4$ (say). We will only consider pairs (a, q) satisfying

$$(4.12) \quad a \in I = \left[\frac{1}{2}, \frac{2}{3}\right], \quad q \in J := \left[\frac{4}{5}, \frac{5}{4}\right].$$

Such pairs will be called *admissible*.

With these notations, we set, for $0 \leq x \leq \pi$:

$$(4.13) \quad F_q(x) = V_a(x) - \frac{m}{n}x = \int_0^x P_{-a}(t) dt - qx,$$

where P_{-a} is the Poisson kernel at $-a$. We have

$$F_q'(x) = \frac{(1 - a^2)}{1 + 2a \cos x + a^2} - q.$$

and the unique (if some) critical point $x_q = x_q(a)$ of F_q in $[0, \pi]$ is given by $P_{-a}(x_q) = q$, that is

$$(4.14) \quad \cos x_q = \frac{1}{q} \frac{1 - a^2}{2a} - \frac{1 + a^2}{2a} =: h_q(a).$$

We now proceed through a series of simple lemmas, and begin by estimates on h_q and x_q .

Lemma 4.14. *There are positive constants $C > 1$ and $\delta \in (0, 1/2)$ such that, for every admissible couple (a, q) , we have:*

$$(4.15) \quad |h_q(a)| \leq 1 - \delta \quad \text{and} \quad |h'_q(a)| \leq C,$$

so, there is one critical point $x_q(a)$ satisfying

$$(4.16) \quad \delta \leq x_q(a) \leq \pi - \delta \quad \text{and} \quad \sin x_q(a) \geq \delta;$$

moreover

$$(4.17) \quad |x'_q(a)| \leq C \quad \text{and} \quad \delta \leq |P'_{-a}(x_q)| \leq C.$$

Proof. We have

$$h_q(a) = \left(\frac{1}{q} \frac{1 - a^2}{2a} \right) + \left(- \frac{1 + a^2}{2a} \right) =: u(a) + v(a),$$

with u and v respectively decreasing and increasing on $[0, 1]$, and with $v \leq 0$, so that we have, for $q \in J$:

$$h_q(a) \leq u\left(\frac{1}{2}\right) = \frac{3}{4q} \leq \frac{15}{16}.$$

Similarly:

$$h_q(a) \geq u\left(\frac{2}{3}\right) + v\left(\frac{1}{2}\right) = \frac{5}{12q} - \frac{5}{4} \geq \frac{1}{3} - \frac{5}{4} = -\frac{11}{12}.$$

Next, $2h'_q(a) = \left(1 - \frac{1}{q}\right) \frac{1}{a^2} - \left(1 + \frac{1}{q}\right)$; hence $|h'_q(a)| \leq C$. So that, writing $x_q = x_q(a) = \arccos h_q(a)$, we get, with another constant $C > 0$,

$$|x'_q(a)| = \frac{|h'_q(a)|}{\sqrt{1 - h_q(a)^2}} \leq C,$$

since $h_q(a)^2 \leq 1 - \delta$. Finally, $1/9 \leq (1 - a)^2 \leq 1 + 2a \cos x_q + a^2 \leq 4$ and since

$$P'_{-a}(x_q) = \frac{2a(1 - a^2) \sin x_q}{(1 + 2a \cos x_q + a^2)^2},$$

we get the final estimates, ending the proof. \square

Back to Proposition 4.13.

We saw in (4.8) that the value of $a_{m,n} := \widehat{T}_a^n(m)$ is given by the formula

$$(4.18) \quad a_{m,n} = \frac{1}{\pi} \Re I_{m,n}.$$

We have the following estimate, whose proof is postponed (recall that $q = m/n$, and $x_q = x_q(a)$).

Proposition 4.15. *We have*

$$(4.19) \quad I_{m,n} = \sqrt{2\pi} n^{-1/2} \frac{e^{i[nF_q(x_q) + \pi/4]}}{\sqrt{|P'_{-a}(x_q)|}} + O(n^{-3/5}),$$

where the O only depends on a , and so is absolute as long as (a, q) is admissible.

Note that $3/5 > 1/2$. We hence have:

$$(4.20) \quad a_{m,n} = \sqrt{\frac{2}{\pi}} n^{-1/2} \frac{\cos[\pi/4 + nF_q(x_q)]}{\sqrt{|P'_{-a}(x_q)|}} + O(n^{-3/5}),$$

It will be convenient to introduce $\varphi_q(a)$, by setting

$$(4.21) \quad F_q(x_q(a)) = \varphi_q(a).$$

Then, since $1/2 + 3/5 = 11/10$, $\cos^2(\pi/4 + x) = \frac{1 - \sin 2x}{2}$ and $|P'_{-a}(x_q)| \geq \delta$ by Lemma 4.14:

$$a_{m,n}^2 = \frac{1}{\pi} n^{-1} \frac{1 - \sin[2n\varphi_q(a)]}{|P'_{-a}(x_q)|} + O(n^{-11/10}),$$

implying, since $|P'_{-a}(x_q)| \leq C$ by Lemma 4.14 (again for (a, q) admissible), and changing δ :

$$(4.22) \quad a_{m,n}^2 \geq \delta n^{-1} (1 - \sin[2n\varphi_q(a)]) + O(n^{-11/10}).$$

We will also need estimates on the derivatives of $\varphi_q(a)$.

Lemma 4.16. *If (a, q) is admissible, then φ_q decreases on I and moreover:*

- 1) $|\varphi'_q(a)| \geq \delta$;
- 2) $|\varphi''_q(a)| \leq C$.

Proof. Note, in passing, that, with $x = x_q(a) \in [0, \pi]$ (thanks to (4.13)):

$$\varphi_q(a) = \int_0^x [P_{-a}(t) - P_{-a}(x)] dt \leq 0,$$

since the integrand is negative. Next, if f and g are real \mathcal{C}^1 -functions and

$$\Phi(a) = \int_0^{f(a)} g(a, t) dt,$$

the chain rule gives

$$\Phi'(a) = f'(a) g(a, f(a)) + \int_0^{f(a)} \frac{\partial g}{\partial a}(a, t) dt.$$

With $g(a, t) = P_{-a}(t)$ and $f(a) = x_q(a)$, we get, remembering that $x_q(a)$ is critical for F_q :

$$\begin{aligned}\varphi'_q(a) &= [P_{-a}(x_q(a)) - q] x'_q(a) + \int_0^{x_q(a)} \frac{\partial P_{-a}}{\partial a}(a, t) dt \\ &= \int_0^{x_q(a)} \frac{\partial P_{-a}}{\partial a}(a, t) dt.\end{aligned}$$

But $P_{-a}(t) = 1 + 2 \sum_{k=1}^{\infty} (-a)^k \cos kt$, so that

$$\varphi'_q(a) = \int_0^{x_q(a)} \left(-2 \sum_{k=1}^{\infty} k (-a)^{k-1} \cos kt \right) dt = \frac{2}{a} \sum_{k=0}^{\infty} (-a)^k \sin [kx_q(a)],$$

that is

$$(4.23) \quad \varphi'_q(a) = \frac{2}{a} \Im \frac{1}{1 + a e^{ix_q(a)}} = \frac{-2 \sin x_q(a)}{1 + 2a \cos x_q(a) + a^2} < 0.$$

Now, (4.16) gives 1).

Since $|x'_q(a)| \leq C$ by Lemma 4.14, the chain rule and (4.23) clearly give the uniform boundedness of $|\varphi''_q(a)|$ when (a, q) is admissible, and this ends the proof. \square

Lemma 4.14 and Lemma 4.16 will now be exploited through a simple variant of the van der Corput inequalities.

Lemma 4.17. *Let $f: [A, B] \rightarrow \mathbb{R}$, with $A < B$, be a C^2 -function satisfying $|f'| \geq \delta$ and $|f''| \leq C$, and let us put $M = \int_A^B e^{inf(x)} dx$. Then*

$$|M| \leq \frac{2}{n\delta} + \frac{C(B-A)}{n\delta^2}.$$

Proof. Write

$$e^{inf} = \frac{(e^{inf})'}{inf'}$$

and integrate by parts to get

$$M = \left[\frac{e^{inf}}{inf'} \right]_A^B - \frac{i}{n} \int_A^B e^{inf(x)} \frac{f''(x)}{[f'(x)]^2} dx =: M_1 + M_2,$$

with $|M_1| \leq \frac{2}{n\delta}$ and $|M_2| \leq \frac{B-A}{n} \frac{C}{\delta^2}$. This gives the result. \square

End of proof of Proposition 4.13. The preceding lemma can be applied with $A = 1/2$, $B = 2/3$, $f = \varphi_q$ and n changed into $2n$, since Lemma 4.16 shows that this f meets the assumptions of Lemma 4.17. This gives us, uniformly with respect to (a, q) admissible:

$$(4.24) \quad \left| \int_I \sin [2n\varphi_q(a)] da \right| \leq \left| \int_I e^{2in\varphi_q(a)} da \right| \leq \frac{C}{n}.$$

Now, integrating (4.22) on I and using (4.24) give for some numerical $\delta \in (0, 1/2)$ (recall that $a_{m,n} = \widehat{T}_a^n(m)$):

$$\int_I |\widehat{T}_a^n(m)|^2 da \geq \delta n^{-1} + O(n^{-2}) + O(n^{-11/10}) \geq (\delta/2) n^{-1}$$

for $n \geq n_0$ and $\alpha^{-1} \leq m/n \leq \alpha$. This ends the proof of Proposition 4.13. \square

Proof of Theorem 4.10. We know, by Proposition 4.2, that C_{T_a} is bounded for all $a \in \mathbb{D}$, and that, thanks to Lemma 4.4,

$$K := \sup_{1/2 \leq a \leq 2/3} \|C_{T_a}\| < +\infty.$$

Matricially, this writes, for all $a \in (1/2, 2/3)$:

$$\left\| \left(\widehat{T}_a^n(m) \sqrt{\frac{\beta_m}{\beta_n}} \right)_{m,n} \right\| \leq K.$$

In particular, for every $n \geq 1$, we have, considering the columns and rows of the previous matrix:

$$\sum_{m=1}^{\infty} |\widehat{T}_a^n(m)|^2 \frac{\beta_m}{\beta_n} \leq K^2, \quad \text{i. e.} \quad \sum_{m=1}^{\infty} |\widehat{T}_a^n(m)|^2 \beta_m \leq K^2 \beta_n,$$

and, for every $m \geq 1$:

$$\sum_{n=1}^{\infty} |\widehat{T}_a^n(m)|^2 \frac{\beta_m}{\beta_n} \leq K^2, \quad \text{i. e.} \quad \sum_{n=1}^{\infty} |\widehat{T}_a^n(m)|^2 \frac{1}{\beta_n} \leq \frac{K^2}{\beta_m}.$$

In particular, for every $n \geq 1$:

$$(4.25) \quad \sum_{(4/5)n \leq j \leq (5/4)n} |\widehat{T}_a^n(j)|^2 \beta_j \leq K^2 \beta_n.$$

and, for every $m \geq 1$:

$$(4.26) \quad \sum_{(4/5)m \leq k \leq (5/4)m} |\widehat{T}_a^k(m)|^2 \frac{1}{\beta_k} \leq \frac{K^2}{\beta_m}.$$

Integrating on $a \in (1/2, 2/3)$, and using Proposition 4.13, we get, from (4.25), for n large enough,

$$(4.27) \quad \frac{\delta}{n} \sum_{(4/5)n \leq j \leq (5/4)n} \beta_j \leq \frac{K^2}{6} \beta_n$$

and, from (4.26), for m large enough, we have both

$$(4.28) \quad \frac{\delta}{m} \sum_{(4/5)m \leq k \leq m} \frac{1}{\beta_k} \leq \frac{5K^2}{24} \frac{1}{\beta_m}$$

and

$$(4.29) \quad \frac{\delta}{m} \sum_{m \leq k \leq (5/4)m} \frac{1}{\beta_k} \leq \frac{5K^2}{24} \frac{1}{\beta_m}$$

Since the harmonic mean (over the sets of integers $[4m/5, m]$ and $[m, 5m/4]$, which have cardinality $\approx n \approx m$) is less than the arithmetical mean, we obtain from (4.28) and (4.29) both

$$(4.30) \quad \beta_m \leq \frac{125}{24\delta} \frac{K^2}{m} \sum_{(4/5)m \leq k \leq m} \beta_k$$

and

$$(4.31) \quad \beta_m \leq \frac{10}{3\delta} \frac{K^2}{m} \sum_{m \leq k \leq (5/4)m} \beta_k$$

Now assume that $n \leq m \leq (5/4)n$. From (4.30), we have

$$\beta_m \lesssim \frac{1}{m} \sum_{(4/5)m \leq k \leq m} \beta_k \lesssim \frac{1}{n} \sum_{(4/5)n \leq k \leq (5/4)n} \beta_k \lesssim \beta_n$$

thanks to (4.27).

From (4.31) and (4.27), we treat the case $(4/5)n \leq m \leq n$ in the same way. We conclude that for some constant $c > 0$, we have for n and m large enough satisfying $(4/5)n \leq m \leq (5/4)n$:

$$(4.32) \quad \beta_m \leq c \beta_n,$$

which means that β is slowly oscillating. \square

Proof of Proposition 4.15. We will use a variant of [21, Lemma 4.6 p. 72] on the van der Corput's version of the stationary phase method. A careful reading of the proof in [21, p. 72] gives the version below, which only needs local estimates on the second derivative F'' , as occurs in our situation. For sake of completeness, we will give a proof, however postponed in the Appendix.

Proposition 4.18 (Stationary phase). *Let F be a real function with continuous derivatives up to the third order on the interval $[A, B]$ and $F'' > 0$ throughout $]A, B[$. Assume that there is a (unique) point c in $]A, B[$ such that $F'(c) = 0$, and that, for some positive numbers λ_2, λ_3 , and η , the following assertions hold:*

- 1) $[c - \eta, c + \eta] \subseteq [A, B]$;
- 2) $F''(x) \geq \lambda_2$ for all $x \in [c - \eta, c + \eta]$;
- 3) $|F'''(x)| \leq \lambda_3$ for all $x \in [A, B]$.

Then:

$$(4.33) \quad \int_A^B e^{iF(x)} dx = \sqrt{2\pi} \frac{e^{i[F(c)+\pi/4]}}{F''(c)^{1/2}} + O\left(\frac{1}{\eta\lambda_2} + \eta^4\lambda_3\right),$$

where the O involves an absolute constant.

We will show that Proposition 4.18 is applicable with $F = nF_q$ and

$$[A, B] = [0, \pi], \quad c = x_q, \quad \lambda_2 = \kappa_0 n, \quad \lambda_3 = C_0 n, \quad \eta = (\lambda_2 \lambda_3)^{-1/5}.$$

The parameter η is chosen in order to make both error terms in Proposition 4.18 equal: $\frac{1}{\eta\lambda_2} = \eta^4\lambda_3$; so:

$$\eta = \kappa n^{-2/5}$$

and

$$(4.34) \quad \frac{1}{\eta\lambda_2} + \eta^4\lambda_3 = \tilde{\kappa} n^{-3/5} = O(n^{-3/5})$$

(with $\kappa = (\kappa_0 C_0)^{-1/5}$ and $\tilde{\kappa} = 2/\kappa_0 \kappa$).

The slight technical difficulty encountered here is that $F_q''(x)$ vanishes at 0 and π . But Proposition 4.18 covers this case. We have

$$F''(x) = nF_q''(x) = nP'_{-a}(x) = 2a(1 - a^2) \frac{\sin x}{(1 + 2a \cos x + a^2)^2} n,$$

and there are some positive (and absolute) constants κ_0 and σ such that

$$(4.35) \quad F''(x) \geq \kappa_0 n = \lambda_2 \quad \text{for } x \in [\sigma, \pi - \sigma].$$

Now (for n large enough), $[x_q - \eta, x_q + \eta] \subseteq [\sigma, \pi - \sigma]$. Hence the assumptions 1) and 2) of Proposition 4.18 are satisfied.

Finally, since $F(x) = nF_q(x) = n[V_a(x) - qx]$, $F''' = nF_q''' = nV_a''' = nP_{-a}''$, we have, for all $x \in [0, \pi]$ and (a, q) admissible:

$$|F'''(x)| \leq C_0 n = \lambda_3,$$

where C_0 is absolute and assertion 3) of Proposition 4.18 holds.

With (4.34) this ends the proof of (4.19), once we remarked that $nV_a'''(x_q) = F'''(x_q)$. \square

5 Boundedness of all composition operators

In this section, we characterize all the sequences β for which all composition operators are bounded on $H^2(\beta)$. The main remaining step is the following theorem.

Theorem 5.1. *Assume that all composition operators C_φ are bounded on $H^2(\beta)$. Then β is essentially decreasing.*

As an immediate consequence, we obtain Theorem 1.2.

Proof of Theorem 1.2. Assume that β is essentially decreasing and slowly oscillating. All composition operators C_ψ with $\psi(0) = 0$ are bounded on $H^2(\beta)$ (see the Introduction or Proposition 3.2). Since β is slowly oscillating, all the composition operators C_{T_a} , with $a \in \mathbb{D}$, are bounded thanks to Theorem 4.7. Now it is very classical that we can get the boundedness of every composition operators. Indeed given a symbol φ , the symbol $\psi = T_a \circ \varphi$ fixes the origin for $a = -\varphi(0)$. Since $C_\varphi = C_\psi \circ C_{T_{-a}}$, the conclusion follows.

Assume that all composition operators are bounded on $H^2(\beta)$, in particular, the C_{T_a} ones are so, and β is slowly oscillating, thanks to Theorem 4.10. It also follows from Theorem 5.1 that β is essentially decreasing. \square

We will use the following elementary, but crucial, lemma.

Lemma 5.2. *Let u be a function analytic in an open neighborhood Ω of $\overline{\mathbb{D}}$. Then, for every $\varepsilon > 0$, there exists an integer $N \geq 1$ such that*

$$(5.1) \quad \sum_{j=Np}^{\infty} |\widehat{u^p}(j)|^2 \leq \varepsilon, \quad \forall p \geq 1.$$

Proof. From Lemma 4.8, we know that there exist some integer $\lambda > 1$ and a constant $b > 0$, such that $|\widehat{u^p}(j)| \leq e^{-bj}$ when $j \geq \lambda p$. Therefore, for any $N \geq \lambda$, we have

$$\sum_{j=Np}^{\infty} |\widehat{u^p}(j)|^2 \leq (1 - e^{-2b})^{-1} e^{-2bNp} \leq (1 - e^{-2b})^{-1} e^{-2bN} \leq \varepsilon,$$

as soon as N is chosen large enough. \square

Proof of Theorem 5.1. Thanks to Theorem 4.10, we know that β is slowly oscillating. Therefore we shall assume that the sequence β is slowly oscillating but is not essentially decreasing. We are going to construct an analytic function $\varphi: \mathbb{D} \rightarrow \mathbb{D}$ such that the composition operator C_φ is not bounded on $H^2(\beta)$. This function φ will be a Blaschke product, of the form

$$\varphi(z) = \prod_{k=1}^{\infty} T_{a_k}(z^{n_k}) = \prod_{k=1}^{\infty} \frac{z^{n_k} + a_k}{1 + a_k z^{n_k}},$$

for a sequence of numbers $a_k \in (0, 1)$ such that $\sum_{k \geq 1} (1 - a_k) < \infty$ and a sequence of positive integers n_k increasing to infinity. We recall that $T_a(z) = \frac{z+a}{1+az}$ for $0 < a < 1$. Then,

$$|T_{a_k}(z^{n_k}) - 1| \leq \frac{2(1 - a_k)}{1 - |z|}$$

and φ is a locally uniformly convergent infinite product. Observe that φ is indeed a convergent Blaschke product, with n_k zeroes of modulus a_k^{1/n_k} , $k = 1, 2, \dots$, and, setting $a_k = e^{-\varepsilon_k}$, $\sum_k n_k(1 - a_k^{1/n_k}) \leq \sum_k n_k(\varepsilon_k/n_k) = \sum_k \varepsilon_k < \infty$.

These sequences will be constructed by induction, together with another sequence of integers $(m_k)_{k \geq 1}$.

Since β is not essentially decreasing, there exist integers $n_1 > m_1 \geq 4$ such that $\beta_{n_1} \geq 2\beta_{m_1}$. We start with

$$a_1 = 1 - \frac{1}{m_1} \geq \frac{3}{4}.$$

Using Lemma 5.2 with $u = T_{a_1}$, we get $N_0 \geq 1$ such that

$$\sum_{j=N_0 m}^{\infty} |\widehat{T_{a_1}^m}(j)|^2 \leq 2^{-15}, \quad \forall m \geq 1.$$

Assume now that we have constructed increasing sequences of integers

$$m_1, m_2, \dots, m_k, \quad n_1, n_2, \dots, n_k, \quad N_0, N_1, \dots, N_{k-1}$$

such that, for $1 \leq l \leq k-1$, we have

$$m_{l+1} \geq 4m_l \quad \text{and} \quad n_{l+1} \geq 4n_l$$

and, for $1 \leq l \leq k$:

$$n_l \geq N_{l-1} m_l \quad \text{and} \quad \beta_{n_l} \geq 2^l \beta_{m_l}$$

and

$$\sum_{j=N_{l-1} m_l}^{\infty} |\widehat{\varphi_l^m}(j)|^2 \leq 2^{-15},$$

where

$$a_l = 1 - \frac{1}{m_l} \quad \text{and} \quad \varphi_l(z) = T_{a_l}(z^{n_l}).$$

We then apply Lemma 5.2 again, to the function $u = u_k = \varphi_1 \cdots \varphi_k$. We get $N_k > N_{k-1}$ such that

$$(5.2) \quad \sum_{j=N_k m}^{\infty} |\widehat{u_k^m}(j)|^2 \leq 2^{-15}, \quad \forall m \geq 1.$$

Since β is not essentially decreasing but is slowly oscillating, there exist $m_{k+1} \geq 4m_k$ and $n_{k+1} \geq 4n_k$ such that

$$n_{k+1} \geq N_k m_{k+1} \quad \text{and} \quad \beta_{n_{k+1}} \geq 2^{k+1} \beta_{m_{k+1}}.$$

We set

$$a_{k+1} = 1 - \frac{1}{m_{k+1}} \quad \text{and} \quad \varphi_{k+1}(z) = T_{a_{k+1}}(z^{n_{k+1}}).$$

That ends the induction.

Now, since $\sum_{k=1}^{\infty} (1-a_k) = \sum_{k=1}^{\infty} \frac{1}{m_k} \leq \sum_{k=1}^{\infty} 4^{-k} = 1/3 < +\infty$, the infinite product $\varphi = \prod_{k \geq 1} \varphi_k$ converges uniformly on compact subsets of \mathbb{D} .

To show that the composition operator C_φ is not bounded on $H^2(\beta)$, it suffices to show that, for some constant $c_1 > 0$, we have, for all $k \geq 2$:

$$(5.3) \quad \sum_{j=n_k}^{2n_k} |\widehat{\varphi^{m_k}}(j)|^2 \geq c_1.$$

Indeed, since β is slowly oscillating, there is a positive constant $\delta < 1$ such that

$$\beta_j \geq \delta \beta_{n_k} \quad \text{for} \quad j = n_k, n_k + 1, \dots, 2n_k.$$

Then, if we set $e_k(z) = z^{m_k}$, we have, since $C_\varphi(e_k) = \varphi^{m_k}$:

$$\frac{\|C_\varphi(e_k)\|_{H^2(\beta)}^2}{\|e_k\|_{H^2(\beta)}^2} \geq \frac{\sum_{j=n_k}^{2n_k} |\widehat{\varphi^{m_k}}(j)|^2 \beta_j}{\beta_{m_k}} \geq \frac{c_1 \delta \beta_{n_k}}{\beta_{m_k}} \geq 2^k c_1 \delta \xrightarrow[k \rightarrow \infty]{} +\infty,$$

so that C_φ is not bounded on $H^2(\beta)$.

We now have to show (5.3). Let us agree to write formally, for an analytic function $f(z) = \sum_{k=0}^{\infty} f_k z^k$ and an arbitrary positive integer p :

$$f(z) = \sum_{k=0}^p f_k z^k + O(z^{p+1}).$$

For that, we set

$$G_k(z) = \prod_{l=k+1}^{\infty} \varphi_l(z) = \prod_{l=k+1}^{\infty} a_l + O(z^{n_{k+1}}).$$

We have, for $k \geq 2$:

$$\varphi(z) = v_k(z) \varphi_k(z) G_k(z),$$

where $v_k = \varphi_1 \cdots \varphi_{k-1}$.

Remark now that, for $0 < a < 1$, we have

$$T_a(z) = a + (1-a^2)z + O(z^2),$$

so

$$\varphi_k(z) = T_{a_k}(z^{n_k}) = a_k + (1 - a_k^2)z^{n_k} + O(z^{2n_k}).$$

Then

$$(5.4) \quad [G_k(z)]^{m_k} = \left(\prod_{l=k+1}^{\infty} a_l \right)^{m_k} + O(z^{n_{k+1}})$$

and

$$(5.5) \quad [\varphi_k(z)]^{m_k} = a_k^{m_k} + (1 - a_k^2)m_k a_k^{m_k-1} z^{n_k} + O(z^{2n_k}).$$

But

$$(5.6) \quad a_k^{m_k-1} = \left(1 - \frac{1}{m_k}\right)^{m_k-1} \geq e^{-1} := c_2$$

and

$$(5.7) \quad (1 - a_k^2)m_k a_k^{m_k-1} \geq (1 - a_k)m_k a_k^{m_k-1} \geq c_2.$$

Moreover, since $1 - x \geq e^{-2x}$ for $0 \leq x \leq 1/2$, we have:

$$(5.8) \quad \begin{aligned} \left(\prod_{l=k+1}^{\infty} a_l \right)^{m_k} &\geq \exp\left(-2\left(\sum_{l=k+1}^{\infty} \frac{1}{m_l}\right)m_k\right) \\ &\geq \exp\left(-2\sum_{l=1}^{\infty} 4^{-l}\right) = \exp(-2/3) := c_3. \end{aligned}$$

Afterwards, by (5.2), we have

$$(5.9) \quad \sum_{j=N_{k-1}m_k}^{\infty} |\widehat{v_k^{m_k}}(j)|^2 \leq 2^{-15}.$$

Set $v_k^{m_k} = g_1 + g_2$, with

$$\begin{cases} g_1(z) = \sum_{j=0}^{N_{k-1}m_k} \widehat{v_k^{m_k}}(j) z^j \\ g_2(z) = \sum_{j>N_{k-1}m_k} \widehat{v_k^{m_k}}(j) z^j. \end{cases}$$

By (5.9), we have, with $\|\cdot\|_2 = \|\cdot\|_{L^2(\mathbb{T})}$:

$$\|g_2\|_2^2 = \sum_{j>N_{k-1}m_k} |\widehat{v_k^{m_k}}(j)|^2 \leq 2^{-15}.$$

Besides, since φ_k is inner as a product of inner functions, we have $|v_k(z)| = 1$ for all $z \in \mathbb{T}$, so

$$\|g_1\|_2^2 = \|v_k\|_2^2 - \|g_2\|_2^2 \geq 1 - 2^{-15}.$$

Now, $\varphi^{m_k} = v_k^{m_k} \varphi_k^{m_k} G_k^{m_k} = F_1 + F_2$, with

$$F_1 = g_1 \varphi_k^{m_k} G_k^{m_k} \quad \text{and} \quad F_2 = g_2 \varphi_k^{m_k} G_k^{m_k}.$$

Using (5.4), (5.5), (5.7) and (5.8), we get

$$\begin{aligned} \sum_{j=n_k}^{n_k+N_{k-1}m_k} |\widehat{F_1}(j)|^2 &= \left(\prod_{l=k+1}^{\infty} a_l \right)^{2m_k} [(1 - a_k^2) m_k a_k^{m_k-1}]^2 \sum_{j=0}^{N_{k-1}m_k} |\widehat{g_1}(j)|^2 \\ &\geq (1 - 2^{-15}) c_2^2 c_3^2. \end{aligned}$$

As

$$\|F_2\|_2^2 \leq \|g_2\|_2^2 \|\varphi_k^{m_k}\|_{\infty}^2 \|G_k^{m_k}\|_{\infty}^2 \leq 2^{-15},$$

we get, using the inequality $|a+b|^2 \geq \frac{1}{2}|a|^2 - |b|^2$:

$$\begin{aligned} \sum_{j=n_k}^{2n_k} |\widehat{\varphi^{m_k}}(j)|^2 &\geq \sum_{j=n_k}^{n_k+N_{k-1}m_k} |\widehat{F_1}(j) + \widehat{F_2}(j)|^2 \\ &\geq \frac{1}{2} \sum_{j=n_k}^{n_k+N_{k-1}m_k} |\widehat{F_1}(j)|^2 - \sum_{j=n_k}^{n_k+N_{k-1}m_k} |\widehat{F_2}(j)|^2 \\ &\geq \frac{1}{2} (1 - 2^{-15}) c_2^2 c_3^2 - 2^{-15} = \frac{1}{2} (1 - 2^{-15}) e^{-10/3} - 2^{-15} \\ &\geq 2^{-9} - 2^{-15} > 0. \end{aligned}$$

The proof is now complete. \square

6 Some results on multipliers

In this section, we give some results on the multipliers on $H^2(\beta)$, which show how the different notions of regularity for β intervene.

The set $\mathcal{M}(H^2(\beta))$ of multipliers of $H^2(\beta)$ is by definition the vector space of functions h analytic on \mathbb{D} and such that $hf \in H^2(\beta)$ for all $f \in H^2(\beta)$. When $h \in \mathcal{M}(H^2(\beta))$, the operator M_h of multiplication by h is bounded on $H^2(\beta)$ by the closed graph theorem. The space $\mathcal{M}(H^2(\beta))$ equipped with the operator norm is a Banach space. We note the obvious property:

$$(6.1) \quad \mathcal{M}(H^2(\beta)) \hookrightarrow H^{\infty} \quad \text{contractively.}$$

Indeed, if $h \in \mathcal{M}(H^2(\beta))$, we easily get for all $w \in \mathbb{D}$:

$$M_h^*(K_w) = \overline{h(w)} K_w;$$

so that taking norms and simplifying, we are left with $|h(w)| \leq \|M_h\|$, showing that $h \in H^{\infty}$ with $\|h\|_{\infty} \leq \|M_h\|$.

Proposition 6.1. *We have $\mathcal{M}(H^2(\beta)) = H^\infty$ isomorphically if and only if β is essentially decreasing.*

Proof. The sufficient condition is proved in [12, beginning of the proof of Proposition 3.16]. For the necessity, we then have $\|M_h\| \approx \|h\|_\infty$ for every $h \in H^\infty$ by the Banach isomorphism theorem. Now, for $m > n$ (recall that $e_n(z) = z^n$):

$$e_m(z) = z^{m-n} z^n = (M_{e_{m-n}} e_n)(z);$$

so, since $\|M_{e_{m-n}}\| \leq C \|e_{m-n}\|_\infty = C$ for some positive constant C :

$$\beta_m = \|e_m\|^2 \leq C^2 \|e_n\|^2 = C^2 \beta_n. \quad \square$$

In [12, Section 3.6], we gave the following notion of *admissible* Hilbert space of analytic functions.

Definition 6.2. *A Hilbert space H of analytic functions on \mathbb{D} , containing the constants, and with reproducing kernels K_a , $a \in \mathbb{D}$, is said admissible if:*

- (i) H^2 is continuously embedded in H ;
- (ii) $\mathcal{M}(H) = H^\infty$;
- (iii) the automorphisms of \mathbb{D} induce bounded composition operators on H ;
- (iv) $\frac{\|K_a\|_H}{\|K_b\|_H} \leq h\left(\frac{1-|b|}{1-|a|}\right)$ for $a, b \in \mathbb{D}$, where $h: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is a non-decreasing function.

We proved in that paper that every weighted Hilbert space $H^2(\beta)$ with β non-increasing is admissible, under the additional hypothesis that the automorphisms of \mathbb{D} induce bounded composition operators. In view of Theorem 4.7, we get the following result.

Proposition 6.3. *Let β be a weight.*

- 1) *If β is essentially decreasing, then we have (i), (ii), (iii) in Definition 6.2.*
- 2) *If β is slowly oscillating, then we have (iv) in Definition 6.2.*

Let us give a different proof from the one in [12].

Proof. 1) Let us assume that β is essentially decreasing. Then item (i) holds, as well as item (ii), by Proposition 6.1. Item (iii) is Theorem 4.7.

2) Now we assume that β is slowly oscillating.

Let $0 < s < r < 1$.

Without loss of generality, we may assume that $r, s \geq 1/2$. It is enough to prove:

$$(6.2) \quad \|K_r\|^2 \leq C \|K_{r^2}\|^2$$

for some constant $C > 1$. Indeed, iteration of (6.2) gives:

$$\|K_r\|^2 \leq C^k \|K_{r^{2^k}}\|^2$$

and if k is the smallest integer such that $r^{2^k} \leq s$, we have $2^{k-1} \log r > \log s$ and $2^k \leq D \frac{1-s}{1-r}$ where D is a numerical constant. Writing $C = 2^\alpha$ with $\alpha > 1$, we obtain:

$$\left(\frac{\|K_r\|}{\|K_s\|} \right)^2 \leq C^k = (2^k)^\alpha \leq D^\alpha \left(\frac{1-s}{1-r} \right)^\alpha.$$

To prove (6.2), we pick some $M > 1$ such that $\beta_{2n} \geq M^{-1}\beta_n$ and $\beta_{2n-1} \geq M^{-1}\beta_n$, for all $n \geq 1$, since β is slowly oscillating. Write $t = r^2$. We have:

$$\|K_r\|^2 = \frac{1}{\beta_0} + \sum_{n=1}^{\infty} \frac{t^{2n}}{\beta_{2n}} + \sum_{n=1}^{\infty} \frac{t^{2n-1}}{\beta_{2n-1}},$$

implying, since $t^{2n-1} \leq 4t^{2n}$:

$$\|K_r\|^2 \leq \frac{1}{\beta_0} + M \sum_{n=1}^{\infty} \frac{t^{2n}}{\beta_n} + 4M \sum_{n=1}^{\infty} \frac{t^{2n}}{\beta_n} \leq 5M \|K_t\|^2. \quad \square$$

The notion of admissible Hilbert space H is useful for the set of conditional multipliers:

$$\mathcal{M}(H, \varphi) = \{w \in H; w(f \circ \varphi) \in H \text{ for all } f \in H\}.$$

As a corollary of [12, Theorem 3.18] we get:

Corollary 6.4. *Let β be essentially decreasing and slowly oscillating. Then:*

- 1) $\mathcal{M}(H^2, \varphi) \subseteq \mathcal{M}(H^2(\beta), \varphi)$;
- 2) $\mathcal{M}(H^2(\beta), \varphi) = H^2(\beta)$ if and only if $\|\varphi\|_\infty < 1$;
- 3) $\mathcal{M}(H^2(\beta), \varphi) = H^\infty$ if and only if φ is a finite Blaschke product.

We add here as another application of our results an answer to a question appearing in Problem 5 in Zorboska's thesis [22].

Theorem 6.5. *Let β be a weight such that $H^2(\beta)$ is disc-automorphism invariant and a symbol φ inducing a compact composition operator on $H^2(\beta)$. Then the Denjoy-Wolff point of φ must be in \mathbb{D} .*

In other words, φ has a fixed point in \mathbb{D} .

In the statement, " $H^2(\beta)$ is disc-automorphism invariant" means that for all the automorphisms T_a , where $a \in \mathbb{D}$, C_{T_a} is bounded on $H^2(\beta)$ (equivalently it is bounded for at least one $a \in \mathbb{D} \setminus \{0\}$).

For the definition of the Denjoy-Wolff point, we refer to [17].

Proof. From Theorem 4.10, we know that β is slowly oscillating, and from Proposition 6.3, we know that

$$(6.3) \quad \frac{\|K_a\|_{H^2(\beta)}}{\|K_b\|_{H^2(\beta)}} \leq h \left(\frac{1-|b|}{1-|a|} \right) \quad \text{for every } a, b \in \mathbb{D},$$

where $h: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is a non-decreasing function.

Now we split the proof into two cases.

If $\sum \frac{1}{\beta_n} < \infty$, then $H^2(\beta) \subset A(\mathbb{D})$ (continuously) thanks to the Cauchy-Schwarz inequality. It follows from a result due to Shapiro [16, Th.2.1] that $\|\varphi\|_\infty < 1$ and the conclusion follows obviously in this case.

If $\sum \frac{1}{\beta_n} = \infty$, then the normalized reproducing kernel $\frac{K_z}{\|K_z\|}$ is weakly converging to 0 when $|z| \rightarrow 1^-$, since $\|K_z\| \rightarrow +\infty$.

Since C_φ is compact, C_φ^* is compact as well and we get

$$\frac{K_{\varphi(z)}}{\|K_z\|} \rightarrow 0 \quad \text{when } |z| \rightarrow 1^-$$

and equivalently

$$\frac{\|K_z\|}{\|K_{\varphi(z)}\|} \rightarrow +\infty \quad \text{when } |z| \rightarrow 1^-.$$

But, from (6.3), we get

$$h\left(\frac{1 - |\varphi(z)|}{1 - |z|}\right) \rightarrow +\infty \quad \text{when } |z| \rightarrow 1^-$$

hence, since h is non-decreasing,

$$\frac{1 - |\varphi(z)|}{1 - |z|} \rightarrow +\infty \quad \text{when } |z| \rightarrow 1^-.$$

By the Denjoy-Wolff theorem [17], the conclusion follows in this case too. \square

7 Appendix

In this appendix, we give the proof of Proposition 4.18.

The following lemma can be found in [14, Lemma 1, page 47].

Lemma 7.1. *Let $F: [u, v] \rightarrow \mathbb{R}$, with $u < v$, be a \mathcal{C}^2 -function with $F'' > 0$, and F' not vanishing on $[u, v]$. Let*

$$J = \int_u^v e^{iF(x)} dx.$$

Then:

- a) *if $F' > 0$ on $[u, v]$, then $|J| \leq \frac{2}{F'(u)}$;*
- b) *if $F' < 0$ on $[u, v]$, then $|J| \leq \frac{2}{|F'(v)|}$.*

Proof of Proposition 4.18. Write now the integral I of Proposition 4.18 on $[A, B]$ as $I = I_1 + I_2 + I_3$ with:

$$I_1 = \int_A^{c-\eta} e^{iF(x)} dx, \quad I_2 = \int_{c-\eta}^{c+\eta} e^{iF(x)} dx, \quad I_3 = \int_{c+\eta}^B e^{iF(x)} dx.$$

Lemma 7.1 with $u = A$ and $v = c - \eta$ implies, since $F' < 0$ on $[A, c - \eta]$:

$$(7.1) \quad |I_1| \leq \frac{2}{|F'(c - \eta)|} \leq \frac{2}{\eta \lambda_2},$$

where, for the last inequality, we just have to write

$$|F'(c - \eta)| = F'(c) - F'(c - \eta) = \eta F''(\xi)$$

for some $\xi \in [c - \eta, c]$ so that $F''(\xi) \geq \lambda_2$.

Similarly, Lemma 7.1 with $u = c + \eta$ and $v = B$ implies

$$(7.2) \quad |I_3| \leq \frac{2}{F'(c + \eta)} \leq \frac{2}{\eta \lambda_2}.$$

We can now estimate I_2 . The Taylor formula shows that

$$F(x) = F(c) + \frac{(x - c)^2}{2} F''(c) + R,$$

with

$$|R| \leq \frac{|x - c|^3}{6} \lambda_3.$$

Hence

$$I_2 = e^{iF(c)} \int_0^\eta 2 \exp\left(\frac{i}{2} x^2 F''(c)\right) dx + S$$

with

$$|S| \leq \lambda_3 \int_0^\eta \frac{x^3}{3} dx = \frac{\eta^4}{12} \lambda_3.$$

Finally, set

$$K = \int_0^\eta 2 \exp\left(\frac{i}{2} x^2 F''(c)\right) dx.$$

We make the change of variable $x = \sqrt{\frac{2}{F''(c)}} \sqrt{t}$. Recall that $\int_0^\infty \frac{e^{it}}{\sqrt{t}} dt = \sqrt{\pi} e^{i\pi/4}$ is the classical Fresnel integral, and that an integration by parts gives, for $m > 0$:

$$\left| \int_m^\infty \frac{e^{it}}{\sqrt{t}} dt \right| \leq \frac{2}{\sqrt{m}}.$$

Therefore, with $m = \frac{\eta^2}{2} F''(c)$:

$$K = \sqrt{\frac{2}{F''(c)}} \int_0^m \frac{e^{it}}{\sqrt{t}} dt = \sqrt{\frac{2\pi}{F''(c)}} e^{i\pi/4} + R_m,$$

with

$$|R_m| \leq C \sqrt{\frac{1}{F''(c)}} \frac{1}{\sqrt{m}} \leq \frac{C}{\eta\lambda_2}.$$

All in all, we proved that

$$(7.3) \quad I_2 = \sqrt{\frac{2\pi}{F''(c)}} \exp [i(F(c) + \pi/4)] + O\left(\frac{1}{\eta\lambda_2} + \eta^4\lambda_3\right).$$

and the same estimate holds for I , thanks to (7.1) and (7.2).

We have hence proved Proposition 4.18. □

Acknowledgements. L. Rodríguez-Piazza is partially supported by the project PGC2018-094215-B-I00 (Spanish Ministerio de Ciencia, Innovación y Universidades, and FEDER funds). Parts of this paper was made when he visited the Université d'Artois in Lens and the Université de Lille in January 2020 and September 2022. It is his pleasure to thank all his colleagues in these universities for their warm welcome.

Parts of this paper were made during an invitation of the second-named and third-named authors by the IMUS at the Universidad de Sevilla; it is their pleasure to thank all people in that University who made this stay possible and very pleasant, and especially Manuel Contreras.

The third-named author was partly supported by the Labex CEMPI (ANR-LABX-0007-01).

This work is also partially supported by the grant ANR-17-CE40-0021 of the French National Research Agency ANR (project Front).

We warmly thank R. Zarouf for useful discussions and informations.

References

- [1] A. A. BORICHEV, K. FOUCHET, R. ZAROUF, On the Fourier coefficients of powers of a Blaschke factor and strongly annular functions, *preprint*, arXiv:2107.00405 (2021).
- [2] I. CHALENDAR, J. R. PARTINGTON, Norm estimates for weighted composition operators on spaces of holomorphic functions, *Complex Anal. Oper. Theory* 8, no. 5 (2014), 1087–1095.
- [3] I. CHALENDAR, J. R. PARTINGTON, Compactness and norm estimates for weighted composition operators on spaces of holomorphic functions, *Harmonic analysis, function theory, operator theory, and their applications*, 81–89, Theta Ser. Adv. Math. 19, Theta, Bucharest (2017).
- [4] C. C. COWEN, An application of Hadamard multiplication to operators on weighted Hardy spaces, *Linear Algebra Appl.* 133 (1990), 21–32.

- [5] P. L. DUREN, *Univalent functions*, Grundlehren der Mathematischen Wissenschaften 259, Springer-Verlag, New York (1983).
- [6] E. A. GALLARDO-GUTIÉRREZ, J. R. PARTINGTON, Norms of composition operators on weighted Hardy spaces, *Israel J. Math.* 196, no. 1 (2013), 273–283.
- [7] D. M. GIRARD, The behavior of the norm of an automorphism of the unit disk, *Pacific J. Math.* 47, no. 2 (1973), 443–456.
- [8] G. M. GOLUZIN, On majorants of subordinate analytic functions I, *Mat. Sbornik, N.S.* 29 (1951), 209–224.
- [9] V. È KACNEL'SON, A remark on canonical factorization in certain spaces of analytic functions (Russian), in: *Investigations on linear operators and the theory of functions III*, edited by N. K. Nikol'skiĭ, *Zap. Nauch. Sem. Leningrad. Otdel. Mat. Inst. Steklov (LOMI)* 30 (1972), 163–164. Translation: *J. Soviet Math.* 4 (1975), no. 2 (1976), 444–445.
- [10] K. KELLAY, P. LEFÈVRE, Compact composition operators on weighted Hilbert spaces of analytic functions, *J. Math. Anal. Appl.* 386 (2012), 718–727.
- [11] T. L. KRIETE, B. D. MACCLUER, A rigidity theorem for composition operators on certain Bergman spaces, *Michigan Math. J.* 42 (1995), 379–386.
- [12] P. LEFÈVRE, D. LI, H. QUEFFÉLEC, L. RODRÍGUEZ-PIAZZA, Comparison of singular numbers of composition operators on different Hilbert spaces of analytic functions, *J. Funct. Anal.* 280, no. 3 (2021), 1–47 (article 108834 – <https://doi.org/10.1016/j.jfa.2020.108834>).
- [13] D. LI, H. QUEFFÉLEC, L. RODRÍGUEZ-PIAZZA, A spectral radius type formula for approximation numbers of composition operators, *J. Funct. Anal.* 267, no. 12 (2014), 4753–4774.
- [14] H. L. MONTGOMERY, *Ten lectures on the interface between analytic number theory and harmonic analysis*, CBMS Regional Conference Series in Mathematics 84, American Mathematical Society, Providence RI (1994).
- [15] E. REICH, An inequality for subordinate analytic functions, *Pacific J. Math.* 4, no. 2 (1954), 259–274.
- [16] J. H. SHAPIRO, Compact composition operators on spaces of boundary regular holomorphic functions, *Proc. Amer. J. Math.* 100, no. 1 (1987), 49–57.
- [17] J. H. SHAPIRO, *Composition operators and classical function theory*, Universitext, Tracts in Mathematics, Springer-Verlag, New York (1993).

- [18] A. SHIELDS, Weighted shift operators and analytic function theory. Topics in operator theory. Math. Surveys, No. 13, Amer. Math. Soc., Providence, R.I. (1974), p. 49–128.
- [19] O. SZEHR, R. ZAROUF, l_p -norms of Fourier coefficients of powers of a Blaschke factor, J. Anal. Math. 140 (2020), 1–30.
- [20] O. SZEHR, R. ZAROUF, A constructive approach to Schaeffer’s conjecture, J. Math. Pures Appl. (9) 146 (2021), 1–30.
- [21] E. C. TITCHMARSCH, The theory of the Riemann Zeta-function, Second edition revised by D. R. Heath-Brown, Oxford Science Publications (1986).
- [22] N. ZORBOSKA, Composition operators on weighted Hardy spaces. Thesis (Ph.D.)–University of Toronto (Canada). 1988.

Pascal Lefèvre

Univ. Artois, UR 2462, Laboratoire de Mathématiques de Lens (LML), F-62 300 LENS, FRANCE
 Faculté Jean Perrin, Rue Jean Souvraz, S.P. 18 F-62 300 LENS, FRANCE
 pascal.lefevre@univ-artois.fr

Daniel Li

Univ. Artois, UR 2462, Laboratoire de Mathématiques de Lens (LML), F-62 300 LENS, FRANCE
 Faculté Jean Perrin, Rue Jean Souvraz, S.P. 18 F-62 300 LENS, FRANCE
 daniel.li@univ-artois.fr

Hervé Queffélec

Univ. Lille Nord de France, USTL, Laboratoire Paul Painlevé U.M.R. CNRS 8524 & Fédération Mathématique des Hauts-de-France FR 2037 CNRS, F-59 655 VILLENEUVE D’ASCQ Cedex, FRANCE
 Herve.Queffelec@univ-lille.fr

Luis Rodríguez-Piazza

Universidad de Sevilla, Facultad de Matemáticas, Departamento de Análisis Matemático & IMUS, Calle Tarfia s/n 41 012 SEVILLA, SPAIN
 piazza@us.es