



HAL
open science

Correlations between oxidative DNA damage and formation of hepatic tumours in two flatfish species from contaminated environments

Ciara Baines, Richard Meitern, Randel Kreitsberg, Jérôme Fort, Jörn Peter Scharsack, Pedro Nogueira, Mathieu Giraudeau, Tuul Sepp

► To cite this version:

Ciara Baines, Richard Meitern, Randel Kreitsberg, Jérôme Fort, Jörn Peter Scharsack, et al.. Correlations between oxidative DNA damage and formation of hepatic tumours in two flatfish species from contaminated environments. *Biology Letters*, 2023, 19 (5), 10.1098/rsbl.2022.0583 . hal-04111579

HAL Id: hal-04111579

<https://hal.science/hal-04111579v1>

Submitted on 27 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Correlations between oxidative DNA damage and formation of hepatic tumours in two flatfish species
2 from contaminated environments

3

4 Ciara Baines^{1,2}, Richard Meitern¹, Randel Kreitsberg¹, Jérôme Fort³, Jörn Peter Scharsack⁴,
5 Pedro Nogueira⁴, Mathieu Giraudeau³, Tuul Sepp¹

6

7 ¹Institute of Ecology and Earth Sciences, University of Tartu, Liivi 2, 50409, Tartu, Estonia

8 ²Estonian Marine Institute, University of Tartu, Mäealuse 14, 12618 Tallinn, Harju County,
9 Estonia

10 ³Littoral, Environnement et Sociétés (LIENSs), UMR7266 CNRS - La Rochelle Université, 2
11 rue Olympe de Gouges, 17000 La Rochelle, France

12 ⁴Thünen Institute of Fisheries Ecology, Herwigstraße 31, 27572 Bremerhaven, Germany

13

14 **Abstract**

15 Many species in aquatic environments face increased exposure to oncogenic pollution due to
16 anthropogenic environmental change which can lead to higher cancer prevalence. The
17 mechanistic relationship connecting environmental pollution and cancer is multifactorial and
18 poorly understood, and the specific mechanisms are so far still uncharacterized. One potential
19 mediator between pollutant exposure and cancer is oxidative damage to DNA. We conducted
20 a study in the field with two flatfish species, European flounder (*Platichthys flesus* L.) and
21 common dab (*Limanda limanda* L.) with overlapping distribution and similar ecological niche,
22 to investigate if the link between oncogenic pollutants and cancer described in ecotoxicological
23 literature could be mediated by oxidative DNA damage. This was not the case for flounders as
24 neither PAH bile metabolites nor metallic trace element concentrations were related to
25 oxidative DNA damage measurements. However, dabs with higher PAH metabolite

26 concentrations did exhibit increased oxidative damage. High oxidative DNA damage also did
27 not predict neoplasm occurrence, rather, healthy individuals tended to have higher oxidative
28 damage measurements compared to fishes with pre-neoplastic tumours. Our analyses showed
29 that flounders had lower concentrations of PAH bile metabolites, suggesting that compared to
30 dab, this species is less exposed or better at eliminating these contaminants.

31 Keywords: Oxidative stress, hepatic tumours, marine contamination, European flounder,
32 *Limanda limanda*, wildlife cancer

33

34 **Introduction**

35 Cancer is a disease that arose with multicellularity, potentially affecting all
36 multicellular organisms [1]. Increasing evidence suggests some species have evolved stronger
37 tumour suppressor mechanisms than others [2,3]. Whilst these mechanisms are the result of
38 species'-specific life-history and/or environmental exposure to natural oncogenic substances
39 [4,5], many species face increasing pressure from anthropogenic activities that could affect
40 cancer prevalence in natural populations [6,7].

41 There is evidence that some species can adapt to increased concentrations of oncogenic
42 pollutants, especially if the selection pressures are high and pollution is present for extended
43 time periods [8,9]. One step towards understanding the mechanisms behind evolved cancer
44 defence mechanisms in natural populations is to focus on specific physiological pathways to
45 explore why cancer prevalence varies between populations and species living in environments
46 contaminated with hazardous chemicals. One potential mediator between pollutant exposure
47 and cancer is oxidative DNA damage, as both organic pollutants [10] and metallic trace
48 elements [11] increase it. This can lead to cancer initiation and progression by increasing DNA
49 mutations or inducing genome instability and cell proliferation [12]. However, relatively little
50 is known about how the amount of oxidative DNA damage impacts cancer risk in pollution-

51 exposed animal populations, as current studies published mostly focus on changes in
52 antioxidants in response to a specific pollutant in laboratory settings [13-15].

53 Flatfish, such as European flounders (*Platichthys flesus* L.) and dabs (*Limanda limanda*
54 *L.*) have been studied for both pollution exposure and hepatic tumours [16, 17]. These benthic
55 species are regularly used in biomonitoring surveys throughout the North Sea and Baltic Sea
56 marine areas, which have been subjected to high concentrations of pollution for over a hundred
57 years [18,19], through atmospheric and riverine inputs, and industrial point-sources (e.g. oil
58 and gas platforms, [20-22]). Histopathological analyses have indicated that dabs have
59 approximately 10 times higher liver neoplasia prevalence than flounders [23-28]. However, the
60 causes behind this species-specific difference are not known.

61 This field study investigates correlations between the formation of hepatic neoplastic
62 lesions, and (a) concentrations of polycyclic aromatic hydrocarbon (PAHs) bile metabolites,
63 (b) liver concentrations of metallic trace elements (Hg, Cd, As, Pb). We also investigate
64 whether oxidative DNA damage (measured via 8- hydroxy-2'-deoxyguanosine, 8-OhdG) acts
65 as a mediator between contaminant exposure and neoplasm occurrence.

66

67 **Methods**

68 **Field Sampling**

69 Fish were collected using a trawler at ten Baltic and North Sea locations (See Table S1 for
70 locations and numbers sampled) in August 2019. Fishes were weighed and measured before
71 being killed by a percussive blow to the head that was followed by destruction of the brain
72 using a surgical knife. Bile and liver samples were flash frozen in liquid nitrogen and stored at
73 -80°C until analysis. A 3 mm liver section for histopathology analysis was stored in 4%
74 formalin for 48 hours, then in 70% ethanol. Otoliths were collected to measure age. Sites were

75 allocated as 'reference' or 'polluted' based on OSPAR/HELCOM data on local anthropogenic
76 pressures (Table S1, [29-31]).

77 Histopathology

78 Liver samples were processed according to Feist et al [32] for histopathological analysis,
79 prepared onto slides and analysed using a Nikon Eclipse 80i microscope. Foci of cellular
80 alteration (FCA) and neoplastic lesions were recorded [32] (see supplementary Table S1 and
81 figures S1, S2 for further details).

82 PAH bile metabolites

83 PAH bile metabolites were measured using fixed fluorescence methods. Bile samples were
84 diluted with 48% ethanol (1:1600) [33] and read using a fluorescence spectrophotometer (BMG
85 Omega Fluostar) measuring fixed wavelength pairs (excitation and emission, respectively).
86 290/380 nm (2-ring PAH); 256/380 nm (3-ring PAH); 341/383 nm (4-ring PAH); and 380/430
87 nm (5-ring PAH) [34]. The results were presented as fluorescence units (FU) [35].

88 Trace metal concentrations

89 Liver samples were freeze-dried for 48 hours and ground to powder. Total Hg concentrations
90 were measured, in duplicate (relative standard deviation for aliquots <10%), in subsamples of
91 ~1mg of homogenized liver using an advanced Hg analyser spectrophotometer (Altec AMA
92 254) [36]. Remaining samples were analysed for trace elements arsenic (As), cadmium (Cd),
93 and lead (Pb). Samples were digested with a mixture of 3 mL HNO₃ and 5 mL HCl Suprapur
94 quality, heated in a microwave oven and diluted to 50 ml with deionized water, and analysed
95 by Inductively Coupled Plasma Atomic Emission Spectrometry (Varian Vista-Pro ICP-AES)
96 and Mass Spectrometry (ICP-MS II Series Thermo Fisher Scientific).

97 Oxidative stress markers

98 Biomarker 8-hydroxy-2'-deoxyguanosine (8-OHdG) was used to measure oxidative lesions
99 formed in nuclear or mitochondrial DNA induced by free radicals. DNA, extracted using the

100 Purelink Genomic DNA Mini Kit (Invitrogen), was quantified using a Qubit 4 fluorometer.
101 Standards were read on the dsDNA HS setting to calibrate the machine. 1µL of sample was
102 added to 199µL of the working solution in triplicates, analysed, and the means calculated in
103 ng/µL.

104 Samples were prepared for ELISA kits (Cell Biolabs Inc, cat number STA-320) following the
105 protocol for cell or tissue DNA samples by converting to single strand DNA, digesting DNA
106 using nuclease P1 (M0660S, New England Biolabs), adding alkaline phosphatase (CIP,
107 M0371S New England Biolabs), centrifuging at 6000g for 5 minutes and collecting
108 supernatants. Plates were washed using BioTek ELx50 microplate washer and absorbance
109 (wavelength 450nm) measured using a spectrophotometer (Biotek Synergy 2). Results were
110 normalised against the DNA concentrations, converted from 1 to 4 strand DNA and presented
111 as ng oxidised DNA bases per mg DNA.

112 Statistical analysis

113 Statistical analyses were run using R 4.0.5[37]. Principle component analysis (PCA), using
114 packages “stats”[38] and “psych”[39], grouped 2 components (see Figures S3-S4) splitting the
115 variables into metals (As, Cd, Pb, Hg) and hydrocarbons (2, 3, 4, 5 ring). Generalised linear
116 models tested whether oxidative stress or tumours could be predicted by trace metals or PAH
117 metabolite concentrations, age, location (Baltic or North Sea), or whether sites were
118 polluted/reference (Table S3). Chi-squared tested for a difference between the number of fishes
119 with or without neoplasms/FCAs (3-levels: no abnormalities, FCA, neoplasm). Finally,
120 ANOVA and post-hoc Tukey tests tested for differences between oxidative stress and tumours.
121 Other packages used for plotting and data reorganisation were ggplot2[40] tidyverse [41],
122 car[42] and various dependencies within them. Additional analysis with subsets of data (fish
123 aged 3+ as younger fish as less likely to develop cancer and sites where both species were

124 present to exclude the influence of site characteristics on the results) are provided as
125 supplementary material alongside all code and data (<https://doi.org/10.5281/zenodo.7846187>).

126

127 **Results**

128 Histopathology analyses indicated that dabs do not have significantly higher tumour prevalence
129 (FCA or neoplasm) than flounders (Figure 1a, Chi-squared =2.81, df=2, p=0.2). This result
130 could, however, be affected by the bias in the sampling, with more younger dabs in the data set
131 compared to flounders. Linear models found that flounders were older and had higher amounts
132 of oxidative DNA damage than dabs, dabs had significantly higher concentrations of PAH
133 metabolites (Table 1). However, these results may be confounded by differences between sites.
134 When we repeated the species comparison, including only the fish from the three sampling
135 sites where both species were sampled, we found that only age was significantly different
136 between dabs and flounders (Table S4). However, all the fishes with the highest contaminant
137 concentrations were dabs (Figure 1b). There was a significantly higher variation (F-test) in
138 metals ($p<0.001$) and hydrocarbons ($p=0.007$) in dabs than flounders, whereas age ($p=0.057$)
139 and log-transformed oxidative DNA damage ($p=0.555$) did not vary significantly between the
140 species.

141 The generalised linear model suggested that neither age, contaminant concentrations, oxidative
142 stress, or sea (North vs Baltic) influenced the likelihood of a fish developing neoplastic/pre-
143 neoplastic lesions (both species). However, there was significantly higher proportion of
144 tumours in fish living in reference sites compared to more polluted sites (Table 2).
145 Additionally, older flounders had more tumours than younger ones. As age is known to
146 significantly affect cancer prevalence, and we had an over-representation of young dabs in our
147 dataset, additional analysis excluding fishes under 3 years is provided in supplementary
148 materials. Results of these models were the same as for the full dataset (Table S5). In order to

149 check that site characteristics were not confounding our results, additional analysis was run
150 with site as a predictor (Supplementary materials: Table S8, Figure S5). The model including
151 site resulted in a singular fit, which suggests preferring a simpler model, as such we chose the
152 models with "polluted/reference" and "sea" for the main results.

153 There was a significantly higher level of oxidative damage in the fishes from the Baltic Sea
154 than the North Sea for both species ($t=-4.511$, $p<0.001$), flounders ($t\text{-value}=-2.305$, $p<0.031$)
155 but age, contaminant concentrations or whether a site was characterised as polluted/reference
156 did not affect oxidative damage (Table S6). In dabs, oxidative damage was significantly higher
157 in the Baltic ($t\text{-value}=-4.198$, $p<0.001$) and in dabs with higher concentrations of PAH
158 metabolites. In the reduced dataset (only fish aged 3+) oxidative damage was significantly
159 higher in the Baltic but the relationship, for dabs, with hydrocarbons (Table S7).

160 An ANOVA model indicated that oxidative damage was significantly higher in healthy dabs
161 compared to those with FCA ($p=0.027$) but not different between healthy individuals and those
162 with neoplasms or an FCA (Figure 2). There were no significant differences for flounders.

163

164 **Discussion**

165 This field study with two flatfish species investigates whether the link between oncogenic
166 contaminants and hepatic tumours, described in ecotoxicological literature, could be mediated
167 by oxidative DNA damage. We did not find evidence for this in flounders, as neither PAH nor
168 metallic trace element concentrations were related to their oxidative DNA damage, whereas in
169 dabs, higher concentrations of PAH metabolites were related to increased oxidative damage. It
170 is possible that including other contaminants (e.g., persistent organic pollutants) could provide
171 a different picture. Furthermore, oxidative DNA damage was not related to neoplasm
172 occurrence in flounders whereas healthy dabs had increased DNA damage compared to dabs
173 with FCAs. When comparing the species with each other (previous studies indicated higher

174 pollution-induced neoplasm prevalence in dabs than flounders [eg.23-28]) we found flounders
175 had more oxidative DNA damage. However, contaminant concentrations varied more in dab
176 compared to flounders suggesting potential differences in either exposure or pollutant removal
177 mechanisms (Figure 1b). Additionally, PAH metabolite concentrations in flounders were lower
178 than in dabs, suggesting more efficient organic pollutant metabolism in flounders. It must be
179 noted, however, that age structure and sample sizes for age groups differed between the species,
180 with more younger dab samples compared to flounders (Table S2). While age was included in
181 the models, this still might confound the results when comparing species. Additionally, it
182 should be noted that there are fewer 'reference' sites than 'polluted', and it is possible that
183 adding additional sampling locations could change these results.

184 Contaminants can induce oxidative stress by either enhanced reactive oxygen species
185 generation and/or weakening of antioxidant/damage repair systems [43]. Fish studies have
186 indicated that environmental contaminants can affect antioxidant levels (organic pollutants
187 [44], heavy metals [45,46]). However, limitations in detecting oxidative stress, through
188 measuring oxidants and antioxidants, occur from time-lagged and hormetic upregulation of
189 protective mechanisms [47]. Therefore, we assessed the oxidative DNA damage (8-OHdG)
190 from the liver tissue of fish exposed to contaminated environments. The levels of 8-OHdG in
191 aquatic species have been studied as a biomarker of oxidative stress caused by contamination,
192 showing increased levels in populations exposed to higher contamination burdens [48,49]. In
193 our study, increased PAH metabolite concentrations in dabs, but not flounders, were related to
194 higher oxidative DNA damage levels, suggesting that PAH metabolism is linked to increased
195 DNA damage in dabs. The different results for dabs and flounders could arise here from the
196 differences in sampling sites, as only 3 of the 10 sampling sites included both species. This
197 result for dabs, whilst unexpected, could be an indication of acclimatization or adaptation
198 processes, resulting from long-term exposure to oncogenic contamination to reduce oxidative

199 DNA damage. However, a study on killifish (*Fundulus heteroclitus*) suggested that most of the
200 DNA damage present following PAH exposure was the result of covalent binding of
201 metabolites to DNA rather than from oxidative damage [50]. Whether this is the case for
202 flounders and dabs remains to be studied. Interestingly, tumours were not related to the levels
203 of oxidative DNA damage in the liver, suggesting other processes must be involved that
204 contribute more to tumour development in wild fish than oxidative DNA damage. Rather, in
205 dabs with no abnormalities detected, there was less oxidative damage than in dabs with an FCA
206 (no difference with neoplastic individuals or in flounders).

207 In this study, older fishes had more oxidative DNA damage than younger ones. Increased DNA
208 damage in older fish have been recorded in dabs using comet assay methods [51]. Additionally,
209 a study on three-spined sticklebacks (*Gasterosteus aculeatus*) found that larger fishes have
210 more DNA damage than smaller ones using comet assay and micronucleus tests [52],
211 potentially resulting from either the higher metabolic need of larger fish, leading to more
212 oxidative DNA damage, or DNA damage accumulation with age resulting from the senescence
213 process.

214 Fishing location had the greatest influence on elevated oxidative damage, demonstrated by
215 Baltic Sea fish exhibiting a significantly higher oxidative damage than North Sea fish. Changes
216 in salinity can cause oxidative stress in fish [53] and a study on olive flounder (*Paralichthys*
217 *olivaceus*) found that hyposaline conditions caused oxidative stress [54] so it is possible that
218 the low saline Baltic environment is increasing oxidative damage in flounders and dabs.

219 Additionally, a report by the European Environment Agency suggests a larger proportion of
220 the Baltic (96%) is polluted than the North Sea area (75%) [55] which could cause increased
221 oxidative damage in the Baltic. Actively removing organic pollutants from the organism is
222 considered a risk factor for cancer, as the by-products of active pollutant metabolism by the
223 cytochrome p450 enzymes (CYPs) lead to carcinogenesis via DNA adduct formation [56,57].

224 In this study, fish caught in ‘reference’ sites had more tumours than those from ‘polluted’ sites.
225 However, it should also be noted that migration of fishes was not accounted for when
226 categorising sites as ‘reference’ or ‘polluted’ and exposure rates could vary between
227 individual’s dependant on migratory distances. It is possible that fish living in heavily polluted
228 environments can adapt/acclimate to cope with contamination [58]. In conclusion, we showed
229 that in an ecosystem where pollution has persisted for a long time, it is not possible to find a
230 clear link between contaminants (PAH and metallic trace elements), oxidative DNA damage,
231 and neoplasms. In the North and Baltic seas, both flounders and dabs have had approximately
232 50 generations to adapt to high concentrations of environmental contamination. We can suggest
233 that both species, but especially flounders, could have gone through evolutionary processes,
234 where stronger cancer defence mechanisms, more efficient pollutant metabolism, and
235 protection against oxidative DNA damage was selected for.

236

237 **Acknowledgements**

238 This work was supported by the Estonian Research council grants PSG653 to Tuul Sepp and
239 PRG1137 to Peeter Hõrak. We are grateful to C. Churlaud and to M. Brault-Favrou from the
240 "Plateforme Analyses Élémentaires" of LIENSs for their assistance during trace element
241 analyses and to the crew of RW Walther Herwig III for all-round help during the fieldwork.

242

243 **References**

244

- 245 1. Trigos AS, Pearson RB, Papenfuss AT, Goode DL. 2018 How the evolution of
246 multicellularity set the stage for cancer. *Br. J. Cancer* **118**, 145-152.
247 (doi.org/10.1038/bjc.2017.398)

- 248 2. Tollis M, Schiffman JD, Boddy AM. 2017 Evolution of cancer suppression as revealed
249 by mammalian comparative genomics. *Curr. Opin. Genet. Dev.* **42**, 40-47.
250 (doi.org/10.1016/j.gde.2016.12.004)
- 251 3. Baines C, Meitern R, Kreitsberg R, Sepp T. 2022 Comparative study of the evolution
252 of cancer gene duplications across fish. *Evol. Appl.* **15**, 1834-1845.
253 (doi.org/10.1111/eva.13481)
- 254 4. Ujvari B, Beckmann C, Biro PA, Arnal A, Tasiemski A, Massol F, Salzet M, Mery F,
255 Boidin-Wichlacz C, Misse D, Renaud F. 2016 Cancer and life-history traits: lessons
256 from host–parasite interactions. *Parasitology* **143**, 533-541.
257 (doi.org/10.1017/S0031182016000147)
- 258 5. Vittecoq M, Giraudeau M, Sepp T, Marcogliese DJ, Klaassen M, Renaud F, Ujvari B,
259 Thomas F. 2018 Turning natural adaptations to oncogenic factors into an ally in the war
260 against cancer. *Evol. Appl.* **11**, 836-844. (doi.org/10.1111/eva.12608)
- 261 6. Giraudeau M, Sepp T, Ujvari B, Ewald PW, Thomas F. 2018 Human activities might
262 influence oncogenic processes in wild animal populations. *Nat. Ecol. Evol.* **2**, 1065-
263 1070. (doi.org/10.1038/s41559-018-0558-7)
- 264 7. Baines C, Lerebours A, Thomas F, Fort J, Kreitsberg R, Gentes S, Meitern R, Saks L,
265 Ujvari B, Giraudeau M, Sepp T. 2021 Linking pollution and cancer in aquatic
266 environments: A review. *Environ. Int.* **149**, 106391.
267 (doi.org/10.1016/j.envint.2021.106391)
- 268 8. Wirgin I, Roy NK, Loftus M, Chambers RC, Franks DG, Hahn ME. 2011 Mechanistic
269 basis of resistance to PCBs in Atlantic tomcod from the Hudson River. *Science* **331**,
270 1322–1325. (DOI: 10.1126/science.1197296)
- 271 9. Reid NM, Proestou DA, Clark BW, Warren WC, Colbourne JK, Shaw JR, Karchner
272 SI, Hahn ME, Nacci D, Oleksiak MF, Crawford DL, Whitehead A. 2016 The genomic

273 landscape of rapid repeated evolutionary adaptation to toxic pollution in wild fish.
274 *Science* **354**, 1305–1308. (DOI: 10.1126/science.aah4993)

275 10. Singh R, Kaur B, Kalina I, Popov TA, Georgieva T, Garte S, Binkova B, Sram RJ,
276 Taioli E, Farmer P B. 2007 Effects of environmental air pollution on endogenous
277 oxidative DNA damage in humans. *Mutat. Res.* **620**, 71–82.
278 (doi.org/10.1016/j.mrfmmm.2007.02.024)

279 11. Hengstler JG, Bolm-Audorff U, Faldum A, Janssen K, Reifenrath M, Götte W, Jung D,
280 Mayer-Popken O, Fuchs J, Gebhard S Bienfait HG. 2003 Occupational exposure to
281 heavy metals: DNA damage induction and DNA repair inhibition prove co-exposures
282 to cadmium, cobalt and lead as more dangerous than hitherto expected. *Carcinogenesis*
283 **24**, 63-73. (doi.org/10.1093/carcin/24.1.63)

284 12. Valavanidis A, Vlachogianni T, Fiotakis C. 2009 8-hydroxy-2'-deoxyguanosine (8-
285 OHdG): a critical biomarker of oxidative stress and carcinogenesis. *J. Environ. Sci.*
286 *Health C: Toxicol.* **27**, 120-139. (doi.org/10.1080/10590500902885684)


287 13. Isaksson C. 2010 Pollution and its impact on wild animals: a meta-analysis on oxidative
288 stress. *EcoHealth* **7**, 342–350. (doi.org/10.1007/s10393-010-0345-7)

289 14. Liu H, Sun P, Liu H, Yang S, Wang L, Wang Z. 2015 Hepatic oxidative stress
290 biomarker responses in freshwater fish *Carassius auratus* exposed to four benzophenone
291 UV filters. *Ecotoxicol. Environ. Saf.* **119**, 116-122.
292 (doi.org/10.1016/j.ecoenv.2015.05.017)

293 15. Tu TY, Hong CY, Sasado T, Kashiwada S, Chen PJ. 2016 Early life exposure to a
294 rodent carcinogen propiconazole fungicide induces oxidative stress and
295 hepatocarcinogenesis in medaka fish. *Aquat. Toxicol.* **170**, 52-61.
296 (doi.org/10.1016/j.aquatox.2015.11.014)

- 297 16. Lerebours A, Stentiford GD, Lyons BP, Bignell JP, Derocles SA, Rotchell JM. 2014
298 Genetic alterations and cancer formation in a European flatfish at sites of different
299 contaminant burdens. *Environ. Sci. Technol.* **48**, 10448-10455
300 (doi.org/10.1021/es502591p)
- 301 17. Williams TD, Turan N, Diab AM, Wu H, Mackenzie C, Bartie KL, Hrydziuszko O,
302 Lyons BP, Stentiford GD, Herbert JM, Abraham JK. 2011 Towards a system level
303 understanding of non-model organisms sampled from the environment: a network
304 biology approach. *PLoS Comput Biol*, **7**, 1002126.
305 (doi.org/10.1371/journal.pcbi.1002126)
- 306 18. HELCOM 2018: State of the Baltic Sea – Second HELCOM holistic assessment 2011-
307 2016. Baltic Sea Environment Proceedings 155.
- 308 19. Carpenter A. 2019 Oil pollution in the North Sea: the impact of governance measures
309 on oil pollution over several decades. *Hydrobiologia*, **845**, 109-127.
310 (doi.org/10.1007/s10750-018-3559-2)
- 311 20. HELCOM. 2021 Inputs of hazardous substances to the Baltic Sea. Baltic Sea
312 Environment Proceedings No.179.
- 313 21. Marappan S, Stokke R, Malinovsky MP, Taylor A. 2022. Assessment of impacts of the
314 offshore oil and gas industry on the marine environment. In: OSPAR, 2023: The 2023
315 Quality Status Report for the North-East Atlantic. OSPAR Commission, London.
- 316 22. Ilyin I, Travnikov O, Rozovskaya O, Strizhkina I. 2022. Trends in Deposition of Heavy
317 Metals to the OSPAR Maritime Area in the period 1990-2019. In: OSPAR, 2023: The
318 2023 Quality Status Report for the North-East Atlantic. OSPAR Commission, London.
- 319 23. Cachot J, Cherel Y, Larcher T, Pfohl-Leszkowicz A, Laroche J, Quiniou L, Morin J,
320 Schmitz J, Burgeot T, Pottier D. 2013 Histopathological lesions and DNA adducts in
321 the liver of European flounder (*Platichthys flesus*) collected in the Seine estuary versus

- 322 two reference estuarine systems on the French Atlantic coast. *Environ. Sci. Poll.*
323 *Res.* **20**, 723-737. (doi.org/10.1007/s11356-012-1287-0)
- 324 24. Lang T, Wosniok W, Baršienė J, Broeg K, Kopecka J, Parkkonen J. 2006 Liver
325 histopathology in Baltic flounder (*Platichthys flesus*) as indicator of biological effects
326 of contaminants. *Mar. Poll. Bulletin.* **53**, 488-496.
327 (doi.org/10.1016/j.marpolbul.2005.11.008)
- 328 25. Stentiford GD, Longshaw M, Lyons BP, Jones G, Green M, Feist SW. 2003
329 Histopathological biomarkers in estuarine fish species for the assessment of biological
330 effects of contaminants. *Mar. Envir. Res.* **55**, 137-159. (doi.org/10.1016/S0141-
331 1136(02)00212-X)
- 332 26. Stentiford GD, Bignell JP, Lyons BP, Feist SW. 2009 Site-specific disease profiles in
333 fish and their use in environmental monitoring. *Mar. Ecol. Progr. Series.* **381**, 1-15.
334 (doi.org/10.3354/meps07947)
- 335 27. Lyons BP, Stentiford GD, Bignell J, Goodsir F, Sivyer DB, Devlin MJ, Lowe D,
336 Beesley A, Pascoe CK, Moore MN, Garnacho E. 2006 A biological effects monitoring
337 survey of Cardigan Bay using flatfish histopathology, cellular biomarkers and sediment
338 bioassays: Findings of the Prince Madog Prize 2003. *Mar. Environ. Res.* **62**, 342-346.
339 (doi.org/10.1016/j.marenvres.2006.04.064)
- 340 28. CEFAS, Marine Environment Monitoring Group. 2004. UK National Marine
341 Monitoring Programme - Second Report (1999-2001). ISBN 0 907545 20 3.
- 342 29. Hylland K, Skei BB, Brunborg G, Lang T, Gubbins MJ, Le Goff J, Burgeot T. 2017
343 DNA damage in dab (*Limanda limanda*) and haddock (*Melanogrammus aeglefinus*)
344 from European seas. *Marine Environmental Research* 124, 54-60.
- 345 30. OSPAR Commission. 2010. OSPAR Quality Status Report 2010.
346 <https://qsr2010.ospar.org/en/index.html>

- 347 31. Zaborska A, Siedlewicz G, Szymczycha B, Dzierzbicka-Głowacka L, Pazdro K. 2019
348 Legacy and emerging pollutants in the Gulf of Gdańsk (southern Baltic Sea) – loads
349 and distribution revisited. *Marine Pollution Bulletin* 139, 238-255
- 350 32. Feist SW, Lang T, Stentiford GD, Köhler AICES. 2004 Biological effects of
351 contaminants: use of liver pathology of the European flatfish dab (*Limanda limanda*
352 L.) and flounder (*Platichthys flesus* L.) for monitoring. *ICES Techniques in Marine*
353 *Environmental Sciences* 38. (doi.org/10.25607/OBP-226)
- 354 33. Aas E, Baussant T, Balk L, Liewenborg B, Andersen OK. 2000 PAH metabolites in
355 bile, cytochrome P4501A and DNA adducts as environmental risk parameters for
356 chronic oil exposure: a laboratory experiment with Atlantic cod. *Aquat. Toxicol.* **51**,
357 241–258. (doi.org/10.1016/S0166-445X(00)00108-9)
- 358 34. Lee RF, Anderson JW. 2005 Significance of cytochrome P450 system responses and
359 levels of bile fluorescent aromatic compounds in marine wildlife following oil spills.
360 *Mar. Pollut. Bull.* **50**, 705–723. (doi.org/10.1016/j.marpolbul.2005.04.036)
- 361 35. Beyer J, Jonsson G, Porte C, Krahn MM, Ariese F. 2010 Analytical methods for
362 determining metabolites of polycyclic aromatic hydrocarbon (PAH) pollutants in fish
363 bile: a review. *Environ. Toxicol. Pharmacol.* **30**, 224–244.
364 (doi.org/10.1016/j.etap.2010.08.004)
- 365 36. Bustamante P, Lahaye V, Durnez C, Churlaud C, Caurant F. 2006 Total and organic
366 Hg concentrations in cephalopods from the North Eastern Atlantic waters: influence of
367 geographical origin and feeding ecology. *Sci. Total Environ.* **368**, 585-596.
- 368 37. R Core Team. 2021 R: A language and environment for statistical computing. R
369 Foundation for Statistical Computing, Vienna, Austria. URL  [https://www.R-](https://www.R-project.org/)
370 [project.org/](https://www.R-project.org/).

- 371 38. Revelle W. 2022 psych: Procedures for Psychological, Psychometric, and Personality
372 Research. Northwestern University, Evanston, Illinois. R package version
373 2.2.9, <https://CRAN.R-project.org/package=psych>.
- 374 39. Harrell Jr FE. 2022. Rms: Regression Modelling strategies. *R package version 6.0*
375 Dept. Biostatist., Vanderbilt Univ., Nashville, TN, USA.
- 376 40. Wickham H, Chang W, Wickham MH. 2016. Create elegant data visualisations using
377 the grammar of graphics. Package ‘ggplot2’ version 3.3.6.
- 378 41. Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, Golemund
379 G, Hayes A, Henry L, Hester J, et al. 2019. Welcome to the Tidyverse. *J Open Source*
380 *Softw.* 4:1686. (DOI: 10.21105/joss.01686)
- 381 42. Fox J, Weisberg S. 2019. An R Companion to Applied Regression, Third edition. Sage,
382 Thousand Oaks CA. <https://socialsciences.mcmaster.ca/jfox/Books/Companion/>.
- 383 43. Lushchak VI. 2016 Contaminant-induced oxidative stress in fish: a mechanistic
384 approach. *Fish Physiol. Biochem.* **42**, 711–747. (doi.org/10.1007/s10695-015-0171-5)
- 385 44. Chinnadurai K, Prema P, Veeramanikandan V, Kumar KR, Nguyen VH, Marraiki N,
386 Zaghoul NS, Balaji P. 2022 Toxicity evaluation and oxidative stress response of
387 fumaronitrile, a persistent organic pollutant (POP) of industrial waste water on tilapia
388 fish (*Oreochromis mossambicus*). *Environ. Res.* **204**, 112030.
389 (doi.org/10.1016/j.envres.2021.112030)
- 390 45. Kumari K, Khare A, Dange S. 2014. The applicability of oxidative stress biomarkers in
391 assessing chromium induced toxicity in the fish *Labeo rohita*. *BioMed Res. Int.* **2014**,
392 782493. (doi.org/10.1155/2014/782493)
- 393 46. Das D, Moniruzzaman M, Sarbajna A, Chakraborty SB. 2017. Effect of heavy metals
394 on tissue-specific antioxidant response in Indian major carps. *Environ. Sci. Pollut. Res.*
395 **24**, 18010–18024. (doi.org/10.1007/s11356-017-9415-5)

- 396 47. Meitern R, Sild E, Kilk K, Porosk R, Hõrak P. 2013 On the methodological limitations
397 of detecting oxidative stress: effects of paraquat on measures of oxidative status in
398 greenfinches. *J. Exp. Biol.* **216**, 2713-2721. (doi.org/10.1242/jeb.087528)
- 399 48. Machella N, Regoli F, Cambria A, Santella RM. 2004 Oxidative damage to DNA: an
400 immunohistochemical approach for detection of 7, 8-dihydro-8-oxodeoxyguanosine in
401 marine organisms. *Mar. Environ. Res.* **58**, 725-729.
402 (doi.org/10.1016/j.marenvres.2004.03.022)
- 403 49. El-Agri AM, Emam MA, Gaber HS, Hassan EA, Hamdy SM. 2022 Integrated use of
404 biomarkers to assess the impact of heavy metal pollution on *Solea aegyptiaca* fish in
405 Lake Qarun. *Environ. Sci. Eur.* **34**, 1-24. (doi.org/10.1186/s12302-022-00656-6)
- 406 50. Jung D, Cho Y, Collins LB, Swenberg JA, Di Giulio RT. 2009. Effects of benzo [a]
407 pyrene on mitochondrial and nuclear DNA damage in Atlantic killifish (*Fundulus*
408 *heteroclitus*) from a creosote-contaminated and reference site. *Aquat. Toxicol.* **95**, 44-
409 51.
- 410 51. Akcha F, Leday G, Pfohl-Leszkowicz A. 2004 Measurement of DNA adducts and
411 strand breaks in dab (*Limanda limanda*) collected in the field: effects of biotic (age,
412 sex) and abiotic (sampling site and period) factors on the extent of DNA
413 damage. *Mutat. Res-Fund. Mol. M.* **552**, 197-207.
414 (doi.org/10.1016/j.mrfmmm.2004.06.020)
- 415 52. Wirzinger G, Weltje L, Gercken J, Sordyl H. 2007 Genotoxic damage in field-collected
416 three-spined sticklebacks (*Gasterosteus aculeatus* L.): a suitable biomonitoring
417 tool?. *Mutat. Res.-Genet. Toxicol.* **628**,19-30.
418 (doi.org/10.1016/j.mrgentox.2006.11.011)
- 419 53. Birnie-Gauvin K, Costantini D, Cooke SJ, Willmore WG. 2017. A comparative and
420 evolutionary approach to oxidative stress in fish: a review. *Fish Fish (Oxf)*, **18** 928-942.

421 54. Lee, D.W., Choi, Y.U., Park, H.S., Park, Y.S. and Choi, C.Y., 2022. Effect of low pH
422 and salinity conditions on the antioxidant response and hepatocyte damage in juvenile
423 olive flounder *Paralichthys olivaceus*. *Mar. Environ. Res.* **175**, 105562.

424 55. European Environment Agency. 2019. Contaminants in Europe's seas. Moving towards
425 a clean, non-toxic marine environment. 61.

426 56. Stegeman JJ, Lech JJ. 1991 Cytochrome P-450 monooxygenase systems in aquatic
427 species: carcinogen metabolism and biomarkers for carcinogen and pollutant exposure.
428 *Environ. Health Perspect.* **90**, 101–109. (doi.org/10.1289/ehp.90-1519513)

429 57. Willett KL, Ganesan S, Patel M, Metzger C, Quiniou S, Waldbieser G, Scheffler B.
430 2006. In vivo and in vitro CYP1B mRNA expression in channel catfish. *Mar. Environ.*
431 *Res.* **62** S332-S336. (doi.org/10.1016/j.marenvres.2006.04.015)

432 58. Wirgin I, Waldman JR. 2004. Resistance to contaminants in North American fish
433 populations. *Mutat. Res-Fund. Mol. M.* **552**, 73-100.

434 59. Sepp T, Baines C, Kreitsberg R, Scharsack JP, Nogueira P, Lang T, Fort J, Sild E,
435 Clarke JT, Meitern R. (2022) ENA archive bioproject PRJEB53201
436 <https://www.ebi.ac.uk/ena/browser/view/PRJEB53201>.

437
438

439 Table 1: Linear model results for a difference between oxidative damage, age and contaminant
440 burdens between species. * represents p-value <0.05, ** represents p<0.01.

	Coefficient	S.E.	Wald Z	Pr(> Z)
Intercept	-18.7452	5.1216	-3.660	0.000252 ***
Oxidative DNA Damage(log)	2.3344	0.8339	2.799	0.005121 **

Metals	-0.5968	0.4578	-1.304	0.1924
Hydrocarbons	-0.9054	0.3677	-2.462	0.0138*
Age	1.1217	0.2451	4.577	<0.0001*
				**

441

442 Table 2: Model results (N=128) of variables influencing the differences in pre-
443 neoplastic/neoplastic tumours between both species and each species individually. *
444 represents p-value <0.05, ** represents p<0.01.

445

		Estimate	Std. Error	t value	Pr(> t)	
Both	(Intercept)	3.11513	4.5926	0.678	0.49759	
	Metals	-0.20913	0.21477	-0.974	0.33019	
	Hydrocarbons	0.12996	0.22643	0.574	0.56601	
	Age	-0.02367	0.1649	-0.144	0.88584	
	Oxidative Stress	-0.56179	0.75214	-0.747	0.45511	
	Sea (North)	1.18170	0.71741	1.647	0.09927	
	Polluted	-1.4411	0.55388	-2.602	0.00927	**
Dab	(Intercept)	9.92253	5.94056	1.670	0.0949	
	Metals	-0.20196	0.22902	-0.882	0.3778	
	Hydrocarbons	0.24261	0.28230	0.859	0.3901	
	Age	-0.03751	0.34256	-0.110	0.9128	
	Oxidative Stress	-1.73457	0.96487	-1.798	0.0722	
	Sea (North)	1.14057	1.03903	1.714	0.2723	
	Polluted	-1.56631	0.75596	-2.142	0.0383	*
Flounder	(Intercept)	-23.9688	12.3031	-1.948	0.0514	
	Metals	-2.0686	1.4950	-1.384	0.1665	
	Hydrocarbons	-0.5449	0.6646	-1.820	0.4123	
	Age	0.9142	0.4362	2.096	0.0361	*
	Oxidative Stress	3.1787	1.9040	1.669	0.0950	
	Sea (North)	1.6869	1.5987	1.055	0.2913	
	Polluted	-3.1224	1.4183	-2.201	0.0277	*

446

447

448

449


450

451 Figure 1: A: The prevalence of cancer and foci of cellular alterations (FCA) in flounder (n=40)

452 and dab (n=88) livers. Individual numbers on bars. B: PCA loadings for metallic trace elements

453 and polycyclic aromatic hydrocarbons (PAHs).

454



455

456

457 Figure 2: Mean (log-transformed) oxidative DNA damage between fish with neoplasms, foci

458 of cellular alterations (FCA) or neither. Asterix (*) represents $p < 0.05$. Bars represent mean \pm

459 standard deviation. Dots are individual data points.

460

