



**HAL**  
open science

## Water vapor in the Martian atmosphere by SPICAM IR/Mars-Express

A. Trokhimovskiy, A. A. Fedorova, O. I. Korablev, Jean-Loup Bertaux, Eric  
Villard, A. A. Rodin

► **To cite this version:**

A. Trokhimovskiy, A. A. Fedorova, O. I. Korablev, Jean-Loup Bertaux, Eric Villard, et al.. Water vapor in the Martian atmosphere by SPICAM IR/Mars-Express. European Planetary Science Congress 2009, Sep 2009, Potsdam (Germany), Germany. pp.568. hal-04113560

**HAL Id: hal-04113560**

**<https://hal.science/hal-04113560v1>**

Submitted on 8 Jun 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Water vapor in the Martian atmosphere by SPICAM IR/Mars-Express

**A.Y. Trokhimovskiy** (1), A.A. Fedorova (1), O.I. Korablev (1), J.-L. Bertaux (2,3), E. Villard (2,3), A.V. Rodin (1)

(1) Space Research Institute (IKI), Moscow, Russia, (2) Service d'Aéronomie du CNRS/IPSL, Verrières-le-Buisson, France, (3) Institut Pierre Simon Laplace, Université de Versailles-Saint-Quentin, Saint-Quentin en Yvelines, France, (trokh@iki.rssi.ru / Fax: +7-495-3332102)

### Introduction

SPICAM experiment along with PFS and OMEGA spectrometers on Mars Express has a capability to sound the water vapor in the atmosphere. The results of H<sub>2</sub>O measurements have been intensively published during last years [1-6]. Here we present the new analysis of SPICAM IR water vapor measurements, covering two Martian years. The near-IR channel of SPICAM experiment on Mars Express spacecraft is a 800-g acousto-optic tunable filter (AOTF)-based spectrometer operating in the spectral range of 1-1.7  $\mu\text{m}$  with resolving power of  $\sim 2000$  [7, 8]. The nadir measurements of H<sub>2</sub>O in the 1.37- $\mu\text{m}$  spectral band is one of the main objectives of the experiment.

### Data treatment

As compared with previous analysis of water vapor presented in [4] we used the spectroscopic database HITRAN2004 [9] instead of HITRAN 2000 and the most recent measurements of the water line-width broadening in CO<sub>2</sub> atmosphere. Latest version HITRAN2008 doesn't have any meaningful changes in water vapour lines.

The new version of the Martian Climate Database V4.2 [10] was adopted for modelling of synthetic spectra and a scenario based on TES MY24 was used.

The spare model of SPICAM IR instrument was recalibrated in June 2007 in Reims, to analyze specifically the sensitivity to the H<sub>2</sub>O vapor band. According to laboratory measurements, a leakage from the AOTF is responsible up to 5% of signal in sharp absorption features, making the apparent depth of the H<sub>2</sub>O band lower.

### Radiative transfer modelling

Sensitivity of retrieval to aerosol scattering and different vertical distributions of aerosol and water vapor was analyzed for H<sub>2</sub>O absorption band at 1.38  $\mu\text{m}$  and 2.56  $\mu\text{m}$  for different dust particles. Dependences of equivalent width of the H<sub>2</sub>O band on the water vapor abundance and aerosol optical depth for different vertical distribution of water vapor and aerosol optical depth are obtained. Calculations of Martian atmospheric dust optical properties are done as well. The aerosol scattering account for analysis of the bulk of SPICAM data is now in progress.

### Results

Here we present the results from January 2004 (L<sub>s</sub>= 330°, MY26) to January 2008 (L<sub>s</sub>= 150°, MY29), i.e. two and a half Martian years, in the assumption of clear atmosphere. The seasonal trend of water vapor obtained by SPICAM IR is consistent with TES results and disagrees with MAWD South pole maximum measurements. The maximum abundance is 50-55 pr.  $\mu\text{m}$  at the North pole (during MY28 data are missing) and 13-16 pr.  $\mu\text{m}$  at the South pole. The northern tropical maximum amounts to 11-14 pr.  $\mu\text{m}$ . The seasonal trend of water vapor obtained by SPICAM IR is consistent for MY27 with TES results [11]. The South Pole maximum for MY28 agrees well with the MAWD South Pole measurements in 1977 [12]. It assumes the same dust conditions and global dust storm happened at MY28 L<sub>s</sub> 270° like during the MAWD observations. The maximum near 30-60S at L<sub>s</sub> 260° relates to Hellas observations. Recent observations of water vapour distribution during the same period by CRISM spectrometer onboard Mars Reconnaissance Orbiter support these results [6]




Figure 1: A seasonal map of the H<sub>2</sub>O distribution by SPICAM for three Martian years

[12] Jakosky, B. M., and C. B. Farmer, (1982), *J. Geophys. Res.*, **87**, B4, 2999-3019.

### Acknowledge

Russian team acknowledges support from RFBR grants 07-02-00850 and 06-02-72563.

### References

- [1] Fouchet, T., (2007), *Icarus* **190**, 32-49.
- [2] Melchiorri, R. (2007), *PSS* **55**, 333-342.
- [3] Encrenaz, Th. (2005), *A&A* **441**, L9-L12.
- [4] Fedorova, A. et al. (2006), *JGR* **111**, DOI:10.1029/2006JE002695.
- [5] Maltagliati, L. et al. (2009), *Icarus*, *in press*.
- [6] Smith, M. et al. (2009), *JGR*, *in press*.
- [7] Bertaux, J.-L. et al. (2006), *JGR* **111**, DOI:10.1029/2006JE002690.
- [8] Korablev, O. et al. (2006), *JGR* **111**, DOI:10.1029/2006JE002696.
- [9] Rothman, L.S. et al. (2005), *JQSRT*, **96**, 139-204.
- [10] Forget, F. et al. (2007), LPICo1353.3098F.
- [11] Smith, M., (2004), *Icarus* **167**, 148-165.