



HAL
open science

Why Do University Students Smoke Tobacco? French Validity of Brief Wisconsin Inventory of Smoking Dependence Motives among (non-) Daily Smokers and Associations with Psychological Variables

Maxime Mauduy, Nicolas Mauny, Helene Beaunieux, Jessica Mange

► To cite this version:

Maxime Mauduy, Nicolas Mauny, Helene Beaunieux, Jessica Mange. Why Do University Students Smoke Tobacco? French Validity of Brief Wisconsin Inventory of Smoking Dependence Motives among (non-) Daily Smokers and Associations with Psychological Variables. 2023. hal-04120073

HAL Id: hal-04120073

<https://hal.science/hal-04120073>

Preprint submitted on 7 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Why Do University Students Smoke Tobacco?

French Validity of Brief Wisconsin Inventory of Smoking Dependence Motives among (non-) Daily Smokers and Associations with Psychological Variables

Maxime Mauduy^{ab}, Nicolas Mauny^a, H  l  ne Beaunieux^a, and Jessica Mange^a

^a*Psychology, University of Caen Normandy, Laboratoire de Psychologie Caen Normandie (LPCN, EA 7452), Caen, France*

^b*University of Lausanne, Facult   des sciences sociales et politiques, 1015 Lausanne, Suisse*

Author note

Supplementary material is available online at

https://osf.io/2zh6c/?view_only=76269d9b547642d6849a291e3ee24212.

Maxime Mauduy, ORCID iD: <https://orcid.org/0000-0002-4604-5314>.

Nicolas Mauny, ORCID iD: <https://orcid.org/0000-0001-5156-3530>

H  l  ne Beaunieux, ORCID iD: <https://orcid.org/0000-0003-1354-6558>

Jessica Mange, ORCID iD: <https://orcid.org/0000-0001-6279-4721>

Correspondence concerning this article should be addressed to Maxime Mauduy, Universit   de Caen Normandie, UNICAEN, Laboratoire de Psychologie Caen Normandie (LPCN, EA 7452), Esplanade de la Paix, 14032 CAEN cedex 5 – France. Email: maxime.mauduy@unicaen.fr. Phone Number: +33659576492

Abstract

Introduction: This study investigates the validity of the Brief Wisconsin Inventory of Smoking Dependence Motives (B-WISDM) among French daily and non-daily university student smokers, and its associations with psychological variables related to smoking, namely attitude, social norms, perceived behavioral control, identity, and psychopathological variables, namely eating disorders, anxiety, and depression.

Methods: We tested three measurement models of the B-WISDM using robust confirmatory factor analysis, measurement invariance for daily vs. non-daily smokers, and predictive and convergent validity using a bootstrap multivariate regression model with tobacco dependence and psychological variables as outcomes.

Results: The results (1) confirmed the B-WISDM dimensionality in 11 first-order intercorrelated factors among university students; (2) showed the B-WISDM measurement invariance for the types of smokers; (3) showed that tobacco dependence is only and positively predicted by primary dependence motives, namely tolerance, craving, and lack of control, which confirm that they are core components of tobacco dependence; and (4) highlighted specific associations between smoking motives and psychological variables, such as smoking identity and perceived behavioral control with some primary dependence motives, social goals with social norms, weight control with eating disorders, and affiliative attachment and affective enhancement with psychopathological variables.

Conclusions: In addition to showing the French B-WISDM good psychometric qualities among university students, this study extends the B-WISDM external validity to non-daily smokers, meaning that tobacco dependence motivations are conceptually identical for (non-)daily student smokers, and shows specific associations of psychological variables with some smoking motives, which is of great interest for inducing smoking reduction.

Keywords: Brief Wisconsin Inventory of Smoking Dependence Motives, University student, Smoking motives, (Non-)Daily smokers, Psychological variables

Introduction

Given the consequences it has on health, tobacco use is considered a global health problem (World Health Organization, 2018). Tobacco dependence, generally defined as the experience of a strong need to smoke (West, 2017), can lead to neuropsychological impairments even among young people (Chamberlain et al., 2012). Therefore, young people, especially university students, constitute a particularly interesting population group for researchers because, they are at risk of developing tobacco dependence in decades to come (Kenford et al., 2005; Schulenberg et al., 2019). In France, 28% of students are smokers (Pasquereau et al., 2017), and among a sample of French student smokers, 15% are already heavily tobacco dependent (Mauduy et al., 2022). Knowing that a basic step for helping people quit smoking consists of identifying their underlying motivations for tobacco dependence, this study focuses on university students' smoking motives.

Drawing on a large body of work suggesting that dependence is a multidimensional and motivational phenomenon (e.g., Colby et al., 2000), Piper et al. (2004) assumed that tobacco dependence could be measured on the basis of the different motivations underlying tobacco use. As a result, the Wisconsin Inventory of Smoking Dependence Motives (WISDM, 68-item, Piper et al., 2004), and later on, its 37-item brief version (the B-WISDM, Smith et al., 2010) were developed. Numerous studies have been conducted on heavy and daily tobacco users to test the psychometric qualities of the WISDM (e.g., Piasecki et al., 2010a, 2010b, 2011; Piper et al., 2008; Shenassa et al., 2009) and the B-WISDM (e.g., Ma et al., 2012; Smith et al., 2010; Tombor et al., 2010). Researchers prefer this shorter version for practical purposes, but also because it shows good psychometric qualities. First, its dimensionality on 11 motivational factors has been confirmed, namely, loss of control, automaticity, craving, tolerance, affiliative attachment, cognitive enhancement, cue exposure/associative processes, social/environmental goals, affective enhancement, taste, and weight control (Pancani et al., 2015). In their study,

Pancani et al. (2015) compared a model with these 11 first-order factors with error covariances to a model with 11 first-order and 2 second-order factors with error covariances. The rationale for testing these two second-order factors lies in the early work of Piper et al. (2008) which highlights two subscales of the WISDM, namely primary dependence motives (PDM) and secondary dependence motives (SDM). PDM represent the core features of advanced tobacco dependence and consist of four scales (loss of control, automaticity, craving, tolerance). SDM represents instrumental motives for tobacco use and is composed of the seven remaining scales. Pancani et al.'s (2015) results support the 11-factor first-order model. Furthermore, these 11 motivational factors have satisfactory internal consistencies (e.g., Pancani et al., 2015; Smith et al., 2010) and are conceptually similar for men and women (gender measurement invariance, Vajer et al., 2011). Finally, the B-WISDM has good predictive validity with measures of tobacco dependence, such as the Cigarette Dependence Scale (CDS; Pancani et al., 2015), the Tobacco Dependence Screener (Vajer et al., 2011), the Fagerström Test for Nicotine Dependence (FTND; Tombor et al., 2010), and biochemical measures (Smith et al., 2010). In sum, the B-WISDM is an essential tool with good psychometric properties for understanding the different motivations underlying tobacco dependence. Nevertheless, there are two elements that limit the identification of these motivations among university students with the B-WISDM.

First, although the psychometric properties of the B-WISDM have always been tested on heavy or daily smokers (e.g., Pancani et al., 2015; Vajer et al., 2011), it has never been tested on university students. This limits the B-WISDM external validity to this specific population (Vajer et al., 2011), as a majority of student smokers are not daily smokers, but rather smoke on occasions and are thus considered as “light” smokers (Levinson et al., 2007; Moran et al., 2004; Thompson et al., 2007). While the original version of the WISDM was used with college student smokers (e.g., Piasecki et al., 2011), and the B-WISDM has already been used to study the characteristics of non-daily smokers (e.g., Scheuermann, Mburu, et al., 2015; Scheuermann,

Nollen, et al., 2015) or tested in a nonclinical sample (Adkison et al., 2016), only the PDM and SDM scores were used, and neither the dimensionality nor the measurement invariance of B-WISDM according to the type of smoker (daily vs. non-daily) were pre-tested. Testing the psychometric qualities of B-WISDM with non-daily smokers is thus necessary to justify its relevance to this particular smoker population.

Second, although the B-WISDM has good psychometric properties in terms of dimensionality, internal consistency and predictive validity with tobacco dependence, researchers have shown little interest in its convergent validity. Several psychological variables associated with tobacco dependence have been identified among university students: attitude towards smoking (i.e., students' evaluation of smoking, e.g., Mao et al., 2009), perceived smoking social norms (i.e., students' perceptions about how significant others approve of smoking, e.g., Riou França et al., 2009), perceived behavioral control to resist smoking (i.e., students' perception of their ability not to smoke, e.g., Jalilian et al., 2016), smoker identity (i.e., the extent to which students view themselves as smokers, e.g., Levinson et al., 2007; Moran et al., 2004), eating disorders (Eisenberg et al., 2011), anxiety (e.g., Bierhoff et al., 2019), and depression (e.g., Schleicher et al., 2009). Hence, on the one hand, the B-WISDM highlights several motivations for tobacco dependence, and, on the other hand, several psychological variables predict tobacco dependence, but little is known about the associations between these smoking dependence motives and psychological variables. We believe these smoking motives to be associated with specific psychological variables, making these latter the psychological matter of the motivations for tobacco dependence. First, we expect both smoking identity and perceived behavioral control to be two core psychological components of the PDM (e.g., Falomir-Pichastor et al., 2020; Lee et al., 2014; Mauduy et al., 2022). Second, we expect the other psychological variables to be more related to instrumental motivations to smoke, namely SDM. Specifically, attitudes would be primarily associated with taste motives, social norms

with social goals motives, eating disorders with weight control motives, and anxiety and depression with coping motives, that is affective and cognitive enhancements, and affiliative attachment. Thus, investigating the associations between the smoking motives and the psychological variables would be of great interest because it would allow us to (1) provide additional evidence of the good psychometric properties of the B-WISDM, (2) better understand the processes underlying tobacco dependence among university students, and (3) better assist university students to quit smoking by focusing prevention efforts on the psychological processes underlying their motivations for smoking.

Thus, the present study aims to test the validity of the B-WISDM among a French university student population. For this purpose, we test the B-WISDM dimensionality in 11 first-order motivational factors, its measurement invariance for types of smokers, namely daily and non-daily smokers, as well as its concurrent validity with tobacco dependence and convergent validity with several psychosocial variables already identified in the literature as being associated with tobacco dependence.

Methods

Procedure and Participants

This study was carried out on a convenience sample of 687 student smokers from the University of Caen Normandy (France). The participants were recruited per mail on their institutional address and were asked if they wished to participate in an online survey on tobacco smoking (via the Limesurvey® application, November 2021). No compensation was provided to the participants.

Ethics

All participants were volunteers and gave their consent before starting the survey. The study was notified to and authorized by the “Commission Nationale de l’Informatique et des Libertés” with the registration number u24- 20171109-01R1. Besides, the participants’ anonymity was guaranteed by the University Information System Direction (DSI). This survey was conducted in full agreement with the Declaration of Helsinki (2008) and the ethical standards set by the Psychology Department, that follows the American Psychological Association Ethical Principles of Psychologists and the Code of Conduct (APA, 2017) for the ethical treatment of human participants.

Measures

Details of all measures and items described below are available on the supplemental online material at https://osf.io/2zh6c/?view_only=76269d9b547642d6849a291e3ee24212.

Demographics and Smoking History

We measured gender, age, academic level, age of smoking onset, smoking parents, cigarette consumption per day, recent attempts to quit smoking (Perski et al., 2018), and motivation to quit smoking (MTSS, Kotz et al., 2013).

French Brief Wisconsin Inventory of Smoking Dependence Motives

The B-WISDM, developed by Smith et al. (2010), was translated into French and then back-translated into English by a professional translation service for verification purposes. The B-WISDM is a questionnaire composed of 37 items rated on a 7-point scale ranging from 1 = *Not true for me at all* to 7 = *Extremely true for me*. It comprised 11 subscales that represent different smoking dependence motivations (Smith et al., 2010; Pancani et al., 2015): loss of control, automaticity, craving, tolerance, affiliative attachment, cognitive enhancement, cue

exposure/associative processes, social/environmental goals, affective enhancement, taste, and weight control (see the *supplemental material* for the items translated).

Concurrent Validity Measures

Tobacco dependence was measured using the validated French-language Cigarette Dependence Scale (CDS-5; Etter, 2005). This 5-item self-report questionnaire provides a continuous score for cigarette addiction. The CDS has shown high internal consistency, a good predictive validity, as well as a high test-retest reliability (e.g., Etter, 2005, 2008; Etter et al., 2003), thus overcoming the psychometric limitations of the FTND.

Convergent Validity Measures

Attitudes, perceived social norms and perceived behavioral control to resist smoking were assessed based on the Theory of Planned Behavior (TPB, Ajzen, 1991; Jalilian et al., 2016; Mauduy et al., 2022). The assessment included *attitude* towards smoking (4-item, Cronbach $\alpha = .92$), perceived social norms for smoking (3-item, Cronbach $\alpha = .81$), and perceived behavioral control to resist smoking (3-item, Cronbach $\alpha = .75$) rated on Likert-type scale scored from 1 = *do not agree* to 5 = *strongly agree*. This scale measuring three factors of the TPB has acceptable fits in French (Mauduy et al., 2022).

Smoker identity was assessed using the Smoker Self-Concept Scale (SSCS; 5-item Likert-type scale scored from 1 = *do not agree* to 5 = *strongly agree* ; Cronbach $\alpha = .85$, Shadel & Mermelstein, 1996). The SSCS assesses the importance of being a smoker for one's self-concept and has shown great predictive and discriminant validities, a good internal consistency (Shadel & Mermelstein, 1996), and excellent fits in French (Mauduy et al., 2022).

Eating disorders were assessed with the French version of the SCOFF questionnaire (Parker et al., 2010). It consists in responding to five questions related to food in a yes/no format. Scores can range from 0 to 5 (i.e., answer yes to all 5 questions), with a score of 2 or higher indicating a likely case of an eating disorder.

Anxiety was measured with the French version of the State-Trait Anxiety Inventory (STAI, Spielberger et al., 1983) which is composed of 20 items (Cronbach $\alpha = .91$) on a 4-point scale ranging from no (1) to yes (4). It has shown good validity and acceptable reliability (Barnes et al., 2002).

Depression was measured with the short French version of the Beck Depression Inventory (BDI, Beck et al., 1988). It consists of 13 items, each composed of four statements reflecting various severity degrees of depression-related symptoms. The BDI has shown high internal consistency, good predictive validity, and high test-retest reliability (Beck et al., 1988; Beck & Steer, 1984).

Statistical Analyses

First, the dimensionality and reliability of the B-WISDM have been tested. Three models derived from the literature were tested using a robust confirmatory factor analysis (CFA). Model 1 contained two first-order correlated factors, namely the PDM and SDM. Model 2 contained 11 first-order correlated factors. Model 3 included 11 first-order correlated factors (as model 2) and 2 second-order factors, namely PDM and SDM. Robust CFA were used to manage the multivariate non-normality of the data (i.e., Kurtosis test > 20 , Mardia, 1974). As standard maximum likelihood estimation assumes multivariate normality, we used the maximum likelihood estimation with robust (Huber-White) standard errors. Five fit indices were used to evaluate each model. The Satorra-Bentler Chi-square (χ^2) test indicate the fit goodness of the model when the value is statistically significant. The robust comparative fit index (CFI) and robust Tucker-Lewis index must be greater than 0.95 to demonstrate a good model fit, but values above 0.90 are still considered as acceptable ((Boateng et al., 2018; Brown, 2006). The robust root mean squared error of approximation (RMSEA) indicates a good model fit when its values are lower than 0.05 and an adequate fit when they are lower than 0.08. The standardized root mean square residual (SRMR) had to be lower than 0.08 (Boateng et al.,

2018). To identify whether one more restricted model has a better fit than another, three criteria have to be met. First, we compared the models using the specialized computation for the Satorra-Bentler χ^2 difference (Satorra & Bentler, 2001). The value of the χ^2 difference must be significant (i.e., $p < .05$). However, since the χ^2 test may be sensitive to the sample size, we also used CFI and RMSEA values. The CFI must show an increase higher than .01 ($CFI_{diff} > .01$, Cheung & Rensvold, 2002), and the RMSEA must show a decrease of over 0.015 ($RMSEA_{diff} < .015$, Chen, 2007). The internal consistencies for the optimal model identified were then calculated using Cronbach's alpha. Values above 0.90 are considered excellent, and values above 0.70 are considered satisfactory (Boateng et al., 2018; Nunnally & Bernstein, 1994).

Second, to test the measurement invariance of types of smokers (daily and non-daily), we conducted a series of multigroup CFAs on the previously identified model. Three nested models with increasing constraints were estimated. First, the configural invariance, which corresponds to a model without constraints, tests whether the dimensionality of the FB-WISDM and the item composition of the factors are identical between groups. A configural invariance between groups indicates that the groups (non-daily or daily smokers) have a similar representation of the underlying construct. Second, the metric invariance, which constrained the factor loadings to be identical in all groups, tests whether the contribution of each item to a factor (as measured by the factor loadings or standardized regression coefficient) is identical in all groups. Metric invariance indicates that the understanding of the content of each item is not influenced by the type of smokers. Third, scalar invariance, which constrained both factor loadings and intercepts to be identical in all groups, tests whether the intercept for each item is not influenced by the groups. Scalar invariance indicates that the differences between groups on the means of measure variables (i.e., items) are due to group differences on the mean of the common latent variable (i.e., factor). Thus, scalar invariance is required to compare groups in terms of latent means. To test measurement invariance, we compared configural to metric

invariance, and then, if metric invariance is retained, we compared metric to scalar invariance. To show measurement invariance, at least one of the three following criteria had to be met. First, the specialized computation for the Satorra-Bentler χ^2 difference between the two models has to be not significant, second, the CFI_{diff} between the two models should be less than or equal to .01, and third, the $RMSEA_{diff}$ between the two models should be less than or equal to .015. It should be noted that, since the Satorra-Bentler χ^2 difference test may be sensitive to the sample size, more attention has been given to the CFI_{diff} and $RMSEA_{diff}$ indicators to evaluate the measurement invariance. Furthermore, we also tested the measurement invariance for gender (men and women) in order to verify that the gender invariance previously shown among adult daily smokers (Vajer et al., 2011) is replicated among university students.

Third, the concurrent and convergent validity were tested using a multivariate regression model that includes all the FB-WISDM dimensions of the identified model as predictors, and tobacco dependence, attitudes, social norms, perceived behavioral control to resist smoking, smoker identity, eating disorders, anxiety, and depression as outcome variables. The Bootstrap method ($N = 1000$) was used to assess the stability of the results.

All statistical analyses were conducted using the R software (version 4.05; R Core Team). The package *mvn* (Korkmaz et al., 2014) was used to assess the multivariate nonnormality, the package *lavaan* (Rosseel, 2012) to test the dimensionality, the measurement invariance, the reliability and concurrent and convergent validities of the FB-WISDM.

Results

Characteristics of the Sample

Table 1 presents the characteristics of the sample. Students were considered current smokers when they indicated that they had already smoked 100 cigarettes in their lifetime and had smoked at least one time in the past 30 days (Adkison et al., 2016; Riou França et al., 2009).

Among the 687 smokers ($M_{age} = 20.06$, $SD = 2.39$), 63.5% were female (2.2% did not choose an answer). On average, they started smoking at age 16.1 years ($SD = 2.03$) and currently smoke 6.34 cigarettes per day ($SD = 16.3$). A majority of participants are non-daily smokers (63.5%). Finally, almost half of the participants reported having made one attempt to quit during the past year (41.4%) and are motivated to quit smoking (42.6%).

Tests of FB-WISDM Dimensionality

Among the 687 participants, 29 had at least one missing data on the FB-WISDM. To address the problem of missing data, we used the full-information maximum likelihood estimation (T. Lee & Shi, 2021).

Robust Confirmatory Factor Analyses

The fit indices of the three tested model are depicted in Table 2. Robust CFAs of Model 1 indicated an inadequate fit with the data, while Model 2 and Model 3 displayed acceptable fit indices. The specialized computation for the Satorra–Bentler χ^2 difference as well as fit indices indicated that Model 2 fitted significantly better the data than Model 1 ($\chi^2_{diff} = 1721.60$, $df_{diff} = 11$, $p < .001$; $CFI_{diff} = 0.271$; $RMSEA_{diff} = -0.061$), indicating that a two-dimension scale , namely, the composite of four subscales related to the PDM and the composite of nine subscales related to the SDM, is not adequate to account for the data. Model 2 showed a significantly better fit than Model 3 on Satorra–Bentler χ^2 difference ($\chi^2_{diff} = 277.91$, $df_{diff} = 43$, $p < .001$) but not on CFI and RMSEA indices ($CFI_{diff} = 0.015$; $RMSEA_{diff} = 0.004$). These results suggest that the 11 first-order dimensions of the FB-WISDM (Model 2) represent the optimal solution, although we cannot reject Model 3. We further compared models 2 and 3 by examining the factor loadings and correlations between factors. No differences were found between the two models for the first-order factors: they displayed standardized factor loadings that are all above 0.60 as well as positive and significant correlations between all factors. However, two elements

weaken the relevance of Model 3. First, the loadings of the second-order factors are different, with PDM values ranging from 0.77 to 0.98 and SDM ranging from 0.27 to 0.86. The dimensionality of the SDM is challenged by the three following sub-scales: social/environmental goals, taste and weight control. Second, discriminant validity is also challenged by the high correlations between the PDM and SDM factors ($r = 0.86$). Thus, we consider Model 2 as the optimal solution to fit the data. The results of the robust CFA for Model 2 are displayed in Table 3.

Cronbach's Alpha

All cronbach's alpha values of the 11 first-order factors were higher than 0.70 (see Table 3), indicating an excellent internal consistency of the FB-WISDM.

Tests of Measurement Invariances through Multiple Group Robust CFA

Multiple group CFA were used to test the measurement invariance (configural, metric, scalar) of the FB-WISDM for types of smokers (daily and non-daily smokers) and gender (men and women).

Measurement Invariance for Types of Smokers

For all three invariance levels of types of smokers, the models have good fit indices (see Table 4). The Satorra–Bentler χ^2 difference between the first model (configural invariance) and the second one (metric invariance) is significant ($\chi^2_{\text{diff}} = 137.80$, $df_{\text{diff}} = 37$, $p < .001$) but CFI and RMSEA values do not differ ($CFI_{\text{diff}} = .005$; $RMSEA_{\text{diff}} = .001$). Then, Satorra–Bentler χ^2 difference between the second and the third model (scalar invariance) is significant ($\chi^2_{\text{diff}} = 92.62$, $df_{\text{diff}} = 26$, $p < .001$), but there was no significant change in fit indices ($CFI_{\text{diff}} = .005$; $RMSEA_{\text{diff}} = .001$). Thus, scalar invariance is not rejected, which means that the factor loadings and intercepts of the 11 first-order factors are invariant between types of smokers. Concretely,

daily and non-daily smokers have a similar understanding of the different smoking motivations measured by the FB-WISDM.

Measurement Invariance for Gender

For all three gender invariance levels, the models have good fit indices (see Table 4). The Satorra–Bentler χ^2 difference between the first model (configural invariance) and the second one (metric invariance) is significant ($\chi^2_{\text{diff}} = 55.11$, $df_{\text{diff}} = 37$, $p = .028$) but the CFI and RMSEA values do not differ. Then, fit indices did not change between the second and the third model (scalar invariance, $\chi^2_{\text{diff}} = 29.96$, $df_{\text{diff}} = 26$, $p = .27$; $CFI_{\text{diff}} = 0.0$; $RMSEA_{\text{diff}} = 0.001$). Thus, scalar invariance is not rejected, which means that the factor loadings and intercepts of the 11 first-order factors are gender invariant. Men and women have a similar understanding of the different motivations for smoking

Tests of FB-WISDM Concurrent and Convergent Validity

The results, displayed in Table 5, support both concurrent and convergent validity of the FB-WISDM. First, tobacco dependence is mainly and positively predicted by PDM, namely, lack of control, craving, and tolerance, and is also predicted by associative processes. Second, attitudes are mainly and positively associated with taste and affective enhancement, and, to a lesser extent, negatively associated with lack of control; social norms are only positively associated with social goals; perceived behavioral control is mainly and negatively associated with craving and associative processes; smoker identity is mainly associated with lack of control and affiliative attachment, and, to a lesser extent, associative processes and tolerance; eating disorders are only associated with weight control; anxiety is mainly associated with affective enhancement, and, to a lesser extent, with affiliative attachment, weight control and negatively associated with tolerance; and depression is mainly associated with affective

enhancement, weight control and affiliative attachment, and, to a lesser extent, negatively associated with taste.

Discussion

This study aimed at testing the psychometric qualities of the B-WISDM on a sample of daily and non-daily university student smokers. The questionnaire had already been validated but only on heavy or daily adult smokers. Overall, the results highlight the good psychometric qualities of the FB-WISDM among university students. In line with previous studies, our results confirm the dimensionality in 11 first-order motivational factors, their satisfactory internal consistencies (Ma e al., 2012; Pancani et al., 2015; Smith et al., 2010; Vajer et al., 2011) and gender invariance (Vajer et al., 2011). On the other hand, and beyond being having carried out the first study to test the validity of the B-WISDM on university students, we have also contributed to the literature by highlighting two new findings that we will discuss.

First, this study highlights the applicability of the B-WISDM to a non-daily, occasional smoking population. Tested only among heavy or daily smokers (e.g., Pancani et al., 2015; Smith et al., 2010; Vajer et al., 2011), the validity of the B-WISDM among occasional smokers was lacking. Based on measurement invariance by types of smokers, this study is the first one to show that motivations underlying tobacco dependence are conceptually identical for daily and non-daily university student smokers. This result, beyond extending the applicability of the B-WISDM to different types of smokers, is of great interest because it would allow us to investigate motivational changes explaining the development of daily smoking in occasional smokers (Kenford et al., 2005; Schulenberg et al., 2019), using the B-WISDM.

Second, this study highlights the convergent validity of the B-WISDM with psychological variables, well-known to be associated with tobacco use and dependence. This result is particularly interesting for prevention purposes, as it potentially makes the B-WISDM a very useful diagnostic tool to identify the underlying psychological reasons for an individual's

smoking, in the first stage of prevention. Then, in the second stage, the B-WISDM would be able to guide prevention actions targeting the psychological variables that are specifically associated with people's reasons for smoking through the use of adapted strategies. In our results, two categories of psychological variables seem to emerge, depending on whether they are more associated with PDM or with SDM. On the one hand, smoker identity and perceived behavioral control to resist smoking are mainly associated with motivations related to lack of control, tolerance and craving, thus accounting for three out of four PDM. These findings are not surprising considering that perceived behavioral control (Lee et al., 2014; Mauduy et al., 2022) and smoker identity (Falomir-Pichastor et al., 2020; Mauduy et al., 2022) are key determinants of tobacco dependence and cessation success. As PDM are the core components of tobacco dependence (Piper et al., 2008), it seems essential for any tobacco dependence prevention program to target these two psychological variables. The development of new prevention strategies, such as the social modeling technique (Burn, 1991; see Webb & Sheeran, 2006 for a meta-analysis) for increasing people's confidence to resist smoking, or the multi-categorization technique (Kang & Bodenhausen, 2015) for encouraging people who smoke to define themselves not only as smokers but also as members of other social groups and then reduce the potential role of the problematic identity in driving behaviors, could be considered for this purpose. On the other hand, attitude towards smoking, perceived smoking social norms, eating disorders, anxiety, and depression are uniquely associated with specific SDM. Thus, our results would suggest targeting positive smoking attitudes among student smokers because of taste and affective enhancement, with for instance the framing technique (Gallagher & Updegraff, 2012), the perceived smoking social norms because of social goals, with the personalized normative feedback technique (Vallentin-Holbech et al., 2018), eating disorders because of weigh control, with dissonance-based prevention programs (see Stice et al., 2019

for a meta-analytic review), and anxiety and depression because of affiliative attachment and affective enhancement, with mindfulness-based interventions (Hofmann & Gómez, 2017).

This study, however, has two main limitations that we must highlight. First, a biochemical measure of tobacco dependence (Bize et al., 2009) or a combination of measures (Hughes et al., 2004) would have allowed us to ensure that the participants are actually smokers. Second, given that our study was conducted with French students and that it is a correlational study, further research could attempt to replicate our results.

In conclusion, after having been validated on heavy smokers or daily smokers in English, Italian and Hungarian participants, this study shows that the B-WISDM has also good psychometric qualities among French daily and non-daily university student smokers. Examining its convergent validity with different psychological variables (attitude, social norms, perceived behavioral control, smoker identity, and eating disorders) has provided a better understanding of the underlying motivations for tobacco dependence and a guide for smoking reduction efforts among university students. Future research must now extend the external validity of the B-WISDM to other populations of non-daily smokers, and investigate the involvement of other psychological variables in understanding motivations underlying tobacco dependence.

Funding

The work reported was supported by RIN Tremplin Grant 19E00906 of Normandie Région, France.

Declaration of Interest Statement

The authors report no conflict of interest.

Data Availability Statement

Data from this study will be provided by the first author upon request.

References

- Adkison, S. E., Rees, V. W., Bansal-Travers, M., Hatsukami, D. K., & O'Connor, R. J. (2016). Psychometric Characteristics of the Brief Wisconsin Inventory of Smoking Dependence Motives Among a Nonclinical Sample of Smokers. *Nicotine & Tobacco Research, 18*(4), 470–476. <https://doi.org/10.1093/ntr/ntv113>
- Ajzen, I. (1991). The theory of planned behavior. *Organizational Behavior and Human Decision Processes, 50*(2), 179–211. [https://doi.org/10.1016/0749-5978\(91\)90020-T](https://doi.org/10.1016/0749-5978(91)90020-T)
- Barnes, L. L. B., Harp, D., & Jung, W. S. (2002). Reliability generalization of scores on the Spielberger state-trait anxiety inventory. *Educational and Psychological Measurement, 62*(4), 603–618.
- Beck, A. T., & Steer, R. A. (1984). Internal consistencies of the original and revised Beck Depression Inventory. *Journal of Clinical Psychology, 40*(6), 1365–1367.
- Beck, A. T., Steer, R. A., & Carbin, M. G. (1988). Psychometric properties of the Beck Depression Inventory: Twenty-five years of evaluation. *Clinical Psychology Review, 8*(1), 77–100. [https://doi.org/10.1016/0272-7358\(88\)90050-5](https://doi.org/10.1016/0272-7358(88)90050-5)
- Bierhoff, J., Haardörfer, R., Windle, M., & Berg, C. J. (2019). Psychological Risk Factors for Alcohol, Cannabis, and Various Tobacco Use among Young Adults: A Longitudinal Analysis. *Substance Use and Misuse, 54*(8), 1365–1375. <https://doi.org/10.1080/10826084.2019.1581220>
- Bize, R., Burnand, B., Mueller, Y., Rège Walther, M., & Cornuz, J. (2009). Biomedical risk assessment as an aid for smoking cessation. In R. Bize (Ed.), *Cochrane Database of Systematic Reviews* (Issue 2). John Wiley & Sons, Ltd. <https://doi.org/10.1002/14651858.CD004705.pub3>
- Boateng, G. O., Neilands, T. B., Frongillo, E. A., Melgar-Quiñonez, H. R., & Young, S. L. (2018). Best Practices for Developing and Validating Scales for Health, Social, and Behavioral Research: A Primer. *Frontiers in Public Health, 6*(June), 1–18. <https://doi.org/10.3389/fpubh.2018.00149>

- Brown, T. A. (2006). *Confirmatory Factor Analysis for Applied Research*. Guilford Publications.
- Burn, S. M. (1991). Social Psychology and the Stimulation of Recycling Behaviors: The Block Leader Approach. *Journal of Applied Social Psychology, 21*(8), 611–629. <https://doi.org/10.1111/j.1559-1816.1991.tb00539.x>
- Chamberlain, S. R., Odlaug, B. L., Schreiber, L. R. N., & Grant, J. E. (2012). Association between Tobacco Smoking and Cognitive Functioning in Young Adults. *American Journal on Addictions, 21*, S14–S19. <https://doi.org/10.1111/j.1521-0391.2012.00290.x>
- Cheung, G. W., & Rensvold, R. B. (2002). Evaluating Goodness-of-Fit Indexes for Testing Measurement Invariance Evaluating Goodness-of-Fit Indexes for Testing Measurement Invariance. *Structural Equation Modeling, 9*(2), 233–255. <https://doi.org/10.1207/S15328007SEM09025>
- Colby, S. M., Tiffany, S. T., Shiffman, S., & Niaura, R. S. (2000). Measuring nicotine dependence among youth: A review of available approaches and instruments. *Drug and Alcohol Dependence, 59*(SUPPL. 1), 23–39. [https://doi.org/10.1016/S0376-8716\(99\)00163-5](https://doi.org/10.1016/S0376-8716(99)00163-5)
- Eisenberg, D., Nicklett, E. J., Roeder, K., & Kirz, N. E. (2011). Eating disorder symptoms among college students: Prevalence, persistence, correlates, and treatment-seeking. *Journal of American College Health, 59*(8), 700–707. <https://doi.org/10.1080/07448481.2010.546461>
- Etter, J.-F. (2005). A comparison of the content-, construct- and predictive validity of the cigarette dependence scale and the Fagerström test for nicotine dependence. *Drug and Alcohol Dependence, 77*(3), 259–268. <https://doi.org/10.1016/j.drugalcdep.2004.08.015>
- Etter, J.-F. (2008). Comparing the validity of the Cigarette Dependence Scale and the Fagerström Test for Nicotine Dependence. *Drug and Alcohol Dependence, 95*(1–2), 152–159. <https://doi.org/10.1016/j.drugalcdep.2008.01.017>
- Etter, J.-F., Houezec, J. Le, & Perneger, T. V. (2003). A self-administered questionnaire to measure

- dependence on cigarettes: The cigarette dependence scale. *Neuropsychopharmacology*, 28(2), 359–370. <https://doi.org/10.1038/sj.npp.1300030>
- Falomir-Pichastor, J. M., Blondé, J., Desrichard, O., Felder, M., Riedo, G., & Folly, L. (2020). Tobacco dependence and smoking cessation: The mediating role of smoker and ex-smoker self-concepts. *Addictive Behaviors*, 102, 106200. <https://doi.org/10.1016/j.addbeh.2019.106200>
- Gallagher, K. M., & Updegraff, J. A. (2012). Health Message Framing Effects on Attitudes, Intentions, and Behavior: A Meta-analytic Review. *Annals of Behavioral Medicine*, 43(1), 101–116. <https://doi.org/10.1007/s12160-011-9308-7>
- Hofmann, S. G., & Gómez, A. F. (2017). Mindfulness-Based Interventions for Anxiety and Depression. *Psychiatric Clinics of North America*, 40(4), 739–749. <https://doi.org/10.1016/j.psc.2017.08.008>
- Hughes, J. R., Oliveto, A. H., Riggs, R., Kenny, M., Liguori, A., Pillitteri, J. L., & MacLaughlin, M. A. (2004). Concordance of different measures of nicotine dependence: Two pilot studies. *Addictive Behaviors*, 29(8), 1527–1539. <https://doi.org/10.1016/j.addbeh.2004.02.031>
- Jalilian, F., Joulaei, H., Mirzaei-Alavijeh, M., Samannezhad, B., Berimvandi, P., Matin, B. K., & Mahboubi, M. (2016). Cognitive factors related to cigarettes smoking among college students: An application of theory of planned. *Social Sciences (Pakistan)*, 11(7), 1189–1193. <https://doi.org/10.3923/sscience.2016.1189.1193>
- Kang, S. K., & Bodenhausen, G. V. (2015). Multiple Identities in Social Perception and Interaction: Challenges and Opportunities. *Annual Review of Psychology*, 66(1), 547–574. <https://doi.org/10.1146/annurev-psych-010814-015025>
- Kenford, S. L., Wetter, D. W., Welsch, S. K., Smith, S. S., Fiore, M. C., & Baker, T. B. (2005). Progression of college-age cigarette samplers: What influences outcome. *Addictive Behaviors*, 30(2), 285–294. <https://doi.org/10.1016/j.addbeh.2004.05.017>

- Korkmaz, S., Goksuluk, D., & Zararsiz, G. (2014). MVN: An R Package for Assessing Multivariate Normality. *The R Journal*, 6(2), 151–162. <https://journal.r-project.org/archive/2014-2/korkmaz-goksuluk-zararsiz.pdf>.
- Kotz, D., Brown, J., & West, R. (2013). Predictive validity of the Motivation To Stop Scale (MTSS): A single-item measure of motivation to stop smoking. *Drug and Alcohol Dependence*, 128(1–2), 15–19. <https://doi.org/10.1016/j.drugalcdep.2012.07.012>
- Lee, H. S., Catley, D., & Harris, K. J. (2014). Improving understanding of the quitting process: Psychological predictors of quit attempts versus smoking cessation maintenance among college students. *Substance Use and Misuse*, 49(10), 1332–1339. <https://doi.org/10.3109/10826084.2014.901386>
- Lee, T., & Shi, D. (2021). A Comparison of Full Information Maximum Likelihood and Multiple Imputation in Structural Equation Modeling With Missing Data. *Psychological Methods*, 26(4), 466–485.
- Levinson, A. H., Campo, S., Gascoigne, J., Jolly, O., Zakharyan, A., & Tran, Z. V. (2007). Smoking, but not smokers: Identity among college students who smoke cigarettes. *Nicotine and Tobacco Research*, 9(8), 845–852. <https://doi.org/10.1080/14622200701484987>
- Ma, J. Z., Li, M. D., & Payne, T. J. (2012). Evaluation of the brief Wisconsin Inventory of smoking dependence motives in African-American and European-American heavy smokers. *Frontiers in Psychiatry*, 3(APR), 1–7. <https://doi.org/10.3389/fpsy.2012.00036>
- Mao, R., Li, X., Stanton, B., Wang, J., Hong, Y., Zhang, H., & Chen, X. (2009). Psychosocial correlates of cigarette smoking among college students in China. *Health Education Research*, 24(1), 105–118. <https://doi.org/10.1093/her/cyn002>
- Mardia, K. V. (1974). Applications of Some Measures of Multivariate Skewness and Kurtosis in Testing Normality and Robustness Studies. *The Indian Journal of Statistics*, 36(2), 115–128.

- Mauduy, M., Mauny, N., & Mange, J. (2022). Tobacco Dependence Among French University Students : A Cluster Analytic Approach to Identifying Distinct Psychological Profiles of Smokers. *Journal of Drug Issues*, 1–21. <https://doi.org/10.1177/00220426221107560>
- Moran, S., Wechsler, H., & Rigotti, N. A. (2004). Social smoking among US college students. *Pediatrics*, 114(4), 1028–1034. <https://doi.org/10.1542/peds.2003-0558-L>
- Nunnally, J. C., & Bernstein, I. (1994). *Psychometric theory* (McGraw-Hil). McGraw-Hill.
- Pancani, L., D'addario, M., Cappelletti, E. R., Greco, A., Monzani, D., & Steca, P. (2015). Smoking behavior: A cross-sectional study to assess the dimensionality of the brief Wisconsin Inventory of Smoking Dependence Motives and identify different typologies among young daily smokers. *Nicotine and Tobacco Research*, 17(1), 98–105. <https://doi.org/10.1093/ntr/ntu143>
- Parker, S. C., Lyons, J., & Bonner, J. (2010). Eating Disorders in Graduate Students: Exploring the SCOFF Questionnaire as a Simple Screening Tool. *Journal of Ame*, 54(2), 103–107. <https://doi.org/10.3200/JACH.54.2.103-107>
- Pasquereau, A., Gautier, A., Andler, R., Guignard, R., Richard, J. B., & Nguyen-Thanh, V. (2017). Tobacco and e-cigarette in France: levels of consumption according to the preliminary results from the 2016. *Health Barometer*.
- Perski, O., Herd, N., Brown, J., & West, R. (2018). Does consistent motivation to stop smoking improve the explanation of recent quit attempts beyond current motivation? A cross-sectional study. *Addictive Behaviors*, 81, 12–16. <https://doi.org/10.1016/j.addbeh.2018.01.037>
- Piasecki, T. M., Piper, M. E., & Baker, T. B. (2010a). Refining the tobacco dependence phenotype using the Wisconsin Inventory of Smoking Dependence Motives: II. Evidence from a laboratory self-administration assay. *Journal of Abnormal Psychology*, 119(3), 513–523. <https://doi.org/10.1037/a0020235>
- Piasecki, T. M., Piper, M. E., & Baker, T. B. (2010b). Tobacco dependence: Insights from

- investigations of self-reported motives. *Current Directions in Psychological Science*, 19(6), 395–401. <https://doi.org/10.1177/0963721410389460>
- Piasecki, T. M., Piper, M. E., Baker, T. B., & Hunt-Carter, E. E. (2011). WISDM primary and secondary dependence motives: Associations with self-monitored motives for smoking in two college samples. *Drug and Alcohol Dependence*, 114(2–3), 207–216. <https://doi.org/10.1016/j.drugalcdep.2010.10.005>
- Piper, M. E., Bolt, D. M., Kim, S.-Y., Japuntich, S. J., Smith, S. S., Niederdeppe, J., Cannon, D. S., & Baker, T. B. (2008). Refining the tobacco dependence phenotype using the Wisconsin Inventory of Smoking Dependence Motives. *Journal of Abnormal Psychology*, 117(4), 747–761. <https://doi.org/10.1037/a0013298>
- Piper, M. E., Piasecki, T. M., Federman, E. B., Bolt, D. M., Smith, S. S., Fiore, M. C., & Baker, T. B. (2004). A Multiple Motives Approach to Tobacco Dependence: The Wisconsin Inventory of Smoking Dependence Motives (WISDM-68). *Journal of Consulting and Clinical Psychology*, 72(2), 139–154. <https://doi.org/10.1037/0022-006X.72.2.139>
- Radzius, A., Moolchan, E. T., Henningfield, J. E., Heishman, S. J., & Gallo, J. J. (2001). A factor analysis of the Fagerstrom Tolerance Questionnaire. *Addictive Behaviors*, 26, 303–310.
- Riou França, L., Dautzenberg, B., Falissard, B., & Reynaud, M. (2009). Are social norms associated with smoking in French university students? A survey report on smoking correlates. *Substance Abuse Treatment, Prevention, and Policy*, 4(4). <https://doi.org/10.1186/1747-597X-4-4>
- Rosseel, Y. (2012). lavaan: An R Package for Structural Equation. *Journal of Statistical Software*, 48(2).
- Satorra, A., & Bentler, P. M. (2001). A scaled difference chi-square test statistic for moment structure analysis. *Psychometrika*, 66(4), 507–514.
- Scheuermann, T. S., Mburu, W. E., Mathur, C., & Ahluwalia, J. S. (2015). Correlates of converted and

native nondaily smoking. *Nicotine and Tobacco Research*, 17(9), 1112–1119.

<https://doi.org/10.1093/ntr/ntu272>

Scheuermann, T. S., Nollen, N. L., Cox, L. S., Reitzel, L. R., Berg, C. J., Guo, H., Resnicow, K., & Ahluwalia, J. S. (2015). Smoking dependence across the levels of cigarette smoking in a multiethnic sample. *Addictive Behaviors*, 43(1), 1–6.

<https://doi.org/10.1016/j.addbeh.2014.11.017>

Schleicher, H., Harris, K., Catley, D., & Nazir, N. (2009). The role of depression and negative affect regulation expectancies in tobacco smoking among college students. *Journal of American College Health*, 57(5), 507–512. <https://doi.org/10.3200/JACH.57.5.507-512>

Schulenberg, J., Johnston, L. D., O'Malley, P., Bachman, J., Miech, R. A., & Patrick, M. E. (2019). *Monitoring the Future national survey results on drug use, 1975-2018: Volume II, college students and adults ages 19-60*. (Ann Arbor). The University of Michigan.

<http://monitoringthefuture.org/pubs.html#monographs>

Shadel, W. G., & Mermelstein, R. (1996). Individual differences in self-concept among smokers attempting to quit: Validation and predictive utility of measures of the smoker self-concept and abstainer self-concept. *Annals of Behavioral Medicine*, 18(3), 151–156.

<https://doi.org/10.1007/BF02883391>

Shenassa, E. D., Graham, A. L., Burdzovic, J. B., & Buka, S. L. (2009). Psychometric properties of the Wisconsin inventory of Smoking Dependence Motives (WISDM-68): A replication and extension. *Nicotine and Tobacco Research*, 11(8), 1002–1010. <https://doi.org/10.1093/ntr/ntp109>

Smith, S. S., Piper, M. E., Bolt, D. M., Fiore, M. C., Wetter, D. W., Cinciripini, P. M., & Baker, T. B. (2010). Development of the Brief Wisconsin Inventory of Smoking Dependence Motives. *Nicotine & Tobacco Research*, 12(5), 489–499. <https://doi.org/10.1093/ntr/ntq032>

Spielberger, C. D., Gorsuch, R. L., Lushene, R., Vagg, P. R., & Jacobs, G. (1983). *Manual for the*

State-Trait Anxiety Inventory. Consulting Psychologists Press.

- Stice, E., Marti, C. N., Shaw, H., & Rohde, P. (2019). Meta-analytic review of dissonance-based eating disorder prevention programs : Intervention , participant , and facilitator features that predict larger effects. *Clinical Psychology Review, 70*(February 2018), 91–107.
<https://doi.org/10.1016/j.cpr.2019.04.004>
- Thompson, B., Coronado, G., Chen, L., Thompson, L. A., Halperin, A., Jaffe, R., McAfee, T., & Zbikowski, S. M. (2007). Prevalence and characteristics of smokers at 30 Pacific Northwest colleges and universities. *Nicotine and Tobacco Research, 9*(3), 429–438.
<https://doi.org/10.1080/14622200701188844>
- Tombor, I., Urbán, R., Berkes, T., & Demetrovics, Z. (2010). Denial of smoking-related risk among pregnant smokers. *Acta Obstetricia et Gynecologica Scandinavica, 89*(4), 524–530.
<https://doi.org/10.3109/00016341003678427>
- Vajer, P., Urbán, R., Tombor, I., Stauder, A., & Kalabay, L. (2011). Psychometric properties and construct validity of the brief wisconsin inventory of smoking dependence motives in an internet-based sample of treatment-seeking Hungarian smokers. *Nicotine and Tobacco Research, 13*(4), 273–281. <https://doi.org/10.1093/ntr/ntq254>
- Vallentin-Holbech, L., Rasmussen, B. M., & Stock, C. (2018). Effects of the social norms intervention The GOOD Life on norm perceptions, binge drinking and alcohol-related harms: A cluster-randomised controlled trial. *Preventive Medicine Reports, 12*(June), 304–311.
<https://doi.org/10.1016/j.pmedr.2018.10.019>
- Webb, T. L., & Sheeran, P. (2006). Does changing behavioral intentions engender behavior change? A meta-analysis of the experimental evidence. *Psychological Bulletin, 132*(2), 249–268.
<https://doi.org/10.1037/0033-2909.132.2.249>
- West, R. (2017). Tobacco smoking: Health impact, prevalence, correlates and interventions.

Psychology and Health, 32(8), 1018–1036. <https://doi.org/10.1080/08870446.2017.1325890>

World Health Organization. (2018). *Who global report on trends in prevalence of tobacco smoking 2000-2025, second edition*.

Tables

Table 1

Sample Characteristics

	<i>M (SD)</i>	<i>Range</i>
Socio demographics		
Females, N (%)	434 (63.17%)	
Males, N (%)	235 (34.21%)	
No answer sex, N (%)	18 (2.62%)	
Age, years	20.06 (2.39)	18 - 35
Smoking-related variables		
First use, year	16.12 (2.03)	10 - 31
Cigarettes per day	6.34 (16.3)	0 - 60
Days of use per month	20.52 (10.38)	1 - 30
CDS –Tobacco dependence	11.72 (4.50)	1 - 24
MTSS – Motivation to quit smoking	2.85 (1.69)	1 - 7
Recent attempts to quit smoking, N (%)	214 (41.39%)	
Smokers Status		
Non-daily, N (%)	436 (63.5%)	
Daily, N (%)	251 (36.5%)	
Psychological variables		
Attitude related to smoking	3.70 (1.08)	1 - 7
Social norms related to smoking	3.86 (1.17)	1 - 7
Perceived behavioral control to resist smoking	3.51 (1.60)	1 - 7
Smoker identity	2.69 (1.64)	1 - 7
SCOFF – Eating Disorders	1.02 (1.18)	1 - 5
STAI-T – Anxiety	50.90 (14.35)	20 - 80
BDI – Depression	9.40 (8.35)	0 - 36

Note. Except for gender, recent attempts to quit, and smokers' status, data show means (standard deviations); CDS: Cigarette Dependence Scale; MTSS: Motivation to stop smoking; SCOFF: Sick Control One Fat Food; STAI-T: State-Trait Anxiety Inventory; BDI: Beck Depression Inventory

Table 2*The Fit Indices for the Three Confirmatory Factor Analysis Models of FB-WISDM*

	Model description	Satorra-Bentler χ^2	df	CFI	TLI	RMSEA	90% Confidence Interval		SRMR
							Lower	Upper	
Model 1	2 first-order factors with error covariances	7291.134	628	0.655	0.634	0.126	0.123	0.129	0.089
Model 2	11 first-order factors with error covariances	2087.628	563	0.926	0.913	0.061	0.058	0.065	0.043
Model 3	11 first-order and 2 second-order factors with error covariances	2442.526	617	0.911	0.903	0.065	0.061	0.068	0.057

Note. CFI = comparative fit index; RMSEA = root mean squared error of approximation; SRMR = standardized root mean square residual; TLI = Tucker–Lewis index.

Table 3*Factor Loadings and Internal Consistencies of the French B-WISDM*

Factor	Estimate	SE	Standard Estimate	Z	p	α
Affiliative Attachment						
Item 11	1.821	0.268	0.821	6.794	< .001	.86
Item 22	1.336	0.191	0.830	6.982	< .001	
Item 26	1.498	0.221	0.844	6.766	< .001	
Automaticity						
Item 1	1.628	0.109	0.776	14.975	< .001	.89
Item 10	1.612	0.103	0.774	15.648	< .001	
Item 14	1.732	0.118	0.900	14.698	< .001	
Item 25	1.519	0.118	0.827	12.845	< .001	
Loss of Control						
Item 2	1.513	0.285	0.872	5.318	< .001	.91
Item 16	1.338	0.258	0.783	5.185	< .001	
Item 21	1.883	0.346	0.911	5.439	< .001	
Item 35	1.547	0.290	0.835	5.342	< .001	
Cognitive Enhancement						
Item 6	1.786	0.066	0.896	27.128	< .001	.92
Item 13	1.818	0.069	0.917	26.205	< .001	
Item 32	1.566	0.067	0.855	23.237	< .001	
Craving						
Item 4	1.678	0.242	0.855	6.928	< .001	.91
Item 17	1.665	0.243	0.851	6.862	< .001	
Item 23	1.723	0.253	0.840	6.823	< .001	
Item 29	1.490	0.217	0.841	6.855	< .001	
Cue Exposure / Associative Processes						
Item 8	1.555	0.173	0.612	8.973	< .001	.73
Item 12	2.022	0.155	0.692	13.022	< .001	
Item 24	2.079	0.167	0.793	12.435	< .001	
Taste						
Item 5	1.505	0.116	0.790	12.956	< .001	.84
Item 15	1.706	0.136	0.857	12.575	< .001	
Item 20	1.519	0.110	0.747	13.771	< .001	
Tolerance						
Item 3	1.433	0.141	0.718	10.176	< .001	.87
Item 28	1.732	0.146	0.863	11.821	< .001	
Item 31	1.555	0.158	0.676	9.841	< .001	
Item 36	1.803	0.148	0.894	12.182	< .001	
Weight Control						
Item 7	1.665	0.095	0.867	17.566	< .001	.88
Item 19	1.073	0.092	0.752	11.615	< .001	
Item 34	1.755	0.100	0.912	17.534	< .001	

Affective Enhancement						
Item 9	1.595	0.251	0.774	6.345	< .001	.82
Item 33	1.690	0.273	0.753	6.199	< .001	
Item 37	1.656	0.263	0.795	6.301	< .001	
Social / Environmental Goals						
Item 18	1.622	0.091	0.866	17.829	< .001	.92
Item 27	1.560	0.092	0.848	17.001	< .001	
Item 30	1.812	0.083	0.965	21.788	< .001	

Note. $N = 687$. The table shows the results of the CFA and internal consistency for Model 2.

SMOKING MOTIVES AMONG UNIVERSITY STUDENTS

Table 4

The Fit Indices for the Three Multigroups Confirmatory Factor Analysis Models of FB-WISDM for Type of Smokers and Gender

	Model description	Satorra-Bentler χ^2	df	CFI	TLI	RMSEA	90% Confidence Interval		SRMR
							Lower	Upper	
Type of smokers (daily vs. non-daily)									
Model 1	Configural Invariance	2644.08	1126	0.919	0.905	0.059	0.056	0.063	0.052
Model 2	Metric Invariance	2766.83	1163	0.914	0.901	0.060	0.057	0.064	0.063
Model 3	Scalar Invariance	2875.13	1189	0.914	0.901	0.060	0.057	0.064	0.063
Gender (men vs. women)									
Model 1	Configural Invariance	2916.475	1126	0.917	0.902	0.065	0.062	0.069	0.049
Model 2	Metric Invariance	2972.161	1163	0.916	0.904	0.065	0.061	0.068	0.054
Model 3	Scalar Invariance	3002.432	1189	0.916	0.906	0.064	0.061	0.067	0.054

Note. CFI = comparative fit index; RMSEA = root mean squared error of approximation; SRMR = standardized root mean square residual; TLI = Tucker–Lewis index.

SMOKING MOTIVES AMONG UNIVERSITY STUDENTS

Table 5

Standardized Coefficients from a Multivariate Regression Model Describing the Association between the French B-WISDM and Tobacco Dependence and Psychological Variables

	Tobacco dependence	Attitude	Social norms	PBC	Smoker identity	Eating disorders	Anxiety	Depression
Adjusted R^2	.761	.319	.317	.520	.447	.211	.247	.282
Loss of Control	.252***	-.199*	.046	-.050	.252***	-.081	-.108	-.113
Craving	.342***	.023	-.004	-.319***	-.013	-.005	.215	.169
Automaticity	.044	-.002	.025	-.048	-.010	.056	-.002	.006
Tolerance	.376***	.045	-.047	.015	.156*	-.030	-.212*	-.123
Affiliative Attachment	-.058*	.047	.025	-.008	.192***	.071	.182*	.234**
Associative Processes	.017	-.003	-.051	-.387***	.123*	.038	.043	-.044
Social Goads	-.004	.001	.538***	.046	-.009	-.019	.028	.057
Taste	.024	.481***	.053	.009	.082	-.049	-.119	-.143*
Weight Control	.035	-.015	.064	.025	-.024	.332***	.129*	.226***
Affective Enhancement	-.082	.181***	.008	-.067	.009	.047	.305***	.269***
Cognitive Enhancement	-.012	.049	.026	.003	.059	-.002	.024	.029

Note. PBC: Perceived Behavioral Control. Cigarettes/day: Number of cigarettes smoked per day. Smoking days/month: Number of smoking days per month. Statistically significant at *** $p < .001$; ** $p < .01$; * $p < .05$