



HAL
open science

A global invariant for path structures and second order differential equations

Elisha Falbel, Jose Miguel Veloso

► **To cite this version:**

Elisha Falbel, Jose Miguel Veloso. A global invariant for path structures and second order differential equations. 2023. hal-04121036

HAL Id: hal-04121036

<https://hal.science/hal-04121036>

Preprint submitted on 12 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A global invariant for path structures and second order differential equations

E. Falbel and J. M. Veloso

Abstract

We study a global invariant for path structures. The invariant is obtained as a secondary invariant from a Cartan connection on a canonical bundle associated to a path structure. It is computed in examples which are defined in terms of reductions of the path structure. In particular we give a formula for this global invariant for second order differential equations defined on a torus T^2 .

1 Introduction

Path structures on a 3-manifold are defined by a choice of contact structure and a decomposition of the contact plane bundle as a direct sum of two line bundles. This structure was thoroughly studied in the 19th century (see in particular [T]) as it appears in the description of second order differential equations and their equivalence under certain transformations (see Section 2 and references [A, IL, BGH]).

In Section 2 we collect definitions and examples. In particular we explain the relation with ordinary second order equations. In the following section we define the most important reductions of path structures. The first one is obtained by fixing a global contact form and it is called strict path structure. There exists a Cartan bundle Y_1 and a connection adapted to that structure (see 2.5) which was used in ([FMMV]) to obtain a classification of compact 3-manifolds with non-compact automorphism group preserving the strict path structure. We recall the construction in Proposition 2.3. The second one, we call enriched path structure following [MM] which were used by Mion-Mouton to classify certain classes of partial-hyperbolic diffeomorphisms of three manifolds. It consists of path structures where we fix a line transverse to the contact distribution. We define an adapted Cartan bundle Y_2 and a canonical connection adapted to this structure (see 2.6 and Proposition 2.5). There exists a natural embedding $Y_1 \rightarrow Y_2$ (Section 2.6.2, Proposition 2.7).

In Section 3 we recall the construction of the Cartan bundle Y and the canonical adapted connection to a path structure on a 3-manifold (see Proposition 3.3). This construction is due to Cartan in [Car]. Although one can find modern treatments of this topic in several references (in particular [IL, BGH]), we include this section for the sake of completeness and because the conventions we use might differ from others. We obtain a natural embedding $Y_2 \rightarrow Y$ (see 3.4, Proposition 3.4) and compute the curvature of the bundle Y in terms of the curvature of Y_2 (see 3.4.1). The formulas are used in the computation of the global invariant

in the next section. We also recall the computations by Cartan of the invariants of a second order differential equation.

In the following section we define the global invariant when Y_2 admits a global section (see Definition 4.2). This construction is inspired by an analogous construction of a Chern-Simons invariant in the case of CR manifolds given in [BE] (see also [CL] for a relative version which does not depend on the existence of a global section). In [FV] we defined a similar invariant for flag structures. Those are manifolds equipped with a decomposition of a complex contact structure defined on the complexified tangent bundle of a 3-manifold. In this paper we restrict the definition to path structures. We obtain the expression of the invariant in terms of a reduction Y_2 or Y_1 of the Cartan bundle Y of the path structure (see Proposition 4.5). We also give a formula of the invariant in the case of a second order differential equation on the torus (Proposition 4.10). It involves an integration of fifth order derivatives of the function defining the ordinary equation in the form $y'' = F(x, y, y')$. We use coordinates in the projective cotangent bundle over a surface as explained in section 4.1. We characterize certain families of differential equations on the torus which have vanishing global invariant in Corollary 4.11. We then compute the invariant for a family of path structures on tight contact structures on the torus (see Proposition 5.3) and characterize those structures with vanishing global invariant, they turn to be flat. Finally we compute the global invariant for homogeneous path structures on $\mathbf{SU}(2)$ (see Proposition 6.1) and identify the flat structure on the sphere where the global invariant is maximal.

The authors thank Martin Mion-Mouton for useful discussions.

2 Path structures in dimension 3

Path geometries are very related to the theory of second order differential equations. See a modern treatment in section 8.6 of [IL] and in [BGH] where the relation to second order differential equations is also explained. Let M be a real three dimensional manifold and TM be its tangent bundle.

Definition 2.1 *A path structure on M is a choice of two sub-bundles T^1 and T^2 in TM such that $T^1 \cap T^2 = \{0\}$ and such that $T^1 \oplus T^2$ is a contact distribution.*

The condition that $T^1 \oplus T^2$ be a contact distribution means that, locally, there exists a one form $\theta \in T^*M$ such that $\ker \theta = T^1 \oplus T^2$ and $d\theta \wedge \theta$ is never zero.

One can choose a contact form θ up to a scalar function. One can interpret this as follows: one has a \mathbb{R}^* -bundle over the manifold given by the choice of θ at each point (one might keep only positive multiples for simplicity). Over this line bundle one defines the tautological form $\omega_x = \pi^*(\theta_{\pi(x)})$. This bundle is trivial if and only if there exists a global contact form θ . If the contact distribution is oriented, then there exists a global contact form. Indeed, using a global metric on the distribution one can define locally a transversal vector to the distribution taking a Lie bracket of orthonormal vectors in the distribution. This defines a global 1-form.

Fix θ and local forms Z^1 and Z^2 defining the lines as above such that $d\theta = Z^1 \wedge Z^2$. There exists global forms Z^1 and Z^2 if and only if there exists global vector fields along the

lines. Clearly, if the contact distribution is oriented, it suffices that there exists a global vector field along one of the foliations by lines.

Local equivalence (also called point equivalence) between path structures happens when there exists a local diffeomorphism which gives a correspondence between the lines defining each structure.

2.1 The flat model space

Flat path geometry is the geometry of real flags in \mathbb{R}^3 . That is the geometry of the space of all couples (p, l) where $p \in \mathbb{R}P^2$ and l is a real projective line containing p . The space of flags is identified to the quotient

$$\mathbf{SL}(3, \mathbb{R})/B$$

where B is the Borel group of all real upper triangular matrices.

The Lie algebra of $\mathbf{SL}(3, \mathbb{R})$ decomposes into the following direct sum of vector subspaces:

$$\mathfrak{sl}(3, \mathbb{R}) = \mathfrak{g}^{-2} \oplus \mathfrak{g}^{-1} \oplus \mathfrak{g}^0 \oplus \mathfrak{g}^1 \oplus \mathfrak{g}^2,$$

where

$$\begin{aligned} \mathfrak{g}^{-2} &= \left\{ \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ z & 0 & 0 \end{pmatrix} \right\}, & \mathfrak{g}^{-1} &= \left\{ \begin{pmatrix} 0 & 0 & 0 \\ x & 0 & 0 \\ 0 & y & 0 \end{pmatrix} \right\}, \\ \mathfrak{g}^0 &= \left\{ \begin{pmatrix} u+v & 0 & 0 \\ 0 & -2v & 0 \\ 0 & 0 & -u+v \end{pmatrix} \right\}, \\ \mathfrak{g}^1 &= \left\{ \begin{pmatrix} 0 & a & 0 \\ 0 & 0 & b \\ 0 & 0 & 0 \end{pmatrix} \right\}, & \mathfrak{g}^2 &= \left\{ \begin{pmatrix} 0 & 0 & c \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \right\}. \end{aligned}$$

That is the graded decomposition of $\mathfrak{sl}(3, \mathbb{R})$ where $\mathfrak{b} = \mathfrak{g}^0 \oplus \mathfrak{g}^1 \oplus \mathfrak{g}^2$ corresponds to upper triangular matrices with null trace. The tangent space of $\mathbf{SL}(3, \mathbb{R})/B$ at $[B]$ is identified to

$$\mathfrak{sl}(3, \mathbb{R})/\mathfrak{b} = \mathfrak{g}^{-2} \oplus \mathfrak{g}^{-1}.$$

2.2 Examples

Example I Consider the Heisenberg group

$$\mathbf{Heis}(3) = \{ (z, t) \mid z \in \mathbb{C}, t \in \mathbb{R} \}$$

with multiplication defined by $(z_1, t_1) \star (z_2, t_2) = (z_1 + z_2, t_1 + t_2 + 2\text{Im } z_1 \overline{z_2})$. The contact form

$$\theta = dt - xdy - ydx$$

is invariant under left multiplications (also called Heisenberg translations). If $\Lambda \subset \mathbf{Heis}(3)$ is a lattice then the quotient $\Lambda \backslash \mathbf{Heis}(3)$ is a circle bundle over the torus with a globally defined contact form.

A lattice Λ determines a lattice $\Gamma \subset \mathbb{C}$ corresponding to projection in the exact sequence

$$0 \rightarrow \mathbb{R} \rightarrow \mathbf{Heis}(3) \rightarrow \mathbb{C} \rightarrow 0.$$

There are many global vector fields in the distribution defined by θ invariant under Λ , it suffices to lift an invariant vector field on \mathbb{C} under Γ . All circle bundles obtained in this way are not trivial and the fibers are transverse to the distribution.

Example II. Here we consider the torus T^3 with coordinates $(x, y, t) \pmod{1}$ and the global contact form

$$\theta_n = \cos(2\pi nt)dx - \sin(2\pi nt)dy.$$

There are two canonical global vector fields on the distribution given by $\frac{\partial}{\partial t}$ and $\sin(2\pi nt)\frac{\partial}{\partial x} + \cos(2\pi nt)\frac{\partial}{\partial y}$. In this example, the fiber given by the coordinate t has tangent space contained in the distribution.

Example III. An homogeneous example is the Lie group $\mathbf{SU}(2)$ with left invariant vector fields X and Y with $Z = [X, Y]$ and cyclic commutation relations. The vector fields X and Y define a path structure on $\mathbf{SU}(2)$.

Example IV. Another homogeneous example is the Lie group $\mathbf{SL}(2, \mathbb{R})$ with left invariant vector fields X and Y with $Z = [X, Y]$ with $[Z, X] = X$ and $[Z, Y] = -Y$ given by generators

$$X = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, Y = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, Z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

The path structure defined by X and Y induces a path structure on the quotient $\Gamma \backslash \mathbf{SL}(2, \mathbb{R})$ by a discrete torsion free subgroup $\Gamma \subset \mathbf{SL}(2, \mathbb{R})$. This structure is invariant under the flow defined by right multiplication by e^{tZ} .

Example V. Let Σ be a surface equipped with a Riemannian metric. The geodesic flow on the unit tangent bundle $T^1\Sigma$ defines a distribution which, together with the distribution defined by the vertical fibers of the projection of the unit tangent bundle on Σ , defines a path structure which is not invariant under the geodesic flow. For $\Sigma = H_{\mathbb{R}}^2$, the hyperbolic space, we obtain $T^1\Sigma = \mathbf{PSL}(2, \mathbb{R})$ with distributions defined by the left invariant distributions $X - Y$ and Z (using the same generators of the Lie algebra as in the previous example).

Example VI Let M be a three manifold equipped with a path structure $D = T^1 \oplus T^2 \subset TM$. Suppose D is orientable and choose a section u of T^1 . Each section v of T^2 such that (u, v) is positive gives rise to a CR structure. Indeed we define $Ju = v$ and $Jv = -u$. The choice of v corresponds to a section of an \mathbb{R}_+^* -bundle over M . Reciprocally given a CR structure on M , defined by $J : D \rightarrow D$, one can associate path structures corresponding to a choice $T^1 \subset D$ and defining then $T^2 = J(T^1)$.

2.3 Path structures and second order differential equations

This is studied since a long time (see [T], [IL] and [BGH]). It turns out that path structures can be obtained putting together second order differential equations in one variable. Indeed, a second order differential equation in one variable is described locally as

$$\frac{d^2y}{dx^2} = F\left(x, y, \frac{dy}{dx}\right).$$

This defines a path structure on a neighborhood of a point in \mathbb{R}^3 with coordinates (x, y, p) :

$$L_1 = \ker\{dp - Fdx\} \cap \ker\{dy - pdx\}, \quad L_2 = \ker dx \cap \ker dy.$$

The contact structure is defined by the form

$$\theta = dy - pdx.$$

Defining the forms $Z^1 = dx$ and $Z^2 = dp - Fdx$, one has that $d\theta = Z^1 \wedge Z^2$.

One can show easily that every path structure is, in fact, locally equivalent to a second order equation. That is, there exists local coordinates such that L_1 and L_2 are defined via a second order ODE as above.

2.4 Reductions of path structures

We will describe two reductions of path geometry corresponding to subgroups $G_1 \subset G_2 \subset SL(3, \mathbb{R})$ where

$$G_1 = \left\{ \begin{pmatrix} a & 0 & 0 \\ \star & \frac{1}{a^2} & 0 \\ \star & \star & a \end{pmatrix} \right\}$$

and

$$G_2 = \left\{ \begin{pmatrix} a & 0 & 0 \\ \star & \frac{1}{ab} & 0 \\ \star & \star & b \end{pmatrix} \right\}.$$

The models are G_1/\mathbb{R}^* and G_2/\mathbb{R}^{*2} and correspond to the Heisenberg group where in the first model we fix a contact form and, in the second, a transverse line to the contact distribution.

Other reductions of the G_2 -structure might occur, namely by choosing other embeddings of \mathbb{R}^* into G_2 . They appear naturally when certain components of the curvature of the Cartan connections on Y_2 or Y are non-vanishing.

We will construct coframe bundles Y_1, Y_2 and a principal bundle Y over M with structure groups $\mathbb{R}^*, \mathbb{R}^{*2}$ and the Borel group B together with Cartan connections and canonical embeddings

$$Y_1 \rightarrow Y_2 \rightarrow Y.$$

They correspond to a strict path structure, an enriched path structure (see next sections for definitions) and finally, a path structures on the manifold M .

2.5 Path structures with a fixed contact form: strict path structures.

In this section we fix a contact form and recall the reduction of the structure group of a path geometry obtained in [FV] where we called the path structure with a fixed contact form a pseudo flag structure. This structure is called strict path structure in [FMMV].

G_1 denotes from now on the subgroup of $\mathbf{SL}(3, \mathbb{R})$ defined by

$$G_1 = \left\{ \begin{pmatrix} a & 0 & 0 \\ x & \frac{1}{a^2} & 0 \\ z & y & a \end{pmatrix} \mid a \in \mathbb{R}^*, (x, y, z) \in \mathbb{R}^3 \right\}$$

and $P_1 \subset G_1$ the subgroup defined by

$$P_1 = \left\{ \begin{pmatrix} a & 0 & 0 \\ 0 & \frac{1}{a^2} & 0 \\ 0 & 0 & a \end{pmatrix} \right\}.$$

Writing the Maurer-Cartan form of G_1 as the matrix

$$\begin{pmatrix} w & 0 & 0 \\ \theta^1 & -2w & 0 \\ \theta & \theta^2 & w \end{pmatrix}$$

one obtains the Maurer-Cartan equations:

$$d\theta + \theta^2 \wedge \theta^1 = 0$$

$$d\theta^1 - 3w \wedge \theta^1 = 0$$

$$d\theta^2 + 3w \wedge \theta^2 = 0$$

$$dw = 0.$$

G_1 is the automorphism group of the canonical left-invariant strict path structure of **Heis**(3), and that its action induces an identification of **Heis**(3) with the homogeneous space $X = G_1/P_1$.

Let M be a three-manifold equipped with a strict path structure (E^1, E^2, θ) defined by two one dimensional bundles E^1 and E^2 and contact form θ . We let R be the associated Reeb vector field (satisfying $\iota_R d\theta = 0$ and $\theta(R) = 1$). Now let $X_1 \in E^1$, $X_2 \in E^2$ be such that $d\theta(X_1, X_2) = 1$. The dual coframe of (X_1, X_2, R) is $(\theta^1, \theta^2, \theta)$, for two 1-forms θ^1 and θ^2 verifying $d\theta = \theta^1 \wedge \theta^2$.

At any point $x \in M$, one can look at the coframes of the form

$$\omega^1 = a^3 \theta^1(x), \quad \omega^2 = \frac{1}{a^3} \theta^2(x), \quad \omega = \theta(x)$$

for $a \in \mathbb{R}^*$.

Definition 2.2 We denote by $p_1 : Y_1 \rightarrow M$ the \mathbb{R}^* -coframe bundle over M given by the set of coframes $(\omega, \omega^1, \omega^2)$ of the above form.

We will denote the tautological forms defined by $\omega^1, \omega^2, \omega$ using the same letters. That is, we write ω^i at the coframe $(\omega^1, \omega^2, \omega)$ to be $p_1^*(\omega^i)$.

Proposition 2.3 There exists a unique Cartan connection on Y_1

$$\pi_1 = \begin{pmatrix} w & 0 & 0 \\ \omega^1 & -2w & 0 \\ \omega & \omega^2 & w \end{pmatrix}$$

such that its curvature form is of the form

$$\Pi_1 = d\pi_1 + \pi_1 \wedge \pi_1 = \begin{pmatrix} dw & 0 & 0 \\ \omega \wedge \tau^1 & -2dw & 0 \\ 0 & -\omega \wedge \tau^2 & dw \end{pmatrix}$$

with $\tau^1 \wedge \omega^2 = \tau^2 \wedge \omega^1 = 0$.

Observe that the condition $\tau^1 \wedge \omega^2 = \tau^2 \wedge \omega^1 = 0$ implies that we may write $\tau^1 = \tau_2^1 \omega^2$ and $\tau^2 = \tau_1^2 \omega^1$. The structure equations are

$$\begin{aligned} d\omega + \omega^2 \wedge \omega^1 &= 0, \\ d\omega^1 - 3w \wedge \omega^1 &= \omega \wedge \tau^1, \\ d\omega^2 + 3w \wedge \omega^2 &= -\omega \wedge \tau^2. \end{aligned}$$

The proof of the proposition is given in [FMMV] and [FV].

Bianchi identities are obtained differentiating the structure equations. They are described in the following equations:

$$dw = C\omega \wedge \omega^1 + D\omega \wedge \omega^2 + S\omega^1 \wedge \omega^2, \quad (1)$$

$$d\tau_2^1 - 6\tau_2^1 w + 3D\omega^1 = \tau_{20}^1 \omega + \tau_{22}^1 \omega^2 \quad (2)$$

$$d\tau_1^2 + 6\tau_1^2 w + 3C\omega^2 = \tau_{10}^2 \omega + \tau_{11}^2 \omega^1 \quad (3)$$

2.6 Path structures with a fixed transverse line: enriched path structures.

In this section we introduce a coframe bundle and a Cartan connection associated to a path structure with a fixed transverse line to the the contact distribution.

The model space is the homogeneous space which is the quotient of the group of lower triangular matrices in $SL(3, \mathbb{R})$ by the subgroup of diagonal matrices. The Maurer-Cartan form is the Lie algebra valued form which can be represented by

$$\pi = \begin{pmatrix} \varphi + w & 0 & 0 \\ \omega^1 & -2w & 0 \\ \omega & \omega^2 & -\varphi + w \end{pmatrix}$$

The Maurer-Cartan equations $d\pi + \pi \wedge \pi = 0$ are given by

$$\begin{aligned} d\omega &= 2\varphi \wedge \omega + \omega^1 \wedge \omega^2 \\ d\omega^1 &= \varphi \wedge \omega^1 + 3w \wedge \omega^1 \\ d\omega^2 &= \varphi \wedge \omega^2 - 3w \wedge \omega^2. \end{aligned}$$

Let M be a three manifold equipped with a path structure $D = T^1 \oplus T^2 \subset TM$. We fix a transverse line L so that $TM = T^1 \oplus T^2 \oplus L$.

We suppose $X_1 \in T^1$, $X_2 \in T^2$ and $X \in L$ is a frame. The dual coframe is θ^1 , θ^2 and θ . Observe that θ is simply a form with $\ker \theta = D$. One can define a coframe bundle defined by all coframes:

$$\omega^1 = a^1 \theta^1, \quad \omega^2 = a^2 \theta^2, \quad \omega = \lambda \theta.$$

where we will suppose, for simplicity, that $a^1, a^2, \lambda > 0$.

A reduction of this coframe bundle is obtained by imposing that each coframe verifies

$$d\omega|_D = (\omega^1 \wedge \omega^2)|_D$$

for an extension of the 1-form such that $\ker \omega = D$. This relation does not depend on the particular extension of a form ω defined at a point because $d\omega|_D(X, Y) = -\omega([X, Y])$ for any vector fields X and Y which are sections of the distribution D .

Definition 2.4 We denote by $p_2 : Y_2 \rightarrow M$ the \mathbb{R}^{*2} -coframe bundle over M given by the set of 1-forms $(\omega, \omega^1, \omega^2)$ defined above. The structure group \mathbb{R}^{*2} acts as follows

$$(\omega', \omega'^1, \omega'^2) = (\omega, \omega^1, \omega^2) \begin{pmatrix} \lambda & 0 & 0 \\ 0 & a^1 & 0 \\ 0 & 0 & a^2 \end{pmatrix}$$

where $\lambda, a^1, a^2 \in \mathbb{R}_+^*$ with $a^1 a^2 = \lambda$.

In order to define a Cartan connection on Y_2 we start taking the tautological forms corresponding to the forms $\omega, \omega^1, \omega^2$, which we will denote by the same letters by abuse of notation.

Using a coframe section $(\theta, \theta^1, \theta^2)$ on M one can express the tautological forms as

$$\omega = \lambda p_2^*(\theta), \quad \omega^1 = a^1 p_2^*(\theta^1), \quad \omega^2 = a^2 p_2^*(\theta^2),$$

with $a^1 a^2 = \lambda$.

We need to define two forms φ and w corresponding to the vertical directions

Observe first that we have

$$d\omega = \frac{d\lambda}{\lambda} \wedge \omega + \omega^1 \wedge \omega^2 \pmod{\omega}$$

and therefore one may write

$$d\omega = 2\varphi \wedge \omega + \omega^1 \wedge \omega^2 \tag{4}$$

where φ restricted to the vertical fiber is $\frac{d\lambda}{2\lambda}$. The form φ is not yet fixed and any other form φ' satisfying the equation satisfies

$$\varphi - \varphi' = s\omega$$

where s is a function on Y_2 .

Differentiating the forms ω^1 and ω^2 we obtain new forms which correspond to the coordinates a^1, a^2 :

$$d\omega^1 = \frac{da^1}{a^1} \wedge \omega^1 + a^1 d\theta^1 \quad \text{and} \quad d\omega^2 = \frac{da^2}{a^2} \wedge \omega^2 + a^2 d\theta^2.$$

Observing that

$$\frac{d\lambda}{\lambda} = \frac{da^1}{a^1} + \frac{da^2}{a^2}$$

we can write

$$\begin{aligned} d\omega^1 &= \frac{d\lambda}{2\lambda} \wedge \omega^1 + \frac{1}{2} \left(\frac{da^1}{a^1} - \frac{da^2}{a^2} \right) \wedge \omega^1 + a^1 d\theta^1 \\ d\omega^2 &= \frac{d\lambda}{2\lambda} \wedge \omega^2 - \frac{1}{2} \left(\frac{da^1}{a^1} - \frac{da^2}{a^2} \right) \wedge \omega^2 + a^2 d\theta^2 \end{aligned}$$

Now we can make the first right hand term of each equation to be $\varphi \wedge \omega^1$ and $\varphi \wedge \omega^2$ respectively by adding terms in $\omega, \omega^1, \omega^2$ to $\frac{d\lambda}{2\lambda}$. The terms in $\omega^1 \wedge \omega^2$ not appearing in these first terms can be absorbed in the second term in each equation. It remains a last term in each equation that we denote by $\omega \wedge \tau^1$ and $-\omega \wedge \tau^2$ respectively. We proved the following:

Lemma 2.1 *There exists forms w, τ^1, τ^2 defined on Y_2 such that*

$$d\omega^1 = \varphi \wedge \omega^1 + 3w \wedge \omega^1 + \omega \wedge \tau^1 \text{ and } d\omega^2 = \varphi \wedge \omega^2 - 3w \wedge \omega^2 - \omega \wedge \tau^2. \quad (5)$$

The forms τ^1 and τ^2 are horizontal, that is, they vanish on vectors tangent to the fibers of $Y_2 \rightarrow M$. Moreover, writing $\omega^1 = a^1\theta^1$, $\omega^2 = a^2\theta^2$, $\omega = \lambda\theta$ for a choice of sections on M , one has $\varphi = \frac{d\lambda}{2\lambda}$ and $6w = \frac{da^1}{a^1} - \frac{da^2}{a^2}$ modulo the tautological forms of the fiber bundle Y_2 .

Let φ', w', τ'^1 and τ'^2 be other forms satisfying equations above. Taking the difference we obtain

$$0 = (\varphi - \varphi') \wedge \omega^1 + 3(w - w') \wedge \omega^1 + \omega \wedge (\tau^1 - \tau'^1)$$

and

$$0 = (\varphi - \varphi') \wedge \omega^2 - 3(w - w') \wedge \omega^2 - \omega \wedge (\tau^2 - \tau'^2)$$

Therefore, as $\varphi - \varphi' = s\omega$, we can write

$$0 = -3\omega^1 \wedge (w - w') + \omega \wedge (s\omega^1 + \tau^1 - \tau'^1)$$

and

$$0 = 3\omega^2 \wedge (w - w') - \omega \wedge (-s\omega^2 + \tau^2 - \tau'^2).$$

By Cartan's lemma we obtain

$$w - w' = a\omega,$$

$$\tau^1 - \tau'^1 = -3a\omega^1 - s\omega^1 + b^1\omega,$$

$$\tau^2 - \tau'^2 = -3a\omega^2 + s\omega^2 + b^2\omega.$$

Now, we can impose that $\tau^1 = \tau_1^1\omega^1 + \tau_2^1\omega^2$ and $\tau^2 = \tau_1^2\omega^1 + \tau_2^2\omega^2$ by choosing convenient b^1 and b^2 (or by simply considering, from the beginning, τ^1 and τ^2 with no terms in ω). Moreover, one can choose unique a and s so that $\tau_1^1 = 0$ and $\tau_2^2 = 0$. We conclude that

Lemma 2.2 *There exists unique forms $\varphi, w, \tau^1, \tau^2$ defined on Y_2 such that*

$$d\omega = 2\varphi \wedge \omega + \omega^1 \wedge \omega^2$$

$$d\omega^1 = \varphi \wedge \omega^1 + 3w \wedge \omega^1 + \omega \wedge \tau^1$$

$$d\omega^2 = \varphi \wedge \omega^2 - 3w \wedge \omega^2 - \omega \wedge \tau^2$$

with $\tau^1 \wedge \omega^2 = \tau^2 \wedge \omega^1 = 0$.

Bianchi identities are obtained differentiating the above equations:

Lemma 2.3 *There exists a 1-form ψ such that*

$$d\varphi = \omega \wedge \psi \quad (6)$$

The form ψ may be chosen satisfying $\psi = A\omega^1 + B\omega^2$ and $d\psi = -2\varphi \wedge \psi + \omega \wedge \alpha$ where A, B are functions on Y_2 and α is a 1-form on Y_2 .

Proof. Differentiating equation $d\omega = 2\varphi \wedge \omega + \omega^1 \wedge \omega^2$ one obtains, using equations 5, that $d\varphi \wedge \omega = 0$, that is,

$$d\varphi = \omega \wedge \psi \quad (7)$$

for a 1-form ψ defined on Y_2 .

Differentiating $d\varphi = \omega \wedge \psi$ one has

$$0 = d\omega \wedge \psi - \omega \wedge d\psi = (2\varphi \wedge \omega + \omega^1 \wedge \omega^2) \wedge \psi - \omega \wedge d\psi = \omega^1 \wedge \omega^2 \wedge \psi - \omega \wedge (d\psi + 2\varphi \wedge \psi).$$

Using Cartan's lemma, $\psi = A\omega^1 + B\omega^2$ modulo ω , and we certainly can choose ψ satisfying $d\varphi = \omega \wedge \psi$ with $\psi = A\omega^1 + B\omega^2$. We conclude that

$$d\psi + 2\varphi \wedge \psi = \omega \wedge \alpha.$$

□

Equation $dd\omega^1 = 0$ gives after simplifications

$$0 = d(\varphi + 3w) \wedge \omega^1 + \omega \wedge \omega^2 (d\tau_2^1 + 2\tau_2^1(\varphi - 3w)). \quad (8)$$

Analogously, $dd\omega^2 = 0$ simplifies to

$$0 = d(\varphi - 3w) \wedge \omega^2 - \omega \wedge \omega^1 (d\tau_1^2 + 2\tau_1^2(\varphi + 3w)). \quad (9)$$

Using the previous lemma we may write

$$dw = C\omega \wedge \omega^1 + D\omega \wedge \omega^2 + S\omega^1 \wedge \omega^2,$$

where C, D and S are functions on Y_2 .

We can represent the equations above as a matrix equation whose entries are differential forms. The forms are disposed in the Lie algebra $\mathfrak{b} \subset \mathfrak{sl}(3, \mathbb{R})$ (the Lie algebra of lower triangular matrices) and we obtain the following Proposition.

Proposition 2.5 *Let Y_2 be the adapted principal bundle constructed above associated to an enriched path structure on a manifold M . Then there exists a unique Cartan's connection with values in \mathfrak{b} :*

$$\pi_2 = \begin{pmatrix} \varphi + w & 0 & 0 \\ \omega^1 & -2w & 0 \\ \omega & \omega^2 & -\varphi + w \end{pmatrix}$$

with curvature:

$$\Pi_2 = d\pi_2 + \pi_2 \wedge \pi_2 = \begin{pmatrix} \omega \wedge \psi + W & 0 & 0 \\ \tau_2^1 \omega \wedge \omega^2 & -2W & 0 \\ 0 & -\tau_1^2 \omega \wedge \omega^1 & -\omega \wedge \psi + W \end{pmatrix} \quad (10)$$

where $W = C\omega \wedge \omega^1 + D\omega \wedge \omega^2 + S\omega^1 \wedge \omega^2$ and $\psi = A\omega^1 + B\omega^2$.

2.6.1 More Bianchi identities

- Substituting the expressions above in equations 8 and 9 we obtain

$$d\tau_2^1 + 2\tau_2^1(\varphi - 3w) + (B + 3D)\omega^1 = \tau_{20}^1\omega + \tau_{22}^1\omega^2. \quad (11)$$

- Analogously we obtain

$$d\tau_1^2 + 2\tau_1^2(\varphi + 3w) - (A - 3C)\omega^2 = \tau_{10}^2\omega + \tau_{11}^2\omega^1. \quad (12)$$

From the last two equations we obtain the following

Proposition 2.6 *If the adapted connection of Y_2 has nul torsion and*

$$dw = S\omega^1 \wedge \omega^2,$$

then $d\varphi = 0$.

- Analogously, $dd\varphi = 0$ simplifies to

$$0 = \omega \wedge \omega^1(dA + 3A(\varphi + w)) + \omega \wedge \omega^2(dB + 3B(\varphi - w))$$

and we obtain

$$dA + 3A(\varphi + w) = A_0\omega + A_1\omega^1 + A_2\omega^2, \quad (13)$$

$$dB + 3B(\varphi - w) = B_0\omega + B_1\omega^1 + B_2\omega^2, \quad (14)$$

with $A_2 = B_1$.

- Also, $ddw = 0$ simplifies to

$$0 = \omega \wedge \omega^1(dC + 3C(\varphi + w)) + \omega \wedge \omega^2(dD + 3D(\varphi - w)) + \omega^1 \wedge \omega^2(dS + 2S\varphi)$$

and we obtain

$$dC + 3C(\varphi + w) = C_0\omega + C_1\omega^1 + C_2\omega^2, \quad (15)$$

$$dD + 3D(\varphi - w) = D_0\omega + D_1\omega^1 + D_2\omega^2, \quad (16)$$

$$dS + 2S\varphi = S_0\omega + S_1\omega^1 + S_2\omega^2, \quad (17)$$

with $C_2 - D_1 + S_0 = 0$.

Lemma 2.4 *If $\tau^1 = \tau^2 = C = D = 0$*

$$d\varphi = 0.$$

Proof. From the last formulae we obtain that ψ is a multiple of ω and the result follows. \square

2.6.2 The embedding $\iota_1 : Y_1 \rightarrow Y_2$

Given a path structure with a fixed contact form ω we obtained first a coframe bundle Y_1 and one can also obtain a canonical transverse direction by considering the Reeb vector field associated to ω . One obtains then a coframe bundle Y_2 of last section.

Given a coframe $(\omega, \omega^1, \omega^2) \in Y_1$ one can view the same coframe as a coframe of Y_2 . This gives an embedding

$$\iota_1 : Y_1 \rightarrow Y_2.$$

By abuse of language we may write the connection forms of Y_1 and Y_2 using the same letters and then obtain:

Proposition 2.7 *There exists a unique embedding $\iota_1 : Y_1 \rightarrow Y_2$ satisfying $\iota_1^*(\omega) = \omega$, $\iota_1^*(\omega^1) = \omega^1$ and $\iota_1^*(\omega^2) = \omega^2$. Moreover, for this embedding, $\iota_1^*(\varphi) = 0$ and $\iota_1^*(w) = w$.*

Proof. If unicity is not satisfied one can obtain the same forms pulling back a different coframe. But from the transformations of the coframe,

$$\begin{aligned}\tilde{\omega} &= \frac{a}{b} \omega \\ \tilde{\omega}^1 &= a^2 b \omega^1 \\ \tilde{\omega}^2 &= \frac{1}{ab^2} \omega^2.\end{aligned}$$

We obtain then that $a = b = 1$ and the embedding is uniquely determined by the conditions.

Comparing the structure equations of both structures we further get the equations $\iota_1^*(\varphi) = 0$ and $\iota_1^*(w) = w$. \square

3 The Cartan connection of a path structure

We review in this section the construction of a Cartan connection. The construction is due to E. Cartan in [Car] and one can read a modern description of it in [IL]. We include this section in order to fix our conventions and describe the embedding of Y_2 into the corresponding fiber bundle associated to a path geometry (see 3.4 and 3.4.1) which will be used to define the global invariant.

The Maurer-Cartan form on $SL(3, \mathbb{R})$ is given by a form with values in the Lie algebra $\mathfrak{sl}(3, \mathbb{R})$:

$$\pi = \begin{pmatrix} \varphi + w & \varphi^2 & \psi \\ \omega^1 & -2w & \varphi^1 \\ \omega & \omega^2 & -\varphi + w \end{pmatrix}$$

satisfying the equation $d\pi + \pi \wedge \pi = 0$. That is

$$\begin{aligned}d\omega &= \omega^1 \wedge \omega^2 + 2\varphi \wedge \omega \\ d\omega^1 &= \varphi \wedge \omega^1 + 3w \wedge \omega^1 + \omega \wedge \varphi^1\end{aligned}$$

$$\begin{aligned}
d\omega^2 &= \varphi \wedge \omega^2 - 3w \wedge \omega^2 - \omega \wedge \varphi^2 \\
dw &= -\frac{1}{2}\varphi^2 \wedge \omega^1 + \frac{1}{2}\varphi^1 \wedge \omega^2 \\
d\varphi &= \omega \wedge \psi - \frac{1}{2}\varphi^2 \wedge \omega^1 - \frac{1}{2}\varphi^1 \wedge \omega^2 \\
d\varphi^1 &= \psi \wedge \omega^1 - \varphi \wedge \varphi^1 + 3w \wedge \varphi^1 \\
d\varphi^2 &= -\psi \wedge \omega^2 - \varphi \wedge \varphi^2 - 3w \wedge \varphi^2 \\
d\psi &= \varphi^1 \wedge \varphi^2 + 2\psi \wedge \varphi.
\end{aligned}$$

3.1 The \mathbb{R}^* -bundle of contact forms and an adapted coframe bundle

We recall the construction of the \mathbb{R}^* -bundle of contact forms. Define E to be the \mathbb{R}^* -bundle of all forms θ on TM such that $\ker \theta = T^1 \oplus T^2$. Remark that this bundle is trivial if and only if there exists a globally defined non-vanishing form θ . Define the set of forms θ^1 and θ^2 on M satisfying

$$\begin{aligned}
\theta^1(T^1) &\neq 0 \quad \text{and} \quad \theta^2(T^2) \neq 0. \\
\ker \theta^1|_{\ker \theta} &= T^2 \quad \text{and} \quad \ker \theta^2|_{\ker \theta} = T^1.
\end{aligned}$$

Fixing one choice, all others are given by $\theta^i = a^i\theta^i + v^i\theta$.

On E we define the tautological form ω . That is $\omega_\theta = \pi^*(\theta)$ where $\pi : E \rightarrow M$ is the natural projection. We also consider the tautological forms defined by the forms θ^1 and θ^2 over the line bundle E . That is, for each $\theta \in E$ we let $\omega_\theta^i = \pi^*(\theta^i)$. At each point $\theta \in E$ we have the family of forms defined on E :

$$\begin{aligned}
\omega' &= \omega \\
\omega'^1 &= a^1\omega^1 + v^1\omega \\
\omega'^2 &= a^2\omega^2 + v^2\omega
\end{aligned}$$

We may, moreover, suppose that

$$d\theta = \theta^1 \wedge \theta^2 \quad \text{modulo } \theta$$

and therefore

$$d\omega = \omega^1 \wedge \omega^2 \quad \text{modulo } \omega.$$

This imposes that $a^1a^2 = 1$.

Those forms vanish on vertical vectors, that is, vectors in the kernel of the map $TE \rightarrow TM$. In order to define non-horizontal 1-forms we let θ be a section of E over M and introduce the coordinate $\lambda \in \mathbb{R}^*$ in E . By abuse of notation, let θ denote the tautological form on the section θ . We write then the tautological form ω over E is

$$\omega_{\lambda\theta} = \lambda\theta.$$

Differentiating this formula we obtain

$$d\omega = 2\varphi \wedge \omega + \omega^1 \wedge \omega^2 \quad (18)$$

where $\varphi = \frac{d\lambda}{2\lambda}$ modulo $\omega, \omega^1, \omega^2$. Here $\frac{d\lambda}{2\lambda}$ is a form intrinsically defined on E up to horizontal forms. We obtain in this way a coframe bundle satisfying equation 18 over E :

$$\begin{aligned} \omega' &= \omega \\ \omega'^1 &= a^1\omega^1 + v^1\omega \\ \omega'^2 &= a^2\omega^2 + v^2\omega \\ \varphi' &= \varphi - \frac{1}{2}a^1v^2\omega^1 + \frac{1}{2}a^2v^1\omega^2 + s\omega \end{aligned}$$

$v^1, v^2, s \in \mathbb{R}$ and $a^1, a^2 \in \mathbb{R}^*$ such that $a^1a^2 = 1$.

Definition 3.1 We denote by Y the coframe bundle $Y \rightarrow E$ given by the set of 1-forms $\omega, \omega^1, \omega^2, \varphi$ as above. Two coframes are related by

$$(\omega', \omega'^1, \omega'^2, \varphi') = (\omega, \omega^1, \omega^2, \varphi) \begin{pmatrix} 1 & v^1 & v^2 & s \\ 0 & a^1 & 0 & -\frac{1}{2}a^1v^2 \\ 0 & 0 & a^2 & \frac{1}{2}a^2v^1 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

where and $s, v^1, v^2 \in \mathbb{R}$ and $a^1, a^2 \in \mathbb{R}^*$ satisfy $a^1a^2 = 1$.

The bundle Y can also be fibered over the manifold M . In order to describe the bundle Y as a principal fiber bundle over M observe that choosing a local section θ of E and forms θ^1 and θ^2 on M such that $d\theta = \theta^1 \wedge \theta^2$ one can write a trivialization of the fiber

$$\begin{aligned} \omega &= \lambda\theta \\ \omega^1 &= a^1\theta^1 + v^1\lambda\theta \\ \omega^2 &= a^2\theta^2 + v^2\lambda\theta \\ \varphi &= \frac{d\lambda}{2\lambda} - \frac{1}{2}a^1v^2\theta^1 + \frac{1}{2}a^2v^1\theta^2 + s\theta, \end{aligned}$$

where $v^1, v^2, s \in \mathbb{R}$ and $a^1, a^2 \in \mathbb{R}^*$ such that $a^1a^2 = \lambda$. Here the coframe $\omega, \omega^1, \omega^2, \varphi$ is seen as composed of tautological forms.

The group H acting on the right of this bundle is

$$H = \left\{ \left(\begin{pmatrix} \lambda & v^1\lambda & v^2\lambda & s \\ 0 & a^1 & 0 & -\frac{1}{2}a^1v^2 \\ 0 & 0 & a^2 & \frac{1}{2}a^2v^1 \\ 0 & 0 & 0 & 1 \end{pmatrix} \text{ where } s, v^1, v^2 \in \mathbb{R} \text{ and } a^1, a^2 \in \mathbb{R}^* \text{ satisfy } a^1a^2 = \lambda \right) \right\}.$$

Consider the homomorphism from the Borel group $B \subset \mathbf{SL}(3, \mathbb{R})$ of upper triangular matrices with determinant one into H

$$j : B \rightarrow H$$

given by

$$\begin{pmatrix} a & c & e \\ 0 & \frac{1}{ab} & f \\ 0 & 0 & b \end{pmatrix} \longrightarrow \begin{pmatrix} \frac{a}{b} & -a^2f & \frac{c}{b} & -eb + \frac{1}{2}acf \\ 0 & a^2b & 0 & -\frac{1}{2}abc \\ 0 & 0 & \frac{1}{ab^2} & -\frac{f}{2b} \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

One verifies that the homomorphism is surjective so that H is isomorphic to the Borel group of upper triangular matrices in $\mathbf{SL}(3, \mathbb{R})$.

Proposition 3.2 *The bundle $Y \rightarrow M$ is a principal bundle with structure group H .*

3.2 Construction of connection forms on the bundle Y

The goal of this section is to review the construction of canonical forms defined on the coframe bundle $Y \rightarrow E$ as in [FV]. They give rise to a Cartan connection on Y with values in $\mathfrak{sl}(3, \mathbb{R})$.

A local section of the coframe bundle over M may be given by three forms

$$\theta, \theta^1, \theta^2$$

satisfying $d\theta = \theta^1 \wedge \theta^2$, with $\ker \theta^1_{|\ker \theta} = T^2$ and $\ker \theta^2_{|\ker \theta} = T^1$. They give coordinates on the cotangent bundle over E . Indeed, at $\lambda\theta \in E$, the coframes of Y are parametrized as follows:

$$\begin{aligned} \omega &= \lambda\theta \\ \omega^i &= a^i\theta^i + v^i\lambda\theta \end{aligned}$$

with $a^1a^2 = \lambda$ and

$$d\omega = 2\varphi \wedge \omega + \omega^1 \wedge \omega^2, \tag{19}$$

where $\varphi = \frac{d\lambda}{2\lambda} \pmod{\omega^1, \omega^2, \omega}$.

Differentiating the forms ω^1 and ω^2 we obtain new forms which correspond to the coordinates a^1, v^1, a^2, v^2 (recall that a^1 and a^2 are not independent):

Lemma 3.1 *There exists linearly independent forms w, φ^1, φ^2 defined on T^*Y such that*

$$d\omega^1 = \varphi \wedge \omega^1 + 3w \wedge \omega^1 + \omega \wedge \varphi^1 \text{ and } d\omega^2 = \varphi \wedge \omega^2 - 3w \wedge \omega^2 - \omega \wedge \varphi^2 \tag{20}$$

with $w = \frac{1}{6}(\frac{da^1}{a^1} - \frac{da^2}{a^2}) \pmod{(\omega, \omega^1, \omega^2)}$ and $\varphi^1 = -dv^1, \varphi^2 = dv^2 \pmod{(\omega, \omega^1, \omega^2)}$.

The coordinate s in the bundle Y is associated to a new form:

Lemma 3.2 *There exists a 1-form ψ such that*

$$d\varphi = \omega \wedge \psi - \frac{1}{2}(\varphi^2 \wedge \omega^1 + \varphi^1 \wedge \omega^2) \tag{21}$$

The forms w, φ^1, φ^2 and ψ are not yet determined. Define

$$\begin{aligned} W &= dw + \frac{1}{2}\omega^2 \wedge \varphi^1 - \frac{1}{2}\omega^1 \wedge \varphi^2 \\ \Phi^1 &= d\varphi^1 + 3\varphi^1 \wedge w + \omega^1 \wedge \psi + \varphi \wedge \varphi^1 \\ \Phi^2 &= d\varphi^2 - 3\varphi^2 \wedge w - \omega^2 \wedge \psi + \varphi \wedge \varphi^2 \end{aligned}$$

Lemma 3.3 *There exists unique forms w, φ^1, φ^2 and ψ such that $W = 0$, $\Phi^1 = Q^1\omega \wedge \omega^2$ and $\Phi^2 = Q^2\omega \wedge \omega^1$ where Q^1 and Q^2 are functions on Y .*

We can represent the structure equations 19, 20, 21 as a matrix equation whose entries are differential forms. The forms are disposed in the Lie algebra $\mathfrak{sl}(3, \mathbb{R})$ and define a Cartan connection on Y .

Proposition 3.3 *There exists a unique Cartan connection $\pi : TY \rightarrow \mathfrak{sl}(3, \mathbb{R})$ defined on Y of the form*

$$\pi = \begin{pmatrix} \varphi + w & \varphi^2 & \psi \\ \omega^1 & -2w & \varphi^1 \\ \omega & \omega^2 & -\varphi + w \end{pmatrix}.$$

such that its curvature satisfies

$$\Pi = d\pi + \pi \wedge \pi = \begin{pmatrix} 0 & \Phi^2 & \Psi \\ 0 & 0 & \Phi^1 \\ 0 & 0 & 0 \end{pmatrix}$$

with $\Phi^1 = Q^1\omega \wedge \omega^2$, $\Phi^2 = Q^2\omega \wedge \omega^1$ and $\Psi = (U_1\omega^1 + U_2\omega^2) \wedge \omega$.

3.3 Curvature forms and Bianchi identities

Curvature forms appear as differentials of connection forms and are used implicitly in order to fix the connection forms.

We recall:

$$W = dw - \frac{1}{2}\omega^2 \wedge \varphi^1 + \frac{1}{2}\omega^1 \wedge \varphi^2 = 0, \quad (22)$$

$$\Phi^1 = d\varphi^1 + 3\varphi^1 \wedge w + \omega^1 \wedge \psi + \varphi \wedge \varphi^1 = Q^1\omega \wedge \omega^2, \quad (23)$$

$$\Phi^2 = d\varphi^2 - 3\varphi^2 \wedge w - \omega^2 \wedge \psi + \varphi \wedge \varphi^2 = Q^2\omega \wedge \omega^1, \quad (24)$$

$$\Psi := d\psi - \varphi^1 \wedge \varphi^2 - 2\varphi \wedge \psi = (U_1\omega^1 + U_2\omega^2) \wedge \omega. \quad (25)$$

where Q^1, Q^2, U^1 and U^2 are functions on Y .

3.3.1

Equation $d(d\varphi^1) = 0$ obtained differentiating Φ^1 above implies

$$dQ^1 - 6Q^1w + 4Q^1\varphi = S^1\omega + U_2\omega^1 + T^1\omega^2, \quad (26)$$

‘where we introduced functions S^1 and T^1 .

3.3.2

Analogously, equation $d(d\varphi^2) = 0$ obtained differentiating Φ^2 above implies

$$dQ^2 + 6Q^2w + 4Q^2\varphi = S^2\omega - U_1\omega^2 + T^2\omega^1, \quad (27)$$

where we introduced new functions S^2 and T^2 .

3.3.3

Equation $d(d\psi) = 0$ obtained from 25 implies

$$dU_1 + 5U_1\varphi + 3U_1w + Q^2\varphi^1 = A\omega + B\omega^1 + C\omega^2 \quad (28)$$

and

$$dU_2 + 5U_2\varphi - 3U_2w - Q^1\varphi^2 = D\omega + C\omega^1 + E\omega^2. \quad (29)$$

3.4 Embedding $\iota_2 : Y_2 \rightarrow Y$

The goal now is to obtain an immersion $\iota_2 : Y_2 \rightarrow Y$. One can construct the bundle Y_2 using the bundle E of contact forms as a first step. Then Y_2 is a coframe bundle over E obtained by the tautological forms $\omega, \omega^1, \omega^2$ corresponding to forms $\theta, \theta^1, \theta^2$ satisfying $d\theta = \theta^1 \wedge \theta^2 + 2\varphi \wedge \omega$ with an appropriate φ .

By abuse of language again as for $\iota_1 : Y_1 \rightarrow Y_2$, we may write the connection forms of Y_1 and Y using the same letters and then obtain:

Proposition 3.4 *There exists a unique embedding $\iota_2 : Y_2 \rightarrow Y$ satisfying $\iota_2^*(\omega) = \omega$, $\iota_2^*(\omega^1) = \omega^1$, $\iota_2^*(\omega^2) = \omega^2$, $\iota_2^*(\varphi) = \varphi$.*

Proof. As Y_2 and Y are both coframe bundles over the line bundle E of all contact forms, we can assume that the embedding projects to the identity map on E . Over E , Y is a coframe bundle with structure group

$$\left\{ \begin{pmatrix} a & c & e \\ 0 & \frac{1}{a^2} & f \\ 0 & 0 & a \end{pmatrix} \right\}.$$

In order to determine the embedding we need to choose functions c, e and f . The diagonal matrix correspond to the fiber of Y_2 and does not need to be fixed. Consider then a map from M to the group above given by

$$h = \begin{pmatrix} 1 & c & e \\ 0 & 1 & f \\ 0 & 0 & 1 \end{pmatrix}.$$

Recall that

$$R_h^* \pi = h^{-1}dh + Ad_{h^{-1}}\pi.$$

We obtain, neglecting the terms of the connection of Y which are not relevant, the following transformation formulae. Remark that the term $h^{-1}dh$ does not appear in the transformation of these components.

$$\begin{aligned}
\tilde{\omega} &= \omega \\
\tilde{\omega}^1 &= \omega^1 - f\omega \\
\tilde{\omega}^2 &= \omega^2 + c\omega \\
\tilde{\varphi} &= \varphi - \frac{1}{2}c\omega^1 - f\omega^2 + \left(\frac{1}{2}cf - e\right)\omega
\end{aligned} \tag{30}$$

The forms ω^1 and ω^2 defined at each point of Y_2 define corresponding forms ω^1 and ω^2 in Y . We observe then that the functions f and c must be zero in order that $\iota_2^*(\omega^1) = \omega^1$, $\iota_2^*(\omega^2) = \omega^2$. Finally the form φ on Y_2 defines a corresponding form on Y and we conclude that $e = 0$ if we impose that $\iota_2^*(\varphi) = \varphi$.

□

3.4.1 The curvature of Y in terms of the curvature of Y_2

We obtain the following equations by pulling back to Y_2 the structure equations on Y through the embedding ι_2 :

$$d\omega = 2\varphi \wedge \omega + \omega^1 \wedge \omega^2$$

$$d\omega^1 = \varphi \wedge \omega^1 + 3\tilde{w} \wedge \omega^1 + \omega \wedge \varphi^1 \tag{31}$$

$$d\omega^2 = \varphi \wedge \omega^2 - 3\tilde{w} \wedge \omega^2 - \omega \wedge \varphi^2 \tag{32}$$

$$d\varphi = \omega \wedge \tilde{\psi} - \frac{1}{2}(\varphi^2 \wedge \omega^1 + \varphi^1 \wedge \omega^2) \tag{33}$$

$$d\tilde{w} = -\frac{1}{2}\varphi^2 \wedge \omega^1 + \frac{1}{2}\varphi^1 \wedge \omega^2 \tag{34}$$

$$d\varphi^1 + 3\varphi^1 \wedge \tilde{w} + \omega^1 \wedge \tilde{\psi} + \varphi \wedge \varphi^1 = Q^1\omega \wedge \omega^2 \tag{35}$$

$$d\varphi^2 - 3\varphi^2 \wedge \tilde{w} - \omega^2 \wedge \tilde{\psi} + \varphi \wedge \varphi^2 = Q^2\omega \wedge \omega^1$$

$$d\tilde{\psi} - \varphi^1 \wedge \varphi^2 - 2\varphi \wedge \tilde{\psi} = (U_1\omega^1 + U_2\omega^2) \wedge \omega.$$

In the formulae above we write the pull back of any form α defined on Y using the same notation α except for the pull backs $\tilde{w} = \iota_2^*w$ and $\tilde{\psi} = \iota_2^*\psi$. We should compare with the structure equations of Y_2 and obtain an expression for Q^1 and Q^2 :

$$d\omega = 2\varphi \wedge \omega + \omega^1 \wedge \omega^2$$

$$d\omega^1 = \varphi \wedge \omega^1 + 3w \wedge \omega^1 + \omega \wedge \tau^1$$

$$d\omega^2 = \varphi \wedge \omega^2 - 3w \wedge \omega^2 - \omega \wedge \tau^2$$

with $\tau^1 \wedge \omega^2 = \tau^2 \wedge \omega^1 = 0$.

Recall also that $d\varphi = \omega \wedge \psi$ with $\psi = A\omega^1 + B\omega^2$ and $dw = C\omega \wedge \omega^1 + D\omega \wedge \omega^2 + S\omega^1 \wedge \omega^2$, where A, B, C, D and S are functions on Y_2 .

- The differences between the pull back equations and the structure equations for $d\omega^1$ and $d\omega^2$ give, respectively,

$$3(\tilde{w} - w) \wedge \omega^1 + \omega \wedge (\varphi^1 - \tau^1) = 0$$

and

$$3(\tilde{w} - w) \wedge \omega^2 + \omega \wedge (\varphi^2 - \tau^2) = 0.$$

Therefore, by Cartan's lemma

$$\tilde{w} - w = m\omega$$

for a function m to be determined and

$$\varphi^1 - \tau^1 = -3m\omega^1 + n\omega \text{ and } \varphi^2 - \tau^2 = -3m\omega^2 + P\omega,$$

where n and P are functions to be determined.

- The difference between the equation $d\varphi = \omega \wedge \psi$ and the pull back equation for $d\varphi$ above is

$$\omega \wedge (\tilde{\psi} - \psi) - \frac{1}{2}(\varphi^2 \wedge \omega^1 + \varphi^1 \wedge \omega^2) = 0.$$

Substituting the expressions for φ^1 and φ^2 obtained in the item above we obtain

$$\omega \wedge (\tilde{\psi} - A\omega^1 - B\omega^2) - \frac{1}{2}((\tau^2 - 3m\omega^2 + P\omega) \wedge \omega^1 + (\tau^1 - 3m\omega^1 + n\omega) \wedge \omega^2) = 0,$$

which simplifies to

$$\omega \wedge (\tilde{\psi} - A\omega^1 - B\omega^2 - \frac{P}{2}\omega^1 - \frac{n}{2}\omega^2) = 0.$$

This implies that

$$\tilde{\psi} = (A + \frac{P}{2})\omega^1 + (B + \frac{n}{2})\omega^2 + q\omega,$$

where q is a function to be determined.

- The difference between the equations for $d\tilde{w}$ and dw is then

$$d(m\omega) = d\tilde{w} - dw = -\frac{1}{2}\varphi^2 \wedge \omega^1 + \frac{1}{2}\varphi^1 \wedge \omega^2 - S\omega^1 \wedge \omega^2 - C\omega \wedge \omega^1 - D\omega \wedge \omega^2.$$

Substituting in this formula the expressions for φ^1 and φ^2 in terms of the enriched structure we obtain

$$dm \wedge \omega + m(2\varphi \wedge \omega + \omega^1 \wedge \omega^2) = (-S - 3m)\omega^1 \wedge \omega^2 + (-\frac{P}{2} - C)\omega \wedge \omega^1 + (\frac{n}{2} - D)\omega \wedge \omega^2.$$

That is,

$$(S + 4m)\omega^1 \wedge \omega^2 + \omega \wedge \left(-dm - 2m\varphi + \left(\frac{P}{2} + C\right)\omega^1 - \left(\frac{n}{2} - D\right)\omega^2 \right) = 0.$$

Therefore

$$m = -S/4$$

and so

$$\frac{1}{4}dS + \frac{1}{2}S\varphi + \left(\frac{P}{2} + C\right)\omega^1 - \left(\frac{n}{2} - D\right)\omega^2 + E\omega = 0$$

where E is a function determined by the derivative of S . Writing $dS + 2S\varphi = S_0\omega + S_1\omega^1 + S_2\omega^2$ and comparing with the above expression, we obtain

$$S_0 = -4E, \quad S_1 = -2(P + 2C), \quad S_2 = -2(-n + 2D). \quad (36)$$

Therefore the functions P and n are determined. It remains to determine the function q .

- Computing $d\varphi^1 = d\left(\frac{3S}{4}\omega^1 + n\omega + \tau^1\right)$ and equating to the structure equation $d\varphi^1 = -3\varphi^1 \wedge \tilde{w} - \omega^1 \wedge \tilde{\psi} - \varphi \wedge \varphi^1 + Q^1\omega \wedge \omega^2$ we obtain, after a computation writing

$$dn + 3n(\varphi - w) = n_0\omega + n_1\omega^1 + n_2\omega^2, \quad (37)$$

$$n_1 = -\tau_2^1\tau_1^2 - q + \frac{9}{16}S^2 - 3E, \quad n_2 = -Q^1 + \frac{3}{2}S\tau_2^1 + \tau_{20}^1. \quad (38)$$

Recalling that $n = S_2/2 + 4D$, S and D are determined by Y_2 , we obtained an expression for Q^1 in terms of the enriched structure. Note also that q is determined by the first equation.

- Analogously, computing $d\varphi^2 = -3\frac{3S}{4}\omega^2 + p\omega + \tau^2$ and equating to the structure equation $d\varphi^2 - 3\varphi^2 \wedge \tilde{w} - \omega^2 \wedge \tilde{\psi} + \varphi \wedge \varphi^2 = Q^2\omega \wedge \omega^1$ we obtain, after a computation, writing $dP + 3P(\varphi + w) = P_0\omega + P_1\omega^1 + P_2\omega^2$,

$$P_1 = -Q^2 - \frac{3}{2}S\tau_1^2 + \tau_{10}^2, \quad P_2 = \tau_2^1\tau_1^2 + q - \frac{9}{16}S^2 - 3E. \quad (39)$$

Recalling that $P = -S_1/2 - 2C$, S and C are determined by Y_2 , we obtained an expression for Q^2 in terms of the enriched structure.

The following proposition follows directly from the computations above.

Proposition 3.5 *Suppose Y_2 with its adapted Cartan connection has null torsion, that is, satisfies $\tau^1 = \tau^2 = 0$ and $dw = S\omega^1 \wedge \omega^2$. Then*

$$Q^2 = \frac{1}{2}S_{11}$$

and

$$Q^1 = -\frac{1}{2}S_{22},$$

where S_{11} is the ω^1 component of the form dS_1 and S_{22} is the ω^2 component of the form dS_2 .

Proof. From Proposition 2.6, null torsion and the condition that $dw = S\omega^1 \wedge \omega^2$ (that is, $C = D = 0$) implies that $P = -S_1/2$ and $n = S_2/2$. The result is therefore implied from the previous formulas. \square

3.5 The embedding $Y_1 \rightarrow Y$

Recall that Y_1 is a coframe bundle of forms $(\theta, \theta^1, \theta^2)$ over M . Choosing a local section, the pullback forms over M are also denoted by $(\theta, \theta^1, \theta^2)$. We recall here an embedding $Y_1 \rightarrow Y$ obtained in [FV].

A section $(\theta, \theta^1, \theta^2)$ of the coframe bundle Y_1 clearly defines a path geometry on M . We obtain then a line bundle E and a principal bundle Y with its associated Cartan connection. Also, $(\theta, \theta^1, \theta^2)$ defines, up to the action by the group of matrices

$$\begin{pmatrix} 1 & c & e \\ 0 & 1 & f \\ 0 & 0 & 1 \end{pmatrix}$$

sections of the tautological forms $(\omega, \omega^1, \omega^2)$ on Y . In order to define a canonical section we use the following

Proposition 3.6 *Let $\theta, \theta^1, \theta^2$ be a coframe section of Y_1 and consider the principal bundle Y defined by this coframe. Then there exists a unique section $s : M \rightarrow Y$ such that $s^*\omega = \theta$, $s^*\omega^1 = \theta^1$, $s^*\omega^2 = \theta^2$ and $s^*\varphi = 0$.*

It is easy to verify that this definition is equivariant with respect to the action G_1 , the one parameter group of the strict contact structure. This defines then the embedding $Y_1 \rightarrow Y$.

3.6 The equivalence problem for a second order differential equation

In this section we recall the treatment by Cartan of the point equivalence between second order differential equations. It is included in order to fix conventions and to compare the invariants defined in the next section.

Recall that for a second order differential equation we define

$$\theta = dy - p dx,$$

and

$$L_1 = \ker\{dp - F dx\} \cap \ker\{dy - p dx\}, \quad L_2 = \ker dx \cap \ker dy.$$

For $Z^1 = dx$ and $Z^2 = dp - F dx$, one has then $d\theta = Z^1 \wedge Z^2$. The general forms defining the lines at each tangent space may be described by

$$\omega^1 = a_1 Z^1, \quad \omega^2 = a_2 Z^2, \quad \omega = a_1 a_2 \theta$$

where a_1, a_2 are non-vanishing positive functions on the manifold, so that we have always

$$2\varphi \wedge \omega + \omega^1 \wedge \omega^2 = d\omega = \left(\frac{da_1}{a_1} + \frac{da_2}{a_2}\right) \wedge \omega + a_1 Z^1 \wedge a_2 Z^2,$$

and we obtain comparing with 4

$$\varphi = \frac{1}{2}\left(\frac{da_1}{a_1} + \frac{da_2}{a_2}\right) + r\omega.$$

One computes

$$(\varphi + 3w) \wedge \omega^1 + \omega \wedge \tau^1 = d\omega^1 = da_1 \wedge Z^1 + a_1 \cdot 0 = \left(\frac{1}{2}\left(\frac{da_1}{a_1} + \frac{da_2}{a_2}\right) + \frac{1}{2}\left(\frac{da_1}{a_1} - \frac{da_2}{a_2}\right)\right) \wedge \omega^1 \quad (40)$$

and obtain

$$3w = \frac{1}{2}\left(\frac{da_1}{a_1} - \frac{da_2}{a_2}\right) - r\omega + s\omega^1, \tau_2^1 = 0.$$

Observe that, if $f(x, y, p)$ then

$$df = \frac{1}{a_1} \frac{df}{dx} \omega^1 + \frac{1}{a_2} f_p \omega^2 + \frac{1}{a_1 a_2} f_y \omega,$$

where $\frac{df}{dx} = f_x + f_y p + f_p F$. Also

$$\begin{aligned} d\omega^2 &= \frac{da_2}{a_2} \wedge \omega^2 + a_2 \left(-\frac{1}{a_2} F_p \omega^2 - \frac{1}{a_1 a_2} F_y \omega\right) \wedge \frac{1}{a_1} \omega^1 \\ &= \left(\frac{1}{2}\left(\frac{da_1}{a_1} + \frac{da_2}{a_2}\right) - \frac{1}{2}\left(\frac{da_1}{a_1} - \frac{da_2}{a_2}\right)\right) \wedge \omega^2 + \frac{1}{a_1} F_p \omega^1 \wedge \omega^2 - \frac{1}{(a_1)^2} F_y \omega \wedge \omega^1 \\ &= (\varphi - 3w - 2r\omega + (s + \frac{1}{a_1} F_p) \omega^1) \wedge \omega^2 - \omega \wedge \frac{F_y}{(a_1)^2} \omega^1 \end{aligned} \quad (41)$$

Comparing with $d\omega^2 = (\varphi - 3w) \wedge \omega^2 - \omega \wedge \tau^2$ we obtain $r = 0, s = -\frac{F_p}{a_1}, \tau_1^2 = \frac{F_y}{a_1^2}$,

$$\varphi = \frac{1}{2}\left(\frac{da_1}{a_1} + \frac{da_2}{a_2}\right), 3w = \frac{1}{2}\left(\frac{da_1}{a_1} - \frac{da_2}{a_2}\right) - \frac{F_p}{a_1} \omega^1.$$

From $d\varphi = 0$, we obtain $\psi = 0$, or $A = B = 0$, and from $\tau_2^1 = 0$ it follows $D = \tau_{20}^1 = \tau_{22}^1 = 0$.

From above we get

$$3dw = -dF_p \wedge \frac{1}{a_1} \omega^1 = -\left(\frac{F_{pp}}{a_2} \omega^2 + \frac{F_{py}}{a_1 a_2} \omega\right) \wedge \frac{1}{a_1} \omega^1$$

and comparing with $dw = C\omega \wedge \omega^1 + S\omega^1 \wedge \omega^2$ we obtain

$$3C = -\frac{F_{py}}{a_1^2 a_2}, \quad 3S = \frac{F_{pp}}{a_1 a_2}.$$

Also

$$d\tau_1^2 = d\left(\frac{F_y}{a_1^2}\right) = -\frac{2}{a_1^2}(\varphi + 3w + \frac{F_p}{a_1} \omega^1) F_y + \frac{1}{a_1^2} \left(\frac{1}{a_1} \frac{dF_y}{dx} \omega^1 + \frac{1}{a_1 a_2} F_{yy} \omega + \frac{1}{a_2} F_{yp} \omega^2\right).$$

Comparing with $d\tau_1^2 = -2\tau_1^2(\varphi + 3w) - 3C\omega^2 + \tau_{10}^2\omega + \tau_{11}^2\omega^1$ we obtain

$$\tau_{11}^2 = \frac{1}{a_1^3}(-2F_p F_y + \frac{dF_y}{dx}), \quad \tau_{10}^2 = \frac{1}{a_1^3 a_2} F_{yy}.$$

Now

$$3dS = -2(3S\varphi) + \frac{1}{a_1 a_2} (\frac{1}{a_1} \frac{dF_{pp}}{dx} \omega^1 + \frac{1}{a_2} F_{ppp} \omega^2 + \frac{1}{a_1 a_2} F_{ppy} \omega)$$

and comparing with $dS = -2S\varphi + S_0\omega + S_1\omega^1 + S_2\omega^2$ we obtain

$$S_1 = \frac{1}{3a_1^2 a_2} \frac{dF_{pp}}{dx}, \quad S_2 = \frac{1}{3a_1 a_2^2} F_{ppp}, \quad S_0 = \frac{1}{3a_1^2 a_2^2} F_{ppy}.$$

It follows from $S_1 = -2P - 4C$ that $6P = \frac{1}{a_1^2 a_2} (4F_{yp} - \frac{dF_{pp}}{dx})$. Then

$$\begin{aligned} 6dP = & -\frac{1}{a_1^2 a_2} (3\varphi + 3w + \frac{F_p}{a_1} \omega^1) (4F_{yp} - \frac{dF_{pp}}{dx}) + \frac{4}{a_1^2 a_2} (\frac{1}{a_1} \frac{dF_{yp}}{dx} \omega^1 + \frac{1}{a_2} F_{ypp} \omega^2 + \frac{1}{a_1 a_2} F_{yyp} \omega) \\ & - \frac{1}{a_1^2 a_2} (\frac{1}{a_1} \frac{d^2 F_{pp}}{dx^2} \omega^1 + \frac{1}{a_2} (\frac{dF_{ppp}}{dx} + F_{ypp} + F_{ppp} F_p) \omega^2 + \frac{1}{a_1 a_2} (\frac{dF_{ypp}}{dx} + F_{ppp} F_y) \omega) \end{aligned}$$

Comparing with $dP = -(3\varphi + 3w)P + P_0\omega + P_1\omega^1 + P_2\omega^2$ we obtain

$$\begin{aligned} P_0 = & \frac{1}{6a_1^3 a_2^2} (4F_{yyp} - \frac{dF_{ypp}}{dx} - F_{ppp} F_y), \quad P_1 = \frac{1}{6a_1^3 a_2} (-4F_p F_{yp} + F_p \frac{dF_{pp}}{dx} + 4\frac{dF_{yp}}{dx} - \frac{d^2 F_{pp}}{dx^2}), \\ P_2 = & \frac{1}{6a_1^2 a_2^2} (4F_{ypp} - \frac{dF_{ppp}}{dx} - F_{ypp} - F_{ppp} F_p). \end{aligned}$$

From $Q^2 = \tau_{10}^2 - \frac{3}{2}S\tau_1^2 - P_1$ it follows

$$Q^2 = \frac{1}{6a_1^3 a_2} (6F_{yy} - 3F_y F_{pp} + 4F_p F_{yp} - F_p \frac{dF_{pp}}{dx} - 4\frac{dF_{yp}}{dx} + \frac{d^2 F_{pp}}{dx^2}). \quad (42)$$

It follows from $S_2 = 2n - 4D$ that $6n = \frac{1}{a_1 a_2^2} F_{ppp}$. Then

$$6dn = -\frac{1}{a_1 a_2^2} (3\varphi - 3w - \frac{F_p}{a_1} \omega^1) F_{ppp} + \frac{1}{a_1 a_2^2} (\frac{1}{a_1} \frac{dF_{ppp}}{dx} \omega^1 + \frac{1}{a_2} F_{pppp} \omega^2 + \frac{1}{a_1 a_2} F_{pppy} \omega).$$

Comparing with $dn = -n(3\varphi - 3w) + n_0\omega + n_1\omega^1 + n_2\omega^2$ we obtain

$$n_0 = \frac{1}{6a_1^2 a_2^3} F_{pppy}, \quad n_1 = \frac{1}{6a_1^2 a_2^2} (\frac{dF_{ppp}}{dx} + F_p F_{ppp}), \quad n_2 = \frac{1}{6a_1 a_2^3} F_{pppp}.$$

From $Q^1 = \tau_{20}^1 + \frac{3}{2}S\tau_2^1 - n_2$ it follows

$$Q^1 = -\frac{1}{6a_1 a_2^3} F_{pppp}. \quad (43)$$

Formulas 43 and 42 are in [Car].

4 A global invariant

In this section we define the global invariant for path structures. It has a very similar definition with the global invariant obtained in [FV] in the context of a structure defined on the complexified tangent space of a 3-manifold. But we make the definition explicit in the case of path structures for the sake of clarity and to adapt differences of conventions with our previous paper.

Define the second Chern class of the bundle Y with connection form π as

$$c_2(Y, \pi) = \frac{1}{8\pi^2} \text{tr}(\Pi \wedge \Pi).$$

$$\begin{pmatrix} 0 & \Phi^2 & \Psi \\ 0 & 0 & \Phi^1 \\ 0 & 0 & 0 \end{pmatrix} \wedge \begin{pmatrix} 0 & \Phi^2 & \Psi \\ 0 & 0 & \Phi^1 \\ 0 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 & \Phi^1 \wedge \Phi^2 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

As $\Phi^1 = Q^1\omega \wedge \omega^2$ and $\Phi^2 = Q^2\omega \wedge \omega^1$ we have $\Pi \wedge \Pi = 0$ and therefore

$$c_2(Y, \pi) = 0.$$

Definition 4.1 *The transgression form is defined as*

$$TC_2(\pi) = \frac{1}{8\pi^2} \left(\text{tr}(\pi \wedge \Pi) + \frac{1}{3} \text{tr}(\pi \wedge \pi \wedge \pi) \right) = \frac{1}{24\pi^2} \text{tr}(\pi \wedge \pi \wedge \pi).$$

Lemma 4.1 *The transgression form is closed, that is, $dTC_2(\pi) = c_2(Y, \pi) = 0$.*

Proof. We compute first, using the expressions of Φ^1 , Φ^2 and Ψ , that

$$\text{tr}(\Pi \wedge \pi) = \Phi^2 \wedge \omega^1 + \Phi^1 \wedge \omega^2 + \Psi \wedge \omega = 0.$$

Differentiating the curvature form we obtain $d\Pi = \Pi \wedge \pi - \pi \wedge \Pi$ and therefore

$$\begin{aligned} 0 &= d \text{tr}(\Pi \wedge \pi) = \text{tr}(d\Pi \wedge \pi + \Pi \wedge d\pi) = \text{tr}((\Pi \wedge \pi - \pi \wedge \Pi) \wedge \pi + \Pi \wedge (\Pi - \pi \wedge \pi)) \\ &= -\text{tr}(\pi \wedge \Pi \wedge \pi). \end{aligned}$$

Note that $\text{tr}(\alpha \wedge \beta) = (-1)^{kl} \text{tr}(\beta \wedge \alpha)$ if α and β are two matrices of forms of degree k and l respectively. Therefore, computing

$$\begin{aligned} \frac{1}{3} d \text{tr}(\pi \wedge \pi \wedge \pi) &= \text{tr}(d\pi \wedge \pi \wedge \pi) = \text{tr}((\Pi - \pi \wedge \pi) \wedge \pi \wedge \pi) \\ &= -\text{tr}(\pi \wedge \pi \wedge \pi \wedge \pi) = 0. \end{aligned}$$

□

Definition 4.2 Suppose that the fiber bundle $Y \rightarrow M$ is trivial and let $s : M \rightarrow Y$ be a section, we define then

$$\mu = \int_M s^* TC_2(\pi) = \frac{1}{24\pi^2} \int_M s^* \text{tr}(\pi \wedge \pi \wedge \pi).$$

In principle that integral depends on the section but the following proposition shows that the integrand

$$s^* TC_2(\pi)$$

defines an element in $H^3(M, \mathbb{R})$ which does not depend on the section.

Proposition 4.3 Suppose s and \tilde{s} are two sections. Then

$$\tilde{s}^* TC_2(\pi) - s^* TC_2(\pi) = -\frac{1}{8\pi^2} d s^* \text{tr}(h^{-1} \pi \wedge dh).$$

where $h : M \rightarrow H$ is a map such that $\tilde{s} = R_h \circ s$.

Proof. Fix the section s . Then there exists a map $h : M \rightarrow H$ such that $\tilde{s} = R_h \circ s$. We have then

$$\tilde{s}^* TC_2(\pi) = \frac{1}{24\pi^2} s^* \text{tr}(R_h^* \pi \wedge R_h^* \pi \wedge R_h^* \pi).$$

From the formula

$$R_h^* \pi = h^{-1} dh + Ad_{h^{-1}} \pi,$$

we obtain

$$\begin{aligned} & \text{tr}(R_h^* \pi \wedge R_h^* \pi \wedge R_h^* \pi) = \\ & \text{tr}(h^{-1} dh \wedge h^{-1} dh \wedge h^{-1} dh + 3h^{-1} dh \wedge h^{-1} \pi \wedge dh + 3h^{-1} \pi \wedge \pi \wedge dh + \pi \wedge \pi \wedge \pi) \\ & = \text{tr}(-h^{-1} dh \wedge dh^{-1} \wedge dh - 3dh^{-1} \wedge \pi \wedge dh + 3h^{-1} \pi \wedge \pi \wedge dh + \pi \wedge \pi \wedge \pi). \end{aligned}$$

Observe that the first term in the right hand side vanishes. Indeed, $dh^{-1} \wedge dh$ is upper triangular with null diagonal. Moreover $h^{-1} dh$ is upper triangular and therefore the Lie algebra valued form also has zero diagonal. Therefore

$$\text{tr}(h^{-1} dh \wedge dh^{-1} \wedge dh) = 0.$$

By the same argument $\text{tr}(h^{-1} \pi \wedge dh) = 0$.

Now we show that

$$d \text{tr}(h^{-1} \pi \wedge dh) = \text{tr}(dh^{-1} \wedge \pi \wedge dh - h^{-1} \pi \wedge \pi \wedge dh).$$

$$\text{Compute } d \text{tr}(h^{-1} \pi \wedge dh) = \text{tr}(dh^{-1} \wedge \pi \wedge dh + h^{-1} d\pi \wedge dh)$$

$$= \text{tr}(dh^{-1} \wedge \pi \wedge dh + h^{-1}(\Pi - \pi \wedge \pi) \wedge dh),$$

which gives, using that $\text{tr}(h^{-1} \Pi \wedge dh) = 0$,

$$d \text{tr}(h^{-1} \pi \wedge dh) = \text{tr}(dh^{-1} \wedge \pi \wedge dh - h^{-1} \pi \wedge \pi \wedge dh).$$

We obtained therefore that

$$\tilde{e}s^*TC_2(\pi) = s^*TC_2(\pi) - \frac{1}{8\pi^2}d s^*\text{tr}(h^{-1}\pi \wedge dh)$$

and this completes the proof of the proposition. □

Let $\mu(t)$ be the invariant defined as a function of the a parameter describing the deformation of the structure on a closed manifold M and define $\delta\mu = \frac{d}{dt}\mu(0)$. One can interpret the flat structures as giving critical points of the global invariant μ through the first variation formula which we refer to [FV] for a proof.

Proposition 4.4

$$\delta\mu = -\frac{1}{4\pi^2} \int_M s^*\text{tr}(\dot{\pi} \wedge \Pi).$$

The global invariant can be computed most easily for a path structure induced by an enriched or strict path structure.

Proposition 4.5 *Let M be an enriched path structure and $Y_2 \rightarrow Y$ be the canonical embedding of the enriched geometry into the induced path geometry. Then*

$$\begin{aligned} 8\pi^2\mu(Y) &= (n_1 - \frac{3}{4}S_0 + 2\tau_2^1\tau_1^2)\omega \wedge \omega^1 \wedge \omega^2 + \omega \wedge \omega^1 \wedge (-2A\varphi - (\frac{3}{2}S_1 + 6C)w) \\ &\quad + \omega \wedge \omega^2(-2B\varphi - (\frac{3}{2}S_2 + 6D)w) - \frac{9}{2}Sw \wedge \omega^1 \wedge \omega^2. \end{aligned}$$

Proof. One compute first the following formula.

$$\frac{1}{3}\text{tr}(\pi \wedge \pi \wedge \pi) = (2\omega \wedge \varphi - \omega^1 \wedge \omega^2) \wedge \psi - \omega \wedge \varphi^1 \wedge \varphi^2 + \omega^1 \wedge (\varphi + 3w) \wedge \varphi^2 + \omega^2 \wedge (\varphi - 3w) \wedge \varphi^1.$$

Therefore using the embedding of $Y_2 \rightarrow Y$ in the previous section we obtain by a computation:

$$\begin{aligned} \frac{1}{3}\text{tr}(\pi \wedge \pi \wedge \pi) &= (n_1 - \frac{3}{4}S_0 + 2\tau_2^1\tau_1^2)\omega \wedge \omega^1 \wedge \omega^2 + \omega \wedge \omega^1 \wedge (-2A\varphi - (\frac{3}{2}S_1 + 6C)w) \\ &\quad + \omega \wedge \omega^2(-2B\varphi - (\frac{3}{2}S_2 + 6D)w) - \frac{9}{2}Sw \wedge \omega^1 \wedge \omega^2. \end{aligned}$$

□

Using the embedding of $Y_1 \rightarrow Y$ in the previous section we obtain by a similar computation:

Proposition 4.6 *Let M be a strict path structure and $Y_1 \rightarrow Y$ be the canonical embedding of the strict geometry into the induced path geometry. Then*

$$8\pi^2\mu(Y) = (n_1 - \frac{3}{4}S_0 + 2\tau_2^1\tau_1^2)\omega \wedge \omega^1 \wedge \omega^2 - w \wedge ((\frac{3}{2}S_1 + 6C)\omega \wedge \omega^1 + (\frac{3}{2}S_2 + 6D)\omega \wedge \omega^2 + \frac{9}{2}S\omega^1 \wedge \omega^2) \tag{44}$$

4.1 The global invariant for second order differential equations on the torus

In this section we obtain formulas for the global invariant in the case of an ordinary differential equation defined on the torus. Recall that the projectivized cotangent bundle $\pi : PT^*S \rightarrow S$ of a surface S is described locally by $(x, y, [p, q])$ where (x, y) are local coordinates on the surface and $pdx + qdy$ is a form at (x, y) . The Liouville form θ on T^*S is defined to be the tautological form $\theta(x, y, -pdx + qdy) = \pi^*(-pdx + qdy)$. It induces a contact distribution on PT^*S , which in the chart $(x, y, p) \rightarrow (x, y, [p, 1])$ is given by the kernel of the form $dy - pdx$. On the chart $(x, y, q) \rightarrow (x, y, [1, q])$ the contact distribution is the kernel of $dx - qdy$. One can also consider, fixing a metric on the surface, the unit cotangent bundle $(T^*)^1S$ which is a double cover of PT^*S .

The fibers of the bundle PT^*S give a canonical field of directions on the Liouville distribution. Observe that, in local coordinates (x, y, p) , it is described by $\ker dx \cap \ker dy = \ker dx \cap \ker(dy - pdx)$. Choosing another direction on the contact distribution amounts to define a form, in local coordinates (x, y, p) , $dp - G(x, y, p)dx$, where $G(x, y, p)$ is a function. On the chart (x, y, q) one writes then

$$d\left(\frac{1}{q}\right) - G(x, y, \frac{1}{q})dx = -\frac{1}{q^2}dq - G(x, y, \frac{1}{q})dx.$$

Therefore, the direction is determined by $dq + G(x, y, \frac{1}{q})q^3dy$ (the contact distribution is $\ker(dx - qdy)$). In order to have a well defined direction we need that the function $G(x, y, \frac{1}{q})q^3$ has a differentiable extension for $q = 0$.

Definition 4.7 *A second order differential equation on a surface S is a path structure on the projective cotangent bundle with contact structure induced by the Liouville form and such that one of the directions is given by the fibers.*

It is convenient to introduce a new coordinate in the fiber $\alpha \in]-\pi, \pi]$ through the formula $p = \tan \alpha/2$. The contact distribution is defined by a globally defined form on the coordinates (x, y, α) :

$$\theta = \cos \alpha/2dy - \sin \alpha/2dx.$$

The fiber direction is defined by the equations $dx = dy = 0$ which can also be described by, defining $\theta^1 = \sin \alpha/2dy + \cos \alpha/2dx$, as $\ker \theta^1 \cap \ker \theta$. The last form, which depends on a choice of a function, is

$$\theta^2 = d\alpha - F(x, y, \alpha)\theta^1.$$

Observe that

$$d\theta = \frac{1}{2}\theta^1 \wedge \theta^2.$$

The relation with the differential equation given on the chart (x, y, p) is given writing

$$dp = \frac{1}{2}(1 + p^2)d\alpha$$

and therefore, as $dy = p dx$ in that chart,

$$\begin{aligned} d\alpha - F(x, y, \alpha)\theta^1 &= \frac{2}{1+p^2} dp - F(x, y, 2 \arctan p)(\sin \alpha/2 dy + \cos \alpha/2 dx) \\ &= \frac{2}{1+p^2} dp - F(x, y, 2 \arctan p)(\sin \alpha/2 \cdot p dx + \cos \alpha/2 dx) \\ &= \frac{2}{1+p^2} dp - F(x, y, 2 \arctan p)(\sin \alpha/2 \cdot \tan \alpha/2 + \cos \alpha/2) dx. \end{aligned}$$

and recalling that $\cos \alpha/2 = \frac{1}{\sqrt{1+p^2}}$,

$$= \frac{2}{1+p^2} dp - F(x, y, 2 \arctan p)(1+p^2)^{1/2} dx.$$

Therefore $2G(x, y, p) = F(x, y, 2 \arctan p)(1+p^2)^{3/2}$.

4.1.1 The strict and enriched structure of a differential equation on the torus

Here we will work with a double cover of the projective cotangent bundle of the torus. We define the path structure associated to a differential equation on the torus through a strict path structure defined by

$$\begin{aligned} \theta &= \cos \alpha dy - \sin \alpha dx. \\ \theta^1 &= \sin \alpha dy + \cos \alpha dx \end{aligned}$$

and

$$\theta^2 = d\alpha - F(x, y, \alpha)\theta^1.$$

Here $F(x, y, \alpha)$ is a function defined on the torus. Observe that

$$d\theta = \theta^1 \wedge \theta^2.$$

In the following we will write, for a function $f : T^3 \rightarrow \mathbb{R}$, defined on the torus,

$$df = f_0 \theta + f_1 \theta^1 + f_2 \theta^2,$$

so that $f_x = -f_0 \sin \alpha + (f_1 - f_2 F) \cos \alpha$, $f_y = f_0 \cos \alpha + (f_1 - f_2 F) \sin \alpha$ and $f_\alpha = f_2$. Compute

$$d\theta^1 = (\theta^2 + F\theta^1) \wedge \theta$$

and

$$d\theta^2 = ((F_0 - F^2)\theta^1 - F\theta^2) \wedge \theta + F_2\theta^1 \wedge \theta^2.$$

Consider now the enriched structure defined by θ, θ^1 and θ^2 and the tautological forms $\omega = a_1 a_2 \theta, \omega^1 = a_1 \theta^1$ and $\omega^2 = a_2 \theta^2$.

We first compute

$$d\omega = 2\varphi \wedge \omega + \omega^1 \wedge \omega^2,$$

where $\varphi = \frac{1}{2}(\frac{da_1}{a_1} + \frac{da_2}{a_2})$. Next we compute $d\omega^1$ and $d\omega^2$:

$$d\omega^1 = \frac{da_1}{a_1} \wedge a_1\theta^1 + a_1(\theta^2 + F\theta^1) \wedge \theta$$

$$d\omega^2 = \frac{da_2}{a_2} \wedge a_2\theta^2 + a_2(((F_0 - F^2)\theta^1 - F\theta^2) \wedge \theta + F_2\theta^1 \wedge \theta^2)$$

Comparing with the structure equations of the enriched structure in 5 we may write

$$d\omega^1 = \varphi \wedge \omega^1 + 3w \wedge \omega^1 + \omega \wedge \tau^1 \quad d\omega^2 = \varphi \wedge \omega^2 - 3w \wedge \omega^2 - \omega \wedge \tau^2. \quad (45)$$

with

$$3w = \frac{1}{2}\left(\frac{da_1}{a_1} - \frac{da_2}{a_2}\right) - \frac{1}{a_1a_2}F\omega - \frac{1}{a_1}F_2\omega^1,$$

$$\tau^1 = -\frac{1}{a_2^2}\omega^2$$

and

$$\tau^2 = \frac{1}{a_1^2}(F_0 - F^2)\omega^1.$$

4.1.2 Curvatures

We compute now $d\varphi = \omega \wedge (A\omega^1 + B\omega^2)$ and $dw = C\omega \wedge \omega^1 + D\omega \wedge \omega^2 + S\omega^1 \wedge \omega^2$ (see Proposition 2.5). From $d\varphi = 0$ we obtain $A = B = 0$. Computing dw and comparing to the formula above we obtain

$$C = \frac{F_1 - F_{20} + FF_2}{3a_1^2a_2}, \quad D = \frac{2F_2}{3a_1a_2^2}, \quad S = \frac{F_{22} - F}{3a_1a_2}.$$

In order to compute the global invariant we need to compute the coefficients S_0, S_1 and S_2 in equation 17 (a Bianchi identity) : $dS + 2S\varphi = S_0\omega + S_1\omega^1 + S_2\omega^2$. One obtains

$$S_0 = \frac{F_{220} - F_0}{3a_1^2a_2^2}, \quad S_1 = \frac{F_{221} - F_1}{3a_1^2a_2}, \quad S_2 = \frac{F_{222} - F_2}{3a_1a_2^2}.$$

Now we use the expressions obtained in section 3.4.1 of the curvatures of Y in terms of the curvature of Y_2 . In order to compute $\mu(Y)$ we need to compute n and its derivatives (see 36 and 37).

We have $2n = S_2 + 4D = \frac{F_{222} + 7F_2}{3a_2^2a_1}$ and compute the left hand of $dn + 3(\varphi - w)n = n_0\omega + n_1\omega^1 + n_2\omega^2$ (formula 37) to obtain then

$$n_0 = \frac{1}{6a_1^2a_2^3} (F(F_{222} + F_2) + F_{2220} + 7F_{20}),$$

$$n_1 = \frac{1}{6a_1^2a_2^2} (F_2(F_{222} + F_2) + F_{2221} + 7F_{21})$$

$$n_2 = \frac{1}{6a_1a_2^3} (F_{2222} + 7F_{22})$$

We use equations 38 and 39 to compute the curvature functions Q^1 and Q^2 . We have $Q^1 = -n_2 + \frac{3}{2}S\tau_2^1 + \tau_{20}^1$ and $Q^2 = -P_1 - \frac{3}{2}S\tau_1^2 + \tau_{10}^2$. For that sake, we compute first the derivatives of the torsion, τ_{20}^1 and τ_{10}^2 , using formulas 11 and 12. Computing the left hand side of the equation $d\tau_2^1 + 2\tau_2^1(\varphi - 3w) + (B + 3D)\omega^1 = \tau_{20}^1\omega + \tau_{22}^1\omega^2$ and comparing to the right hand side we obtain

$$\tau_{20}^1 = -\frac{2F}{a_1a_2^3}.$$

Analogously, computing the left hand side of the equation $d\tau_1^2 + 2\tau_1^2(\varphi + 3w) - (A - 3C)\omega^2 = \tau_{10}^2\omega + \tau_{11}^2\omega^1$ we obtain

$$\tau_{10}^2 = \frac{F_{00} - 4FF_0 + 2F^3}{a_1^3a_2}.$$

Proposition 4.8 *Given a (local) differential equation as a path structure induced by the forms θ, θ^1 and θ^2 as above one computes the curvature functions in terms of the enriched structure:*

$$Q^1 = -\frac{1}{6a_1a_2^3}(F_{2222} + 10F_{22} + 9F)$$

and

$$Q^2 = \frac{1}{6a_1^3a_2}(F_2F_{221} + 3F_2F_1^2 - 4F_2F_{20} + 4FF_2^2 + F_{2211} + 6F_1F_{11} - 4F_{201} + 4F_2F_1 + 4FF_{21} - 3F_{22}F_0 + 3F_{22}F^2 - 21FF_0 + 9F^3 + 6F_{00}).$$

Proof. Recall from formula 36 that $P = -S_1/2 - 2C$ and we write $dP + 3P(\varphi + w) = P_0\omega + P_1\omega^1 + P_2\omega^2$. Computing the left side and comparing the right side we obtain the expression of P_1 which we use in the formulas above. \square

The following proposition describes locally differential equations satisfying $Q^1 = 0$.

Proposition 4.9 *Differential equations on an open subset with coordinates (x, y) given by $\theta = \cos \alpha dy - \sin \alpha dx$, $\theta^1 = \sin \alpha dy + \cos \alpha dx$ and $\theta^2 = d\alpha - F(x, y, \alpha)\theta^1$ satisfy*

$$Q^1 = 0$$

if and only if

$$F(x, y, \alpha) = A(x, y) \cos \alpha + B(x, y) \sin \alpha + C(x, y) \cos 3\alpha + D(x, y) \sin 3\alpha$$

where A, B, C and D are functions on x and y .

Proof. Observe that $Q^1 = 0$ is equivalent to $F_{2222} + 10F_{22} + 9F = F_{\alpha\alpha\alpha\alpha} + 10F_{\alpha\alpha} + 9F = 0$. The only solutions to this linear equation are of the form above. \square

Using the coordinates (x, y, p) as above where the differential equation is described as $dp - G(x, y, p)dx = 0$ the condition $Q^1 = 0$ implies that $G(x, y, p)$ is at most a third order polynomial in p with coefficients functions of x and y (see [A]).

4.1.3 The global invariant

We are ready now to use Proposition 4.5 to determine the global invariant:

Proposition 4.10 *Let M be an enriched path structure defined by an ordinary differential equation of second order on the torus with strict structure defined by the forms θ, θ^1 and θ^2 as above. Let $Y_2 \rightarrow Y$ be the canonical embedding of the enriched geometry into the induced path geometry whose connection is π . Then*

$$8\pi^2 s^*(TC_2(\pi)) = \frac{1}{12}(-12F_\alpha^2 + 2(F_{\alpha\alpha x} \cos \alpha + F_{\alpha\alpha y} \sin \alpha + F_{\alpha\alpha\alpha} F) + 14(F_{\alpha x} \cos \alpha + F_{\alpha y} \sin \alpha + F_{\alpha\alpha} F) - 3(-F_{\alpha\alpha x} \sin \alpha + F_{\alpha\alpha y} \cos \alpha) + 3F_\alpha - 24(-F_x \sin \alpha + F_y \cos \alpha) + 18F^2 + 6FF_{\alpha\alpha})\theta \wedge \theta^1 \wedge \theta^2.$$

and

$$8\pi^2 \mu(Y) = 8\pi^2 \int_M s^*(TC_2(\pi)) = \frac{1}{12} \int_M (-32F_\alpha^2 + 2F_{\alpha\alpha}^2 + 18F^2)\theta \wedge \theta^1 \wedge \theta^2.$$

Proof. The terms in Proposition 4.5 were all computed before. A substitution of these terms in the formula gives the first formula. The second formula is obtained by integration by parts.

□

Corollary 4.11 *Let M be equipped with a path structure defined by an ordinary differential equation of second order on the torus with strict structure defined by the forms θ, θ^1 and θ^2 . Let Y be the canonical Cartan bundle with its associated Cartan connection.*

1. *If $\theta^2 = d\alpha - F(x, y)\theta^1$ (the function F does not depend on α). Then $\mu(Y) = 0$ if and only if $F = 0$.*
2. *$Q^1(Y) = 0$ and $\mu(Y) = 0$ if and only if $F = 0$.*

Proof. Clearly, if $F = 0$ then $\mu(Y) = 0$ and $Q^1 = 0$. If F does not depend on α then the invariant becomes

$$\frac{4}{3} \int_M F^2 \theta \wedge \theta^1 \wedge \theta^2,$$

which is zero only if $F = 0$. Suppose now that $Q^1(Y) = 0$ and $\mu(Y) = 0$. Observe that the integral formula for the invariant, by an integration by part and a slight rearrangement, may be written as

$$\frac{1}{12} \int_M (-12F_\alpha^2 + 2F_{\alpha\alpha\alpha} F + 20F_{\alpha\alpha} F + 18F^2)\theta \wedge \theta^1 \wedge \theta^2.$$

Using the expression of Q^1 given in 4.8 and the hypothesis $Q^1 = 0$, we obtain $F_{2222} + 10F_{22} + 9F = 0$, and therefore

$$2F_{\alpha\alpha\alpha} F + 20F_{\alpha\alpha} F + 18F^2 = 0.$$

Therefore

$$8\pi^2\mu(Y) = - \int_M F_\alpha^2 \theta \wedge \theta^1 \wedge \theta^2.$$

We observe therefore that if $\mu(Y) = 0$ then F_α should be null. But if F does not depend on α it should be null by the first part. \square

5 Path structures on a torus

We recall example **II** which is the torus T^3 with coordinates $(x, y, t) \pmod{1}$ and the global contact form, for a fixed $n \in \mathbb{Z}^*$,

$$\theta = \cos(2\pi nt)dx - \sin(2\pi nt)dy.$$

It was proven independently by E. Giroux and Y. Kanda that the contact structures defined by these contact forms classify all tight structures on T^3 (see [?]). We will show here that for each of these contact structures one can define a flat path structure.

There are two canonical global vector fields on the distribution given by $X_1 = \frac{\partial}{\partial t}$ and $X_2 = \sin(2\pi nt)\frac{\partial}{\partial x} + \cos(2\pi nt)\frac{\partial}{\partial y}$.

We define

$$\theta^1 = -2\pi n dt, \quad \theta^2 = \sin(2\pi nt)dx + \cos(2\pi nt)dy,$$

so that $d\theta = \theta^1 \wedge \theta^2$ and we define the strict path structure defined by these forms. We compute

$$d\theta^1 = 0, \quad d\theta^2 = -\theta^1 \wedge \theta.$$

Comparing now with the enriched path connection we obtain

$$\varphi = \frac{1}{2}\left(\frac{da_1}{a_1} + \frac{da_2}{a_2}\right), \quad 3w = \frac{1}{2}\left(\frac{da_1}{a_1} - \frac{da_2}{a_2}\right), \quad \tau_2^1 = 0, \quad \tau_1^2 = -\frac{1}{a_1^2}$$

and therefore $d\varphi = dw = d\tau_2^1 = d\tau_1^2 + 2\tau_1^2(\varphi + 3w) = 0$. It follows that $A = B = C = D = S = \tau_{20}^1 = \tau_{10}^2 = 0$, and it follows from formulas 38 and 39 that $Q^1 = Q^2 = 0$.

We proved:

Lemma 5.1 *The path structures defined by the forms $\theta^1, \theta^2, \theta$ on T^3 are flat.*

We now define a new strict path structure by fixing the contact form θ and changing θ^1 and θ^2 by a constant matrix:

$$\begin{pmatrix} \theta^{1'} \\ \theta^{2'} \end{pmatrix} = \begin{pmatrix} a & b \\ c & f \end{pmatrix} \begin{pmatrix} \theta^1 \\ \theta^2 \end{pmatrix} \quad \text{or} \quad \begin{pmatrix} \theta^1 \\ \theta^2 \end{pmatrix} = \begin{pmatrix} f & -b \\ -c & a \end{pmatrix} \begin{pmatrix} \theta^{1'} \\ \theta^{2'} \end{pmatrix},$$

where

$$\det \begin{pmatrix} a & b \\ c & f \end{pmatrix} = 1.$$

We compute $d\theta^{1'} = ad\theta^1 + bd\theta^2 = -b\theta^1 \wedge \theta = -bf\theta^{1'} \wedge \theta + b^2\theta^{2'} \wedge \theta$ and $d\theta^{2'} = cd\theta^1 + fd\theta^2 = -f\theta^1 \wedge \theta = -f^2\theta^{1'} \wedge \theta + bf\theta^{2'} \wedge \theta$. In order to compute the enriched connection we need to find $\varphi', w', \tau^{1'}, \tau^{2'}$ satisfying

$$d\omega^{1'} = (\varphi' + 3w') \wedge \omega^{1'} + \omega' \wedge \tau_2^{1'} \omega^{2'}, \quad d\omega^{2'} = (\varphi' - 3w') \wedge \omega^{2'} - \omega' \wedge \tau_1^{2'} \omega^{1'}.$$

Comparing with the structure equations and observing $\omega^{i'} = a_i \theta^{i'}$ we obtain

$$\begin{aligned} \varphi' &= \frac{1}{2} \left(\frac{da_1}{a_1} + \frac{da_2}{a_2} \right), \quad 3w' = \frac{1}{2} \left(\frac{da_1}{a_1} - \frac{da_2}{a_2} \right) + \frac{bf}{a_1 a_2} \omega, \\ \tau_2^{1'} &= -\frac{b^2}{a_2^2} \theta^{2'} \quad \text{and} \quad \tau_1^{2'} = -\frac{f^2}{a_1^2} \theta^{1'}. \end{aligned}$$

Then $d\varphi = 0$, $3dw = \frac{bf}{a_1 a_2} \omega^{1'} \wedge \omega^{2'}$, and it follows $A = B = C = D = 0$ and $3S = \frac{bf}{a_1 a_2}$. Also

$$d\tau_2^{1'} = -2\tau_2^{1'}(\varphi - 3w) + \frac{2b^3 f}{a_2^3 a_1} \omega, \quad d\tau_1^{2'} = -2\tau_1^{2'}(\varphi + 3w) - \frac{2bf^3}{a_1^3 a_2} \omega,$$

and $\tau_{20}^{1'} = \frac{2b^3 f}{a_2^3 a_1}$, $\tau_{10}^{2'} = -\frac{2bf^3}{a_1^3 a_2}$. At last $dS = -2S\varphi$ and we get $S_0 = S_1 = S_2 = 0$. Then $P = n = 0$, and we obtain from formulas 38 and 39 that

$$Q^1 = \frac{3}{2} \frac{b^3 f}{a_1 a_2^3}, \quad Q^2 = -\frac{3}{2} \frac{bf^3}{a_1^3 a_2}.$$

We proved

Lemma 5.2 *The path structures defined by the forms $\theta^{1'}, \theta^{2'}, \theta$ on T^3 have curvatures $Q^1 = \frac{3}{2}b^3 f$, $Q^2 = -\frac{3}{2}bf^3$ (computed through a section on the torus).*

Note that the path structure is flat if and only if the one of the torsions $\tau^{1'}$ or $\tau^{2'}$ are zero and this happens if the direction defined by $\frac{\partial}{\partial t}$ is one of the line bundles contained in the contact bundle of the path structure. The couple (b, f) is determined up to a sign by the curvatures Q^1 and Q^2 .

The global invariant is given in the next Proposition.

Proposition 5.3 *Let $T^3_n(a, b, c, d, f)$ as the path structure on the torus defined as above. Then the global invariant is*

$$\mu(T^3_n(a, b, c, d, f)) = \frac{3n}{8\pi} (bf)^2.$$

Proof. This is a direct computation using the formula for the global invariant (see formula 44):

$$\int_{T^3} s^* TC_2(\pi) = \int_{T^3} \frac{1}{8\pi^2} (2\tau_2^{1'} \tau_1^{2'} \theta \wedge \theta^1 \wedge \theta^2 - \frac{9}{2} w \wedge \theta^1 \wedge \theta^2) = \int_{T^3} \frac{1}{8\pi^2} \frac{3}{2} b^2 f^2 \theta \wedge \theta^1 \wedge \theta^2.$$

Therefore

$$\mu(T^3_n(a, b, c, d, f)) = \int_{T^3} \frac{3}{16\pi^2} (bf)^2 \theta d\theta = \frac{3n}{8\pi} (bf)^2.$$

□

Note that the global invariant is null if and only if the path structure is flat.

6 Invariant path structures on $\mathbf{SU}(2)$

Tight contact structures on S^3 are all contactomorphic (see [E]). In this section we explicit homogeneous strict path structures on $\mathbf{SU}(2)$ which are carried by a fixed left invariant tight contact structure.

Let α, β, γ be a basis of left invariant 1-forms defined on $\mathbf{SU}(2)$ with

$$d\alpha = -\beta \wedge \gamma, \quad d\beta = -\gamma \wedge \alpha, \quad d\gamma = -\alpha \wedge \beta$$

A strict path structure on $\mathbf{SU}(2)$ is given by fixing the contact form $\theta = \gamma$ and the line fields $E^1 = \ker \alpha \cap \ker \theta$ and $E^2 = \ker \beta \cap \ker \theta$.

We define strict path structures by choosing a map from $\mathbf{SU}(2)$ to $\mathbf{SL}(2, \mathbb{R})$:

$$\theta = \gamma, \quad Z^1 = r_1\beta + r_2\alpha, \quad Z^2 = s_1\beta + s_2\alpha,$$

with $r_1s_2 - r_2s_1 = 1$. Then

$$d\theta = Z^1 \wedge Z^2.$$

In the case the map $\mathbf{SU}(2) \rightarrow \mathbf{SL}(2, \mathbb{R})$ is constant, from $\beta = s_2Z^1 - r_2Z^2$ and $\alpha = -s_1Z^1 + r_1Z^2$, we obtain

$$dZ^1 = r_1d\beta + r_2d\alpha = \theta \wedge (xZ^1 + yZ^2)$$

and analogously,

$$dZ^2 = \theta \wedge (zZ^1 - xZ^2),$$

where

$$x = r_1s_1 + r_2s_2, \quad y = -(r_1^2 + r_2^2), \quad z = s_1^2 + s_2^2.$$

Observe that $x^2 + yz = -1$. Then for a enriched path structure with coframes obtained from the tautological forms $\omega = a_1a_2\theta$, $\omega^1 = a_1Z^1$ and $\omega^2 = a_2Z^2$ we obtain

$$d\omega^1 = \left(\frac{da_1}{a_1} + x\theta\right) \wedge \omega^1 + a_1y\theta \wedge Z^2.$$

$$d\omega^2 = \left(\frac{da_2}{a_2} - x\theta\right) \wedge \omega^2 + a_2z\theta \wedge Z^1$$

From Proposition 2.3 we have

$$\varphi = \frac{1}{2}\left(\frac{da_1}{a_1} + \frac{da_2}{a_2}\right), \quad 3w = \frac{1}{2}\left(\frac{da_1}{a_1} - \frac{da_2}{a_2}\right) + \frac{x}{a_1a_2}\omega,$$

$$\tau_2^1 = \frac{y}{a_2^2}, \quad \tau_1^2 = -\frac{z}{a_1^2}.$$

and therefore

$$d\varphi = 0, \quad 3dw = d(x\theta) = \frac{x}{a_1a_2}\omega^1 \wedge \omega^2$$

so that $S = \frac{x}{3a_1a_2}$, $A = B = C = D = 0$.

From

$$\begin{aligned} d\tau_2^1 &= -2\frac{da_2}{a_2}\frac{y}{a_2^2} = -2\tau_2^1(\varphi - 3w) - 2\frac{xy}{a_1a_2^3}\omega \\ d\tau_1^2 &= 2\frac{da_1}{a_1}\frac{z}{a_1^2} = -2\tau_1^2(\varphi + 3w) - 2\frac{xz}{a_2a_1^3}\omega \end{aligned}$$

we obtain

$$\tau_{20}^1 = -2\frac{xy}{a_1a_2^3}, \quad \tau_{10}^2 = -2\frac{xz}{a_2a_1^3}$$

From

$$dS = -\frac{x}{3a_1a_2}\left(\frac{da_1}{a_1} + \frac{da_2}{a_2}\right) = -2\varphi S$$

we obtain $S_0 = S_1 = S_2 = 0$, and $P = n = 0$.

It follows from formulas 38 and 39 that $Q^1 = \tau_{20}^1 + \frac{3}{2}S\tau_2^1$ and $Q^2 = \tau_{10}^2 - \frac{3}{2}S\tau_1^2$, therefore

$$Q^1 = -\frac{xy}{a_1a_2^3}$$

and

$$Q^2 = -\frac{xz}{a_1^3a_2}.$$

Observe that y and z never vanish. We conclude that the invariant strict structure on $\mathbf{SU}(2)$ is a flat path structure if and only if $x = 0$. This can be interpreted, because $x = r_1s_1 + r_2s_2$, as the strict structures such that the directions E^1 and E^2 are perpendicular for the canonical metric defined by the forms α and β .

Proposition 6.1 *Define strict path structures on $\mathbf{SU}(2)$ by choosing a constant map from $\mathbf{SU}(2)$ to $\mathbf{SL}(2, \mathbb{R})$:*

$$\theta = \gamma, \quad Z^1 = r_1\beta + r_2\alpha, \quad Z^2 = s_1\beta + s_2\alpha,$$

with $r_1s_2 - r_2s_1 = 1$. Let $x = r_1s_1 + r_2s_2$. Then the global invariant of the induced path structure is

$$\mu(\mathbf{SU}(2)(r_1, r_2, s_1, s_2)) = -\frac{1}{2} - \frac{3}{8}x^2.$$

Proof. We compute, using formula 44, the global invariant for the family of structures defined on $\mathbf{SU}(2)$. We have from above that $x = r_1s_1 + r_2s_2$, $y = -(r_1^2 + r_2^2)$, $z = s_1^2 + s_2^2$ and that $x^2 + yz = -1$. Then it follows

$$\begin{aligned} \int_{\mathbf{SU}(2)} s^*TC_2(\pi) &= \int_{\mathbf{SU}(2)} \frac{1}{8\pi^2} (2\tau_2^1\tau_1^2\theta - \frac{9}{2}Sw) \wedge \theta^1 \wedge \theta^2 = - \int_{\mathbf{SU}(2)} \frac{1}{8\pi^2} (2yz + \frac{1}{2}x^2) \gamma \wedge \beta \wedge \alpha \\ &= \int_{\mathbf{SU}(2)} \frac{1}{8\pi^2} (-2 - \frac{3}{2}x^2) \gamma \wedge \beta \wedge \alpha. \end{aligned}$$

We use then that $\int_{\mathbf{SU}(2)} \gamma \wedge \beta \wedge \alpha = 2\pi^2$. □

Observe that the invariant is never null for this family even in the case of a flat path structure (which happens when $x = 0$). Also the critical point of the invariant along this family is a maximal at $x = 0$, at a flat structure, and it is equal to $-\frac{1}{2}$.

References

- [A] Arnold, V. : Chapitres supplémentaires de la théorie des équations différentielles ordinaires. Mir, Moscow, 1980.
- [B] Bryant, R. : Élie Cartan and geometric duality. Preprint 1998.
- [BGH] Bryant, R. , Griffiths, P. , Hsu, L. : Toward a geometry of differential equations. Geometry, topology, and physics, 1-76, Conf. Proc. Lecture Notes Geom. Topology, IV, Int. Press, Cambridge, MA, 1995.
- [BE] Burns, D. , Epstein, C. L. A global invariant for three-dimensional CR-manifolds. Invent. Math. 92 (1988), no. 2, 333-348.
- [Ca] Cartan, E. : Sur les variétés à connexion projective. Bull. Soc. Math. France 52 (1924), 205-241.
- [Car] Cartan, E.: Les espaces généralisés et l'intégration de certaines classes d'équations différentielles, C. R. Acad. Sci. 206 (1938), 1689-1693.
- [CL] Cheng, J. H. , Lee, J. M. : The Burns-Epstein invariant and deformation of CR structures. Duke Math. J. 60 (1990), no. 1, 221-254.
- [E] Eliashberg, Y. : Contact 3-manifolds twenty years since J. Martinet's work, Ann. Inst Fourier, 42(1992), 165-192.
- [FMMV] Falbel, E.; Mion-Mouton, M.; Veloso, J. M. Cartan connections and path structures with large automorphism groups. Internat. J. Math. 32 (2021), no. 12,
- [FV] Falbel, E. , Veloso, J. M. : Flag structures on real 3-manifolds. Geom. Dedicata 209 (2020), 149-176.
- [IL] Ivey, T. A. , Landsberg, J. M. : Cartan for beginners: differential geometry via moving frames and exterior differential systems. Graduate Studies in Mathematics, 61. American Mathematical Society, Providence, RI, 2003.
- [MM] Mion-Mouton, M : Partially hyperbolic diffeomorphisms and Lagrangian contact structures. Ergodic Theory Dynam. Systems 42 (2022), no. 8, 2583-2629.
- [T] Tresse, A. M. : Détermination des Invariants Ponctuels de l'équation Différentielle Ordinaire du Second Ordre $y'' = \omega(x, y, y')$, Preisschriften der Furstlichen Jablonowskischen Gesellschaft XXXII, Leipzig, 1896.

E. FALBEL
INSTITUT DE MATHÉMATIQUES
DE JUSSIEU-PARIS RIVE GAUCHE
CNRS UMR 7586 AND INRIA EPI-OURAGAN
SORBONNE UNIVERSITÉ, FACULTÉ DES SCIENCES
4, PLACE JUSSIEU 75252 PARIS CEDEX 05, FRANCE
elisha.falbel@imj-prg.fr

J. M. VELOSO
FACULDADE DE MATEMÁTICA - ICEN
UNIVERSIDADE FEDERAL DO PARÁ
66059 - BELÉM- PA - BRAZIL
veloso@ufpa.br