



HAL
open science

Interactions of human drug transporters with chemical additives present in plastics: Potential consequences for toxicokinetics and health

Valentin Tastet, M Le Vee, Arnaud Bruyère, Olivier Fardel

► **To cite this version:**

Valentin Tastet, M Le Vee, Arnaud Bruyère, Olivier Fardel. Interactions of human drug transporters with chemical additives present in plastics: Potential consequences for toxicokinetics and health. *Environmental Pollution*, 2023, 331 (Pt 2), pp.121882. 10.1016/j.envpol.2023.121882 . hal-04121151

HAL Id: hal-04121151

<https://hal.science/hal-04121151v1>


Submitted on 27 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21

**Interactions of human drug transporters with chemical additives present in plastics:
Potential consequences for toxicokinetics and health**

Valentin Tastet¹, Marc Le Vée¹, Arnaud Bruyère¹ and Olivier Fardel²

¹Univ Rennes, Inserm, EHESP, Irset (Institut de recherche en santé, environnement et travail)
- UMR_S 1085, F-35000 Rennes, France

²Univ Rennes, CHU Rennes, Inserm, EHESP, Irset (Institut de recherche en santé,
environnement et travail) - UMR_S 1085, F-35000 Rennes, France

Corresponding author: Olivier Fardel, Irset, Faculté de Pharmacie, 2 Avenue Pr Léon
Bernard, 35043 Rennes, France. E-mail: olivier.fardel@univ-rennes.fr

22 Abstract

23 Human membrane drug transporters are recognized as major actors of pharmacokinetics; they
24 also handle endogenous compounds, including hormones and metabolites. Chemical additives
25 present in plastics interact with human drug transporters, which may have consequences for the
26 toxicokinetics and toxicity of these widely-distributed environmental and/or dietary pollutants,
27 to which humans are highly exposed. The present review summarizes key findings about this
28 topic. *In vitro* assays have demonstrated that various plastic additives, including bisphenols,
29 phthalates, brominated flame retardants, poly-alkyl phenols and per- and poly-fluoroalkyl
30 substances, can inhibit the activities of solute carrier uptake transporters and/or ATP-binding
31 cassette efflux pumps. Some are substrates for transporters or can regulate their expression. The
32 relatively low human concentration of plastic additives from environmental or dietary exposure
33 is a key parameter to consider to appreciate the *in vivo* relevance of plasticizer-transporter
34 interactions and their consequences for human toxicokinetics and toxicity of plastic additives,
35 although even low concentrations of pollutants (in the nM range) may have clinical effects.
36 Existing data about interactions of plastic additives with drug transporters remain somewhat
37 sparse and incomplete. A more systematic characterization of plasticizer-transporter
38 relationships is needed. The potential effects of chemical additive mixtures towards transporter
39 activities and the identification of transporter substrates among plasticizers, as well as their
40 interactions with transporters of emerging relevance deserve particular attention. A better
41 understanding of the human toxicokinetics of plastic additives may help to fully integrate the
42 possible contribution of transporters to the absorption, distribution, metabolism and excretion
43 of plastics-related chemicals, as well as to their deleterious effects towards human health.

44 Key-words

45 Drug transporters, plastic additives, pollutants, toxicity, toxicokinetics.

47 **1. Introduction**

48 Human drug transporters are membrane proteins, implicated in the uptake or secretion
49 of drugs at the plasma membrane (Petzinger and Geyer, 2006). They segregate into ATP-
50 binding cassette (ABC) transporter and solute carrier (SLC) superfamilies (Roberts, 2021).
51 Drug transporters are located in main organs/tissues implicated in drug absorption, disposition
52 and/or elimination, such as the intestine, the liver, the kidney and blood-tissue barriers,
53 including the blood-brain barrier, the blood-testis barrier and the placental barrier (Giacomini
54 and Huang, 2013). Drug transporters thus play a key role in pharmacokinetics, and are notably
55 involved in the phases of absorption, distribution and excretion of drugs (Liu, 2019). Moreover,
56 drugs are not only substrates for transporters, but can also act as potential inhibitors, which can
57 lead to drug-drug interactions (DDIs) when the pharmacokinetics features of a drug, *i.e.*, the
58 “victim”, are impaired, due to inhibition of its membrane transport by a co-administrated drug,
59 *i.e.*, the “perpetrator” (König et al., 2013). Blockage of transporter activity can additionally
60 result in altered transport of endogenous substrates, thereby triggering adverse drug effects
61 (Zolk and Fromm, 2011). Endogenous substrates for transporters include hormones, signalling
62 molecules, nutrients, neuromediators, metabolites, nucleosides, and bile acids (Nigam, 2015).
63 Some of them have been proposed as clinical biomarkers for transporters-related DDI
64 evaluation (Chu et al., 2018). It is noteworthy that inhibition of hormone transport may
65 represent a contributing mechanism to endocrine disruption, an adverse effect of various
66 chemical pollutants (Lisco et al., 2022). In addition to inhibition of their activity, changes in the
67 expression of drug transporters, caused by drug treatment or pathophysiological situations such
68 as inflammation, can impair pharmacokinetics and be the source of DDIs or toxicity (Brouwer
69 et al., 2022; Cressman et al., 2012). Genetic polymorphisms of certain transporters can also
70 cause pharmacokinetics variability among human populations (Yee et al., 2018). For all these
71 reasons, transporters are now fully recognized by the pharmacology community and by
72 regulatory agencies, as major actors of drug disposition in the body (Giacomini et al., 2010).

73 Consequently, from a regulatory standpoint, interactions of novel small drugs with main drug
74 transporters have to be studied and characterized during their pharmaceutical development (Lee
75 et al., 2017). This has to be performed using human, and not animal, transporter models, owing
76 to the known interspecies differences in drug transporter activity and expression (Hammer et
77 al., 2021; Morse et al., 2021; Wang et al., 2015).

78 In addition to drugs, other chemicals, especially pollutants of public health concern, are
79 likely to interact with human drug transporters (Clerbaux et al., 2019; Nicklisch and Hamdoun,
80 2020), which may have important consequences for chemical risk assessment (Clerbaux et al.,
81 2018; Darney et al., 2020). In contrast to drugs, the handling of pollutants by human drug
82 transporters and their potential inhibitory effects remain rather poorly studied (Fardel et al.,
83 2012). The exact implications of these potential interactions for pollutant toxicokinetics or
84 toxicity are consequently poorly understood (Abu-Qare et al., 2003). The proof of concept for
85 the interaction of environmental chemicals with human drug transporters was, however,
86 provided more than twenty five years ago, through the demonstration that various pesticides
87 belonging to different chemical classes can inhibit the ABC transporter P-glycoprotein (P-gp,
88 *ABCB1*) (Bain and LeBlanc, 1996). Pesticides have since been shown to interact with other
89 ABC transporters and with SLCs, as recently reviewed (Chedik et al., 2018; Guéniche et al.,
90 2020). Transporters notably contribute to the pharmacokinetics, and moreover to the toxicity of
91 certain pesticides, such as those of the herbicide paraquat (Wang et al., 2021). Besides
92 pesticides, chemicals belonging to various classes of environment, food or drinking water
93 contaminants also interfere with human transporters (Nicklisch and Hamdoun, 2020). This is
94 notably the case for chemical additives present in plastics (Bruyere et al., 2017; Kim et al.,
95 2007), known to also interact with xenobiotic transporters of aquatic organisms (Anselmo et
96 al., 2012; Horion et al., 2015; Keiter et al., 2016; Nicklisch et al., 2021; Popovic et al., 2014).
97 More than 400 of these plastics-related chemicals, basically used for enhancing polymer

98 properties and prolonging their life, are in use (Gunaalan et al., 2020). The most common plastic
99 additives are phthalates (used as plasticizers), bisphenols (used as
100 monomers/plasticizers/antioxidants), brominated flame retardants (BFRs) such as
101 polybrominated diphenyls ethers (PBDE) or tetrabromobisphenol A (TBBPA, 4,4'-(propane-
102 2,2-diyl)bis(2,6-dibromophenol)), alkylphenols (used as antioxidants/plasticizers), and per- and
103 poly-fluoroalkyl substances (PFASs) (used as coating agents) (Barhoumi et al., 2022; Groh et
104 al., 2019; Hahladakis et al., 2018; Hermabessiere et al., 2017; Wiesinger et al., 2021). Because
105 these additives are usually not chemically bound to plastics, they can migrate out of them and
106 are consequently widely released in the environment, notably when plastics degrade
107 (Hermabessiere et al., 2017). They are, in particular, found in food and drinking water, where
108 they can originate from plastic packaging or bottling, and are also closely associated with
109 nanoplastics and microplastics, which are now recognized as major emerging environmental
110 pollutants (Sangkham et al., 2022). Humans are thus highly exposed to these plastic additives
111 through ingestion, inhalation and dermal contact. This is problematic because of the established
112 deleterious effects of many of these additives; for example, bisphenols, phthalates and BFRs
113 cause endocrine disruption, fertility problems, metabolic diseases and cancers (Benjamin et al.,
114 2017; Campanale et al., 2020; Feiteiro et al., 2021; Ma et al., 2019; Tarafdar et al., 2022). Plastic
115 additives are therefore chemicals of considerable concern with respect to the human exposome
116 and the public health. This led us to survey the interactions of plastic additives with human drug
117 transporters in the present review, with a special emphasis on possible consequences for human
118 exposure, toxicokinetics and health. The plastic additives considered are mainly those for which
119 interactions with at least one human drug transporter are documented in the literature, according
120 to a search of the PubMed database (<https://pubmed.ncbi.nlm.nih.gov/>), as of 08/31/2022; these
121 plastic additives are listed in Table 1.

122 **2. Brief overview of drug transporters**

123 2.1. ABC transporters

124 Plasma membrane drug transporters belonging to the ABC protein family mediate
125 primary active efflux of drugs, through hydrolysis of ATP via their intrinsic ATPase activity
126 (Schinkel and Jonker, 2003). They comprise P-gp, encoded by the multidrug resistance gene 1
127 (*MDR1*), multidrug resistance-associated proteins (MRPs) such as MRP1 (*ABCC1*), MRP2
128 (*ABCC2*), MRP3 (*ABCC3*) and MRP4 (*ABCC4*), breast cancer resistance protein
129 (BCRP/*ABCG2*), as well as the hepatic bile salt export pump (BSEP/*ABCB11*). The main types
130 of substrates for these efflux pumps are given in Table 2. It is noteworthy that most of these
131 transporters, particularly P-gp, MRP1, MRP2 and BCRP, usually exhibit broad and sometimes
132 overlapping substrate specificity. Several of them, especially P-gp, have been historically
133 implicated in multidrug resistance of cancer cells (Amawi et al., 2019), thus illustrating the
134 protective effect of these efflux pumps towards xenobiotics (Leslie et al., 2005).

135 The tissue location of ABC transporters is shown in Fig. 1. P-gp and BCRP are notably
136 found at the brush-border membrane of enterocytes, where they actively expel their substrates
137 into the digestive lumen, thereby preventing or limiting their intestinal absorption (Müller et
138 al., 2017). These pumps are also present at the luminal pole of brain microvessel endothelial
139 cells, at the apical pole of syncytiotrophoblasts and at the basolateral pole of Sertoli cells, where
140 they contribute to the blood–brain, blood-placenta and blood-testis barriers, respectively (Liu
141 and Liu, 2019; Miller, 2015; Mruk et al., 2011). P-gp and BCRP, as well as MRP2 and BSEP,
142 are additionally found at the canalicular pole of hepatocytes, where they secrete their substrates
143 into the bile canaliculus lumen (Jetter and Kullak-Ublick, 2020); in this context, drug-mediated
144 inhibition of BSEP transport has been associated with cholestasis (Chatterjee and Annaert,
145 2018). In the kidney, P-gp, BCRP and MRP2 are located at the apical pole of proximal tubular
146 cells, where they contribute to tubular secretion of drugs (Lee and Kim, 2004).

147 2.2. SLC transporters

148 SLCs mediate secondary active transport or facilitated diffusion of drugs (Zhou and Shu,
149 2022). They belong mainly to the *SLCO*, *SLC22*, and *SLC47* transporter subfamilies. The list
150 of these transporters and their main types of substrates are indicated in Table
151 2. *SLCO* transporters, also known as organic anion transporting polypeptides (OATPs), include
152 three members recognized as important drug transporters, *i.e.*, OATP1B1 (*SLCO1B1*),
153 OATP1B3 (*SLCO1B3*) and OATP2B1 (*SLCO2B1*) (Roth et al., 2012). *SLC22* drug transporters
154 correspond to organic cation transporters (OCTs) such as OCT1 (*SLC22A1*) and OCT2
155 (*SLC22A2*), and organic anion transporters (OATs) like OAT1 (*SLC22A6*), OAT2 (*SLC22A7*),
156 OAT3 (*SLC22A8*) and OAT4 (*SLC22A11*) (Ciarimboli, 2008; Koepsell, 2020; Roth et al.,
157 2012). *SLC47* transporters are multidrug and toxin extrusion (MATE) proteins and comprise
158 two members, *i.e.*, MATE1 (*SLC47A1*) and MATE2 (*SLC47A2*) (Nies et al., 2016). Similarly
159 to ABC transporters, there is a high degree of overlapping substrates between SLC transporters,
160 notably between OCTs and MATEs and between OATPs and OATs (Table 2).

161 OATPs, OCTs and OATs are notably expressed at the basolateral side of hepatocytes
162 and kidney proximal tubular cells (Fig. 1) and primarily act as uptake drug transporters enabling
163 the entry of drugs from blood into cells (Ciarimboli, 2008; Roth et al., 2012). By contrast,
164 MATE1 and MATE2-K, the functional splice variant of MATE2, function as pH-dependent
165 efflux transporters, at the apical surface of hepatocytes or proximal tubular cells (Koepsell,
166 2020) (Fig. 1). OAT4 is implicated in drug reabsorption from the primitive urine at the apical
167 pole of proximal tubular cells and also in compound uptake into the placenta from the foetus
168 blood at the basal pole of syncytiotrophoblasts (Rizwan and Burckhardt, 2007). SLC
169 transporters also include the proton-coupled peptide transporter (PEPT) 1 (*SLC15A1*), involved
170 in absorption of peptidomimetic drugs at the apical pole of enterocytes (Inui et al., 2000), the
171 sodium-taurocholate cotransporting polypeptide (NTCP/*SLC10A1*) and the apical sodium-
172 dependent bile acid transporter (ASBT/*SLC10A2*), implicated in bile acid uptake at the

173 basolateral pole of hepatocytes and the apical pole of enterocytes, respectively (Meier and
174 Stieger, 2002), and the human organic solute transporter (OST) α/β (*SLC51A/SLC51B*),
175 implicated in the efflux of bile acids at the basolateral pole of enterocytes and hepatocytes
176 (Beaudoin et al., 2020).

177 **3. Modulation of transporter activities by plastic additives**

178 Direct modulation of human transporter activities by plastic additives have been studied
179 using conventional *in vitro* methods, based on vesicular or cellular assays (Brouwer et al.,
180 2013). Vesicular transport assays, used mainly for ABC transporters, rely upon the ATP-
181 dependent translocation of reference substrates into the vesicular compartment of inside-out
182 plasma membrane vesicles prepared from cells expressing the transporter of interest (Volpe,
183 2016); the inhibition or stimulation of transporter activity is visualized by the decreased or
184 increased accumulation of the reference substrate into the vesicular compartment in response
185 to the chemical tested for its transporter activity modulating potential. Cellular assays, used for
186 ABC and SLC transporters, correspond to uptake or efflux of reference substrates in, or from,
187 recombinant cells overexpressing the transporter of interest, or to bidirectional transport of
188 reference substrates across transporter-expressing polarized cells (Jani and Krajcsi, 2014); the
189 potential transporter modulating activity of compounds is demonstrated by the alteration of
190 reference substrate uptake, efflux or transcellular passage.

191 Modulations of ABC transporter activities by plastic additives are summarized in Table
192 3. Plastic additives usually act as inhibitors of ABC transporter activities or fail to interfere with
193 them. Only bisphenol A (BPA, 4,4'-(propane-2,2-diyl)diphenol) was reported to stimulate the
194 activity of an ABC transporter, *i.e.*, that of P-gp (Table 3), which is consistent with the fact that
195 most ABC transporter modulators behave as inhibitors (Cui et al., 2015). It is, however,
196 noteworthy that the stimulating effect of BPA towards P-gp activity observed in cellular assays
197 (Jin and Audus, 2005) has not been confirmed in vesicular assays (Dankers et al., 2013), which

198 may reflect differential sensitivities of cellular and vesicular assays to modulators. Other
199 discrepancies have been observed between cellular and vesicular assays notably with
200 diethylhexylphthalate (DEHP, bis(2-ethylhexyl) benzene-1,2-dicarboxylate) which acts as a P-
201 gp inhibitor in cellular assays (Kim et al., 2007) but not in vesicular assays (Dankers et al.,
202 2013), and the BFR 2,2',4,4'-tetrabromodiphenyl ether (BDE-47, 3,5-dibromo-2-(2,4-
203 dibromophenoxy)phenol), which inhibits BCRP in vesicular assays but not in cellular assays
204 (Marchitti et al., 2017). Overall, among plastic additives, phthalates like DEHP and its
205 metabolite monoethylhexylphthalate (MEHP, 2-(2-ethylhexoxycarbonyl)benzoate) interact
206 poorly with ABC transporter activity; they notably fail to inhibit MRP1, MRP4 or BCRP
207 activity (Table 3). By contrast, the BFR TBBPA, as well as the PFASs perfluorooctanoic acid
208 (PFOA, 2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-pentadecafluorooctanoic acid) and perfluorooctane
209 sulfonate (PFOS, 1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptadecafluorooctane-1-sulfonic acid),
210 appear to be potent inhibitors of ABC transporters as they block P-gp, MRP1, MRP4 and BCRP
211 activities (Table 3); PFOS additionally inhibits BSEP and MRP2. Among ABC transporters,
212 BCRP is the most sensitive to plastic additives, as it is inhibited by BPA, BFRs and PFASs
213 (Table 3). It is noteworthy that the concentrations of plastic additives required to inhibit ABC
214 transporters are usually in the 10-100 μM range (Table 3); only the alkylphenol 4-nonylphenol
215 (NP) blocks P-gp activity when used at low concentrations in the 0.1-1 μM range (Doo et al.,
216 2005).

217 Modulations of SLC drug transporter activities by plastic additives are summarized in
218 Table 4. Some additives, *e.g.*, bisphenols used at 100 μM , stimulate the activity of SLCs; this
219 is the case for bisphenol F (BPF, 4-[(4-hydroxyphenyl)methyl]phenol), proposed as a substitute
220 for BPA, which increases the activity of OCT2, OATP1B3, and OAT1, whereas BPA stimulates
221 only that of OAT1 (Table 4). Inhibition of SLC activities by plastic additives is nevertheless a
222 more common situation (Table 4). It occurs notably for TBBPA which blocks OCT1, OCT2,

223 OATP1B1, OATP1B3, MATE1, OAT3 and NTCP activities, but fails to alter MATE2-K and
224 OAT1 activities. Other BFRs like BDE-47 and PFASs target OATPs and/or OATs (Table 4),
225 whereas effects of phthalates towards SLC transporters have not yet been investigated. The
226 concentrations of bisphenols, BFRs or PFAS acting on SLC transporter activities are often in
227 the 10-100 μM range (Table 4), as reported above for those acting on ABC transporters. Low
228 concentrations of TBBPA (in the 0.5-5 μM range), however, inhibit OAT3 and NTCP activity,
229 whereas inhibition of OATP activities is caused by much lower concentrations of BDE (in the
230 100 pM-100 nM range) (Table 4).

231 The mechanisms by which plastic additives modulate transporter activities remain to be
232 clarified. Competitive or non-competitive interactions with drug binding sites on transporter
233 proteins are likely involved, as proposed for drugs inhibiting transporters (Wigler and Patterson,
234 1993). The fact that BPA has been demonstrated to inhibit OAT3-mediated transport of estrone
235 3-sulfate in a competitive manner (Bruyere et al., 2017) and that PFOA, perfluorononanoic acid
236 (PFNA, 2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,9-heptadecafluorononanoic acid) and perfluorodecanoic
237 acid (PFDA, 2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,10-nonadecafluorodecanoic acid)
238 competitively block OATP2B1-mediated uptake of sulfobromophthalein in Caco-2 cells
239 (Kimura et al., 2020) fully supports this conclusion. BDEs probably also inhibit OATPs by
240 competitive mechanisms because they are high affinity ligands for these SLCs (Pacyniak et al.,
241 2010) (Table 5).

242 **4. Plastic additives as substrates for transporters**

243 Handling of plastic additives by human drug transporters has mainly been investigated
244 using conventional *in vitro* methods addressing whether a compound is a substrate for a plasma
245 membrane transporter (Jani and Krajcsi, 2014). Such methods correspond to (i) cellular assays
246 based on the uptake, efflux or transcellular passage of the chemicals in, from or across
247 transporter-expressing cells, in the presence or absence of reference inhibitors for the

248 considered transporter, (ii) vesicular assays, based on the transport of the compounds into the
249 intravesicular compartment of transporter-expressing inside-outside membrane vesicles and
250 (iii) ATPase assays, which indirectly measure ABC drug transporter activity through
251 quantifying the stimulation of ATPase activity by substrates in plasma membrane preparations
252 from ABC transporter-expressing cells. Additionally, the perfused human placenta model has
253 been used to demonstrate that bisphenol S (BPS, 4-(4-hydroxyphenyl)sulfonylphenol) is
254 transported by P-gp (Grandin et al., 2019).

255 The plastic additives which are either substrates or non substrates for human ABC and
256 SLC transporters are listed in Table 5. With respect to P-gp, DEHP and alkyphenols such as
257 nonylphenol ethoxylate (NPE, 2-(4-nonylphenoxy)ethanol), but not its metabolite NP, are
258 substrates. For BPA, conflicting results have been obtained with respect to P-gp. BPA has thus
259 been shown to be transported by P-gp across human intestinal Caco-2 cells (Yoshikawa et al.,
260 2002), but not across P-gp-overexpressing MDCK cells (Dankers et al., 2013); moreover, BPA
261 failed to stimulate ATPase activity of P-gp (Mazur et al., 2012). Further studies are therefore
262 required to more precisely determine whether BPA is a substrate for P-gp. With respect to other
263 ABC transporters, MRP2, MRP3 and BCRP transport BPA, whereas PFOA is a substrate for
264 BCRP (Table 5). Various BDEs, including BDE-47, have been demonstrated to be transported
265 by OATP1B1, OATP1B3 and OATP2B1, with rather low K_m values (around 0.3-1.9 μM)
266 (Table 5), indicating that they are high affinity substrates for OATPs. Whereas OATs transport
267 PFOA with K_m values in the 50-310 μM range. Other PFASs such as PFOS have additionally
268 been found to be substrates for the bile salt transporters NTCP, ASBT and OST α/β (Table 5).
269 Finally, the trans-epithelial transport of various BDEs across Caco-2 cells has been shown to
270 be altered by various transporter inhibitors, suggesting that several transporters, *i.e.*, P-gp,
271 MRPs, BCRP and OCTs, may participate in the transcellular passage of these chemicals, even

272 if passive diffusion may predominate (Yu et al., 2017). Further studies are required to confirm
273 whether BDEs are substrates for these transporters in intestinal cells.

274 **5. Regulation of human drug transporter expression by plastic additives**

275 Regulations of human drug transporter expression by plastic additives have been mostly
276 studied using cellular *in vitro* models and are listed in Table 6. These regulations, either
277 induction or repression of transporter expression, depend on the nature of both the transporter
278 and the plastic additive. For P-gp, chronic exposure (48 h) of placental BeWo cells to 10 μM
279 BPA results in enhanced expression of the efflux pump, but in a rather modest manner (55%
280 increase) (Jin and Audus, 2005). By contrast, BPA exposure at 80 μM for 24 h fails to induce
281 P-gp expression in human hepatic HepG2 cells, despite an increase in *MDR1* mRNA levels
282 (Hanet et al., 2008). These contradicting results may reflect a cell type-dependent regulation of
283 P-gp by BPA. This chemical also increases the activity of the *MDR1* gene promoter in an
284 haplotype-dependent manner (Speidel et al., 2018). Phthalates such as DEHP used at 10 μM
285 for 24 h-72 h enhance P-gp expression in colon cancer and sarcoma cells (Angelini et al., 2011;
286 Chen et al., 2018; Takeshita et al., 2006), which is associated with increased resistance to
287 chemotherapeutics drugs (Chen et al., 2018). P-gp induction and multidrug resistance also occur
288 in response to chronic exposure to 100 nM DEHP for three months in breast cancer MDA-MB-
289 231 cells (Jadhao et al., 2021). P-gp expression has additionally been shown to be induced in a
290 rat epididymal epithelial cell line in response to treatment by NP (20 μM) for 24 h (Jones and
291 Cyr, 2011). Effects of PFASs towards P-gp expression remain to be characterized, whereas the
292 bile salt efflux pump BSEP has been shown to be repressed by both PFOA and PFOS in human
293 hepatic HepaRG cells (Table 6).

294 With respect to MRPs, MRP1, MRP2 and MRP3, but not MRP4, have been
295 demonstrated to be up-regulated by BPA (80 μM) in human hepatic HepG2 cells (Hanet et al.,
296 2008) (Table 6). By contrast, DEHP tends to decrease Mrp2 protein levels in rat liver (Johnson

297 and Klaassen, 2002), whereas it induces expression of Mdr2 (*Abcb4*), involved in canalicular
298 secretion of phospholipids, in the mouse liver (Miranda et al., 1997). MRP3 mRNA expression
299 is repressed by the PFASs PFOA and PFOS, but is increased by the PFASs PFNA and PFDA
300 in HepaRG cells, thus highlighting that regulation of this ABC transporter depends on the nature
301 of the PFAS. BCRP is repressed in human term placental explant cultures exposed to 1 nM
302 BPA; NP has a similar effect. BPA also reduces BCRP expression in placental BeWo cells,
303 when used, however, at very high concentration (100 mM) (Table 6). PFNA and PFDA also
304 repress BCRP expression in human hepatic HepaRG cells. On the other hand, PFOA enhances
305 expression of Bcrp in the liver, but not in the kidney, of mice (Eldasher et al., 2013); this
306 suggests that regulation of hepatic BCRP by PFAS may be complex and species-dependent.

307 Regarding the regulation of SLC transporter expression, PFASs are the only plastic
308 additives that have been shown to play a role. They notably decrease mRNA expression of
309 hepatic OATP1B1, OATP2B1 and NTCP, whereas they increase that of OST β (Table 6).

310 The mechanisms of transporter expression regulation by plastic additives remain to be
311 fully characterized. Drug-sensing receptors, such as the pregnane-X-receptor (PXR), the aryl
312 hydrocarbon receptor (AhR) or the constitutive androstane receptor (CAR), may be implicated
313 because (i) such receptors regulate expression of various transporters, including P-gp, MRP2
314 and BCRP (Tirona, 2011), and (ii) plastic additives can activate them, notably CAR is activated
315 by PFASs and BDE-47 and PXR is activated by BDE-47 and BPA (Abe et al., 2017; Grimaldi
316 et al., 2019; Sueyoshi et al., 2014). Plastic additives-related oxidative stress and subsequent
317 activation of the antioxidant nuclear factor erythroid-derived 2-like 2 (Nrf2) pathway may also
318 be contributing factors, notably for BDEs (Dunnick et al., 2018). Additionally, repression of
319 BCRP protein expression in human term placenta explant cultures and BeWo cells by BPA
320 likely involves the estrogenic-like effects of this chemical (Cao et al., 2022; Sieppi et al., 2016).

321 **6. Consequences for toxicokinetics and human health**

322 6.1 Consequences for plastic additives altering transporter activity and/or expression

323 The possible consequences of the alteration of drug transporter activity and/or
324 expression by plastic additives may be drawn from what is now well-established regarding
325 drug-mediated modulation of transporters (Giacomini and Huang, 2013). Plastic additives
326 acting as transporter modulators, considered as “perpetrators”, are likely to impair membrane
327 passage of transporter substrates, considered as “victims”, and thus possibly impair their
328 pharmacokinetics features, *i.e.*, intestinal absorption, distribution and liver or kidney
329 elimination. For drugs or pollutants as “victims”, this may ultimately result in toxicity (if plasma
330 victim concentrations/ area under the curve (AUC) of plasma victim concentration versus time
331 are increased) or lack of effect (if plasma victim concentration/AUC of victims are decreased)
332 (Fig. 2). This may theoretically result in drug-pollutant or pollutant-pollutant interactions. On
333 the other hand, if the victim substrates are endogenous substrates such as, for example, bile
334 acids, toxicity, such as cholestasis due to inhibition of bile acid transport, may develop
335 (Chatterjee and Annaert, 2018). Altered transport of hormones as substrates may also contribute
336 to endocrine disruptive effects, as already suggested for inhibition of testosterone precursor
337 secretion by TBBPA in Leydig cells (Dankers et al., 2013).

338 The potential drug-pollutant or pollutant-pollutant interactions as well as pollutant
339 toxicity due to an abrogation of endogenous substrate transport have not yet been reported for
340 plastic additives, even for those inhibiting transporter activity or expression. This may likely be
341 due to the fact that human plasma concentrations of plastic additives, such as BPA, phthalates,
342 alkylphenols, BDEs, TBBPA and PFASs, due to environmental exposure or food intake, are
343 usually low, *i.e.*, in the nM or pM range (Asimakopoulos et al., 2012; Huang et al., 2014; Li et
344 al., 2020; Teeguarden et al., 2013; Wang et al., 2012; Wang et al., 2019). Their unbound
345 fraction, which is the one pharmacologically active towards transporters according to the free
346 drug theory (Summerfield et al., 2022), may even be much lower, but it is not commonly

347 considered in studies investigating serum pollutant concentrations. By contrast, the
348 concentrations of plastic additives required to modulate transporter activity and/or expression
349 are mostly in the 10-100 μM range according to *in vitro* studies (Table 3, Table 4 and Table 6).
350 *In vivo* inhibition of most drug transporters might therefore be unlikely to occur in response to
351 environmental exposure or food intake of plastic additives. This conclusion notably applies to
352 inhibition of SLC activities by BPA. Indeed, when applying the criterion initially defined by
353 the International Transporter Consortium for predicting *in vivo* inhibition of SLC transporters,
354 based on the ratio between the unbound maximum plasma concentration (C_{max}) of the drug
355 perpetrator and its IC_{50} value (which has to be ≥ 0.1 to consider *in vivo* inhibition of SLCs)
356 (Giacomini et al., 2010), human plasma concentrations of BPA (less than 50 nM) are very
357 unlikely to block the activity of OCT1, MATE1, OATP1B1, OATP1B3 or OAT3, as previously
358 reported (Bruyere et al., 2017). However, this general lack of *in vivo* modulation of SLCs by
359 plastic additives may be challenged for those exhibiting potent *in vitro* inhibitory effects
360 towards transporters or those regulating transporter expression when used at low
361 concentrations. Thus, BDE-47 at 100 pM inhibits OATP1B1 activity *in vitro* by approximately
362 50% (Pacyniak et al., 2010), thus suggesting that the IC_{50} value of this BDE towards this OATP
363 is around 100 pM. With total BDE-47 concentrations observed in human plasma in the 500-
364 1000 pM range (Wang et al., 2012), the $C_{\text{max}}/\text{IC}_{50}$ ratio is in the 5-10 range, indicating that
365 OATP1B1 activity may be inhibited in humans by circulating BDE-47 and that BDE-47 may
366 consequently potentially interfere with OATP1B1 substrates like statins. Nevertheless, this
367 conclusion can be questioned by considering what is the unbound maximum plasma
368 concentration of BDE-47, or even the unbound maximum plasma concentration of BDE-47 at
369 the liver inlet, instead of the total plasma concentration of BDE-47, if taking into account the
370 hepatic location of OATP1B1. With respect to regulation of transporter expression, repression
371 of BCRP by 1 nM of BPA in placenta explants (Sieppi et al., 2016) may be hypothesized to

372 occur *in vivo*, because human plasma concentration of BPA in pregnant women is around 28.5
373 nM (Yang et al., 2021), even though the limit of the free concentration of BPA has to be
374 considered. Through repressing placental BCRP expression, BPA may alter the function of the
375 blood placenta barrier and may thereby decrease the protection of the fetus against toxicant
376 substrates of BCRP, such as carcinogenic heterocyclic aromatic amines (Myllynen et al., 2008),
377 thus indicating a possible pollutant-pollutant interaction.

378 For P-gp and BCRP, it is noteworthy that their intestinal inhibition is predicted to
379 potentially occur if the ratio between the perpetrator maximal gastrointestinal concentration
380 (calculated as the ingested dose of perpetrator in a volume of 250 mL) and the IC₅₀ value is ≥
381 10 (Giacomini et al., 2010). The daily dietary intake of plastic additives has therefore to be
382 considered when calculating theoretical concentrations to evaluate inhibitory effects towards
383 P-gp and/or BCRP. With respect to BPA, TBBPA and BDEs, they can be estimated to be
384 approximately 2.38 µg, 70 ng and 75.4 ng, respectively, for a standard adult male weighing 70
385 kg (Colnot et al., 2014; Domingo et al., 2008; Huang et al., 2017), leading to gut concentrations
386 in the 1-100 nM range, which is lower than the concentrations needed to inhibit P-gp or BCRP
387 *in vitro* (in the 500 nM-100 µM range) (Table 3). Similarly, and despite daily DEHP intakes
388 being higher, *i.e.*, up to 840 µg (Cheng et al., 2016), the DEHP concentrations in the gut lumen,
389 up to 8.6 µM, remain much lower than the 30.1 mM DEHP concentration active on P-gp *in*
390 *vitro*. Overall, these data suggest that inhibition of intestinal P-gp or BCRP by plastic additives
391 in response to common contamination by the diet is rather unlikely. The risk of inhibition
392 remains to be considered, however, for highly-exposed subjects, notably in the case of exposure
393 to a mixture of pollutants, including plastic additives, whose inhibitory effects towards P-gp
394 and/or BCRP may be additive or synergic, as already demonstrated for inhibition of P-gp by
395 mixtures of marine pollutants (Nicklisch et al., 2016).

396 6.2 Consequences for plastic additives that are transporter substrates

397 Handling of plastic additives by transporters is likely to contribute to the various steps
398 of their toxicokinetics. In particular, transport by P-gp or BCRP at the apical pole of enterocytes
399 may limit their intestinal absorption, especially since saturation of these intestinal pumps is
400 unlikely to occur, owing to the relative low concentrations of plastic additives in the gut lumen,
401 as already discussed above. Transport of plastic additives by P-gp and/or BCRP at the blood-
402 brain barrier may contribute to limit their entry into the brain and prevent any neurotoxicity. A
403 similar conclusion also applies to handling by P-gp and/or BCRP at the blood-placenta barrier
404 and protection of the fetus. Globally, transport of plastic additives by ABC efflux pumps can
405 be considered as a general protective mechanism against the potential toxicity of these
406 compounds.

407 SLC transporters may play a key-role in liver and kidney uptake of their plastic additive
408 substrates. This is notably the case for BDE-47, BDE-99 and BDE-153, recognized as high
409 affinity substrates for OATPs, which probably explains their liver-specific accumulation
410 (Pacyniak et al., 2010). For PFOA, its transport by OAT1 and OAT3 likely contributes to its
411 renal secretion (Nakagawa et al., 2008) whereas its transport by OAT4 probably mediates its
412 tubular reabsorption, thus resulting in the final detection of only a small amount of PFOA in
413 urine in humans (Nakagawa et al., 2009). PFOS has additionally been demonstrated to be a
414 substrate for intestinal and hepatic uptake transporters of bile salts, *i.e.*, ASBT and NTCP,
415 which likely participate to its enterohepatic circulation and, furthermore, to its long half-life
416 and its hepatic accumulation (Zhao et al., 2015).

417 In the case of plastic additive substrates of transporters, especially for those with a
418 marked lipophilic nature, it is noteworthy that passive diffusion probably also contributes to
419 their passage across the plasma membrane. The net transport is therefore the sum of active and
420 passive transport, as already suggested for drugs (Sugano et al., 2010). For SLC transporters,
421 the uptake velocity is consequently the addition of transporter-mediated saturable transport and

422 of transporter-independent non-saturable passage. As illustrated in Fig. 3, for low
423 concentrations of plastic additives, relevant to those occurring in response to environmental
424 exposures, the fraction of saturable transporter-dependent passage across the plasma membrane
425 may be preponderant, thus highlighting the theoretical importance of transporters for the
426 toxicokinetics of these environmental contaminants.

427 **7. Conclusion and perspectives**

428 The studies summarized in this review clearly indicate that various plastic additives can
429 interact with drug transporters, as inhibitors of their activity, as their substrates or as modulators
430 of their expression. However, these regulatory effects have been mostly demonstrated *in vitro*
431 and their *in vivo* relevance remains a crucial point to determine. The rather low concentrations
432 of plastic additives in humans in response to environmental exposure (when compared to those
433 of drugs administered to patients) probably limits the *in vivo* effects of plastic additives as
434 single agents towards the activity or expression of transporters. However, a caveat to this
435 conclusion is the fact that humans are commonly exposed to mixtures of pollutants, including
436 plastic additives, whose inhibitory effects towards transporters may add up or synergise, as
437 already discussed above for marine pollutants inhibiting P-gp (Nicklisch et al., 2016), or as
438 demonstrated for a binary mixture of heterocyclic amines blocking OAT3 (Sayyed et al., 2017)
439 or for the inhibitory effects of pesticides towards P-gp activity (Pivčević and Zaja, 2006). The
440 study of potential inhibitory effects of mixtures of pollutants, such as mixtures of plastic
441 additives or mixtures of plastic additives and other pollutants, towards drug transporters is
442 therefore warranted. In this context, the recently described inhibitory effects of microplastics
443 towards ABC transporters (Koepsell, 2020) is particularly noteworthy because they are likely
444 to add up with those of plasticizers that are commonly released from plastic microparticles or
445 nanoparticles (Gulizia et al., 2023). Moreover, although plastic additive concentrations are
446 usually rather low in plasma of exposed humans, the specific accumulation of some of them in

447 certain tissues cannot be ruled out (Cimmino et al., 2020), potentially resulting in intra-tissular
448 concentrations of plastic additives reaching levels active on transporter activity or expression.
449 A better characterization of the human toxicokinetics of plastic additives, including predictions
450 of their intra-tissular concentrations and evaluation of their unbound form, in response to
451 environmental exposure, may therefore be useful to apprehend the possible *in vivo* relevance of
452 their effects on drug transporter activity and/or expression and their putative toxicity. This may
453 be done through physiologically-based toxicokinetic (PBTK) approaches, already developed
454 for plasticizers (Jeong et al., 2020; Sarigiannis et al., 2016). Other *in silico* methods, including
455 those based on quantitative structure-activity relationships (QSAR) or molecular docking (Ai
456 et al., 2015), may also be useful for predicting interactions of plastic additives with drug
457 transporters, especially for emerging plastic additives such as organophosphate flame retardants
458 for which data are currently lacking. The recent use of molecular docking to determine the
459 binding affinities of various chemicals, including PFASs, to P-gp, BCRP and MRP1, fully
460 supports this methodology (Barhoumi et al., 2022). New approaches to study clinical inhibition
461 of transporter activities by measuring altered pharmacokinetics of endogenous substrates,
462 reported notably for OATP1B1 and OAT inhibition by drugs (Neuvonen et al., 2021; Willemin
463 et al., 2021), may additionally be effective to study putative inhibitory effects of plasticizers
464 in humans.

465 The handling of plastic additives by transporters and their pharmacokinetics
466 consequences have been determined for some additives, such as BDEs and PFOA. This fully
467 validates the concept that transporters can participate in the pharmacokinetics of plastic
468 additives. However, data about the nature of plastic additives, and of chemical pollutants
469 overall, as substrates of drug transporters, remain rather scarce. An extensive analysis of the
470 potential handling of plastic additives, including emergent ones, by drug transporters is
471 therefore warranted. It is noteworthy that the traditional approach for such analyses, *i.e.*,

472 comparing the uptake of putative substrates in cellular or vesicular models overexpressing or
473 not the transporter of interest and in the absence or presence of specific transporter inhibitors,
474 requires the precise quantification of cellular or vesicular accumulation of plastic additives
475 through sensitive liquid chromatography-tandem mass spectrometry (LC-MS/MS). This
476 method can, however, be extremely consuming in time and/or money, and may be difficult to
477 integrate into high throughput processes. Experimental approaches such as *trans*-stimulation or
478 competitive counterflow assays (Schäfer et al., 2018; Severance et al., 2017) may be interesting
479 alternatives. Indeed, these assays, based on the ability of candidate substrates to *trans*-stimulate
480 uptake or efflux of a reference tracer substrate, do not require measurement of intracellular
481 levels of candidate substrates such as plastic additives and may be applied to large series of
482 chemicals when using a fluorescent dye as a tracer substrate; this is noteworthy for plastic
483 additives, of which there are more than 400. The relatively low sensitivity of the *trans*-
484 stimulation assay can, nevertheless, be limiting (Lefèvre et al., 2021). In addition to identifying
485 the plastic additives that are substrates for transporters, the possible effect of transporter
486 polymorphisms towards plasticizer toxicokinetics, as was reported for drugs (Yee et al., 2018),
487 will need to be investigated. This implication of transporter polymorphism could apply for
488 environmental chemicals overall (Darney et al., 2020), and it may help to identify individual
489 susceptibility factors to pollutants, including plastic additives.

490 Finally, it is noteworthy that the list of membrane transporters of interest for drug
491 pharmacokinetics and toxicity is rather evolutive and emerging transporters of clinical
492 significance, including plasma membrane transporters of vitamins or lysosomal and
493 mitochondrial transporters, likely merit attention for drug candidates (Chu et al., 2018;
494 Giacomini et al., 2022; Yee and Giacomini, 2021). These emerging transporters are also
495 potentially targeted by pollutants and may therefore be considered in the future for plastic
496 additives. Amino-acid transporters, such as aspartate transporters (*SLC1A2* and *SLC1A3*)

497 regulated by BPA in human hepatic HepG2 cells (Jiménez-Torres et al., 2020), as well as
498 glucose transporters, norepinephrine transporter (*SLC6A2*) and thyroid hormone transporters,
499 impacted by BPA or phthalates (Bereketoglu and Pradhan, 2022; Li et al., 2016; Quesnot et al.,
500 2014; Sakurai et al., 2004; Toyohira et al., 2003), may additionally be of interest.

501

502 **Author contribution statement**

503 Valentin Tastet - conceptualization, writing; Marc Le Vée - reviewing; Arnaud Bruyère -
504 reviewing; Olivier Fardel - conceptualization, writing, editing.

505 **Declaration of competing interest**

506 The authors declare that they have no competing financial interests or personal relationships
507 that could have appeared to influence the work reported in this paper.

508 **Data availability**

509 No data was used for the research described in the article.

510 **Acknowledgments**

511 This study has received funding from the European Union's Horizon Europe research and
512 innovation program under Grant Agreement N° 101057014 (European Partnership for the
513 Assessment of Risks from Chemicals) and from the French Agency for Food, Environmental
514 and Occupational Health & Safety (ANSES) under Grant Agreement N° 2020/01/124. The
515 authors thank Dr Catherine Lavau for proofreading the manuscript.

516

517

518

519

520

521

522

523 **References**

524 Abe, T., Takahashi, M., Kano, M., Amaike, Y., Ishii, C., Maeda, K., Kudoh, Y., Morishita,
525 T., Hosaka, T., Sasaki, T., Kodama, S., Matsuzawa, A., Kojima, H., Yoshinari, K., 2017.
526 Activation of nuclear receptor CAR by an environmental pollutant perfluorooctanoic acid. *Arch*
527 *Toxicol* 91, 2365-2374.

528 Abu-Qare, A.W., Elmasry, E., Abou-Donia, M.B., 2003. A role for P-glycoprotein in
529 environmental toxicology. *J Toxicol Environ Health B Crit Rev* 6, 279-288.

530 Ai, N., Fan, X., Ekins, S., 2015. In silico methods for predicting drug-drug interactions
531 with cytochrome P-450s, transporters and beyond. *Adv Drug Deliv Rev* 86, 46-60.

532 Amawi, H., Sim, H.M., Tiwari, A.K., Ambudkar, S.V., Shukla, S., 2019. ABC Transporter-
533 Mediated Multidrug-Resistant Cancer. *Adv Exp Med Biol* 1141, 549-580.

534 Angelini, A., Centurione, L., Sancilio, S., Castellani, M.L., Conti, P., Di Ilio, C., Porreca,
535 E., Cuccurullo, F., Di Pietro, R., 2011. The effect of the plasticizer diethylhexyl phthalate on
536 transport activity and expression of P-glycoprotein in parental and doxo-resistant human
537 sarcoma cell lines. *J Biol Regul Homeost Agents* 25, 203-211.

538 Anselmo, H.M., van den Berg, J.H., Rietjens, I.M., Murk, A.J., 2012. Inhibition of cellular
539 efflux pumps involved in multi xenobiotic resistance (MXR) in echinoid larvae as a possible
540 mode of action for increased ecotoxicological risk of mixtures. *Ecotoxicology* 21, 2276-2287.

541 Asimakopoulos, A.G., Thomaidis, N.S., Koupparis, M.A., 2012. Recent trends in
542 biomonitoring of bisphenol A, 4-t-octylphenol, and 4-nonylphenol. *Toxicol Lett* 210, 141-154.

543 Bain, L.J., LeBlanc, G.A., 1996. Interaction of structurally diverse pesticides with the
544 human MDR1 gene product P-glycoprotein. *Toxicol Appl Pharmacol* 141, 288-298.

545 Barhoumi, B., Sander, S.G., Tolosa, I., 2022. A review on per- and polyfluorinated alkyl
546 substances (PFASs) in microplastic and food-contact materials. *Environ Res* 206, 112595.

547 Beaudoin, J.J., Brouwer, K.L.R., Malinen, M.M., 2020. Novel insights into the organic
548 solute transporter alpha/beta, OST α/β : From the bench to the bedside. *Pharmacol Ther* 211,
549 107542.

550 Behr, A.C., Kwiatkowski, A., Ståhlman, M., Schmidt, F.F., Luckert, C., Braeuning, A.,
551 Buhrke, T., 2020. Impairment of bile acid metabolism by perfluorooctanoic acid (PFOA) and
552 perfluorooctanesulfonic acid (PFOS) in human HepaRG hepatoma cells. *Arch Toxicol* 94,
553 1673-1686.

554 Benjamin, S., Masai, E., Kamimura, N., Takahashi, K., Anderson, R.C., Faisal, P.A., 2017.
555 Phthalates impact human health: Epidemiological evidences and plausible mechanism of
556 action. *J Hazard Mater* 340, 360-383.

557 Bereketoglu, C., Pradhan, A., 2022. Plasticizers: negative impacts on the thyroid hormone
558 system. *Environ Sci Pollut Res Int* 29, 38912-38927.

- 559 Brouwer, K.L., Keppler, D., Hoffmaster, K.A., Bow, D.A., Cheng, Y., Lai, Y., Palm, J.E.,
560 Stieger, B., Evers, R., 2013. In vitro methods to support transporter evaluation in drug discovery
561 and development. *Clin Pharmacol Ther* 94, 95-112.
- 562 Brouwer, K.L.R., Evers, R., Hayden, E., Hu, S., Li, C.Y., Meyer Zu Schwabedissen, H.E.,
563 Neuhoﬀ, S., Oswald, S., Piquette-Miller, M., Saran, C., Sjöstedt, N., Sprowl, J.A., Stahl, S.H.,
564 Yue, W., 2022. Regulation of Drug Transport Proteins-From Mechanisms to Clinical Impact:
565 A White Paper on Behalf of the International Transporter Consortium. *Clin Pharmacol Ther*.
- 566 Bruyere, A., Hubert, C., Le Vee, M., Chedik, L., Sayyed, K., Stieger, B., Denizot, C.,
567 Parmentier, Y., Fardel, O., 2017. Inhibition of SLC drug transporter activities by environmental
568 bisphenols. *Toxicol In Vitro* 40, 34-44.
- 569 Burckhardt, G., 2012. Drug transport by Organic Anion Transporters (OATs). *Pharmacol*
570 *Ther* 136, 106-130.
- 571 Campanale, C., Massarelli, C., Savino, I., Locaputo, V., Uricchio, V.F., 2020. A Detailed
572 Review Study on Potential Effects of Microplastics and Additives of Concern on Human
573 Health. *Int J Environ Res Public Health* 17.
- 574 Cao, Y., Chen, Z., Zhang, M., Shi, L., Qin, S., Lv, D., Li, D., Ma, L., Zhang, Y., 2022.
575 Maternal exposure to bisphenol A induces fetal growth restriction via upregulating the
576 expression of estrogen receptors. *Chemosphere* 287, 132244.
- 577 Charuk, M.H., Grey, A.A., Reithmeier, R.A., 1998. Identification of the synthetic
578 surfactant nonylphenol ethoxylate: a P-glycoprotein substrate in human urine. *Am J Physiol*
579 274, F1127-1139.
- 580 Chatterjee, S., Annaert, P., 2018. Drug-induced Cholestasis: Mechanisms, Models, and
581 Markers. *Curr Drug Metab* 19, 808-818.
- 582 Chedik, L., Bruyere, A., Bacle, A., Potin, S., Le Vée, M., Fardel, O., 2018. Interactions of
583 pesticides with membrane drug transporters: implications for toxicokinetics and toxicity. *Expert*
584 *Opin Drug Metab Toxicol* 14, 739-752.
- 585 Chen, H.P., Lee, Y.K., Huang, S.Y., Shi, P.C., Hsu, P.C., Chang, C.F., 2018. Phthalate
586 exposure promotes chemotherapeutic drug resistance in colon cancer cells. *Oncotarget* 9,
587 13167-13180.
- 588 Cheng, Z., Li, H.H., Wang, H.S., Zhu, X.M., Sthiannopkao, S., Kim, K.W., Yasin, M.S.M.,
589 Hashim, J.H., Wong, M.H., 2016. Dietary exposure and human risk assessment of phthalate
590 esters based on total diet study in Cambodia. *Environ Res* 150, 423-430.
- 591 Choi, Y.H., Yu, A.M., 2014. ABC transporters in multidrug resistance and
592 pharmacokinetics, and strategies for drug development. *Curr Pharm Des* 20, 793-807.
- 593 Chu, X., Liao, M., Shen, H., Yoshida, K., Zur, A.A., Arya, V., Galetin, A., Giacomini,
594 K.M., Hanna, I., Kusuhara, H., Lai, Y., Rodrigues, D., Sugiyama, Y., Zamek-Gliszczyński,
595 M.J., Zhang, L., 2018. Clinical Probes and Endogenous Biomarkers as Substrates for
596 Transporter Drug-Drug Interaction Evaluation: Perspectives From the International Transporter
597 Consortium. *Clin Pharmacol Ther* 104, 836-864.
- 598 Ciarimboli, G., 2008. Organic cation transporters. *Xenobiotica* 38, 936-971.

- 599 Cimmino, I., Fiory, F., Perruolo, G., Miele, C., Beguinot, F., Formisano, P., Oriente, F.,
600 2020. Potential Mechanisms of Bisphenol A (BPA) Contributing to Human Disease. *Int J Mol*
601 *Sci* 21.
- 602 Clerbaux, L.A., Coecke, S., Lumen, A., Kliment, T., Worth, A.P., Paini, A., 2018.
603 Capturing the applicability of in vitro-in silico membrane transporter data in chemical risk
604 assessment and biomedical research. *Sci Total Environ* 645, 97-108.
- 605 Clerbaux, L.A., Paini, A., Lumen, A., Osman-Ponchet, H., Worth, A.P., Fardel, O., 2019.
606 Membrane transporter data to support kinetically-informed chemical risk assessment using non-
607 animal methods: Scientific and regulatory perspectives. *Environ Int* 126, 659-671.
- 608 Colnot, T., Kacew, S., Dekant, W., 2014. Mammalian toxicology and human exposures to
609 the flame retardant 2,2',6,6'-tetrabromo-4,4'-isopropylidenediphenol (TBBPA): implications
610 for risk assessment. *Arch Toxicol* 88, 553-573.
- 611 Cressman, A.M., Petrovic, V., Piquette-Miller, M., 2012. Inflammation-mediated changes
612 in drug transporter expression/activity: implications for therapeutic drug response. *Expert Rev*
613 *Clin Pharmacol* 5, 69-89.
- 614 Cui, H., Zhang, A.J., Chen, M., Liu, J.J., 2015. ABC Transporter Inhibitors in Reversing
615 Multidrug Resistance to Chemotherapy. *Curr Drug Targets* 16, 1356-1371.
- 616 Dankers, A.C., Roelofs, M.J., Piersma, A.H., Sweep, F.C., Russel, F.G., van den Berg, M.,
617 van Duursen, M.B., Masereeuw, R., 2013. Endocrine disruptors differentially target ATP-
618 binding cassette transporters in the blood-testis barrier and affect Leydig cell testosterone
619 secretion in vitro. *Toxicol Sci* 136, 382-391.
- 620 Darney, K., Turco, L., Buratti, F.M., Di Consiglio, E., Vichi, S., Roudot, A.C., Béchaux,
621 C., Testai, E., Dorne, J., Lautz, L.S., 2020. Human variability in influx and efflux transporters
622 in relation to uncertainty factors for chemical risk assessment. *Food Chem Toxicol* 140, 111305.
- 623 Domingo, J.L., Martí-Cid, R., Castell, V., Llobet, J.M., 2008. Human exposure to PBDEs
624 through the diet in Catalonia, Spain: temporal trend. A review of recent literature on dietary
625 PBDE intake. *Toxicology* 248, 25-32.
- 626 Doo, M.H., Li, H., Jang, H.I., Song, I.S., Chung, S.J., Shim, C.K., 2005. Effect of
627 nonylphenol ethoxylates (NPEs) on barrier functions of epithelial cell membranes: opening of
628 tight junctions and competitive inhibition of P-gp-mediated efflux. *Int J Pharm* 302, 145-153.
- 629 Dunnick, J.K., Shockley, K.R., Pandiri, A.R., Kissling, G.E., Gerrish, K.E., Ton, T.V.,
630 Wilson, R.E., Brar, S.S., Brix, A.E., Waidyanatha, S., Mutlu, E., Morgan, D.L., 2018. PBDE-
631 47 and PBDE mixture (DE-71) toxicities and liver transcriptomic changes at PND 22 after in
632 utero/postnatal exposure in the rat. *Arch Toxicol* 92, 3415-3433.
- 633 Eldasher, L.M., Wen, X., Little, M.S., Bircsak, K.M., Yacovino, L.L., Aleksunes, L.M.,
634 2013. Hepatic and renal Bcrp transporter expression in mice treated with perfluorooctanoic
635 acid. *Toxicology* 306, 108-113.
- 636 Engdahl, E., van Schijndel, M.D.M., Voulgaris, D., Di Criscio, M., Ramsbottom, K.A.,
637 Rigden, D.J., Herland, A., Rüegg, J., 2021. Bisphenol A Inhibits the Transporter Function of

- 638 the Blood-Brain Barrier by Directly Interacting with the ABC Transporter Breast Cancer
639 Resistance Protein (BCRP). *Int J Mol Sci* 22.
- 640 Fardel, O., Kolasa, E., Le Vee, M., 2012. Environmental chemicals as substrates, inhibitors
641 or inducers of drug transporters: implication for toxicokinetics, toxicity and pharmacokinetics.
642 *Expert Opin Drug Metab Toxicol* 8, 29-46.
- 643 Feiteiro, J., Mariana, M., Cairrão, E., 2021. Health toxicity effects of brominated flame
644 retardants: From environmental to human exposure. *Environ Pollut* 285, 117475.
- 645 Giacomini, K.M., Huang, S.M., 2013. Transporters in drug development and clinical
646 pharmacology. *Clin Pharmacol Ther* 94, 3-9.
- 647 Giacomini, K.M., Huang, S.M., Tweedie, D.J., Benet, L.Z., Brouwer, K.L., Chu, X.,
648 Dahlin, A., Evers, R., Fischer, V., Hillgren, K.M., Hoffmaster, K.A., Ishikawa, T., Keppler, D.,
649 Kim, R.B., Lee, C.A., Niemi, M., Polli, J.W., Sugiyama, Y., Swaan, P.W., Ware, J.A., Wright,
650 S.H., Yee, S.W., Zamek-Gliszczynski, M.J., Zhang, L., 2010. Membrane transporters in drug
651 development. *Nat Rev Drug Discov* 9, 215-236.
- 652 Giacomini, K.M., Yee, S.W., Koleske, M.L., Zou, L., Matsson, P., Chen, E.C., Kroetz,
653 D.L., Miller, M.A., Gozalpour, E., Chu, X., 2022. New and Emerging Research on Solute
654 Carrier and ATP Binding Cassette Transporters in Drug Discovery and Development: Outlook
655 From the International Transporter Consortium. *Clin Pharmacol Ther* 112, 540-561.
- 656 Gottesman, M.M., Pastan, I., 1993. Biochemistry of multidrug resistance mediated by the
657 multidrug transporter. *Annu Rev Biochem* 62, 385-427.
- 658 Grandin, F.C., Lacroix, M.Z., Gayrard, V., Viguié, C., Mila, H., de Place, A., Vayssière,
659 C., Morin, M., Corbett, J., Gayrard, C., Gely, C.A., Toutain, P.L., Picard-Hagen, N., 2019. Is
660 bisphenol S a safer alternative to bisphenol A in terms of potential fetal exposure ? Placental
661 transfer across the perfused human placenta. *Chemosphere* 221, 471-478.
- 662 Grimaldi, M., Boulahtouf, A., Toporova, L., Balaguer, P., 2019. Functional profiling of
663 bisphenols for nuclear receptors. *Toxicology* 420, 39-45.
- 664 Groh, K.J., Backhaus, T., Carney-Almroth, B., Geueke, B., Inostroza, P.A., Lennquist, A.,
665 Leslie, H.A., Maffini, M., Slunge, D., Trasande, L., Warhurst, A.M., Muncke, J., 2019.
666 Overview of known plastic packaging-associated chemicals and their hazards. *Sci Total*
667 *Environ* 651, 3253-3268.
- 668 Guéniche, N., Bruyere, A., Le Vée, M., Fardel, O., 2020. Implication of human drug
669 transporters to toxicokinetics and toxicity of pesticides. *Pest Manag Sci* 76, 18-25.
- 670 Gulizia, A.M., Patel, K., Philippa, B., Motti, C.A., van Herwerden, L., Vamvounis, G.,
671 2023. Understanding plasticiser leaching from polystyrene microplastics. *Sci Total Environ*
672 857, 159099.
- 673 Gunaalan, K., Fabbri, E., Capolupo, M., 2020. The hidden threat of plastic leachates: A
674 critical review on their impacts on aquatic organisms. *Water Res* 184, 116170.
- 675 Hahladakis, J.N., Velis, C.A., Weber, R., Iacovidou, E., Purnell, P., 2018. An overview of
676 chemical additives present in plastics: Migration, release, fate and environmental impact during
677 their use, disposal and recycling. *J Hazard Mater* 344, 179-199.

- 678 Hammer, H., Schmidt, F., Marx-Stoelting, P., Pötz, O., Braeuning, A., 2021. Cross-species
679 analysis of hepatic cytochrome P450 and transport protein expression. *Arch Toxicol* 95, 117-
680 133.
- 681 Hanet, N., Lancon, A., Delmas, D., Jannin, B., Chagnon, M.C., Cherkaoui-Malki, M.,
682 Latruffe, N., Artur, Y., Heydel, J.M., 2008. Effects of endocrine disruptors on genes associated
683 with 17beta-estradiol metabolism and excretion. *Steroids* 73, 1242-1251.
- 684 Hermabessiere, L., Dehaut, A., Paul-Pont, I., Lacroix, C., Jezequel, R., Soudant, P., Duflos,
685 G., 2017. Occurrence and effects of plastic additives on marine environments and organisms:
686 A review. *Chemosphere* 182, 781-793.
- 687 Horion, S., Thomé, J.P., Gismondi, É., 2015. Changes in antitoxic defense systems of the
688 freshwater amphipod *Gammarus pulex* exposed to BDE-47 and BDE-99. *Ecotoxicology* 24,
689 959-966.
- 690 Huang, F., Wen, S., Li, J., Zhong, Y., Zhao, Y., Wu, Y., 2014. The human body burden of
691 polybrominated diphenyl ethers and their relationships with thyroid hormones in the general
692 population in Northern China. *Sci Total Environ* 466-467, 609-615.
- 693 Huang, R.P., Liu, Z.H., Yuan, S.F., Yin, H., Dang, Z., Wu, P.X., 2017. Worldwide human
694 daily intakes of bisphenol A (BPA) estimated from global urinary concentration data (2000-
695 2016) and its risk analysis. *Environ Pollut* 230, 143-152.
- 696 Inui, K., Terada, T., Masuda, S., Saito, H., 2000. Physiological and pharmacological
697 implications of peptide transporters, PEPT1 and PEPT2. *Nephrol Dial Transplant* 15 Suppl 6,
698 11-13.
- 699 Jadhao, M., Tsai, E.M., Yang, H.C., Chen, Y.F., Liang, S.S., Wang, T.N., Teng, Y.N.,
700 Huang, H.W., Wang, L.F., Chiu, C.C., 2021. The Long-Term DEHP Exposure Confers
701 Multidrug Resistance of Triple-Negative Breast Cancer Cells through ABC Transporters and
702 Intracellular ROS. *Antioxidants (Basel)* 10.
- 703 Jani, M., Krajcsi, P., 2014. In vitro methods in drug transporter interaction assessment.
704 *Drug Discov Today Technol* 12, e105-112.
- 705 Jeong, S.H., Jang, J.H., Cho, H.Y., Lee, Y.B., 2020. Risk assessment for humans using
706 physiologically based pharmacokinetic model of diethyl phthalate and its major metabolite,
707 monoethyl phthalate. *Arch Toxicol* 94, 2377-2400.
- 708 Jetter, A., Kullak-Ublick, G.A., 2020. Drugs and hepatic transporters: A review. *Pharmacol*
709 *Res* 154, 104234.
- 710 Jiménez-Torres, C., Hernández-Kelly, L.C., Najimi, M., Ortega, A., 2020. Bisphenol A
711 exposure disrupts aspartate transport in HepG2 cells. *J Biochem Mol Toxicol* 34, e22516.
- 712 Jin, H., Audus, K.L., 2005. Effect of bisphenol A on drug efflux in BeWo, a human
713 trophoblast-like cell line. *Placenta* 26 Suppl A, S96-s103.
- 714 Johnson, D.R., Klaassen, C.D., 2002. Regulation of rat multidrug resistance protein 2 by
715 classes of prototypical microsomal enzyme inducers that activate distinct transcription
716 pathways. *Toxicol Sci* 67, 182-189.

- 717 Jones, S.R., Cyr, D.G., 2011. Regulation and characterization of the ATP-binding cassette
718 transporter-B1 in the epididymis and epididymal spermatozoa of the rat. *Toxicol Sci* 119, 369-
719 379.
- 720 Kalliokoski, A., Niemi, M., 2009. Impact of OATP transporters on pharmacokinetics. *Br J*
721 *Pharmacol* 158, 693-705.
- 722 Keiter, S., Burkhardt-Medicke, K., Wellner, P., Kais, B., Färber, H., Skutlarek, D.,
723 Engwall, M., Braunbeck, T., Keiter, S.H., Luckenbach, T., 2016. Does perfluorooctane
724 sulfonate (PFOS) act as chemosensitizer in zebrafish embryos? *Sci Total Environ* 548-549, 317-
725 324.
- 726 Kim, J.H., Yun, J., Sohng, J.K., Cha, J.M., Choi, B.C., Jeon, H.J., Kim, S.H., Choi, C.H.,
727 2007. Di(2-ethylhexyl)phthalate leached from medical PVC devices serves as a substrate and
728 inhibitor for the P-glycoprotein. *Environ Toxicol Pharmacol* 23, 272-278.
- 729 Kimura, O., Fujii, Y., Haraguchi, K., Kato, Y., Ohta, C., Koga, N., Endo, T., 2017. Uptake
730 of perfluorooctanoic acid by Caco-2 cells: Involvement of organic anion transporting
731 polypeptides. *Toxicology Letters* 277, 18-23.
- 732 Kimura, O., Fujii, Y., Haraguchi, K., Kato, Y., Ohta, C., Koga, N., Endo, T., 2020. Effects
733 of perfluoroalkyl carboxylic acids on the uptake of sulfobromophthalein via organic anion
734 transporting polypeptides in human intestinal Caco-2 cells. *Biochem Biophys Res Commun* 524, 100807.
- 735 Koepsell, H., 2020. Organic Cation Transporters in Health and Disease. *Pharmacol Rev*
736 72, 253-319.
- 737 König, J., Müller, F., Fromm, M.F., 2013. Transporters and drug-drug interactions:
738 important determinants of drug disposition and effects. *Pharmacol Rev* 65, 944-966.
- 739 Lee, S.C., Arya, V., Yang, X., Volpe, D.A., Zhang, L., 2017. Evaluation of transporters in
740 drug development: Current status and contemporary issues. *Adv Drug Deliv Rev* 116, 100-118.
- 741 Lee, W., Kim, R.B., 2004. Transporters and renal drug elimination. *Annu Rev Pharmacol*
742 *Toxicol* 44, 137-166.
- 743 Lefèvre, C.R., Le Vée, M., Gaubert, S., Jouan, E., Bruyere, A., Moreau, C., Fardel, O.,
744 2021. Substrate-Dependent Trans-Stimulation of Organic Cation Transporter 2 Activity. *Int J*
745 *Mol Sci* 22.
- 746 Leslie, E.M., Deeley, R.G., Cole, S.P., 2005. Multidrug resistance proteins: role of P-
747 glycoprotein, MRP1, MRP2, and BCRP (ABCG2) in tissue defense. *Toxicol Appl Pharmacol*
748 204, 216-237.
- 749 Li, A., Zhuang, T., Shi, W., Liang, Y., Liao, C., Song, M., Jiang, G., 2020. Serum
750 concentration of bisphenol analogues in pregnant women in China. *Sci Total Environ* 707,
751 136100.
- 752 Li, J., Wang, Y., Fang, F., Chen, D., Gao, Y., Liu, J., Gao, R., Wang, J., Xiao, H., 2016.
753 Bisphenol A disrupts glucose transport and neurophysiological role of IR/IRS/AKT/GSK3 β
754 axis in the brain of male mice. *Environ Toxicol Pharmacol* 43, 7-12.

- 755 Lim, J.J., Suh, Y., Faustman, E.M., Cui, J.Y., 2021. Regulation of transporters by
756 perfluorinated carboxylic acids in HepaRG cells. *Drug Metab Dispos.*
- 757 Lisco, G., Giagulli, V.A., Iovino, M., Guastamacchia, E., Pergola, G., Triggiani, V., 2022.
758 Endocrine-Disrupting Chemicals: Introduction to the Theme. *Endocr Metab Immune Disord*
759 *Drug Targets* 22, 677-685.
- 760 Liu, L., Liu, X., 2019. Contributions of Drug Transporters to Blood-Placental Barrier. *Adv*
761 *Exp Med Biol* 1141, 505-548.
- 762 Liu, X., 2019. Overview: Role of Drug Transporters in Drug Disposition and Its Clinical
763 Significance. *Adv Exp Med Biol* 1141, 1-12.
- 764 Loo, T.W., Clarke, D.M., 1998. Nonylphenol ethoxylates, but not nonylphenol, are
765 substrates of the human multidrug resistance P-glycoprotein. *Biochem Biophys Res Commun*
766 247, 478-480.
- 767 Ma, Y., Liu, H., Wu, J., Yuan, L., Wang, Y., Du, X., Wang, R., Marwa, P.W., Petlulu, P.,
768 Chen, X., Zhang, H., 2019. The adverse health effects of bisphenol A and related toxicity
769 mechanisms. *Environ Res* 176, 108575.
- 770 Mao, Q., Unadkat, J.D., 2015. Role of the breast cancer resistance protein (BCRP/ABCG2)
771 in drug transport--an update. *Aaps j* 17, 65-82.
- 772 Marchitti, S.A., Mazur, C.S., Dillingham, C.M., Rawat, S., Sharma, A., Zastre, J., Kenneke,
773 J.F., 2017. Inhibition of the Human ABC Efflux Transporters P-gp and BCRP by the BDE-47
774 Hydroxylated Metabolite 6-OH-BDE-47: Considerations for Human Exposure. *Toxicol Sci*
775 155, 270-282.
- 776 Mazur, C.S., Marchitti, S.A., Dimova, M., Kenneke, J.F., Lumen, A., Fisher, J., 2012.
777 Human and rat ABC transporter efflux of bisphenol a and bisphenol a glucuronide: interspecies
778 comparison and implications for pharmacokinetic assessment. *Toxicol Sci* 128, 317-325.
- 779 Meier, P.J., Stieger, B., 2002. Bile salt transporters. *Annu Rev Physiol* 64, 635-661.
- 780 Miller, D.S., 2015. Regulation of ABC transporters at the blood-brain barrier. *Clin*
781 *Pharmacol Ther* 97, 395-403.
- 782 Miranda, S., Vollrath, V., Wielandt, A.M., Loyola, G., Bronfman, M., Chianale, J., 1997.
783 Overexpression of mdr2 gene by peroxisome proliferators in the mouse liver. *J Hepatol* 26,
784 1331-1339.
- 785 Morse, B.L., Fallon, J.K., Kolar, A., Hogan, A.T., Smith, P.C., Hillgren, K.M., 2021.
786 Comparison of Hepatic Transporter Tissue Expression in Rodents and Interspecies Hepatic
787 OCT1 Activity. *Aaps j* 23, 58.
- 788 Mruk, D.D., Su, L., Cheng, C.Y., 2011. Emerging role for drug transporters at the blood-
789 testis barrier. *Trends Pharmacol Sci* 32, 99-106.
- 790 Müller, J., Keiser, M., Drozdik, M., Oswald, S., 2017. Expression, regulation and function
791 of intestinal drug transporters: an update. *Biol Chem* 398, 175-192.
- 792 Myllynen, P., Kummu, M., Kangas, T., Ilves, M., Immonen, E., Rysä, J., Pirilä, R.,
793 Lastumäki, A., Vähäkangas, K.H., 2008. ABCG2/BCRP decreases the transfer of a food-born

- 794 chemical carcinogen, 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) in perfused
795 term human placenta. *Toxicol Appl Pharmacol* 232, 210-217.
- 796 Nakagawa, H., Hirata, T., Terada, T., Jutabha, P., Miura, D., Harada, K.H., Inoue, K.,
797 Anzai, N., Endou, H., Inui, K., Kanai, Y., Koizumi, A., 2008. Roles of organic anion
798 transporters in the renal excretion of perfluorooctanoic acid. *Basic Clin Pharmacol Toxicol* 103,
799 1-8.
- 800 Nakagawa, H., Terada, T., Harada, K.H., Hitomi, T., Inoue, K., Inui, K., Koizumi, A.,
801 2009. Human organic anion transporter hOAT4 is a transporter of perfluorooctanoic acid. *Basic*
802 *Clin Pharmacol Toxicol* 105, 136-138.
- 803 Neuvonen, M., Tornio, A., Hirvensalo, P., Backman, J.T., Niemi, M., 2021. Performance
804 of Plasma Coproporphyrin I and III as OATP1B1 Biomarkers in Humans. *Clin Pharmacol Ther*
805 110, 1622-1632.
- 806 Nicklisch, S.C., Rees, S.D., McGrath, A.P., Gökirmak, T., Bonito, L.T., Vermeer, L.M.,
807 Cregger, C., Loewen, G., Sandin, S., Chang, G., Hamdoun, A., 2016. Global marine pollutants
808 inhibit P-glycoprotein: Environmental levels, inhibitory effects, and cocrystal structure. *Sci*
809 *Adv* 2, e1600001.
- 810 Nicklisch, S.C.T., Hamdoun, A., 2020. Disruption of small molecule transporter systems
811 by Transporter-Interfering Chemicals (TICs). *FEBS Lett* 594, 4158-4185.
- 812 Nicklisch, S.C.T., Pouv, A.K., Rees, S.D., McGrath, A.P., Chang, G., Hamdoun, A., 2021.
813 Transporter-interfering chemicals inhibit P-glycoprotein of yellowfin tuna (*Thunnus albacares*).
814 *Comp Biochem Physiol C Toxicol Pharmacol* 248, 109101.
- 815 Nies, A.T., Damme, K., Kruck, S., Schaeffeler, E., Schwab, M., 2016. Structure and
816 function of multidrug and toxin extrusion proteins (MATEs) and their relevance to drug therapy
817 and personalized medicine. *Arch Toxicol* 90, 1555-1584.
- 818 Nies, A.T., Keppler, D., 2007. The apical conjugate efflux pump ABCC2 (MRP2). *Pflugers*
819 *Arch* 453, 643-659.
- 820 Nigam, S.K., 2015. What do drug transporters really do? *Nat Rev Drug Discov* 14, 29-44.
- 821 Pacyniak, E., Roth, M., Hagenbuch, B., Guo, G.L., 2010. Mechanism of polybrominated
822 diphenyl ether uptake into the liver: PBDE congeners are substrates of human hepatic OATP
823 transporters. *Toxicol Sci* 115, 344-353.
- 824 Petzinger, E., Geyer, J., 2006. Drug transporters in pharmacokinetics. *Naunyn*
825 *Schmiedebergs Arch Pharmacol* 372, 465-475.
- 826 Pivčević, B., Zaja, R., 2006. Pesticides and their binary combinations as P-glycoprotein
827 inhibitors in NIH 3T3/MDR1 cells. *Environ Toxicol Pharmacol* 22, 268-276.
- 828 Popovic, M., Zaja, R., Fent, K., Smital, T., 2014. Interaction of environmental
829 contaminants with zebrafish organic anion transporting polypeptide, Oatp1d1 (Slco1d1).
830 *Toxicol Appl Pharmacol* 280, 149-158.
- 831 Quesnot, N., Bucher, S., Fromenty, B., Robin, M.A., 2014. Modulation of metabolizing
832 enzymes by bisphenol a in human and animal models. *Chem Res Toxicol* 27, 1463-1473.

- 833 Rizwan, A.N., Burckhardt, G., 2007. Organic anion transporters of the SLC22 family:
834 biopharmaceutical, physiological, and pathological roles. *Pharm Res* 24, 450-470.
- 835 Roberts, A.G., 2021. The Structure and Mechanism of Drug Transporters. *Methods Mol*
836 *Biol* 2342, 193-234.
- 837 Roth, M., Obaidat, A., Hagenbuch, B., 2012. OATPs, OATs and OCTs: the organic anion
838 and cation transporters of the SLCO and SLC22A gene superfamilies. *Br J Pharmacol* 165,
839 1260-1287.
- 840 Russel, F.G., Koenderink, J.B., Masereeuw, R., 2008. Multidrug resistance protein 4
841 (MRP4/ABCC4): a versatile efflux transporter for drugs and signalling molecules. *Trends*
842 *Pharmacol Sci* 29, 200-207.
- 843 Sakurai, K., Kawazuma, M., Adachi, T., Harigaya, T., Saito, Y., Hashimoto, N., Mori, C.,
844 2004. Bisphenol A affects glucose transport in mouse 3T3-F442A adipocytes. *Br J Pharmacol*
845 141, 209-214.
- 846 Sangkham, S., Faikhaw, O., Munkong, N., Sakunkoo, P., Arunlertaree, C., Chavali, M.,
847 Mousazadeh, M., Tiwari, A., 2022. A review on microplastics and nanoplastics in the
848 environment: Their occurrence, exposure routes, toxic studies, and potential effects on human
849 health. *Mar Pollut Bull* 181, 113832.
- 850 Sarigiannis, D.A., Karakitsios, S.P., Handakas, E., Simou, K., Solomou, E., Gotti, A., 2016.
851 Integrated exposure and risk characterization of bisphenol-A in Europe. *Food Chem Toxicol*
852 98, 134-147.
- 853 Sayyed, K., Le Vee, M., Abdel-Razzak, Z., Fardel, O., 2017. Inhibition of organic anion
854 transporter (OAT) activity by cigarette smoke condensate. *Toxicol In Vitro* 44, 27-35.
- 855 Schäfer, A.M., Bock, T., Meyer Zu Schwabedissen, H.E., 2018. Establishment and
856 Validation of Competitive Counterflow as a Method To Detect Substrates of the Organic Anion
857 Transporting Polypeptide 2B1. *Mol Pharm* 15, 5501-5513.
- 858 Schinkel, A.H., Jonker, J.W., 2003. Mammalian drug efflux transporters of the ATP
859 binding cassette (ABC) family: an overview. *Adv Drug Deliv Rev* 55, 3-29.
- 860 Severance, A.C., Sandoval, P.J., Wright, S.H., 2017. Correlation between Apparent
861 Substrate Affinity and OCT2 Transport Turnover. *J Pharmacol Exp Ther* 362, 405-412.
- 862 Shen, H., Lai, Y., Rodrigues, A.D., 2017. Organic Anion Transporter 2: An Enigmatic
863 Human Solute Carrier. *Drug Metab Dispos* 45, 228-236.
- 864 Sieppi, E., Vähäkangas, K., Rautio, A., Ietta, F., Paulesu, L., Myllynen, P., 2016. The
865 xenoestrogens, bisphenol A and para-nonylphenol, decrease the expression of the ABCG2
866 transporter protein in human term placental explant cultures. *Mol Cell Endocrinol* 429, 41-49.
- 867 Speidel, J.T., Xu, M., Abdel-Rahman, S.Z., 2018. Bisphenol A (BPA) and bisphenol S
868 (BPS) alter the promoter activity of the ABCB1 gene encoding P-glycoprotein in the human
869 placenta in a haplotype-dependent manner. *Toxicol Appl Pharmacol* 359, 47-54.

- 870 Sueyoshi, T., Li, L., Wang, H., Moore, R., Kodavanti, P.R., Lehmler, H.J., Negishi, M.,
871 Birnbaum, L.S., 2014. Flame retardant BDE-47 effectively activates nuclear receptor CAR in
872 human primary hepatocytes. *Toxicol Sci* 137, 292-302.
- 873 Sugano, K., Kansy, M., Artursson, P., Avdeef, A., Bendels, S., Di, L., Ecker, G.F., Faller,
874 B., Fischer, H., Gerebtzoff, G., Lennernaes, H., Senner, F., 2010. Coexistence of passive and
875 carrier-mediated processes in drug transport. *Nat Rev Drug Discov* 9, 597-614.
- 876 Summerfield, S.G., Yates, J.W.T., Fairman, D.A., 2022. Free Drug Theory - No Longer
877 Just a Hypothesis? *Pharm Res* 39, 213-222.
- 878 Takeshita, A., Inagaki, K., Igarashi-Migitaka, J., Ozawa, Y., Koibuchi, N., 2006. The
879 endocrine disrupting chemical, diethylhexyl phthalate, activates MDR1 gene expression in
880 human colon cancer LS174T cells. *J Endocrinol* 190, 897-902.
- 881 Tarafdar, A., Sirohi, R., Balakumaran, P.A., Reshmy, R., Madhavan, A., Sindhu, R., Binod,
882 P., Kumar, Y., Kumar, D., Sim, S.J., 2022. The hazardous threat of Bisphenol A: Toxicity,
883 detection and remediation. *J Hazard Mater* 423, 127097.
- 884 Teeguarden, J., Hanson-Drury, S., Fisher, J.W., Doerge, D.R., 2013. Are typical human
885 serum BPA concentrations measurable and sufficient to be estrogenic in the general population?
886 *Food Chem Toxicol* 62, 949-963.
- 887 Tirona, R.G., 2011. Molecular mechanisms of drug transporter regulation. *Handb Exp*
888 *Pharmacol*, 373-402.
- 889 Toyohira, Y., Utsunomiya, K., Ueno, S., Minami, K., Uezono, Y., Yoshimura, R., Tsutsui,
890 M., Izumi, F., Yanagihara, N., 2003. Inhibition of the norepinephrine transporter function in
891 cultured bovine adrenal medullary cells by bisphenol A. *Biochem Pharmacol* 65, 2049-2054.
- 892 van der Schoor, L.W., Verkade, H.J., Kuipers, F., Jonker, J.W., 2015. New insights in the
893 biology of ABC transporters ABCC2 and ABCC3: impact on drug disposition. *Expert Opin*
894 *Drug Metab Toxicol* 11, 273-293.
- 895 Volpe, D.A., 2016. Transporter assays as useful in vitro tools in drug discovery and
896 development. *Expert Opin Drug Discov* 11, 91-103.
- 897 Wang, C., Lin, Z., Dong, Q., Lin, Z., Lin, K., Wang, J., Huang, J., Huang, X., He, Y.,
898 Huang, C., Yang, D., Huang, C., 2012. Polybrominated diphenyl ethers (PBDEs) in human
899 serum from Southeast China. *Ecotoxicol Environ Saf* 78, 206-211.
- 900 Wang, L., Prasad, B., Salphati, L., Chu, X., Gupta, A., Hop, C.E., Evers, R., Unadkat, J.D.,
901 2015. Interspecies variability in expression of hepatobiliary transporters across human, dog,
902 monkey, and rat as determined by quantitative proteomics. *Drug Metab Dispos* 43, 367-374.
- 903 Wang, X., Wang, X., Zhu, Y., Chen, X., 2021. ADME/T-based strategies for paraquat
904 detoxification: Transporters and enzymes. *Environ Pollut* 291, 118137.
- 905 Wang, Y., Zhu, H., Kannan, K., 2019. A Review of Biomonitoring of Phthalate Exposures.
906 *Toxics* 7.
- 907 Wiesinger, H., Wang, Z., Hellweg, S., 2021. Deep Dive into Plastic Monomers, Additives,
908 and Processing Aids. *Environ Sci Technol* 55, 9339-9351.

- 909 Wigler, P.W., Patterson, F.K., 1993. Inhibition of the multidrug resistance efflux pump.
910 *Biochim Biophys Acta* 1154, 173-181.
- 911 Willemin, M.E., Van Der Made, T.K., Pijpers, I., Dillen, L., Kunze, A., Jonkers, S.,
912 Steemans, K., Tuytelaars, A., Jacobs, F., Monshouwer, M., Scotcher, D., Rostami-Hodjegan,
913 A., Galetin, A., Snoeys, J., 2021. Clinical Investigation on Endogenous Biomarkers to Predict
914 Strong OAT-Mediated Drug-Drug Interactions. *Clin Pharmacokinet* 60, 1187-1199.
- 915 Yang, C.H., Glover, K.P., Han, X., 2010. Characterization of cellular uptake of
916 perfluorooctanoate via organic anion-transporting polypeptide 1A2, organic anion transporter
917 4, and urate transporter 1 for their potential roles in mediating human renal reabsorption of
918 perfluorocarboxylates. *Toxicol Sci* 117, 294-302.
- 919 Yang, J., Wang, H., Du, H., Xu, L., Liu, S., Yi, J., Chen, Y., Jiang, Q., He, G., 2021. Serum
920 Bisphenol A, glucose homeostasis, and gestational diabetes mellitus in Chinese pregnant
921 women: a prospective study. *Environ Sci Pollut Res Int* 28, 12546-12554.
- 922 Yee, S.W., Brackman, D.J., Ennis, E.A., Sugiyama, Y., Kamdem, L.K., Blanchard, R.,
923 Galetin, A., Zhang, L., Giacomini, K.M., 2018. Influence of Transporter Polymorphisms on
924 Drug Disposition and Response: A Perspective From the International Transporter Consortium.
925 *Clin Pharmacol Ther* 104, 803-817.
- 926 Yee, S.W., Giacomini, K.M., 2021. Emerging Roles of the Human Solute Carrier 22
927 Family. *Drug Metab Dispos* 50, 1193-1210.
- 928 Yoshikawa, Y., Hayashi, A., Inai, M., Matsushita, A., Shibata, N., Takada, K., 2002.
929 Permeability characteristics of endocrine-disrupting chemicals using an in vitro cell culture
930 model, Caco-2 cells. *Curr Drug Metab* 3, 551-557.
- 931 Yu, Y., Wang, M., Zhang, K., Yang, D., Zhong, Y., An, J., Lei, B., Zhang, X., 2017. The
932 transepithelial transport mechanism of polybrominated diphenyl ethers in human intestine
933 determined using a Caco-2 cell monolayer. *Environ Res* 154, 93-100.
- 934 Zhao, W., Zitzow, J.D., Ehresman, D.J., Chang, S.C., Butenhoff, J.L., Forster, J.,
935 Hagenbuch, B., 2015. Na⁺/Taurocholate Cotransporting Polypeptide and Apical Sodium-
936 Dependent Bile Acid Transporter Are Involved in the Disposition of Perfluoroalkyl Sulfonates
937 in Humans and Rats. *Toxicol Sci* 146, 363-373.
- 938 Zhou, S., Shu, Y., 2022. Transcriptional Regulation of Solute Carrier (SLC) Drug
939 Transporters. *Drug Metab Dispos*.
- 940 Zolk, O., Fromm, M.F., 2011. Transporter-mediated drug uptake and efflux: important
941 determinants of adverse drug reactions. *Clin Pharmacol Ther* 89, 798-805.
- 942
- 943
- 944
- 945

946

947 Table 1. Plastic additives considered in this review.

Chemical class	Function	Name
Phthalates	Plasticizers	Di(2-ethylhexyl) phthalate (DEHP) Mono(2-ethylhexyl) phthalate (MEHP) (metabolite of DEHP) Butylbenzyl phthalate (BBP) Diethyl phthalate (DEP) Dibutyl phthalate (DBP)
Bisphenols	Monomers/ plasticizers/ antioxidants	Bisphenol A (BPA) BPA glucuronide (metabolite of BPA) Bisphenol F (BPF) Bisphenol S (BPS)
Brominated flame retardants (BFRs)	Flame retardants	Polybrominated diphenyl ethers (PBDE): BDE-47, BDE-99, BDE-100, BDE-153, 6-hydroxy-BDE-47 (metabolite of BDE-47) Tetrabromobisphenol A (TBBPA)
Alkylphenols	Plasticizers/antioxidants	tert-Butylphenol (tBP) tert-Octylphenol (tOP) Nonylphenol ethoxylate (NPE) Nonylphenol (NP) (metabolite of NPE)
Per- and poly-fluorinated alkyl substances (PFASs)	Coating agents	Perfluorobutane sulfonate (PFBS) Perfluorooctanoic acid (PFOA) Perfluorononanoic acid (PFNA) Perfluorodecanoic acid (PFDA) Perfluorohexane sulfonate (PFHxS) Perfluorooctane sulfonate (PFOS)

948

949

950

951

952

953

954 Table 2. Main types of substrates for human drug transporters.

Subfamily	Transporter	Type of substrates	Reference
ABCB	P-gp (<i>MDR1/ABCB1</i>)	Neutral or positively charged hydrophobic drugs (Example: etoposide, digoxin, doxorubicin, fexofenadine, paclitaxel, vinblastine)	(Gottesman and Pastan, 1993)
	BSEP (<i>ABCB11</i>)	Bile acids (Example: taurocholate)	(Jetter and Kullak-Ublick, 2020)
ABCC	MRP1 (<i>ABCC1</i>)	Hydrophobic and anionic drugs; glucuronide and glutathione conjugates; leukotriene C4 (Example: ciprofloxacin, doxorubicin, etoposide, irinotecan, ritonavir, saquinavir, vincristine)	(Choi and Yu, 2014)
	MRP2 (<i>ABCC2</i>)	Anionic drugs; glucuronide, glutathione and sulfate conjugates; sulphated bile salts; leukotriene C4 (Example: adefovir, ampicillin, bilirubin glucuronides, ritonavir, sequinavir)	(Nies and Keppler, 2007)
	MRP3 (<i>ABCC3</i>)	Anionic drugs; glucuronide conjugates; bile salts (Example: morphine-3-glucuronide, estradiol-17 β - glucuronide, taurocholate)	(van der Schoor et al., 2015)
	MRP4 (<i>ABCC4</i>)	Anionic drugs; antiviral nucleoside analogues; signalling molecules (Example: adefovir, cAMP, cGMP, furosemide, tenofovir)	(Russel et al., 2008)
ABCG	BCRP (<i>ABCG2</i>)	Hydrophobic drugs; hydrophilic conjugated organic anions (Example: cimetidine, doxorubicin, estrone 3-sulfate, mitoxantrone, prazosine, rosuvastatin, uric acid)	(Mao and Unadkat, 2015)
SLC10	NTCP (<i>SLC10A1</i>) ASBT (<i>SLC10A2</i>)	Bile acids (Example: taurocholate)	(Meier and Stieger, 2002)
SLC15	PEPT1 (<i>SLC15A1</i>)	Peptidomimetic drugs (Example: β -lactam antibiotics)	(Inui et al., 2000)
SLCO	OATP1B1 (<i>SLCO1B1</i>)	Anionic drugs; bile acids; bilirubin; steroid and thyroid hormones (Example: bromosulphophthalein, estrone 3-sulfate, statins, valsartan)	(Kalliokoski and Niemi, 2009)
	OATP1B3 (<i>SLCO1B3</i>)	Anionic drugs; bile acids; bilirubin; steroid and thyroid hormones (Example: cholecystokinin, estrone 3 sulfate, statins)	
	OATP2B1 (<i>SLCO2B1</i>)	Anionic drugs; bile acids; steroid hormones (Example: estrone 3-sulfate, fexofenadine, statins)	
SLC22	OCT1 (<i>SLC22A1</i>) OCT2 (<i>SLC22A2</i>)	Cationic drugs; monoamine neurotransmitters (Example: acyclovir, cimetidine, cisplatin, metformin, oxaliplatin, choline, dopamine)	(Koepsell, 2020)
	OAT1 (<i>SLC22A6</i>) OAT3 (<i>SLC22A8</i>)	Anionic drugs; cyclic nucleotides; dicarboxylic acids (Example : adefovir, zidovudine, ciprofloxacin, cephaloridin, methotrexate, para-aminohippurate, pravastatin)	(Burckhardt, 2012)
	OAT2 (<i>SLC22A8</i>)	Anionic drugs; cyclic nucleotides; prostaglandins; steroid hormones (Example: bumetanide, cGMP, creatinine, diclofenac, erythromycin, estrone 3-sulfate, uric acid)	(Shen et al., 2017)
SLC47	MATE1 (<i>SLC47A1</i>) MATE2-K (<i>SLC47A2</i>)	Cationic drugs (Example: cimetidine, creatinine, ganciclovir, metformin, oxaliplatin)	(Koepsell, 2020)
SLC51	OST α (<i>SLC51A</i>) OST β (<i>SLC51B</i>)	Bile acids; conjugated steroids (Example: taurocholate, estrone 3-sulfate, digoxin)	(Beaudoin et al., 2020)

955 Note: P-gp - P-glycoprotein; *MDR* - multidrug resistance gene; BSEP - bile salt export pump; MRP - multidrug
956 resistance-associated protein; BCRP - breast cancer resistance protein; NTCP - sodium-taurocholate
957 cotransporting polypeptide; ASBT - apical sodium-dependent bile acid transporter; PEPT - peptide transporter;
958 OATP - organic anion transporting polypeptide; OCT - organic cation transporter; OAT - organic anion
959 transporter; MATE - multidrug and toxin extrusion protein; OST - organic solute transporter.

961

962 Table 3. Modulation of human ABC drug transporter activities by plastic additives.

Transporter	Chemical class	Transporter activity		
		Stimulation	Inhibition	Lack of effect
P-gp	Bisphenols	BPA (100 nM ^{CA}) (Jin and Audus, 2005)		BPA (100 μM ^{VA}) (Dankers et al., 2013)
	Phthalates		DEHP (30.1 mM ^{CA}) (Kim et al., 2007) DEP (16.1 mM ^{CA}) (Kim et al., 2007) DBP (32.1 mM ^{CA}) (Kim et al., 2007)	DEHP (100 μM ^{VA}) (Dankers et al., 2013) MEHP (100 μM ^{VA}) (Dankers et al., 2013)
	BFRs		BDE-47 (50 μM ^{VA}) (Nicklisch et al., 2016) 6-OH-BDE-47 (IC ₅₀ =11.4 μM ^{VA} , 50 μM ^{CA}) (Marchitti et al., 2017) TBBPA (IC ₅₀ =22.9 μM ^{VA}) (Dankers et al., 2013)	PBDE-47 (300 μM ^{VA} , 100 μM ^{CA}) (Marchitti et al., 2017) PBDE-100 (50 μM ^{VA}) (Nicklisch et al., 2016)
	Alkylphenols		NPE (0.1-1 μM ^{CA}) (Doo et al., 2005)	
	PFASs		PFOA (100 μM ^{VA}) (Dankers et al., 2013) PFOS (100 μM ^{VA}) (Dankers et al., 2013)	
BSEP	PFASs		PFOS (100 μM ^{VA}) (Zhao et al., 2015)	PFBS (100 μM ^{VA}) (Zhao et al., 2015) PFHxS (100 μM ^{VA}) (Zhao et al., 2015)
MRP1	Bisphenols			BPA (100 μM ^{VA}) (Dankers et al., 2013)
	Phthalates			DEHP (100 μM ^{VA}) (Dankers et al., 2013) MEHP (100 μM ^{VA}) (Dankers et al., 2013)
	BFRs		TBBPA (100 μM ^{VA}) (Dankers et al., 2013)	
	PFASs		PFOA (100 μM ^{VA}) (Dankers et al., 2013) PFOS (100 μM ^{VA}) (Dankers et al., 2013)	
MRP2	PFASs		PFHxS (100 μM ^{VA}) (Zhao et al., 2015) PFOS (100 μM ^{VA}) (Zhao et al., 2015)	PFBS (100 μM ^{VA}) (Zhao et al., 2015)
MRP4	Bisphenols			BPA (100 μM ^{VA}) (Dankers et al., 2013)
	Phthalates			DEHP (100 μM ^{VA}) (Dankers et al., 2013) MEHP (100 μM ^{VA}) (Dankers et al., 2013)
	BFRs		TBBPA (IC ₅₀ =24.0 μM ^{VA}) (Dankers et al., 2013)	
	PFASs		PFOA (100 μM ^{VA}) (Dankers et al., 2013) PFOS (100 μM ^{VA}) (Dankers et al., 2013)	
BCRP	Bisphenols		BPA (500 nM ^{CA} , 100 μM ^{VA}) (Dankers et al., 2013; Engdahl et al., 2021)	
	Phthalates			DEHP (100 μM ^{VA}) (Dankers et al., 2013) MEHP (100 μM ^{VA}) (Dankers et al., 2013)
	BFRs		BDE-47 (IC ₅₀ =45.7 μM ^{VA}) (Marchitti et al., 2017) 6-hydroxy-BDE-47 (IC ₅₀ =9.4 μM ^{VA} , 50 μM ^{CA}) (Marchitti et al., 2017) TBBPA (100 μM ^{VA}) (Dankers et al., 2013)	BDE-47 (100 μM ^{CA}) (Marchitti et al., 2017)
	PFASs		PFOA (5-100 μM ^{VA}) (Dankers et al., 2013; Eldasher et al., 2013) PFOS (100 μM ^{VA}) (Dankers et al., 2013) PFHxS (100 μM ^{VA}) (Zhao et al., 2015)	

963 Note: Abbreviations for plastic additives and transporters are those defined in Table 1 and Table 2, respectively; CA - cellular
 964 assay; VA - vesicular assay; IC₅₀ - half maximal inhibitory concentration.

965

966

967

968

969

970

971

972

973

974

975 Table 4. Modulation of human SLC drug transporter activities by plastic additives.

Transporter	Chemical class	Transporter activity		
		Stimulation	Inhibition	Lack of effect
OCT1	Bisphenols	BPS (100 μ M) (Bruyere et al., 2017)	BPA (IC ₅₀ =39.0 μ M) (Bruyere et al., 2017) BPF (100 μ M) (Bruyere et al., 2017)	
	BFRs		TBBPA (IC ₅₀ =37.5 μ M) (Bruyere et al., 2017)	
OCT2	Bisphenols	BPF (100 μ M) (Bruyere et al., 2017) BPS (100 μ M) (Bruyere et al., 2017)		BPA (100 μ M) (Bruyere et al., 2017)
	BFRs		TBBPA (100 μ M) (Bruyere et al., 2017)	
MATE1	Bisphenols		BPA (IC ₅₀ =73.5 μ M) (Bruyere et al., 2017) BPF (100 μ M) (Bruyere et al., 2017) BPS (100 μ M) (Bruyere et al., 2017)	
	BFRs		TBBPA (IC ₅₀ =23.1 μ M) (Bruyere et al., 2017)	
MATE2-K	Bisphenols			BPA (100 μ M) (Bruyere et al., 2017) BPF (100 μ M) (Bruyere et al., 2017) BPS (100 μ M) (Bruyere et al., 2017)
	BFRs			TBBPA (100 μ M) (Bruyere et al., 2017)
OATP1B1	Bisphenols		BPA (IC ₅₀ =18.7 μ M) (Bruyere et al., 2017) BPF (IC ₅₀ =13.4 μ M) (Bruyere et al., 2017)	BPS (100 μ M) (Bruyere et al., 2017)
	BFRs		TBBPA (IC ₅₀ =0.6 μ M) (Bruyere et al., 2017) BDE-47 (100 pM) (Pacyniak et al., 2010) BDE-99 (100 nM) (Pacyniak et al., 2010) BDE-153 (100 nM) (Pacyniak et al., 2010)	
OATP1B3	Bisphenols	BPF (100 μ M) (Bruyere et al., 2017)	BPA (100 μ M) (Bruyere et al., 2017)	BPS (100 μ M) (Bruyere et al., 2017)
	BFRs		TBBPA (IC ₅₀ =12.3 μ M) (Bruyere et al., 2017) BDE-47 (100 pM) (Pacyniak et al., 2010) BDE-99 (100 nM) (Pacyniak et al., 2010) BDE-153 (100 nM) (Pacyniak et al., 2010)	
OATP2B1	BFRs		BDE-47 (100 pM) (Pacyniak et al., 2010) BDE-99 (100 nM) (Pacyniak et al., 2010) BDE-153 (100 nM) (Pacyniak et al., 2010)	
	PFASs		PFOA (Ki=62.2 μ M) (Kimura et al., 2020) PFNA (Ki=35.3 μ M) (Kimura et al., 2020) PFDA (Ki=43.2 μ M) (Kimura et al., 2020)	
OAT1	Bisphenols	BPA (100 μ M) (Bruyere et al., 2017) BPF (100 μ M) (Bruyere et al., 2017)	BPS (100 μ M) (Bruyere et al., 2017)	
	BFRs			TBBPA (100 μ M) (Bruyere et al., 2017)
	PFASs		PFOA (10 μ M) (Nakagawa et al., 2008)	
OAT2	PFASs		PFOA (10 μ M) (Nakagawa et al., 2008)	
OAT3	Bisphenols		BPA (IC ₅₀ =9.2 μ M) (Bruyere et al., 2017) BPF (IC ₅₀ =26.8 μ M) (Bruyere et al., 2017) BPS (IC ₅₀ =23.3 μ M) (Bruyere et al., 2017)	
	BFRs		TBBPA (IC ₅₀ =0.5 μ M) (Bruyere et al., 2017)	
	PFASs		PFOA (10 μ M) (Nakagawa et al., 2008)	
NTCP	Bisphenols		BPA (100 μ M) (Bruyere et al., 2017) BPF (100 μ M) (Bruyere et al., 2017)	BPS (100 μ M) (Bruyere et al., 2017)
	BFRs		TBBPA (IC ₅₀ =4.5 μ M) (Bruyere et al., 2017) PFBS (10 μ M) (Zhao et al., 2015) PFHxS (10 μ M) (Zhao et al., 2015) PFOS (10 μ M) (Zhao et al., 2015)	

976 Note: Abbreviations for plastic additives and transporters are those defined in Table 1 and Table 2, respectively; IC₅₀ - half
 977 maximal inhibitory concentration; Ki - inhibition constant.

978

979

980

981 Table 5. Plastic additives as substrates for human ABC and SLC drug transporters.

Transporter	Plastic additive		
	Chemical class	Substrate	Not substrate
P-gp	Bisphenols	BPA ^{CA} (Yoshikawa et al., 2002)	BPA ^{ATA, CA} (Dankers et al., 2013; Mazur et al., 2012) BPA-Glucuronide ^{ATA} (Mazur et al., 2012) BPS ^{Perfused human placenta} (Grandin et al., 2019)
	Phthalates	DEHP ^{CA, ATA} (Kim et al., 2007; Yoshikawa et al., 2002)	BBP ^{CA} (Yoshikawa et al., 2002) DBP ^{CA, ATA} (Kim et al., 2007; Yoshikawa et al., 2002) DEP ^{CA, ATA} (Kim et al., 2007)
	BFRs		TBBPA ^{CA} (Dankers et al., 2013)
	Alkylphenols	tBP ^{CA} (Yoshikawa et al., 2002) tOP ^{CA} (Yoshikawa et al., 2002) NPE ^{ATA, CA} (Charuk et al., 1998; Loo and Clarke, 1998)	NP ^{CA} (Loo and Clarke, 1998)
	PFASs		PFOA ^{CA, ATA} (Dankers et al., 2013)
MRP2	Bisphenols	BPA ^{ATA} (Mazur et al., 2012)	BPA-Glucuronide ^{ATA} (Mazur et al., 2012)
MRP3	Bisphenols	BPA ^{ATA} (Mazur et al., 2012) BPA-Glucuronide ^{ATA} (Mazur et al., 2012)	
BCRP	Bisphenols	BPA ^{ATA, CA} (Dankers et al., 2013; Mazur et al., 2012)	BPA-Glucuronide ^{ATA} (Mazur et al., 2012) TBBPA ^{CA} (Dankers et al., 2013)
	BFRs		TBBPA ^{CA} (Dankers et al., 2013)
	PFASs	PFOA ^{CA} (Dankers et al., 2013)	
OATP1B1	BFRs	BDE-47 (Km=0.31 μM^{CA}) (Pacyniak et al., 2010) BDE-99 (Km=0.91 μM^{CA}) (Pacyniak et al., 2010) BDE-153 (Km=1.91 μM^{CA}) (Pacyniak et al., 2010)	
OATP2B1	BFRs	BDE-47 (Km=0.41 μM^{CA}) (Pacyniak et al., 2010) BDE-99 (Km=0.70 μM^{CA}) (Pacyniak et al., 2010) BDE-153 (Km=1.66 μM^{CA}) (Pacyniak et al., 2010)	
	PFASs	PFOA (Km=8.3 μM^{CA}) (Kimura et al., 2017)	
OATP1B3	BFRs	BDE-47 (Km=0.81 μM^{CA}) (Pacyniak et al., 2010) BDE-99 (Km=0.87 μM^{CA}) (Pacyniak et al., 2010) BDE-153 (Km=0.65 μM^{CA}) (Pacyniak et al., 2010)	
OAT1	PFASs	PFOA (Km=48.0 μM^{CA}) (Nakagawa et al., 2008)	
OAT3	Bisphenols		BPA ^{CA} (Bruyere et al., 2017)
	PFASs	PFOA (Km=49.1 μM^{CA}) (Nakagawa et al., 2008)	
OAT4	PFASs	PFOA (Km=310.3 μM^{CA}) (Nakagawa et al., 2009; Yang et al., 2010)	
NTCP	PFASs	PFBS (Km=39.6 μM^{CA}) (Zhao et al., 2015) PFHxS (Km=112 μM^{CA}) (Zhao et al., 2015) PFOS (Km=130 μM^{CA}) (Zhao et al., 2015)	
ASBT	PFASs	PFOS ^{CA} (Zhao et al., 2015)	PFBS ^{CA} (Zhao et al., 2015) PFHxS ^{CA} (Zhao et al., 2015)
OST α/β	PFASs	PFBS ^{CA} (Zhao et al., 2015) PFHxS ^{CA} (Zhao et al., 2015) PFOS ^{CA} (Zhao et al., 2015)	

982 Note: Abbreviations for plastic additives and transporters are those defined in Table 1 and Table 2, respectively; CA - cellular
 983 assay; ATA - vesicular ATPase assay; Km - Michaelis constant.


984

985

986 Table 6. Regulation of human drug transporter expression by plastic additives.

Transporter	Chemical class (Plastic additive)	Regulation	Reference
P-gp/ <i>MDR1</i>	Bisphenols	Induction of <i>MDR1</i> promoter activity by BPA (50 nM/acute exposure/15 min or 3 nM/chronic exposure/12 days), BPS (0.3 nM/chronic exposure/12 days) or a mixture BPA/BPS (5 nM BPA/1.5 nM BPS/chronic exposure/12 days) in an haplotype-dependent manner. Repression of <i>MDR1</i> promoter activity by BPS (0.5 nM/acute exposure/15 min) in an haplotype-dependent manner.	(Speidel et al., 2018)
		Induction of <i>MDR1</i> mRNA expression, but not that of P-gp, by BPA (80 μ M/24 h) in human hepatic HepG2 cells.	(Hanet et al., 2008)
		Modest induction of P-gp expression by BPA (10 μ M/48 h) in placental BeWo cells.	(Jin and Audus, 2005)
	Phthalates	Induction of P-gp expression by DEHP (10 μ M/24-72 h) and MEHP (10 μ M/24-72 h) in cancer colon cell lines.	(Chen et al., 2018)
		Induction of P-gp expression by DEHP (6 and 12 μ M/24 h) in sarcoma cells.	(Angelini et al., 2011)
		Induction of <i>MDR1</i> mRNA expression by DEHP (1-10 μ M/24 h) in human cancer colon cells.	(Takeshita et al., 2006)
		Induction of <i>MDR1</i> mRNA and P-gp expression in breast cancer MDA-MB-231 cells by chronic exposure to DEHP (100 nM/three months).	(Jadhaio et al., 2021)
Alkylphenols	No apparent induction of P-gp expression in human intestinal Caco-2 cells exposed to NPE (up to 1 μ M/72 h).	(Doo et al., 2005)	
BSEP	PFASs	Repression of BSEP mRNA expression by PFOA (250 and 500 μ M/24 h) and PFOS (25-100 μ M/24 h) in human hepatic HepaRG cells.	(Behr et al., 2020)
MRP1	Bisphenols	Induction of MRP1 mRNA expression by BPA (80 μ M/24 h) in HepG2 cells.	(Hanet et al., 2008)
MRP2	Bisphenols	Induction of MRP2 mRNA and protein expression by BPA (80 μ M/24 h) in HepG2 cells.	(Hanet et al., 2008)
	PFASs	Repression of MRP2 mRNA expression by PFOA (250 and 500 μ M/24 h) and PFOS (25 μ M/24 h) in HepaRG cells.	(Behr et al., 2020)
MRP3	Bisphenols	Induction of MRP3 mRNA and protein expression by BPA (80 μ M/24 h) in HepG2 cells.	(Hanet et al., 2008)
	PFASs	Induction of MRP3 mRNA expression by PFNA (45 μ M/24 h) and PFDA (45 μ M/24 h), but repression by PFOA (500 μ M/24 h) and PFOS (100 μ M/24 h) in HepaRG cells.	(Behr et al., 2020; Lim et al., 2021)
MRP4	Bisphenols	Lack of effects of BPA (80 μ M/24 h) towards MRP4 mRNA expression in human hepatic HepG2 cells.	(Hanet et al., 2008)
BCRP	Bisphenols	Repression of BCRP protein expression by BPA (1 nM/48 h) in human term placental explant cultures.	(Sieppi et al., 2016)
		Repression of BCRP mRNA and protein expression by BPA (100 mM/48 h) in BeWo cells.	(Cao et al., 2022)
	Alkylphenols	Repression of BCRP protein expression by NP (1 nM/48 h) in human term placental explant cultures.	(Sieppi et al., 2016)
	PFASs	Induction of BCRP mRNA expression by PFNA (45 μ M/24 h) and PFDA (45 μ M/24 h) in HepaRG cells.	(Lim et al., 2021)
OATP1B1	PFASs	Repression of OATP1B1 mRNA expression by PFOA (250 and 500 μ M/24 h) and PFOS (100 μ M/24 h) in HepaRG cells.	(Behr et al., 2020)
OATP2B1	PFASs	Repression of OATP2B1 mRNA expression by PFDA (45 μ M/24 h) in HepaRG cells.	(Lim et al., 2021)
MATE2-K	PFASs	Induction of MATE2-K mRNA expression by PFDA (45 μ M/24 h) in HepaRG cells.	(Lim et al., 2021)
NTCP	PFASs	Repression of NTCP mRNA expression by PFDA (45 μ M/24 h), PFOA (10-500 μ M/24 h) and PFOS (10-100 μ M/24 h) in HepaRG cells.	(Behr et al., 2020; Lim et al., 2021)
OST β	PFAS	Induction of OST β mRNA expression by PFOA (50-250 μ M/24 h) and PFOS (25-100 μ M/24 h) in HepaRG cells.	(Behr et al., 2020)

987 Note: Abbreviations for plastic additives and transporters are those defined in Table 1 and Table 2, respectively.



989


990

991 Fig. 1. Schematic representation of human drug transporter expression. Red and green arrows

992 correspond to ATP-binding cassette (ABC) and solute carrier (SLC) transporters, respectively.

993

994



995

996

997 Fig. 2. Schematic representation of the possible deleterious consequences of drug transporter

998 inhibition by a plastic additive, considered here as a perpetrator.


999

1000

1001

1002

1003



1004

1005

1006 Fig. 3. Relative contribution of SLC transporter-mediated uptake and transporter-unrelated
1007 passive diffusion across the plasma membrane for plastic additives. At low plastic additive
1008 concentrations (usually the case for these pollutants in response to environmental or dietary
1009 exposure), the relative contribution of transporter-mediated uptake velocity to total uptake
1010 velocity is maximal.

1011

1012

Highlights

- Human drug transporters are key actors of pharmacokinetics for drugs and pollutants
- Various plastic additives inhibit *in vitro* activities of human drug transporters
- Some plasticizers regulate human drug transporter expression in cellular systems
- Plastic additives can behave as substrates for human drug transporters
- *In vivo* relevance of plasticizers-transporters interactions remains to elucidate

Journal Pre-proof

Author statement

This paper, entitled “Interactions of human drug transporters with chemical additives present in plastics: Potential consequences for toxicokinetics and health”, by Tastet et al., has been approved by all the authors and is not being considered elsewhere. This paper does not involve the use of human subjects or animal experiments.

On behalf of all co-authors,



Olivier Fardel

Journal Pre-proof

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Journal Pre-proof