



HAL
open science

On the limit problem arising in the kinetic derivation of the Cahn-Hilliard equation

Charles Elbar, Benoît Perthame, Jakub Skrzeczkowski

► **To cite this version:**

Charles Elbar, Benoît Perthame, Jakub Skrzeczkowski. On the limit problem arising in the kinetic derivation of the Cahn-Hilliard equation. 2023. hal-04125162

HAL Id: hal-04125162

<https://hal.science/hal-04125162>

Preprint submitted on 12 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON THE LIMIT PROBLEM ARISING IN THE KINETIC DERIVATION OF THE CAHN-HILLIARD EQUATION

CHARLES ELBAR, BENOÎT PERTHAME, AND JAKUB SKRZECZKOWSKI

ABSTRACT. The non-local degenerate Cahn-Hilliard equation is derived from the Vlasov equation with long range attraction. We study the local limit as the delocalization parameter converges to 0. The difficulty arises from the degeneracy which requires compactness estimates, but all necessary a priori estimates can be obtained only on the nonlocal quantities yielding almost no information on the limiting solution itself. We introduce a novel condition on the nonlocal kernel which allows us to exploit the available nonlocal a priori estimates. The condition, satisfied by most of the kernels appearing in the applications, can be of independent interest. Our approach is flexible and systems can be treated as well.

1. INTRODUCTION

We consider the nonlocal PDE

$$\partial_t \rho_\varepsilon - \Delta \rho_\varepsilon + \operatorname{div}(\rho_\varepsilon \nabla \Delta(\rho_\varepsilon * \omega_\varepsilon * \omega_\varepsilon)) = 0 \quad (1.1)$$

arising in the kinetic derivation of the degenerate Cahn-Hilliard equation via hydrodynamic limit as in our recent paper [13] which makes the formal approach of Takata and Noguchi [20] fully rigorous. Furthermore, PDEs of the form (1.1) appear in the numerical analysis of the particle method as explained below. Concerning the notation, $\varepsilon > 0$ is a small parameter, $\rho_\varepsilon : \mathbb{R}^d \times [0, T] \rightarrow [0, \infty)$ is the solution while

$$\omega_\varepsilon(x) = \frac{1}{\varepsilon^d} \omega\left(\frac{x}{\varepsilon}\right)$$

2020 *Mathematics Subject Classification.* 35B40, 35D30, 35K25, 35K55.

Jakub Skrzeczkowski was supported by National Science Center, Poland through project no. 2019/35/N/ST1/03459.

is a usual mollification kernel (one can think of ω being smooth, nonnegative and compactly supported function but more general assumption will be presented below). The remaining question is whether one can consider the limit $\varepsilon \rightarrow 0$ in (1.1) and arrive at

$$\partial_t \rho - \Delta \rho + \operatorname{div}(\rho \nabla \Delta \rho) = 0. \quad (1.2)$$

A natural strategy for such problems are so-called commutator estimates, originally studied in the context of renormalized solutions to transport equation [12] and conservation of energy for weak solutions of Euler equation [8] or more general hyperbolic systems [1, 16]. The commutator estimates control the error resulting from the fact that the mollification operator does not commute with nonlinear functions. In our case, they cannot be applied directly, as we do not have sufficient estimates on $\{\rho_\varepsilon\}$. Indeed, both energy and entropy identities for (1.1) yield estimates on non-local quantities $\rho_\varepsilon * \omega_\varepsilon$, $\nabla \rho_\varepsilon * \omega_\varepsilon$ and $\Delta \rho_\varepsilon * \omega_\varepsilon$ (see Section 3). In particular, we do not even know that $\{\rho_\varepsilon\}$ is uniformly bounded in $L^2_{\text{loc}}((0, T) \times \mathbb{R}^d)$ (except in dimension $d = 2$). In Appendix A we show that for a fairly arbitrary kernel (up to some weak growth assumptions at infinity), the direct commutator estimates work in dimension $d = 2$. The same is true if one knows a priori that $\{\rho_\varepsilon\}$ is uniformly integrable in $L^2((0, T) \times \mathbb{R}^d)$ (or even $L^2_{\text{loc}}((0, T) \times \mathbb{R}^d)$ if ω is compactly supported).

Being not satisfied as we cannot cover the physical dimension $d = 3$, our approach is to look for an appropriate assumption on ω so that we can exploit more the aforementioned nonlocal quantities. We were inspired by the recent work on Landau equation [4, 5] where the authors consider the kernel $\omega(x) = e^{-(1+|x|^2)^{1/2}}$ which satisfies the crucial estimate

$$|x| |\nabla \omega(x)| \leq C \omega(x). \quad (1.3)$$

Such inequalities allow to transform difficult-to-estimate terms by better understood ones, for instance $\varepsilon \rho_\varepsilon * |\nabla \omega_\varepsilon|$ can be estimated pointwisely by $\rho_\varepsilon * \omega_\varepsilon$ due to nonnegativity of ρ_ε and ω_ε . Nevertheless, (1.3) is fairly restrictive - for instance, it excludes compactly supported kernels. In our setting, the following assumption turns out to be successful:

Assumption 1.1. We assume that $\omega : \mathbb{R}^d \rightarrow [0, \infty)$ is a smooth function such that $\int_{\mathbb{R}^d} \omega(x) dx = 1$ and $\omega(x) = \omega(-x)$. Moreover, there exists an integrable kernel $f : \mathbb{R}^d \rightarrow [0, \infty)$ such that for all $x \in \mathbb{R}^d$

$$(|x| + |x|^2) |\nabla \omega(x)| \leq C \omega * f(x). \quad (1.4)$$

Furthermore, we assume that ω has sufficient decay at $+\infty$:

$$\lim_{R \rightarrow \infty} \sup_{|x|=R} |x|^d \omega(x) = 0. \quad (1.5)$$

This covers the case of Gaussian $\omega(x) = \frac{1}{(2\pi)^{d/2}} e^{-|x|^2/2}$ and any nonnegative, compactly supported kernel by choosing $f = \omega$, see Lemmas 2.1 and 2.2. Moreover, we can cover the kernel $\omega(x) = e^{-(1+|x|^2)^{1/2}}$ by choosing f more carefully, see Lemma 2.3.

For the initial condition we suppose that $\rho(t=0) = \rho^0$ where ρ^0 satisfies

$$\rho^0 \geq 0, \quad \rho^0 \in L^1(\mathbb{R}^d) \cap H^1(\mathbb{R}^d), \quad \rho^0 |\log \rho^0| \in L^1(\mathbb{R}^d), \quad |x|^2 \rho^0 \in L^1(\mathbb{R}^d). \quad (1.6)$$

Our main result reads:

Theorem 1.2. *Let $\{\rho_\varepsilon\}_\varepsilon$ be a sequence of solutions to (1.1) with initial condition ρ^0 satisfying (1.6). Then, up to a subsequence not relabeled, $\rho_\varepsilon \rightarrow \rho$ in $L^p(0, T; L^1(\mathbb{R}^d))$ for all $p \in [1, \infty)$ where ρ is a weak solution of the degenerate Cahn-Hilliard equation (1.2).*

Our methods are quite flexible and they allow to study the same question for systems of the type

$$\partial_t \rho_\varepsilon^i - \Delta \rho_\varepsilon^i - \operatorname{div} \left(\rho_\varepsilon^i \nabla \Delta \sum_{j=1}^N K_\varepsilon^{i,j} * \rho_\varepsilon^j \right) = 0, \quad i = 1, \dots, N, \quad (1.7)$$

under some additional structural assumptions. This is discussed in Section 5 (see Theorem 5.1).

To conclude the introduction, let us mention that similar problems have been studied in the literature for the porous media equation. Up to our knowledge, the first result of this type was obtained by Lions and Mas-Gallic [18] for the PDE

$$\partial_t \rho_\varepsilon = \operatorname{div}(\rho_\varepsilon \nabla \rho_\varepsilon * \omega_\varepsilon * \omega_\varepsilon)$$

Then, the cases of cross-diffusion systems and general nonlinear diffusion equations has been considered in [7] and [2, 17], respectively. These problems are motivated by the numerical algorithms called particle methods. More precisely, consider N particles moving according to the system of ODEs

$$X_i'(t) = -\frac{1}{N} \sum_{j \neq i} \nabla W(X_i(t) - X_j(t)).$$

Then, the empirical measure $\mu^N(t) = \frac{1}{N} \sum_{i=1}^N \delta_{X_i(t)}$ solves in the sense of distributions

$$\partial_t \mu^N - \operatorname{div}(\mu^N \nabla \mu^N * W) = 0$$

so in the limit $N \rightarrow \infty$

$$\partial_t \mu - \operatorname{div}(\mu \nabla \mu * W) = 0.$$

If $W \xrightarrow{*} \delta_0$, we recover the porous media equation. For numerical experiments based on this method we refer to [18].

Let us also comment that (1.1) could be called the nonlocal Cahn-Hilliard equation but it should not be confused with the nonlocal effect in the following PDE

$$\partial_t \rho_\varepsilon = \operatorname{div} (\rho_\varepsilon \nabla (B_\varepsilon[\rho_\varepsilon] + F'(\rho_\varepsilon))), \quad (1.8)$$

where F is the potential and $B_\varepsilon[u]$ is a nonlocal operator approximating $-\Delta u$, i.e.

$$B_\varepsilon[u](x) = \frac{1}{\varepsilon^2} (u(x) - \omega_\varepsilon * u(x)) = \frac{1}{\varepsilon^2} \int_{\mathbb{T}^d} \omega_\varepsilon(y) (u(x) - u(x-y)) \, dy.$$

The equation was obtained by Giacomini and Lebowitz [15] as a derivation of the degenerate Cahn-Hilliard equation

$$\partial_t \rho = \operatorname{div} (\rho \nabla (-\Delta \rho + F'(\rho))), \quad (1.9)$$

proposed in [3] to model the dynamics of phase separation in binary mixtures. The question of passing to the limit from (1.8) to (1.9) was addressed only recently in [14] for a single equation and in [6] for a system. This problem is fairly different from (1.1) as energy and entropy yields strong compactness of $\{\rho_\varepsilon\}$ and $\{\nabla \rho_\varepsilon\}$ rather than their mollifications $\{\rho_\varepsilon * \omega_\varepsilon\}$ and $\{\nabla \rho_\varepsilon * \omega_\varepsilon\}$ as in the case of (1.1). We also remark that the same problem was studied in the context of the nondegenerate Cahn-Hilliard equation [9–11, 19]

$$\partial_t \rho_\varepsilon = \operatorname{div} \nabla \mu_\varepsilon, \quad \mu_\varepsilon = B_\varepsilon[\rho_\varepsilon] + F'(\rho_\varepsilon).$$

Here, one obtains immediately an estimate on $\{\nabla \mu_\varepsilon\}$ (by multiplying by μ_ε) which greatly simplifies identification of the limits. Nevertheless, we point out that in [9–11, 19] the difficulty is rather the low regularity of the potential and the kernel.

The structure of the paper is as follows. In Section 2 we show that Assumption 1.1 is satisfied by a wide class of kernels. In Section 3 we gather the a priori estimates necessary for the proof of the main result, Theorem 1.2, which is proved in Section 4. In the last Section 5, we show how the result can be extended to systems. Finally,

Appendix A is dedicated to the proof of Theorem 1.2 in dimension 2 for a broader class of kernels.

2. EXAMPLES OF KERNELS SATISFYING ASSUMPTION 1.1

Three particular classes of kernels are usually found in the literature and we show they satisfy Assumption 1.1. In fact, in all of those examples, we only need to verify condition (1.4).

Lemma 2.1. *Let $\omega : \mathbb{R}^d \rightarrow [0, \infty)$ be a smooth function such that $\int_{\mathbb{R}^d} \omega(x) dx = 1$. Suppose that ω is supported on the unit ball $\{x \in \mathbb{R}^d : |x| \leq 1\}$ and $\omega > 0$ on the interior $\{x \in \mathbb{R}^d : |x| < 1\}$. Then, ω satisfies (1.4) with $f = \omega$.*

Proof. To prove (1.4), we only need to consider $|x| \leq 1$. By smoothness and compact support of ω , there exists a constant such that $(|x| + |x|^2) |\nabla \omega(x)| \leq C$ and it remains to prove that $\inf_{|x| \leq 1} \omega * \omega(x) > 0$. For any $|x| \leq 1$, we see from the formula $\omega * \omega(x) = \int_{|y| \leq 1} \omega(x - y) \omega(y) dy$ that $\omega * \omega(x) > 0$. As any continuous function attains its infimum on a compact set, the conclusion follows. \square

Lemma 2.2. *Let $\omega(x) = \frac{1}{(2\pi)^{d/2}} e^{-|x|^2/2}$. Then, ω satisfies (1.4) with $f = \omega$.*

Proof. For Gaussians, we know that

$$\omega * \omega(x) = \frac{1}{(2\pi)^d} e^{-|x|^2/4} \int_{\mathbb{R}^d} e^{-\left|\frac{x}{\sqrt{2}} - \sqrt{2}y\right|^2/2} dy = \frac{1}{(2\pi)^{d/2}} e^{-|x|^2/4}.$$

Therefore, since the function $(|x|^2 + |x|^3) e^{-|x|^2/4}$ is globally bounded, we find

$$(|x| + |x|^2) |\nabla \omega(x)| \leq C (|x|^2 + |x|^3) e^{-|x|^2/2} \leq C e^{-|x|^2/4} = C \omega * \omega(x).$$

\square

Lemma 2.3. *Let $\omega(x) = e^{-(1+|x|^2)^{1/2}}$. Then, ω satisfies (1.4) with $f(x) = e^{-(1+|x|^2/3)^{1/2}}$.*

Proof. We need to estimate the convolution $\omega * f$ from below so we need to estimate the expression $\sqrt{1 + |x - y|^2} + \sqrt{1 + |y|^2/3}$ from above. Using $\sqrt{x} + \sqrt{y} \leq \sqrt{2(x + y)}$ and $\sqrt{x + y} \leq \sqrt{x} + \sqrt{y}$, we have

$$\begin{aligned} \sqrt{1 + |x - y|^2} + \sqrt{1 + |y|^2/3} &\leq \sqrt{2} \sqrt{2 + |x|^2 - 2x \cdot y + \frac{4}{3}|y|^2} \\ &\leq \sqrt{2} \sqrt{\frac{1 + |x|^2}{4} + \left| \frac{\sqrt{3}}{2}x - \frac{2}{\sqrt{3}}y \right|^2 + \frac{7}{4}} \\ &\leq \frac{\sqrt{2}}{2} \sqrt{1 + |x|^2} + \sqrt{2} \sqrt{\left| \frac{\sqrt{3}}{2}x - \frac{2}{\sqrt{3}}y \right|^2 + \frac{7}{4}}. \end{aligned}$$

Note that the integral $\int_{\mathbb{R}^d} e^{-\sqrt{2} \sqrt{\left| \frac{\sqrt{3}}{2}x - \frac{2}{\sqrt{3}}y \right|^2 + \frac{7}{4}}} dy$ is a constant independent of x (by a change of variables). Therefore,

$$e^{-\frac{\sqrt{2}}{2} \sqrt{1 + |x|^2}} \leq C \omega * f(x).$$

We conclude by observing that the function $(|x| + |x|^2) e^{-(1 - \frac{\sqrt{2}}{2}) \sqrt{1 + |x|^2}}$ is globally bounded

$$\begin{aligned} (|x| + |x|^2) |\nabla \omega(x)| &\leq C (|x| + |x|^2) e^{-\sqrt{1 + |x|^2}} = \\ &= C (|x| + |x|^2) e^{-(1 - \frac{\sqrt{2}}{2}) \sqrt{1 + |x|^2}} e^{-\frac{\sqrt{2}}{2} \sqrt{1 + |x|^2}} \leq C e^{-\frac{\sqrt{2}}{2} \sqrt{1 + |x|^2}} \leq C \omega * f(x). \end{aligned}$$

□

3. UNIFORM ESTIMATES FOR (1.1) AND COMPACTNESS

The first immediate estimate is the conservation of mass. Integrating the equation in space we obtain an $L^\infty(0, T; L^1(\mathbb{R}^d))$ control on the solution. Moreover, the nonlocal equation (1.1) comes with an energy/entropy structure. Defining

$$E_\varepsilon[\rho] = \int_{\mathbb{R}^d} \frac{|\nabla \rho * \omega_\varepsilon|^2}{2} dx, \quad \Phi[\rho] = \int_{\mathbb{R}^d} \rho \log(\rho) dx, \quad (3.1)$$

we obtain the dissipation equalities:

$$\frac{dE_\varepsilon[\rho]}{dt} + \int_{\mathbb{R}^d} |\Delta \rho * \omega_\varepsilon|^2 dx + \int_{\Omega} \rho |\nabla \Delta \rho * \omega_\varepsilon * \omega_\varepsilon|^2 dx = 0, \quad (3.2)$$

$$\frac{d\Phi[\rho]}{dt} + \int_{\mathbb{R}^d} \frac{|\nabla \rho|^2}{\rho} dx + \int_{\mathbb{R}^d} |\Delta \rho * \omega_\varepsilon|^2 dx = 0. \quad (3.3)$$

Of course one has to be careful with the entropy equality, as $\rho \log(\rho)$ can be negative when ρ is small and one needs to show that its negative part is integrable.

Proposition 3.1. *Suppose the initial condition ρ^0 satisfies (1.6). Then, there exists a unique nonnegative weak solution to (1.1) satisfying the following bounds, uniformly with respect to ε :*

- (A) $\{\rho_\varepsilon\}_\varepsilon \in L^\infty(0, T; L^1(\mathbb{R}^d) \cap L \log L(\mathbb{R}^d))$,
- (B) $\{\partial_t \rho_\varepsilon\}_\varepsilon \in L^2(0, T; H^{-k}(\mathbb{R}^d))$ for some k ,
- (C) $\{\sqrt{\rho_\varepsilon} \nabla \Delta \rho_\varepsilon * \omega_\varepsilon * \omega_\varepsilon\}_\varepsilon \in L^2((0, T) \times \mathbb{R}^d)$,
- (D) $\{\rho_\varepsilon * \omega_\varepsilon\}_\varepsilon \in L^\infty(0, T; H^1(\mathbb{R}^d)) \cap L^2(0, T; H^2(\mathbb{R}^d))$,
- (E) $\{|x|^2 \rho_\varepsilon\}_\varepsilon \in L^\infty(0, T; L^1(\mathbb{R}^d))$,
- (F) $\{\nabla \sqrt{\rho_\varepsilon}\}_\varepsilon \in L^2(0, T; L^2(\mathbb{R}^d))$,
- (G) $\{\nabla \rho_\varepsilon\}_\varepsilon \in L^2(0, T; L^1(\mathbb{R}^d))$.

Moreover, we can extract a subsequence such that

$$\rho_\varepsilon \rightarrow \rho \text{ strongly in } L^p(0, T; L^1(\mathbb{R}^d)), \quad p < \infty, \quad (3.4)$$

$$\rho_\varepsilon * \omega_\varepsilon \rightharpoonup \rho \text{ weakly in } L^\infty(0, T; H^1(\mathbb{R}^d)) \cap L^2(0, T; H^2(\mathbb{R}^d)), \quad (3.5)$$

$$\rho_\varepsilon * \omega_\varepsilon \rightarrow \rho \text{ a.e. and strongly in } L^2(0, T; H_{loc}^1(\mathbb{R}^d)). \quad (3.6)$$

Proof of Proposition 3.1. The existence and uniqueness of solutions is a classical matter as (1.1) is an advection-diffusion equation with smooth advection (as ω_ε is smooth). The L^1 bound in (A) is a consequence of mass conservation. (C) follows directly from (3.2). Estimate (B) is a consequence of (A), (C), Equation (1.1) and splitting

$$\rho_\varepsilon \nabla \Delta \rho_\varepsilon * \omega_\varepsilon * \omega_\varepsilon = \sqrt{\rho_\varepsilon} \sqrt{\rho_\varepsilon} \nabla \Delta \rho_\varepsilon * \omega_\varepsilon * \omega_\varepsilon.$$

To prove (D), we first deduce from (3.2) bounds on $\nabla \rho_\varepsilon * \omega_\varepsilon$ and $\Delta \rho_\varepsilon * \omega_\varepsilon$. Then, the $L^\infty(0, T; L^2(\mathbb{R}^d))$ bound on $\{\rho_\varepsilon * \omega_\varepsilon\}_\varepsilon$ follows from bounds on $\{\rho_\varepsilon * \omega_\varepsilon\}_\varepsilon$ in $L^\infty(0, T; L^1(\mathbb{R}^d))$, $\{\nabla \rho_\varepsilon * \omega_\varepsilon\}_\varepsilon$ in $L^\infty(0, T; L^2(\mathbb{R}^d))$ and from the Gagliardo-Nirenberg inequality). The estimate in $L^2(0, T; H^2(\mathbb{R}^d))$ is a consequence of the second-order regularizing property of the operator Δ on the whole space.

To see (E), we multiply Equation (1.1) by $|x|^2$ and obtain after integration by parts

$$\frac{d}{dt} \int_{\mathbb{R}^d} |x|^2 \rho_\varepsilon dx = 2 \int_{\mathbb{R}^d} \rho_\varepsilon dx + 2 \int_{\mathbb{R}^d} \sqrt{\rho_\varepsilon} x \cdot \sqrt{\rho_\varepsilon} \nabla \Delta \rho_\varepsilon * \omega_\varepsilon * \omega_\varepsilon dx.$$

Using (A), (C), the Cauchy-Schwarz inequality and the Gronwall lemma, we obtain (E).

It remains to prove the second part of estimate (A) namely the $L \log(L)$ bound on ρ_ε . A small difficulty is that the negative part of $\rho_\varepsilon \log(\rho_\varepsilon)$ might not be integrable on the whole space. Nevertheless, as in [17], one can prove that $\rho_\varepsilon |\log \rho_\varepsilon|_-$ is uniformly bounded in $L^\infty(0, T; L^1(\mathbb{R}^d))$ by splitting \mathbb{R}^d for $\{x : |\rho_\varepsilon| \leq e^{-|x|^2}\}$, $\{x : |\rho_\varepsilon| > e^{-|x|^2}\}$ and applying tail estimate (E). Hence, we can use (3.3) and deduce Estimate (F). Estimate (G) follows from (A) and (F) by writing $\nabla \rho_\varepsilon = \frac{\nabla \rho_\varepsilon}{\sqrt{\rho_\varepsilon}} \sqrt{\rho_\varepsilon}$.

The convergences (3.4)–(3.6) are a consequence of the Lions-Aubin lemma and the Banach-Alaoglu theorem where convergence (3.4) has been upgraded from a local to a global one by the tail estimate (E). \square

4. THE PROOF OF THE MAIN RESULT

We only need to study the term $\int_0^T \int_{\mathbb{R}^d} \nabla \varphi \rho_\varepsilon \nabla \Delta(\rho_\varepsilon * \omega_\varepsilon * \omega_\varepsilon) dx dt$, for test functions $\varphi \in C_c^\infty([0, T] \times \mathbb{R}^d)$. Using the properties of the mollifiers

$$\int_0^T \int_{\mathbb{R}^d} \nabla \varphi \rho_\varepsilon \nabla \Delta(\rho_\varepsilon * \omega_\varepsilon * \omega_\varepsilon) dx dt = - \int_0^T \int_{\mathbb{R}^d} (\nabla \varphi \rho_\varepsilon) * \nabla \omega_\varepsilon \Delta(\rho_\varepsilon * \omega_\varepsilon) dx dt.$$

Thanks to the weak convergence (3.6), we only need to prove the following strong convergence result

$$(\nabla\varphi\rho_\varepsilon) * \nabla\omega_\varepsilon \rightarrow \nabla\varphi \cdot \nabla\rho + \Delta\varphi\rho = \operatorname{div}(\nabla\varphi\rho) \text{ strongly in } L^2((0, T) \times \mathbb{R}^d). \quad (4.1)$$

We write $\nabla\varphi(y) = \nabla\varphi(y) - \nabla\varphi(x) + \nabla\varphi(x)$ which results in two terms:

$$\begin{aligned} (\nabla\varphi\rho_\varepsilon) * \nabla\omega_\varepsilon(x) &= \int_{\mathbb{R}^d} \nabla\varphi(y)\rho_\varepsilon(y) \cdot \nabla\omega_\varepsilon(x-y) \, dy = \nabla\varphi(x) \cdot \nabla\rho_\varepsilon * \omega_\varepsilon + \\ &\quad + \int_{\mathbb{R}^d} (\nabla\varphi(y) - \nabla\varphi(x))\rho_\varepsilon(y) \cdot \nabla\omega_\varepsilon(x-y) \, dy =: I + J. \end{aligned}$$

According to (3.6), the term I converges strongly in $L^2((0, T) \times \mathbb{R}^d)$ (note that φ is compactly supported). The rest of the proof is devoted to the analysis of the term J .

By Taylor's expansion

$$\nabla\varphi(y) - \nabla\varphi(x) = D^2\varphi(x)(y-x) + R(x, y)$$

where the term R satisfies $|R| \leq C|y-x|^2$. We split $J = J_1 + J_2$, where

$$J_1 = \int_{\mathbb{R}^d} D^2\varphi(x)(y-x)\rho_\varepsilon(y)\nabla\omega_\varepsilon(x-y) \, dy, \quad J_2 = \int_{\mathbb{R}^d} R(x, y)\rho_\varepsilon(y)\nabla\omega_\varepsilon(x-y) \, dy.$$

Term J_1 . We prove that $\lim_{\varepsilon \rightarrow 0} J_1 = \Delta\varphi\rho$ in $L^2((0, T) \times \mathbb{R}^d)$. Since φ is compactly supported, it is sufficient to prove that

$$\int_{\mathbb{R}^d} \rho_\varepsilon(y)(x_i - y_i)\partial_j\omega_\varepsilon(x-y) \, dy \rightarrow -\rho(x)\delta_{i,j} \quad \text{in } L^2_{\text{loc}}((0, T) \times \mathbb{R}^d). \quad (4.2)$$

Assertion (4.2) will be obtained by proving convergence in $L^1((0, T) \times \mathbb{R}^d)$ and uniform boundedness in $L^p((0, T) \times \mathbb{R}^d)$ for some $p > 2$. Concerning the convergence in $L^1((0, T) \times \mathbb{R}^d)$, we first change variables

$$\int_{\mathbb{R}^d} \rho_\varepsilon(y)(x_i - y_i)\partial_j\omega_\varepsilon(x-y) \, dy = \int_{\mathbb{R}^d} \rho_\varepsilon(x - \varepsilon z)z_i\partial_j\omega(z) \, dz$$

We can estimate in $L^1((0, T) \times \mathbb{R}^d)$ the difference

$$\begin{aligned} & \int_0^T \int_{\mathbb{R}^d} \left| \int_{\mathbb{R}^d} (\rho_\varepsilon(x - \varepsilon z) - \rho_\varepsilon(x)) z_i \partial_j \omega(z) dz \right| dx dt \leq \\ & \leq \int_0^T \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_0^1 \varepsilon |\nabla \rho_\varepsilon(x - \varepsilon s z)| |z|^2 |\nabla \omega(z)| ds dz dx dt \leq \varepsilon \|\nabla \rho_\varepsilon\|_{L^1_{t,x}} \| |z|^2 \nabla \omega \|_{L^1_x}, \end{aligned}$$

where integrability of $|z|^2 \nabla \omega(z)$ is a consequence of assumption (1.4):

$$\int_{\mathbb{R}^d} |z|^2 |\nabla \omega(z)| dz \leq C \int_{\mathbb{R}^d} \omega * f(z) dz = C \|\omega\|_{L^1} \|f\|_{L^1} \leq C.$$

Therefore, it is sufficient to study the term $\rho_\varepsilon(x) \int_{\mathbb{R}^d} z_i \partial_j \omega(z) dz$ which equals $-\rho_\varepsilon(x)$ because

$$\int_{\mathbb{R}^d} z_i \partial_j \omega(z) dz = -\delta_{i,j} \int_{\mathbb{R}^d} \omega(z) dz = -\delta_{i,j},$$

where the boundary term vanishes thanks to (1.5). The conclusion follows because ρ_ε is strongly convergent in $L^1((0, T) \times \mathbb{R}^d)$, cf. (3.4).

Concerning the uniform boundedness in $L^p((0, T) \times \mathbb{R}^d)$ with $p > 2$, by nonnegativity of ρ_ε , definition of $\omega_\varepsilon(x) = \frac{1}{\varepsilon^d} \omega\left(\frac{x}{\varepsilon}\right)$ and assumption (1.4),

$$\begin{aligned} \left| \int_{\mathbb{R}^d} \rho_\varepsilon(y) (x_i - y_i) \partial_j \omega_\varepsilon(x - y) dy \right| &= \left| \int_{\mathbb{R}^d} \rho_\varepsilon(y) \frac{(x_i - y_i)}{\varepsilon^{d+1}} \partial_j \omega\left(\frac{x - y}{\varepsilon}\right) dy \right| \leq \\ &\leq C \int_{\mathbb{R}^d} \rho_\varepsilon(y) \frac{1}{\varepsilon^d} \omega * f\left(\frac{x - y}{\varepsilon}\right) dy. \end{aligned} \quad (4.3)$$

A change of variables shows that

$$\begin{aligned} \frac{1}{\varepsilon^d} \omega * f\left(\frac{x - y}{\varepsilon}\right) &= \frac{1}{\varepsilon^d} \int_{\mathbb{R}^d} \omega(z) f\left(\frac{x - y}{\varepsilon} - z\right) dz = \\ &= \frac{1}{\varepsilon^{2d}} \int_{\mathbb{R}^d} \omega\left(\frac{z}{\varepsilon}\right) f\left(\frac{x - y - z}{\varepsilon}\right) dz = \omega_\varepsilon * f_\varepsilon(x - y), \end{aligned} \quad (4.4)$$

where $f_\varepsilon(x) := \frac{1}{\varepsilon^d} f\left(\frac{x}{\varepsilon}\right)$. Due to (4.3),

$$\left| \int_{\mathbb{R}^d} \rho_\varepsilon(y) (x_i - y_i) \partial_j \omega_\varepsilon(x - y) dy \right| \leq C \rho_\varepsilon * \omega_\varepsilon * f_\varepsilon(x).$$

Note that by the Gagliardo-Nirenberg inequality and uniform bound in $L^\infty(0, T; H^1(\mathbb{R}^d))$, $\{\rho_\varepsilon * \omega_\varepsilon\}_\varepsilon$ is bounded in $L^\infty(0, T; L^{\frac{2d}{d-2}}(\mathbb{R}^d))$ where $\frac{2d}{d-2} > 2$. The same is true for $\{\rho_\varepsilon * \omega_\varepsilon * f_\varepsilon\}_\varepsilon$ by the Young convolutional inequality. The conclusion follows.

Term J_2 . We prove that $\lim_{\varepsilon \rightarrow 0} J_2 = 0$. By $|R| \leq C|y - x|^2$, it is sufficient to prove

$$\int_{\mathbb{R}^d} \rho_\varepsilon(y) |x - y|^2 |\nabla \omega_\varepsilon(x - y)| dy \rightarrow 0 \quad \text{in } L^2((0, T) \times \mathbb{R}^d).$$

Again, we want to use assumption (1.4). By definition of ω_ε :

$$\begin{aligned} \int_{\mathbb{R}^d} \rho_\varepsilon(y) |x - y|^2 |\nabla \omega_\varepsilon(x - y)| dy &= \varepsilon \int_{\mathbb{R}^d} \rho_\varepsilon(y) \frac{1}{\varepsilon^d} \left| \frac{x - y}{\varepsilon} \right|^2 \left| \nabla \omega \left(\frac{x - y}{\varepsilon} \right) \right| dy \leq \\ &\leq \varepsilon \int_{\mathbb{R}^d} \rho_\varepsilon(y) \frac{1}{\varepsilon^d} \omega * f \left(\frac{x - y}{\varepsilon} \right) = \varepsilon \rho_\varepsilon * \omega_\varepsilon * f_\varepsilon(x), \end{aligned}$$

where in the last line we applied (4.4). By the Young convolutional inequality, $\rho_\varepsilon * \omega_\varepsilon * f_\varepsilon$ on the (RHS) is bounded in $L^2((0, T) \times \mathbb{R}^d)$ so the conclusion follows.

5. EXTENSION TO SYSTEMS

Motivated by [17], we consider the system of N equations

$$\partial_t \rho_\varepsilon^i - \Delta \rho_\varepsilon^i + \operatorname{div} \left(\rho_\varepsilon^i \nabla \Delta \sum_{j=1}^N K_\varepsilon^{i,j} * \rho_\varepsilon^j \right) = 0, \quad (5.1)$$

where $1 \leq i \leq N$ and the kernels $K_\varepsilon^{i,j}$ are of the form

$$K_\varepsilon^{i,j} = \sum_{k=1}^N \alpha^{i,k} \alpha^{j,k} \omega_\varepsilon^i * \omega_\varepsilon^j,$$

where the ω^i are kernels satisfying Assumption 1.1. The coefficients $\{\alpha^{i,k}\}$ form a matrix A and we assume it is invertible. Under these assumptions, for any set of functions η_1, \dots, η_N :

$$\sum_{i,j=1}^N \int_{\mathbb{R}^d} \eta^i K_\varepsilon^{i,j} * \eta^j = \int_{\mathbb{R}^d} \sum_{k=1}^N \left(\sum_{i=1}^N \alpha^{i,k} \eta^i * \omega_\varepsilon^i \right)^2$$

so that

$$\tilde{C} \|A^{-1}\|^2 \int_{\mathbb{R}^d} \sum_{i=1}^N (\eta^i * \omega_\varepsilon^i)^2 \geq \sum_{i,j=1}^N \int_{\mathbb{R}^d} \eta^i K_\varepsilon^{i,j} * \eta^j \geq C \|A^{-1}\|^2 \int_{\mathbb{R}^d} \sum_{i=1}^N (\eta^i * \omega_\varepsilon^i)^2. \quad (5.2)$$

Theorem 5.1. *Let $\{\rho_\varepsilon^i\}_\varepsilon$ be a sequence of solutions to (5.1) with initial condition $\rho^{0,i}$ satisfying (1.6). Then, for $i = 1, \dots, N$, and for a subsequence not relabeled, $\rho_\varepsilon^i \rightarrow \rho^i$ in $L^p(0, T; L^1(\mathbb{R}^d))$ for all $p \in [1, \infty)$ where ρ^i is a weak solution of*

$$\partial_t \rho^i - \Delta \rho^i + \operatorname{div} \left(\rho^i \nabla \Delta \sum_{j=1}^N K^{i,j} \rho^j \right) = 0, \quad K^{i,j} = \sum_{k=1}^N \alpha^{i,k} \alpha^{j,k}.$$

We first extend the uniform bounds in Proposition 3.1 to the case of system (5.1).

Proposition 5.2. *Suppose that for all $i = 1, \dots, N$, the initial conditions $\rho^{0,i}$ satisfy (1.6). Then, the nonnegative solution to (5.1) satisfies the following bounds, uniformly with respect to ε :*

- (A) $\{\rho_\varepsilon^i\}_\varepsilon \in L^\infty(0, T; L^1(\mathbb{R}^d) \cap L \log L(\mathbb{R}^d))$,
- (B) $\{\partial_t \rho_\varepsilon^i\}_\varepsilon \in L^2(0, T; H^{-k}(\mathbb{R}^d))$ for some k ,
- (C) $\{\sqrt{\rho_\varepsilon^i} \nabla \Delta \sum_{j=1}^N K_\varepsilon^{i,j} * \rho_\varepsilon^j\}_\varepsilon \in L^2((0, T) \times \mathbb{R}^d)$,
- (D) $\{\rho_\varepsilon^i * \omega_\varepsilon^i\}_\varepsilon \in L^\infty(0, T; H^1(\mathbb{R}^d)) \cap L^2(0, T; H^2(\mathbb{R}^d))$,
- (E) $\{|x|^2 \rho_\varepsilon^i\}_\varepsilon \in L^\infty(0, T; L^1(\mathbb{R}^d))$,
- (F) $\{\nabla \sqrt{\rho_\varepsilon^i}\}_\varepsilon \in L^2(0, T; L^2(\mathbb{R}^d))$,
- (G) $\{\nabla \rho_\varepsilon^i\}_\varepsilon \in L^2(0, T; L^1(\mathbb{R}^d))$,

Moreover, we can extract a subsequence such that for all $i = 1, \dots, N$

$$\rho_\varepsilon^i \rightarrow \rho^i \text{ strongly in } L^p(0, T; L^1(\mathbb{R}^d)), \quad p < \infty \quad (5.3)$$

$$\rho_\varepsilon^i * \omega_\varepsilon^i \rightharpoonup \rho^i \text{ weakly in } L^\infty(0, T; H^1(\mathbb{R}^d)) \cap L^2(0, T; H^2(\mathbb{R}^d)), \quad (5.4)$$

$$\rho_\varepsilon^i * \omega_\varepsilon^i \rightarrow \rho^i \text{ a.e. and strongly in } L^2(0, T; H_{loc}^1(\mathbb{R}^d)). \quad (5.5)$$

Proof. The proof is almost the same as the proof of Proposition 3.1. The only difficulty is to obtain the energy and entropy identities corresponding to (3.2) and (3.3), respectively.

Concerning the energy, we multiply (5.1) with $\Delta \sum_{j=1}^N K^{i,j} * \rho_\varepsilon^j$, integrating in space and summing up for $i = 1, \dots, N$ yields

$$\begin{aligned} & \frac{C \|A^{-1}\|^2}{2} \sum_{i=1}^N \int_{\mathbb{R}^d} |\nabla \rho_\varepsilon^i * \omega_\varepsilon^i(t, x)|^2 dx + C \|A^{-1}\|^2 \sum_{i=1}^N \int_0^t \int_{\mathbb{R}^d} |\Delta \rho_\varepsilon^i * \omega^i|^2 dx ds + \\ & + \sum_{i=1}^N \int_0^t \int_{\mathbb{R}^d} \rho_\varepsilon^i \left| \nabla \Delta \sum_{j=1}^N K_\varepsilon^{i,j} * \rho_\varepsilon^j \right|^2 dx ds \leq \frac{\tilde{C} \|A\|^2}{2} \sum_{i=1}^N \int_{\mathbb{R}^d} |\nabla \rho^{i,0}|^2 dx. \end{aligned}$$

This identity implies (C) and (D). Estimate (B) follows from the PDE (5.1) and (C). Thanks to (C), we also deduce (E).

Concerning the entropy, we multiply (5.1) with $\log \rho_\varepsilon^i$, integrate in space and sum up to obtain

$$\sum_{i=1}^N \partial_t \int_{\mathbb{R}^d} \rho_\varepsilon^i (\log \rho_\varepsilon^i - 1) dx + \sum_{i=1}^N \int_{\mathbb{R}^d} \frac{|\nabla \rho_\varepsilon^i|^2}{\rho_\varepsilon^i} dx + \sum_{i,j=1}^N \int_{\mathbb{R}^d} K^{i,j} * \Delta \rho_\varepsilon^j \Delta \rho_\varepsilon^i dx = 0.$$

Applying (5.2) with $\eta_i = \Delta \rho_\varepsilon^i$, we deduce

$$\sum_{i=1}^N \partial_t \int_{\mathbb{R}^d} \rho_\varepsilon^i (\log \rho_\varepsilon^i - 1) dx + \sum_{i=1}^N \int_{\mathbb{R}^d} \frac{|\nabla \rho_\varepsilon^i|^2}{\rho_\varepsilon^i} dx + C \|A^{-1}\| \sum_{i=1}^N \int_{\mathbb{R}^d} |\Delta \rho_\varepsilon^i * \omega_\varepsilon^i|^2 dx \leq 0.$$

As in the proof of Proposition 3.1, one may check that $\rho_\varepsilon^i |\log \rho_\varepsilon^i|_-$ is uniformly bounded in $L^\infty(0, T; L^1(\mathbb{R}^d))$ which implies (A), (F) and (G). The convergences (5.3)–(5.5) easily follow from the estimates. \square

Proof of Theorem 5.1. By linearity, we only need to explain how to pass to the limit in the term

$$\int_0^T \int_{\mathbb{R}^d} \nabla \varphi \rho_\varepsilon^i \nabla \Delta (\rho_\varepsilon^j * \omega_\varepsilon^j * \omega_\varepsilon^i) dx dt = - \int_0^T \int_{\mathbb{R}^d} (\nabla \varphi \rho_\varepsilon^i) * \nabla \omega_\varepsilon^i \Delta (\rho_\varepsilon^j * \omega_\varepsilon^j) dx dt.$$

However, in the proof of Theorem 1.2, we proved that

$$(\nabla\varphi\rho_\varepsilon^i) * \nabla\omega_\varepsilon^i \rightarrow \nabla\varphi \cdot \nabla\rho^i + \Delta\varphi\rho^i \text{ strongly in } L^2((0, T) \times \mathbb{R}^d),$$

see (4.1). Thanks to the weak convergence $\Delta(\rho_\varepsilon^j * \omega_\varepsilon^j) \rightharpoonup \Delta\rho_\varepsilon^j$ in (5.5), we conclude the proof. \square

APPENDIX A. PROOF OF THE CONVERGENCE FOR GENERAL KERNELS AND $d = 2$

In dimension $d = 2$ another proof of the main result uses weaker assumptions, namely

$$d = 2, \quad y\omega(y) \in L^1(\mathbb{R}^d), \quad y\nabla\omega(y) \in L^2(\mathbb{R}^d).$$

As in the main proof, we only need to study term $\int_0^T \int_{\mathbb{R}^d} \nabla\varphi\rho_\varepsilon \nabla\Delta(\rho_\varepsilon * \omega_\varepsilon * \omega_\varepsilon) dx dt$, where $\varphi \in C_c^\infty([0, T] \times \mathbb{R}^d)$. Integrating by parts

$$\int_0^T \int_{\mathbb{R}^d} \nabla\varphi\rho_\varepsilon \nabla\Delta(\rho_\varepsilon * \omega_\varepsilon * \omega_\varepsilon) dx dt = - \int_0^T \int_{\mathbb{R}^d} \operatorname{div}(\nabla\varphi\rho_\varepsilon) * \omega_\varepsilon \Delta(\rho_\varepsilon * \omega_\varepsilon) dx dt$$

According to the a priori estimate (D), we need to prove that $\operatorname{div}(\nabla\varphi\rho_\varepsilon) * \omega_\varepsilon$ converges strongly in $L^2((0, T) \times \mathbb{R}^d)$. We introduce the truncation operator

$$T_M(\rho) = \begin{cases} \rho & \text{if } \rho \leq M, \\ M & \text{if } \rho > M, \end{cases}$$

so that splitting $\rho_\varepsilon = \rho_\varepsilon - T_M(\rho_\varepsilon) + T_M(\rho_\varepsilon)$ we have

$$\begin{aligned} \operatorname{div}(\nabla\varphi\rho_\varepsilon) * \omega_\varepsilon &= \operatorname{div}(\nabla\varphi T_M(\rho_\varepsilon)) * \omega_\varepsilon + \operatorname{div}(\nabla\varphi(\rho_\varepsilon - T_M(\rho_\varepsilon))) * \omega_\varepsilon \\ &= (\Delta\varphi T_M(\rho_\varepsilon)) * \omega_\varepsilon + (\nabla\varphi \cdot \nabla\rho_\varepsilon \mathbf{1}_{\rho_\varepsilon \leq M}) * \omega_\varepsilon + (\nabla\varphi(\rho_\varepsilon - T_M(\rho_\varepsilon))) * \nabla\omega_\varepsilon \\ &=: I_1 + I_2 + I_3. \end{aligned}$$

The parameter M will be chosen later in terms of ε so that $M \rightarrow \infty$ as $\varepsilon \rightarrow 0$.

Term I_1 . We write

$$\begin{aligned} I_1(t, x) &= \int_{\mathbb{R}^d} (\Delta\varphi(x-y) - \Delta\varphi(x)) T_M(\rho_\varepsilon)(x-y) \omega_\varepsilon(y) \, dy + \Delta\varphi(T_M(\rho_\varepsilon)) * \omega_\varepsilon \\ &= I_1^A + I_1^B. \end{aligned}$$

As $|\Delta\varphi(x-y) - \Delta\varphi(x)| \leq C|y|$, we can estimate

$$\|I_1^A\|_{L_{t,x}^2} \leq \sqrt{M} \|\sqrt{\rho_\varepsilon}\|_{L_{t,x}^2} \| |y| \omega_\varepsilon(y) \|_{L^1} \leq \varepsilon \sqrt{M} \|\sqrt{\rho_\varepsilon}\|_{L_{t,x}^2} \| |y| \omega(y) \|_{L^1}$$

so that $\|I_1^A\|_{L_{t,x}^2} \leq C\varepsilon\sqrt{M}$. Furthermore, note that the term I_1^B is compact in $L^2((0, T) \times \mathbb{R}^d)$ whenever $M \rightarrow \infty$, $\varepsilon \rightarrow 0$. To see this, first note that it is sufficient to establish local compactness as φ is compactly supported. The latter can be proved by the Vitali theorem: we have convergence in measure (even in $L_{\text{loc}}^1((0, T) \times \mathbb{R}^d)$) of $T_M(\rho_\varepsilon) * \omega_\varepsilon$ and uniform integrability thanks to the pointwise estimate

$$0 \leq T_M(\rho_\varepsilon) * \omega_\varepsilon \leq \rho_\varepsilon * \omega_\varepsilon$$

since $\rho_\varepsilon * \omega_\varepsilon$ is compact in $L_{\text{loc}}^2((0, T) \times \mathbb{R}^d)$. We conclude that

$$I_1^B \rightarrow \Delta\varphi \rho \text{ in } L^2((0, T) \times \mathbb{R}^d) \quad \text{when } \varepsilon \rightarrow 0, M \rightarrow \infty. \quad (\text{A.1})$$

Term I_2 . We have

$$\begin{aligned} I_2(t, x) &= \int_{\mathbb{R}^d} (\nabla\varphi(x-y) - \nabla\varphi(x)) \nabla\rho_\varepsilon(x-y) \mathbf{1}_{\rho_\varepsilon(x-y) \leq M} \omega_\varepsilon(y) \, dy \\ &\quad + \nabla\varphi \cdot (\nabla\rho_\varepsilon \mathbf{1}_{\rho_\varepsilon \leq M}) * \omega_\varepsilon =: I_2^A + I_2^B. \end{aligned}$$

As $(\nabla\varphi(x-y) - \nabla\varphi(x)) \leq C|y|$ and $|\nabla\rho_\varepsilon(x-y)| \mathbf{1}_{\rho_\varepsilon(x-y) \leq M} \leq \sqrt{M} \frac{|\nabla\rho_\varepsilon(x-y)|}{\sqrt{\rho_\varepsilon(x-y)}}$, we can estimate the term I_2^A as follows

$$\|I_2^A\|_{L_{t,x}^2} \leq \sqrt{M} \left\| \frac{\nabla\rho_\varepsilon}{\sqrt{\rho_\varepsilon}} \right\|_{L_{t,x}^2} \| |y| \omega_\varepsilon(y) \|_{L^1} \leq \varepsilon \sqrt{M} \left\| \frac{\nabla\rho_\varepsilon}{\sqrt{\rho_\varepsilon}} \right\|_{L_{t,x}^2} \| |y| \omega(y) \|_{L^1}$$

so that $|I_2^A| \leq C\varepsilon\sqrt{M}$ according to estimate (F).

Term I_3 . We write

$$\begin{aligned} I_3(t, x) &= \int_{\mathbb{R}^d} (\nabla\varphi(x-y) - \nabla\varphi(x)) (\rho_\varepsilon(x-y) - T_M(\rho_\varepsilon(x-y))) \nabla\omega_\varepsilon(y) \, dy \\ &\quad + \nabla\varphi(\rho_\varepsilon - T_M(\rho_\varepsilon)) * \nabla\omega_\varepsilon = I_3^A + I_3^B. \end{aligned}$$

We observe that $|\nabla\varphi(x-y) - \nabla\varphi(x)| \leq C|y|$ and $|\rho_\varepsilon - T_M(\rho_\varepsilon)| \leq 2\rho_\varepsilon \mathbf{1}_{\rho_\varepsilon \geq M}$ so the term I_3^A can be estimated as

$$\|I_3^A\|_{L^2_{t,x}} \leq C \|(\rho_\varepsilon \mathbf{1}_{\rho_\varepsilon \geq M}) * (|y| |\nabla\omega_\varepsilon(y)|)\|_{L^2_{t,x}}.$$

By the Gagliardo-Nirenberg-Sobolev inequality, we get that $\{\rho_\varepsilon\}$ is uniformly bounded in $L^2((0, T) \times \mathbb{R}^d)$. Therefore,

$$\|I_3^A\|_{L^2_{t,x}} \leq C \|\rho_\varepsilon \mathbf{1}_{\rho_\varepsilon \geq M}\|_{L^2_t L^1_x} \| |y| |\nabla\omega_\varepsilon(y)| \|_{L^2} \leq C \|\rho_\varepsilon\|_{L^2_{t,x}} \|\mathbf{1}_{\rho_\varepsilon \geq M}\|_{L^\infty_t L^2_x} \| |y| |\nabla\omega_\varepsilon(y)| \|_{L^2}.$$

It remains to estimate $\|\mathbf{1}_{\rho_\varepsilon \geq M}\|_{L^\infty_t L^2_x}$ and $\| |y| |\nabla\omega_\varepsilon(y)| \|_{L^2}$. We have

$$\|\mathbf{1}_{\rho_\varepsilon \geq M}\|_{L^\infty_t L^2_x} \leq \sup_{t \in (0, T)} \left(\int_{\mathbb{R}^d} \frac{\rho_\varepsilon \log \rho_\varepsilon}{M \log(M)} \, dx \right)^{1/2} \leq \frac{C}{M^{1/2} \log^{1/2} M},$$

$$\| |y| |\nabla\omega_\varepsilon(y)| \|_{L^2}^2 \leq \int_{\mathbb{R}^d} \frac{1}{\varepsilon^{2d+2}} |y|^2 \left| \nabla\omega\left(\frac{y}{\varepsilon}\right) \right|^2 \, dy = \frac{1}{\varepsilon^d} \int_{\mathbb{R}^d} |y|^2 |\nabla\omega(y)|^2 \, dy \leq \frac{C}{\varepsilon^2},$$

since $d = 2$ and using mass conservation (A). We conclude that

$$\|I_3^A\|_{L^2_{t,x}} \leq \frac{C}{\varepsilon M^{1/2} \log^{1/2} M}.$$

The conclusion. Note that the terms I_2^B and I_3^B combine to

$$I_2^B + I_3^B = \nabla\varphi \nabla(\rho_\varepsilon * \omega_\varepsilon)$$

which is compact in $L^2((0, T) \times \mathbb{R}^d)$ and converges to $\nabla\varphi \nabla\rho$. Therefore,

$$\operatorname{div}(\nabla\varphi \rho_\varepsilon) * \omega_\varepsilon = \Delta\varphi(T_M(\rho_\varepsilon)) * \omega_\varepsilon + \nabla\varphi \nabla(\rho_\varepsilon * \omega_\varepsilon) + R, \quad (\text{A.2})$$

where the first two terms are compact in $L^2((0, T) \times \mathbb{R}^d)$ (see also (A.1)) while

$$\|R\|_{L^2_{t,x}} \leq C \varepsilon \sqrt{M} + \frac{C}{\varepsilon \sqrt{M} \log^{1/2}(M)}.$$

The conclusion follows by choosing M such that $\varepsilon^2 M \log^{1/2} M = 1$.

Remark A.1. In arbitrary dimension d , if we knew that $\{\rho_\varepsilon\}_\varepsilon$ is uniformly integrable in $L^2((0, T) \times \mathbb{R}^d)$, i.e.

$$\lim_{\varepsilon \rightarrow 0} \|\rho_\varepsilon \mathbb{1}_{\rho_\varepsilon > \frac{1}{\varepsilon}}\|_{L^2_{t,x}} = 0, \quad (\text{A.3})$$

we could conclude in an easier way. Indeed, assuming that $y \nabla \omega(y) \in L^1(\mathbb{R}^d)$, one can estimate

$$\|I_3^A\|_{L^2_{t,x}} \leq \|\psi \rho_\varepsilon \mathbb{1}_{\rho_\varepsilon \geq M}\|_{L^2_{t,x}} \| |y| \nabla \omega_\varepsilon(y) \|_{L^1} \leq C \|\psi \rho_\varepsilon \mathbb{1}_{\rho_\varepsilon \geq M}\|_{L^2_{t,x}}.$$

Choosing $M = \frac{1}{\varepsilon}$, we conclude. The condition (A.3) can be relaxed to be satisfied locally when ω is compactly supported. We stress that we do not have any a priori estimate implying (A.3).

REFERENCES

- [1] C. Bardos, P. Gwiazda, A. Świerczewska Gwiazda, E. S. Titi, and E. Wiedemann. Onsager's conjecture in bounded domains for the conservation of entropy and other companion laws. *Proc. A.*, 475(2230):20190289, 18, 2019.
- [2] M. Burger and A. Esposito. Porous medium equation and cross-diffusion systems as limit of nonlocal interaction. *arXiv preprint arXiv:2202.05030*, 2022.
- [3] J. W. Cahn and J. E. Hilliard. Free energy of a nonuniform system. I. Interfacial free energy. *The Journal of Chemical Physics*, 28(2):258–267, 1958.
- [4] J. A. Carrillo, M. G. Delgadino, L. Desvillettes, and J. S. Wu. The Landau equation as a gradient flow. *arXiv preprint arXiv:2007.08591*, to appear in *Analysis & PDE*, 2020.
- [5] J. A. Carrillo, M. G. Delgadino, and J. Wu. Convergence of a particle method for a regularized spatially homogeneous Landau equation. *Mathematical Models and Methods in Applied Sciences (in press)*, 2022.
- [6] J. A. Carrillo, C. Elbar, and J. Skrzeczkowski. Degenerate Cahn-Hilliard systems: From nonlocal to local. *arXiv preprint arXiv:2303.11929*, 2023.
- [7] J. A. Carrillo, A. Esposito, and J. S.-H. Wu. Nonlocal approximation of nonlinear diffusion equations. *arXiv preprint arXiv:2302.08248*, 2023.

- [8] P. Constantin, W. E, and E. S. Titi. Onsager’s conjecture on the energy conservation for solutions of Euler’s equation. *Comm. Math. Phys.*, 165(1):207–209, 1994.
- [9] E. Davoli, H. Ranetbauer, L. Scarpa, and L. Trussardi. Degenerate nonlocal Cahn-Hilliard equations: well-posedness, regularity and local asymptotics. *Ann. Inst. H. Poincaré C Anal. Non Linéaire*, 37(3):627–651, 2020.
- [10] E. Davoli, L. Scarpa, and L. Trussardi. Local asymptotics for nonlocal convective Cahn-Hilliard equations with $W^{1,1}$ kernel and singular potential. *J. Differential Equations*, 289:35–58, 2021.
- [11] E. Davoli, L. Scarpa, and L. Trussardi. Nonlocal-to-local convergence of Cahn-Hilliard equations: Neumann boundary conditions and viscosity terms. *Arch. Ration. Mech. Anal.*, 239(1):117–149, 2021.
- [12] R. J. DiPerna and P.-L. Lions. Ordinary differential equations, transport theory and Sobolev spaces. *Invent. Math.*, 98(3):511–547, 1989.
- [13] C. Elbar, M. Mason, B. Perthame, and J. Skrzeczkowski. From Vlasov equation to degenerate nonlocal Cahn-Hilliard equation. *arXiv preprint arXiv:2208.01026*, to appear in *Communications in Mathematical Physics*, 2022.
- [14] C. Elbar and J. Skrzeczkowski. Degenerate Cahn-Hilliard equation: From nonlocal to local. *J. Differential Equations*, 364:576–611, 2023.
- [15] G. Giacomin and J. L. Lebowitz. Phase segregation dynamics in particle systems with long range interactions. I. Macroscopic limits. *J. Statist. Phys.*, 87(1-2):37–61, 1997.
- [16] P. Gwiazda, M. Michálek, and A. Świerczewska Gwiazda. A note on weak solutions of conservation laws and energy/entropy conservation. *Arch. Ration. Mech. Anal.*, 229(3):1223–1238, 2018.
- [17] S. Hecht, M. Doumic, B. Perthame, and D. Peurichard. Multispecies cross-diffusions: from a nonlocal mean-field to a porous medium system without self-diffusion. *HAL preprint hal-04108050*, May 2023.
- [18] P.-L. Lions and S. Mas-Gallic. Une méthode particulière déterministe pour des équations diffusives non linéaires. *C. R. Acad. Sci. Paris Sér. I Math.*, 332(4):369–376, 2001.
- [19] S. Melchionna, H. Ranetbauer, L. Scarpa, and L. Trussardi. From nonlocal to local Cahn-Hilliard equation. *Adv. Math. Sci. Appl.*, 28(2):197–211, 2019.
- [20] S. Takata and T. Noguchi. A simple kinetic model for the phase transition of the van der Waals fluid. *Journal of Statistical Physics*, 172(3):880–903, 2018.

Charles Elbar: SORBONNE UNIVERSITÉ, LABORATOIRE JACQUES-LOUIS LIONS (LJLL), F-75005
PARIS, FRANCE

Email address: `charles.elbar@sorbonne-universite.fr`

Benoît Perthame: SORBONNE UNIVERSITÉ, CNRS, UNIVERSITÉ DE PARIS, INRIA, LABORATOIRE
JACQUES-LOUIS LIONS UMR7598, F-75005 PARIS

Email address: `Benoit.Perthame@sorbonne-universite.fr`

Jakub Skrzeczkowski: INSTITUTE OF MATHEMATICS OF POLISH ACADEMY OF SCIENCES; FACULTY
OF MATHEMATICS, INFORMATICS AND MECHANICS, UNIVERSITY OF WARSAW, POLAND

Email address: `jakub.skrzeczowski@student.uw.edu.pl`