



HAL
open science

Erratum for Global Sensitivity Analysis: a novel generation of mighty estimators based on rank statistics

Fabrice Gamboa, Thierry Klein, Agnès Lagnoux, Paul Rochet

► To cite this version:

Fabrice Gamboa, Thierry Klein, Agnès Lagnoux, Paul Rochet. Erratum for Global Sensitivity Analysis: a novel generation of mighty estimators based on rank statistics. 2023. hal-04125285

HAL Id: hal-04125285

<https://hal.science/hal-04125285v1>

Preprint submitted on 13 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Global Sensitivity Analysis: a novel generation of mighty estimators based on rank statistics

Fabrice Gamboa¹, Thierry Klein², Agnès Lagnoux³, and Paul Rochet⁴

¹Institut de Mathématiques de Toulouse and ANITI; UMR5219. Université de Toulouse; CNRS. UT3, F-31062 Toulouse, France.

²Institut de Mathématiques de Toulouse; UMR5219. Université de Toulouse; ENAC - Ecole Nationale de l'Aviation Civile, Université de Toulouse, France

³Institut de Mathématiques de Toulouse; UMR5219. Université de Toulouse; CNRS. UT2J, F-31058 Toulouse, France.

⁴ENAC - Ecole Nationale de l'Aviation Civile, Université de Toulouse, France.

June 13, 2023

Abstract

Correction to the proof of Lemma B.2 in Global Sensitivity Analysis: a novel generation of mighty estimators based on rank estimators. F. Gamboa, P. Gremaud, T. Klein, and A. Lagnoux. Bernoulli. (2021) Vol. 28, n°4, pp. 2345-2374.

Key words Global sensitivity analysis, Cramér-von-Mises distance, Pick-Freeze method, Chatterjee's coefficient of correlation, Sobol indices estimation.

AMS subject classification 62G05, 62G20, 62G30.

The purpose of this note is twofold. First, we correct a mistake in the expression of the limiting variance Σ_C that appears in [1, Lemma B.2]. Secondly, we provide simpler expressions for Σ_B in [1, Lemma B.1] and Σ_C . In particular, the expression of Σ_C does no longer depend on the first derivative of the model f . Furthermore, the updated formula of the limiting variance is now consistent with the limiting variance σ_{rank}^2 obtained in [2] by other means.

1 Computation of the asymptotic covariance matrix

Σ_B

In [1, Section B.2], we obtained:

$$\Sigma_B^{1,1} = \mathbb{E}[\text{Var}(YY'|X)] + 2\mathbb{E}[\text{Cov}(YY', YY''|X)],$$

where we remind that $Y = f(X, W)$, $Y' = f(X, W')$, and $Y'' = f(X, W'')$ with W' and W'' independent copies of W . Let us introduce the notation

$$\phi(x) = \mathbb{E}[Y|X = x] \quad \text{and} \quad \sigma^2(x) = \text{Var}(Y|X = x).$$

Then, by conditional independence, we get

$$\text{Var}(YY'|X) = \mathbb{E}[(YY')^2|X] - \mathbb{E}[YY'|X]^2 = \mathbb{E}[Y^2|X]^2 - \mathbb{E}[Y|X]^4 = \sigma^4(X) + 2\sigma^2(X)\phi^2(X)$$

so that

$$\mathbb{E}[\text{Var}(YY'|X)] = \mathbb{E}[\sigma^4(X)] + 2\mathbb{E}[\sigma^2(X)\phi^2(X)].$$

Analogously,

$$\mathbb{E}[\text{Cov}(YY', YY''|X)] = \mathbb{E}[\mathbb{E}[Y^2Y'Y''|X]] - \mathbb{E}[\mathbb{E}[YY'|X]\mathbb{E}[YY''|X]] = \mathbb{E}[\sigma^2(X)\phi^2(X)].$$

Finally,

$$\Sigma_B^{1,1} = \mathbb{E}[\sigma^4(X)] + 4\mathbb{E}[\sigma^2(X)\phi^2(X)].$$

Similarly, the other terms are

$$\begin{aligned} \Sigma_B^{2,2} &= \mathbb{E}[\text{Var}(Y|X)] = \mathbb{E}[\sigma^2(X)], \\ \Sigma_B^{3,3} &= \mathbb{E}[\text{Var}(Y^2|X)] = \mathbb{E}[\phi_4(X)] - \mathbb{E}[\phi_2(X)^2], \\ \Sigma_B^{1,2} &= \Sigma_B^{2,1} = 2\mathbb{E}[\text{Cov}(YY', Y|X)] = 2\mathbb{E}[\sigma^2(X)\phi(X)], \\ \Sigma_B^{1,3} &= \Sigma_B^{3,1} = 2\mathbb{E}[\text{Cov}(YY', Y^2|X)] = 2\mathbb{E}[\phi_3(X)\phi(X)] - 2\mathbb{E}[\phi_2(X)\phi^2(X)], \\ \Sigma_B^{2,3} &= \Sigma_B^{3,2} = \mathbb{E}[\text{Cov}(Y, Y^2|X)] = \mathbb{E}[\phi_3(X)] - \mathbb{E}[\phi(X)\phi_2(X)], \end{aligned}$$

where $\phi_i(x) = \mathbb{E}[Y^i|X = x]$ for $i = 2, 3, 4$.

2 Computation of the asymptotic covariance matrix

Σ_C

In [1, Section B.3], we obtained

$$\begin{aligned} \Sigma_C^{1,1} &= \int \psi_{100}(x_1, x'_1, x''_1, \chi_1) \psi_{100}(x_2, x'_2, x''_2, \chi_2) x_1 \wedge x_2 \wedge x'_1 \wedge x'_2 \wedge x''_1 \wedge x''_2 \\ &\quad \times dQ(x_1, x'_1, x''_1, \chi_1) dQ(x_2, x'_2, x''_2, \chi_2) - \left(\int \psi_{100}(x, x', x'', \chi) x \wedge x' \wedge x'' dQ(x, x', x'', \chi) \right)^2 \end{aligned}$$

leading to the corrected expression of $\Sigma_C^{1,1}$;

$$\begin{aligned} \Sigma_C^{1,1} &= \mathbb{E}[(Y_1 + Y_1'')(Y_2 + Y_2'') f_x(X_1, W_1') f_x(X_2, W_2')(X_1 \wedge X_2)] - \mathbb{E}[(Y + Y'') f_x(X, W') X]^2 \\ &= 4\mathbb{E}[Y_1 Y_2 f_x(X_1, W_1') f_x(X_2, W_2')(X_1 \wedge X_2)] - 4\mathbb{E}[Y f_x(X, W') X]^2, \end{aligned}$$

where we remind that $Y = f(X, W)$, $Y' = f(X, W')$, and $Y'' = f(X, W'')$ with W' and W'' independent copies of W (and analogously for Y_1 and Y_2). Hence, we remark that $\Sigma_C^{1,1} = \Sigma_C^{3,3}$. Considering the second term in $\Sigma_C^{1,1}$, we get

$$\begin{aligned}\mathbb{E}[Y f_x(X, W')X] &= \mathbb{E}[f(X, W)f_x(X, W')X] = \mathbb{E}[\mathbb{E}[f(X, W)f_x(X, W')X|X]] \\ &= \mathbb{E}[X\mathbb{E}[f(X, W)|X]\mathbb{E}[f_x(X, W')|X]] \quad \text{by conditional independence} \\ &= \frac{1}{2}\mathbb{E}[X\phi(X)\phi'(X)] = \frac{1}{2}(\phi^2(1) - \mathbb{E}[\phi^2(X)]) \quad \text{by integration by parts.}\end{aligned}$$

Similarly,

$$4\mathbb{E}[Y_1 f_x(X_1, W'_1)Y_2 f_x(X_2, W'_2)(X_1 \wedge X_2)] = \phi^4(1) - 2\phi^2(1)\mathbb{E}[\phi^2(X)] + \mathbb{E}[\phi^4(X)]$$

that yields

$$\Sigma_C^{1,1} = \phi^4(1) - 2\phi^2(1)\mathbb{E}[\phi^2(X)] + \mathbb{E}[\phi^4(X)] - (\phi^2(1) - \mathbb{E}[\phi^2(X)])^2 = \text{Var}(\phi^2(X)).$$

Finally, the remaining terms of Σ_C (after correction) are:

$$\begin{aligned}\Sigma_C^{2,2} &= \mathbb{E}[f_x(X_1, W_1)f_x(X_2, W_2)(X_1 \wedge X_2)] - \mathbb{E}[f_x(X, W)X]^2 = \text{Var}(\phi(X)) \\ \Sigma_C^{1,2} &= \mathbb{E}[(Y_1 + Y_1'')f_x(X_1, W'_1)f_x(X_2, W_2)(X_1 \wedge X_2)] - \mathbb{E}[(Y + Y'')f_x(X, W')X]\mathbb{E}[f_x(X, W)X] \\ &= 2\mathbb{E}[Y_1 f_x(X_1, W'_1)f_x(X_2, W_2)(X_1 \wedge X_2)] - 2\mathbb{E}[Y f_x(X, W')X]\mathbb{E}[f_x(X, W)X] \\ &= \text{Cov}(\phi(X), \phi^2(X)) \\ \Sigma_C^{1,3} &= 2\mathbb{E}[(Y_1 + Y_1'')f_x(X_1, W'_1)Y_2 f_x(X_2, W'_2)(X_1 \wedge X_2)] - 2\mathbb{E}[(Y + Y'')f_x(X, W')X]\mathbb{E}[Y f_x(X, W')X] \\ &= 4\mathbb{E}[Y_1 f_x(X_1, W'_1)Y_2 f_x(X_2, W'_2)(X_1 \wedge X_2)] - 4\mathbb{E}[Y f_x(X, W')X]^2 = \Sigma_C^{1,1} \\ \Sigma_C^{2,3} &= 2\mathbb{E}[f_x(X_1, W_1)Y_2 f_x(X_2, W'_2)(X_1 \wedge X_2)] - 2\mathbb{E}[f_x(X, W)X]\mathbb{E}[Y f_x(X, W')X] = \Sigma_C^{1,2}.\end{aligned}$$

Evidently, $\Sigma_C^{2,1} = \Sigma_C^{1,2}$, $\Sigma_C^{3,1} = \Sigma_C^{1,3}$, and $\Sigma_C^{3,2} = \Sigma_C^{2,3}$.

References

- [1] F. Gamboa, P. Gremaud, T. Klein, and A. Lagnoux. Global sensitivity analysis: A novel generation of mighty estimators based on rank statistics. *Bernoulli*, 28(4):2345–2374, 2022.
- [2] T. Klein and P. Rochet. Efficiency of the averaged rank-based estimator for first order Sobol index inference. working paper or preprint, June 2023.