



HAL
open science

Gain of function SCN1A disease-causing variants: Expanding the phenotypic spectrum and functional studies guiding the choice of effective antiseizure medication

Sara Matricardi, Sandrine Cestèle, Marina Trivisano, Benedetta Kassabian,
Nathalie Leroudier, Roberta Vittorini, Margherita Nosadini, Elisabetta
Cesaroni, Sabrina Siliquini, Cristina Marinaccio, et al.

► **To cite this version:**

Sara Matricardi, Sandrine Cestèle, Marina Trivisano, Benedetta Kassabian, Nathalie Leroudier, et al..
Gain of function SCN1A disease-causing variants: Expanding the phenotypic spectrum and functional
studies guiding the choice of effective antiseizure medication. *Epilepsia*, 2023, 64 (5), pp.1331-1347.
10.1111/epi.17509 . hal-04127124

HAL Id: hal-04127124

<https://hal.science/hal-04127124v1>

Submitted on 13 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Gain of function SCN1A disease-causing variants: Expanding the phenotypic spectrum and functional studies guiding the choice of effective antiseizure medication

Journal:	<i>Epilepsia</i>
Manuscript ID	EPI-00494-2022.R2
Manuscript Type:	Research Article
Date Submitted by the Author:	19-Nov-2022
Complete List of Authors:	<p>Matricardi, Sara; AOU Ospedali Riuniti di Ancona, Neuropsychiatry Cestele, Sandrine; Institut de Pharmacologie Moléculaire et Cellulaire, CNRS UMR6097 and University of Nice-Sophia Antipolice</p> <p>Trivisano, Marina; Children's Hospital Bambino Gesù, Department of Neuroscience</p> <p>Kassabian, Benedetta; University of Padua, Department of Neurosciences</p> <p>Leroudier, Nathalie ; Universite Cote d'Azur, CNRS UMR7275, Institute of Molecular and Cellular Pharmacology (IPMC)</p> <p>Vittorini, Roberta; 6Child and adolescence neuropsychiatry Service, Department of child pathology and cure, Children hospital "Regina Margherita"</p> <p>Nosadini, Margherita; The Children's Hospital at Westmead, Neuroimmunology group</p> <p>Cesaroni, Elisabetta; AOU Ospedali Riuniti di Ancona, Neuropsychiatry</p> <p>Siliquini, Sabrina; Paediatric Hospital G Salesi, Neurology Unit and Neurogenetics Laboratories</p> <p>Marinaccio, Cristina; 6Child and adolescence neuropsychiatry Service, Department of child pathology and cure, Children hospital "Regina Margherita"</p> <p>Longaretti, Francesca ; Child and adolescence neuropsychiatry service, S. Croce e Carle Hospital,</p> <p>Podestà, Barbara; Santa Croce e Carle Hospital, Department of Pediatric Neurology</p> <p>Operto, Francesca; Child and Adolescent Neuropsychiatry, Medical School, University of Salerno, Child and Adolescent Neuropsychiatry</p> <p>Luisi, Concetta; University of Padua, Department of Neurosciences</p> <p>Sartori, Stefano; Department of Woman and Child Health, Pediatric Neurology and Neurophysiology Unit;</p> <p>Boniver, Clementina ; Department of Woman and Child Health, Pediatric Neurology and Neurophysiology Unit</p> <p>Specchio, Nicola; Bambino Gesu Pediatric Hospital, Department of Neuroscience</p> <p>Vigevano, Federico; Bambino Gesu Children's Hospital, Neurology</p> <p>Marini, Carla; Paediatric Hospital G Salesi, Neurology Unit and Neurogenetics Laboratories; Child Neurology and Psychiatric Unit, Salesi Pediatric Hospital</p> <p>Mantegazza, Massimo; Universite Cote d'Azur, CNRS UMR7275, Institute of Molecular and Cellular Pharmacology (IPMC);</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Key Words:	gain of function, focal epilepsy, Sodium channel Blockers, SCN1A gene

SCHOLARONE™
Manuscripts

1
2
3 **GAIN OF FUNCTION *SCN1A* DISEASE-CAUSING VARIANTS: EXPANDING THE**
4 **PHENOTYPIC SPECTRUM AND FUNCTIONAL STUDIES GUIDING THE CHOICE OF**
5 **EFFECTIVE ANTISEIZURE MEDICATION**
6
7
8
9

10 **Sara Matricardi^{1,2*}, Sandrine Cestèle^{3,4*}, Marina Trivisano⁵, Benedetta Kassabian⁶, Nathalie**
11 **Leroudier^{3,4}, Roberta Vittorini⁷, Margherita Nosadini⁸, Elisabetta Cesaroni¹, Sabrina**
12 **Siliquini¹, Cristina Marinaccio⁷, Francesca Longaretti⁹, Barbara Podestà⁹, Francesca Felicia**
13 **Operto¹⁰, Concetta Luisi^{5,6}, Stefano Sartori⁸, Clementina Boniver⁸, Nicola Specchio⁵, Federico**
14 **Vigevano¹¹, Carla Marini^{1#}, Massimo Mantegazza^{4,5,12#}.**

15
16
17
18
19 ¹Child Neurology and Psychiatry Unit, Children's Hospital "G. Salesi", Ospedali Riuniti Ancona,
20 Italy

21
22 ²Department of Pediatrics, University of Chieti, Italy

23
24 ³Université Cote d'Azur, Valbonne-Sophia Antipolis, France

25
26 ⁴CNRS, Institute of Molecular and Cellular Pharmacology (IPMC), Valbonne-Sophia Antipolis,
27 France

28
29 ⁵Rare and Complex Epilepsy Unit, Department of Neuroscience, Bambino Gesù Children's
30 Hospital, IRCCS, Full Member of European Reference Network EpiCARE, Rome, Italy

31
32 ⁶Neurology Unit, Department of Neuroscience, University of Padua, Padua, Italy

33
34 ⁷Child and adolescence neuropsychiatry Service, Department of child pathology and cure, Children
35 hospital "Regina Margherita", Turin, Italy

36
37 ⁸Pediatric Neurology and Neurophysiology Unit, Department of Women and Children's Health,
38 University of Padua, Italy

39
40 ⁹Child and adolescence neuropsychiatry service, S. Croce e Carle Hospital, Cuneo, Italy

41
42 ¹⁰Child and Adolescent Neuropsychiatry Unit, Department of Medicine, Surgery and Dentistry
43 University of Salerno, Italy

44
45
46 ¹¹Neurology Unit, Department of Neuroscience, Bambino Gesù Children's Hospital, IRCCS, Full
47 Member of European Reference Network EpiCARE, Rome, Italy.

48
49 ¹²Inserm, Valbonne-Sophia Antipolis, France

50
51
52
53 ***Authors equally contributed**

54 Sara Matricardi, ORCID ID: <https://orcid.org/0000-0002-4403-6342>

55 Sandrine Cestèle, ORCID ID: <https://orcid.org/0000-0002-5982-1562>

56
57
58
59
60 **#Co-corresponding authors**

1
2
3 Carla Marini, ORCID ID: <https://orcid.org/0000-0002-9212-2691>

4 Child Neurology and Psychiatry Unit, Children's Hospital "G. Salesi", Ospedali Riuniti Ancona, Italy

5
6 Email: carla.marini@ospedaliriuniti.marche.it
7
8
9

10 Massimo Mantegazza, ORCID ID: <https://orcid.org/0000-0002-1070-7929>

11 Université Cote d'Azur, Valbonne-Sophia Antipolis, France; CNRS, Institute of Molecular and
12 Cellular Pharmacology (IPMC), Valbonne-Sophia Antipolis, France; Inserm, Valbonne-Sophia
13 Antipolis, France.

14
15 Email: mantegazza@ipmc.cnrs.fr
16
17
18
19

20 **Co-author emails:**

21 sara.matricardi@yahoo.it
22 cestele@ipmc.cnrs.fr
23 marina.trivisano@opbg.net
24 benedettakassabian@gmail.com
25 leroudier@ipmc.cnrs.fr
26 roberta.vittorini@unito.it
27 margherita.nosadini@gmail.com
28 elisabetta.cesaroni@ospedaliriuniti.marche.it
29 sabrina.siliquini@ospedaliriuniti.marche.it
30 cmarinaccio@cittadellasalute.to.it
31 longaretti.f@ospedale.cuneo.it
32 podesta.b@ospedale.cuneo.it
33 opertofrancesca@gmail.com
34 concetta.luisi@unipd.it
35 stefano.sartori@unipd.it
36 clementinaboniverconte@gmail.com
37 nicola.specchio@opbg.net
38 federico.vigevano@opbg.net
39 mantegazza@ipmc.cnrs.fr
40 carla.marini@ospedaliriuniti.marche.it
41
42
43
44
45
46
47

48 Abstract word count: 283

49 **Main article word count: 4836**

50
51 Number of tables: 3

52
53 Number of figures: 3

54
55 Number of Supplemental Tables: 1

56 **Number of Supplemental Figures: 3**

57
58 **Number of references: 58**
59
60

ABSTRACT

Objectives: To refine the spectrum of *SCN1A*-epileptic disorders other than Dravet syndrome (DS) and genetic epilepsy with febrile seizures plus (GEFS+) and optimize anti-seizure management by correlating phenotype-genotype relationship and functional consequences of *SCN1A* variants in a cohort of patients. We also performed a literature review including individuals with *SCN1A* variants causing non-DS and non-GEFS+ phenotypes and compared the features of the two cohorts.

Results: Sixteen probands were ascertained via a national collaborative network, nine (56%) with de novo pathogenic variants causing developmental and epileptic encephalopathy (DEE) with seizure onset at a median age of 2 months and severe intellectual disability. Seven subjects (54%), five with inherited and two with de novo variants, manifested focal epilepsies (FE) with mild or no intellectual disability. Sodium-channel-blockers never worsened seizures, and 50% of patients experienced long periods of seizure freedom. We found 13 *SCN1A* missense variants, eight of them were novel and never reported. Functional studies of three representative variants showed a gain of channel function. The literature review led to the identification of 44 individuals with *SCN1A* variants and non-DS, non-GEFS+ phenotypes. The comparison with our cohort highlighted that DEE phenotypes are a common feature.

Significance: The boundaries of *SCN1A*-disorders are wide and still expanding. In our cohort, more than 50% of patients manifested focal epilepsies, which are thus a frequent feature of *SCN1A* pathogenic variants beyond DS and GEFS+. *SCN1A* testing should therefore be included in the diagnostic workup of pediatric, familial and non-familial, focal epilepsies. Alternative, non-DS/non-GEFS+ phenotypes might be associated with gain of channel function, and sodium-channel-blockers could control seizures by counteracting excessive channel function. Functional analysis evaluating the consequences of pathogenic *SCN1A* variants is thus relevant to tailor the appropriate ASM.

1
2
3 **Keywords:** *SCN1A* gene, Gain of Function, Focal Epilepsy, Sodium channel Blockers
4

5 **Key Points:**
6

- 7
- 8 - The boundaries of *SCN1A*-disorders are wide and still expanding
 - 9
 - 10 - Focal epilepsies and other DEEs should be included in the spectrum of *SCN1A*-disorders
 - 11
 - 12 - Functional studies of *SCN1A* missense variants associated with these alternative phenotypes
13 show a moderate gain of channel function
14
 - 15 - Sodium-channel-blockers could control seizures by counteracting excessive channel
16 function
17
 - 18 - Functional studies for evaluating the consequences of *SCN1A* pathogenic variants are
19 relevant to tailor the appropriate ASM
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1. INTRODUCTION

The *SCN1A* gene encoding the voltage-gated sodium channel Nav1.1 is the target of thousands of pathogenic variants responsible for a broad spectrum of neurological phenotypes.¹⁻⁵ Dravet syndrome (DS) remains the prototypic developmental and epileptic encephalopathy (DEE) linked to de novo *SCN1A* pathogenic variants. Less than 10% of variants are inherited from unaffected or mildly affected parents whose phenotypes fall within the genetic epilepsy with febrile seizures plus (GEFS+) spectrum.⁶ Rare patients carrying *SCN1A* pathogenic variants have been classified as early-onset DEE with movement disorders and, more recently, neonatal presentation with arthrogryposis.⁷⁻¹⁰ In addition to epilepsy, *SCN1A* variants have also been linked to paroxysmal non-epileptic attacks, including hemiplegic migraine and neurodevelopmental disorders.^{5,11} The most common underlying epileptogenic mechanism is related to loss-of-function (LoF) of Nav1.1, leading to hypoexcitability of GABAergic neurons and reduced inhibition.^{1,2} This *SCN1A* haploinsufficiency¹² makes sodium-channel blocker (SCB) drugs contraindicated in these patients.¹³ The clinical evidence that some DS patients have increased seizure frequency when treated with SCBs supports the experimental hypothesis.^{14,15} *SCN1A* pathogenic variants have also been found to cause Nav1.1 gain-of-function (GoF) in patients with hemiplegic migraine and in several patients with neonatal and early-onset DEE.^{10,16-19} Overall, current knowledge suggests that when the pathogenic variants cause Nav1.1 GoF, the phenotype is a ‘non-DS and non-GEFS+ disorder’, and SCBs could be effective medications in these patients, counteracting excess function of Nav1.1 and possible neuronal hyperexcitability.¹⁰ Our study aims to assess the phenotypes associated with *SCN1A* pathogenic variants alternative to DS and GEFS+ and to analyze the effect of treatments with antiseizure medications (ASMs), including SCBs, in 16 newly diagnosed and 44 previously reported patients. To gain insight into pathological mechanisms, we also performed functional analysis of three variants representative of the phenotypic spectrum of this cohort.

2. MATERIALS & METHODS

2.1 Study design and data collection.

A national multicenter retrospective observational cohort study was performed by selecting patients carrying *SCN1A* pathogenic variants presenting with phenotypes other than DS or GEFS+. Patients with a medical history of prolonged, focal clonic (hemiclonic) or myoclonic seizures triggered by fever/illness in the first year of life supposed to have a DS phenotype were excluded. Furthermore, patients with typical febrile seizures that continue beyond the age of 6 years, evolving into afebrile seizures, supposed to have a GEFS+ phenotype were excluded.²⁰

Patients were collected by contacting tertiary epilepsy centers in Italy. Clinical and genetic data were collected from medical records using a specific form. Data were stored in a dataset sheet for analysis. EEG recordings and neuroimaging were reviewed by the expert epileptologist and neuroradiologist of the referral center.

We assessed: age, gender, family history, age at seizure onset, seizure type and frequency, episodes of status epilepticus, triggering factors, seizure off-set, duration of follow-up from seizure onset, EEG and MRI findings, treatment with ASMs and response to treatment, neurological findings, psychomotor development, and additional features.

Seizure types were classified according to the 2017 Operational Classification of the International League Against Epilepsy,²¹ and seizure frequency was reported as daily, weekly, monthly, or sporadic. Seizure classification for each patient was based on a combined description of events from parents and caregivers, home video, and ictal video-EEG. Parents and caregivers compiled seizure diaries allowing determination of seizure frequency, worsening, and improvements.

We compared the median number of seizures in the 3 months preceding the introduction of SCBs with the median number of seizures following the drug introduction and the achievement of the therapeutic dose. Treatment with ASMs was classified as “effective” when leading to seizure freedom or to greater than 75% seizure reduction, “mildly effective” between 50 to 75%, “no

1
2
3 significant changes” with a reduction between 0-50%, and “non-effective” when seizures persisted
4
5 at the same frequency or worsened.
6

7
8 Psychomotor development was defined as normal or delayed with mild, moderate, or severe
9
10 impairment according to neurological examination and evaluation of main motor and cognitive
11
12 milestones, including independent walking, speech development, and interactive skills.
13
14

15 16 17 **2.2 Genetic analysis.**

18
19 *SCN1A* disease-causing variants were identified through NGS epilepsy gene panels from different
20
21 laboratories. We included in the study only patients whose variants were classified as “pathogenic”
22
23 (Class-5) and “likely pathogenic” (Class-4), according to criteria detailed by Antoniadis et al.,²² and
24
25 based on the Classification following the Association of Clinical Genetics Science Practice
26
27 Guidelines.²³
28
29

30 31 32 33 **2.3 Functional study.**

34
35 We used the cDNA of the shorter splice variant isoform (-11 aa) of the human $Na_v1.1$ channel alpha
36
37 subunit (GenBank database accession no. NM_006920.4). which we subcloned into the pCDM8
38
39 vector to minimize rearrangements²⁷ and that we used in numerous other studies^{18,19,24-26}. The
40
41 variants p.T162I, p.T1501A, and p.R1892L were introduced by site-directed with the Quick Change
42
43 Lightning Kit (Stratagene). The entire open reading frame was sequenced after each amplification
44
45 to avoid spurious mutations. We expressed wild-type (WT) or mutant $hNa_v1.1$ in the cell line tsA-
46
47 201 (Sigma-Aldrich 96121229), which was maintained and transiently transfected with $CaPO_4$.²⁵
48
49 Cells were co-transfected with the pCDM8- $hNa_v1.1$ vector and a reporter vector expressing Yellow
50
51 Fluorescent Protein (pEYFP-N1; Clontech) to identify the transfected cells for electrophysiological
52
53 recordings. For some experiments, we co-transfected the human clones of $\beta 1$ and $\beta 2$ accessory
54
55 subunits at equimolar ratio with the wild-type or mutant $hNav1.1$, using the bicistronic plasmids
56
57
58
59
60

1
2
3 pIRES-YFP-h β 1 and pIRES-CFP-h β 2, to express with the same plasmid both the protein of interest
4 and the fluorescent protein as reported^{18,25,26}.
5
6

7 Sodium currents were recorded using the whole-cell configuration of the patch-clamp technique, as
8 previously reported.^{18,19,24-26} We started the recordings 5 min after establishing the whole-cell
9 configuration (t_0), to allow a complete dialysis of the cytoplasm and stabilize the initial time-
10 dependent shift of the voltage dependences; the different protocols were run in all the conditions at
11 the same time points from t_0 . Cells were recorded at room temperature (20-24°C) using a MultiClamp
12 700A amplifier and pClamp 10.2 software (Axon Instruments/Molecular devices). Signals were
13 filtered at 10 kHz and sampled at 50 kHz. Electrode capacitance and series resistance were
14 compensated during the experiment. Pipette resistance was 2-2.5 M Ω and maximal accepted voltage-
15 clamp error was less than 2.5 mV. The remaining transient and leakage currents were eliminated
16 using a P/4 subtraction paradigm. Recording solutions were (in mM): external solution 150 NaCl, 1
17 MgCl₂, 1.5 CaCl₂ and 10 HEPES (pH 7.4 with NaOH); internal pipette solution 105 CsF, 35 NaCl,
18 10 EGTA, 10 HEPES and (pH 7.4 with CsOH). Recordings were not corrected for junction potentials.
19 Voltage dependence of activation was studied by applying test pulses of 100-ms from -110 to +60
20 mV from a holding potential at -120 mV. Voltage dependence of inactivation was studied with a 100-
21 ms prepulse at different potentials followed by a test pulse at -10 mV. Conductance-voltage curves
22 were derived from current-voltage (I-V) curves according to $G=I/(V-V_r)$, where I is the peak current,
23 V is the test voltage, and V_r is the apparent observed reversal potential for tsA-201. The voltage
24 dependence of activation and of inactivation were fit to Boltzmann relationships in the form
25 $y=1/(1+\exp((V_{1/2}-V)/k))$, where y is normalized G_{Na} or I_{Na} , $V_{1/2}$ is the voltage of half-maximal activation
26 (V_a) or inactivation (V_h) and k is a slope factor; for the inactivation curve, we included a baseline for
27 taking into account incomplete inactivation.¹ The kinetics of current decay was quantified at -10mV,
28 fitting the first 10 ms of the decay with a single exponential function. Recovery from fast inactivation
29 was studied using a test pulse at 0 mV followed by repolarization at -80 mV of different duration and
30 a test pulse to 0 mV. The persistent sodium current (I_{NaP}) was quantified about 5 minutes after
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 obtaining the whole-cell configuration as the mean current between 40 and 50 ms after the beginning
4 of the voltage step and expressed as a percentage of the maximal transient current (I_{NaT}). Action
5 potential clamp recordings were performed using as voltage stimulus a GABA-ergic neuronal
6 discharge recorded from fast spiking banded cell injecting a 1s-long depolarizing current step as in
7 Chever et al.²⁸; the instantaneous firing frequency of the discharge was between 106 and 82Hz. The
8 intersweep interval was 8s for all the protocols. Data were analyzed with pClamp 10.2 (Axon
9 Instruments/Molecular devices) and Origin2021 (OriginLab).

10
11 For statistical analysis, normality was tested with the Kolmogorov-Smirnov test, homogeneity of
12 variance with the Levene's Test, and statistical comparisons were performed with one-way ANOVA
13 followed by Tukey post-hoc test. $P < 0.05$ was considered significant.

14
15
16
17
18
19
20
21
22
23
24
25
26
27
28 **2.4 Literature review:** PubMed search using the term “*SCN1A* gene mutations or pathogenic
29 variants” and included in the analysis only studies reporting patient-relevant information and
30 genetic findings, including epileptic phenotypes other than DS and GEFS+. Additional patients with
31 phenotypes other than DS and GEFS+ reported in other studies lacking clinical information were
32 excluded. Clinical and genetic data were collected in a separate dataset (Table S1 in Supplemental
33 Material).

2.5 Standard Protocol Approvals and Patient Consents

34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
This study was performed according to ethical principles for medical research involving human
subjects stated in the Declaration of Helsinki. Patients or their legally authorized representatives
consented to participate in the study. The study was approved by Marche Region Ethics Committee
(CERM).

3. RESULTS

We enrolled 16 newly diagnosed patients (8 females and 8 males), with a median age of 10.2 years (interquartile range [IQR]: 17.5 months-19 years; 8 months–43 years) (table 1).

Eleven patients had de novo *SCN1A* pathogenic variants leading to neonatal/early infantile DEE with seizure onset at a median age of 2 months (IQR: 2 days–2.5 months; 1 day–5 months) in nine (patients 2, 3, 5, 8, 9, 10, 11, 12, 16), and infantile/childhood focal epilepsy (FE) in two, with seizure onset at 9 and 36 months (patients 4, 7). Four patients presented familial pathogenic variants causing childhood-onset FE (median 7.5 years; IQR: 4.5-10.5 years; 4-11 years), and an 18-month child with early infantile-onset FE carried a variant inherited from his affected father (Figure S1 in Supplemental Material).

The median follow-up after seizure onset was 7.7 years (IQR: 17 months–14.5 years; 5.5 months–33 years).

3.1 De novo *SCN1A* pathogenic variants

- Nine patients with neonatal and early infantile DEE presented with focal motor (9/9; 100%), tonic (8/9; 88.8%), myoclonic (8/9; 88.8%), and bilateral tonic-clonic seizures (9/9; 100%). Over time other seizure types were epileptic spasms (4/9; 44.4%), focal dysautonomic (4/9; 44.4%), and migrating seizures (2/9; 22.2%). Four had episodes of status epilepticus (SE) (44.4%). One patient died of sudden unexplained death in epilepsy (SUDEP) at 27 years. Four of them also had hyperkinetic movement disorder. All patients had daily seizures at onset; over time, seizures persisted but with periods of reduced frequency or seizure freedom in 77.7%.

VEEG recordings revealed poorly organized background activity (BA) in most (6/9; 66.6%), with focal/multifocal discharges in 77.7% (7/9), migrating pattern in 22.2% (2/9), and diffuse epileptiform abnormalities in 22.2% (2/9).

Brain MRI was unrevealing in 66.5% (6/9), delayed myelination was present in 22.2% (2/9), while one patient with a longer follow-up showed progressive brain atrophy (patient 11).

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
- Two patients with infantile/childhood FE presented with predominant focal motor seizures with a clonic component, usually of alternating side involvement, both had focal to bilateral seizures. Onset was characterized by daily/weekly seizures, while sporadic events persisted over time with long periods of better control. VEEG recording was characterized by focal/multifocal epileptiform discharges. Neuroimaging was unrevealing in both.

3.2 Familial SCN1A pathogenic variants

- 15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
- The 18-month child (patient 1) carrying a familial p.V157I variant manifested focal seizures at one month of age and global delay with hypotonia. Seizures were rapidly controlled with phenobarbital (PB) which was withdrawn at 12 months. At 19 months, he was seizure-free off medication, able to walk, presented mild diffuse hypotonia, good non-verbal communication skills, and speech characterized by a limited number of single words. The repeated MRI was normal. He inherited the variant from his 34-year-old father, who had 3 seizures in the first year of life and is now healthy with normal cognition and motor function. *SCN1A* variant was also identified in the paternal uncle, who had few seizures in infancy; he is in his late twenties and leads a normal functioning, working life. Blood sample from additional family members was not available.
 - Patient 6, harboring the p.N1338H variant inherited from her healthy father, manifested with refractory focal seizures at 11 years, both spontaneous and reflex to tactile stimuli causing frequent falls; repeated brain MRI revealed a possible blurring over the left frontal-mesial lobe.
 - Two siblings and their father, harboring the p.R1892L variant, presented with childhood-onset sporadic focal motor seizures and auditory aura. VEEG recording revealed focal epileptiform abnormalities. Brain MRI was normal in all. NGS epilepsy gene panel excluded the presence of an LGI gene mutation.

58 None of the 16 patients presented classical or prolonged febrile seizures.

3.3 Treatment options and response

1
2
3 In this cohort, all patients but one received SCBs, which were effective or mildly effective in 73%
4 of them. Carbamazepine (CBZ) was administered to twelve patients, and it was mildly effective in
5 four with DEE and effective in five with childhood FE who gained seizure freedom for long
6 periods, with sporadic relapses related to low plasma levels. Lacosamide (LCM) was administered
7 to four patients, and three DEE were responders. Other effective treatments were VPA (5/11) and
8 ACTH (2/4). One familial patient (patient 1) was seizure-free without medication. At the age of 25
9 years, patient 6 underwent stereo-EEG recording of focal seizures followed by left frontal-mesial
10 lobectomy. Histology revealed only gliosis. Following the surgical procedure, the patient continued
11 to have seizures at a reduced frequency and without falls; when CBZ was added, she had long
12 periods of seizure freedom.
13
14
15
16
17
18
19
20
21
22
23
24

25 26 **3.4 Development and additional features**

27
28 Developmental delay (DD) was evident in the nine patients with neonatal/early infantile DEE who
29 had periods of regression resulting in a moderate to severe disability at the last follow-up. The two
30 patients with infantile/childhood FE and de novo *SCN1A* variants showed mild motor DD and
31 borderline cognitive functioning. The two siblings and their father (patients 13-15) with childhood
32 FE had normal cognitive functioning. The adult patient carrying a familial *SCN1A* variant (patient
33 6), despite a long history of focal refractory epilepsy and frontal lobectomy, had normal cognitive
34 functioning. The familial patient (patient 1) with early-infantile FE at 18 months had a mild DD.
35
36 Four patients with neonatal/early infantile DEE (4/9; 44.4%) developed an early hyperkinetic
37 movement disorder (MD), comprising hyperkinesia with perioral and upper limbs involvement,
38 dystonic posturing, myoclonic jerks, and chorea. Additional features in DEE patients were
39 hypotonia (3/9; 33.3%), pyramidal signs and spastic quadriplegia (4/9; 44.4%).
40
41
42
43
44
45
46
47
48
49
50
51
52

53 The three patients with familial childhood FE also suffered from migraine without aura.

54 55 56 **3.5 Genetic analysis**

1
2
3 The study includes thirteen *SCN1A* missense variants, ten are de novo, and three are inherited;
4
5 eleven occur singularly, whereas two are recurrent, including the inherited variant identified in the
6
7 affected siblings and father.
8

9
10 The p.R1636Q variant, detected in two, was previously reported in several other patients with
11
12 variable phenotypes, including DS,^{29,30} Lennox-Gastaut Syndrome,³¹ and DEE, myoclonic seizures,
13
14 dystonia, and spasticity.^{10,32} At the same position but with a different amino-acid change, the
15
16 p.R1636P variant detected in one of our patients was reported in a proband with EIMFS.³³ The
17
18 p.T226M variant, detected in one, was associated with DEE in several other patients.^{9,10} The
19
20 p.A420D variant identified in one of our patients was reported by Butler et al.,³² without phenotypic
21
22 description. The p.T162I variant detected in one of our patients was associated with DS.³⁴ The
23
24 p.N1338H variant was not previously reported, but a frameshift variant at position p.N1338 is
25
26 associated with DS.³⁵ As expected, the frameshift mutation causing LoF is associated with a more
27
28 severe, DS-like phenotype. Eight variants are novel and never reported.
29
30
31
32
33
34

35 **3.6 Functional study**

36
37 Whole-cell patch-clamp experiments were performed for variants p.T162I (neonatal DEE), p.T1501A
38
39 (infantile FE), and p.R1892L (childhood FE). The mean values of the functional properties are shown
40
41 in table 2. Figure 1 (A-L) displays representative current traces of WT and mutant channels elicited
42
43 with depolarizing voltage steps. WT and mutants had similar maximal current density consistent with
44
45 a similar plasma-membrane targeting (figure 1E). **IV curves showed similar features for WT and**
46
47 **mutants, but for p.T162I, whose current started to activate at more negative potentials (Figure S2 in**
48
49 **Supplemental Material). Consistently,** the activation curve of p.T162I was negatively shifted by
50
51 6.6mV, **indicating** a GoF (figure 1F-G). The slope of the activation curves of the mutants was similar
52
53 to **that of the WT** (figure 1H). The inactivation curve of p.T1501A and p.R1892L was positively
54
55 shifted by 9.5mV and 7.7mV respectively (figure 1I-J), consistent with a GoF, which is larger for
56
57 p.T1501A than p.R1892L. The inactivation curve of p.T162I was not shifted, but it was steeper than
58
59
60

1
2
3 that of the WT, although this modification does not have clear functional relevance (figure 1I-J-K).
4
5 The variant p.T1501A induced a faster recovery from fast inactivation, with 70% reduction in the
6
7 time constant of recovery at -80 mV (figure 1L), which is consistent with a GoF. We evaluated the
8
9 kinetics of activation and of decay of the current elicited with a depolarizing step to -10 mV (figure
10
11 2A-E). Activation kinetics was not modified. The exponential fit of the decay showed that the time
12
13 constant of p.T1501A is 1.53-fold slower (figure 2D; table 2), consistent with a GoF. The variant
14
15 p.T1501A was the only one modifying the amplitude of the persistent component of the current
16
17 (I_{NaP}), inducing a 3.94-fold increase (figure 2B, E), which is consistent with a GoF. To reproduce
18
19 neuronal dynamic conditions during a series of action potentials and to evaluate the overall effect of
20
21 the variants, we performed action potential-clamp experiments recording Na^+ action currents in tsA-
22
23 201 cells, using as voltage command the action potential discharge of a fast spiking GABAergic
24
25 neuron (figure 3A-E). These experiments evaluated the overall effect of the variants, with the
26
27 potential to identify dysfunctions that were not evident or not included (e.g., slow inactivation) in the
28
29 classical parameters, which we previously characterized applying voltage steps. Our action potential
30
31 clamp data correlated with the results obtained with the classical voltage step protocols. The
32
33 amplitude of the first action current was not statistically significant, but p.T1501A showed a strong
34
35 trend toward an increase, whereas p.T162I and p.R1892L showed a looser trend (figure 2F; table 2).
36
37 The effect was more prominent at the end of the discharge because p.T1501A and p.T162I showed a
38
39 7.05 and 2.35-fold increase, respectively, and p.R1892L showed a strong trend towards an increase
40
41 (figure 3G; table 2). **Action potential-clamp experiments performed with cells co-expressing with the
42
43 ~~alpha-subunits~~ $\beta 1$ and $\beta 2$ accessory subunits showed similar features (Figure S3 in Supplemental
44
45 Material), consistent with previous studies^{18,25,26,36}. Our functional study was performed in tsA-201
46
47 cells. Further studies are warranted to better elucidate the effect of GoF $Na_v 1.1$ epilepsy variants on
48
49 neuronal excitability including exploiting gene targeted animal models. A neuronal cell background
50
51 could modify functional effects, increasing them, possibly in a variant-specific manner. However, in
52
53 our previous studies ~~comparing~~ functional properties in tsA-201 cells and cortical neurons in culture,
54
55
56
57
58
59
60**

we observed that a neuronal cell background was able to rescue some folding/trafficking defective $Na_v1.1$ mutants increasing current density/membrane targeting, but did not significantly modify the effect that $Na_v1.1$ variants had on gating properties^{18,19,37}. Thus, since the GoF $Na_v1.1$ epilepsy variants that we studied do not reduce current density/membrane targeting, it is likely that their overall functional effect is similar in a neuronal background.

3.5 Literature review

We identified 44 unrelated subjects (18 females; 26 males) from ten previous studies (Table S1).

Clinical details were available for re-analysis for 42 probands with phenotypes other than DS and GEFS+ (Table 3).^{7-10,38-42} Forty patients carried *SCN1A* de novo missense variants, and the remaining four had de novo deletions of chromosome 2q24.3.

Seventeen patients had neonatal DEE, associated with hyperkinetic movement disorder in seven.

Eleven patients also had arthrogyriposis, multiple contractures, hip dysplasia and talipes, two

additional subjects who died in utero showed prenatal detection of arthrogyriposis. Twenty-five

patients presented with early infantile DEE, with and without MD. Multiple seizure types persisted over time and were refractory to all prescribed ASMs. Treatment details were reported for thirty-

eight patients. Commonly prescribed ASMs included BZP (73.6%), PB (55.2%), LEV (65.7%),

TPM (52.6%), VPA (63.1%), STP (38.3%), VGB (20.5%), cannabidiol (23.6%), fenfluramine

(13.1%), acetazolamide (11.7%), zonisamide (8.8%), ACTH (8.8%), perampanel (5.8%),

pyridoxine (5.8%), gabapentin (7.9%), rufinamide (5.8%), ESM (2.9%), felbamate (2.9%), cannabis

oil (2.9%), and bromide (2.9%). Thirty-two patients received more than one SCB (15 CBZ, 16

PHT, 9 LTG, 11 oxcarbazepine, 3 LCM) without seizure aggravation. Carbamazepine was effective

in 46.6% (7/15; two neonatal DEE, and five early infantile DEE), PHT in 50% (8/16; three neonatal

DEE, five early infantile DEE), LTG in 33.3% (3/9; early infantile DEE), oxcarbazepine in 63.6%

(7/11; 5 neonatal DEE, 2 early infantile DEE), LCM in 100% (3/3; neonatal DEE)

4. DISCUSSION

This study, including clinical and functional analysis of 16 novel patients and re-analysis of 44 previously reported subjects carrying *SCN1A* disease-causing variants with GoF effect, refines the clinical spectrum beyond the classical DS and GEFS+, increases the number of patients with *SCN1A*-related phenotypes, and describes their response to ASMs.

4.1 *SCN1A* and DEE phenotypes

Nine subjects, all carrying *de novo* variants, manifested a DEE. They exhibited early infantile focal clonic, tonic, myoclonic, and bilateral tonic-clonic seizures. Four patients also had spasms or focal dysautonomic seizures, and two presented migrating seizures. Unlike DS, these patients had multiple seizures per day at onset and concomitant cognitive impairment. Indeed, motor development and cognitive functions were impaired, ranging from mild to severe. Impaired cognition was present from the first weeks of life, confirming that *SCN1A* disease-causing variants underlie seizures and abnormal development. However, seizures, especially when very frequent and associated with an abnormal EEG, worsen motor and cognitive development. Our patient carrying the p.R1636Q variant presented a mild delay at the onset, when seizures were controlled with CBZ. At 14 months, she went through an ‘encephalopathic phase’, seizures reappeared, and EEG evolved to a continuous spike-wave pattern during sleep and wakefulness, accompanied by a decline of cognitive skills. Treatment with steroids ended the ‘encephalopathic phase’, leading to the re-acquisition of motor and cognitive abilities.

Approximately 40% of DEE patients developed an early MD, including hyperkinesia with perioral and upper limbs involvement, dystonic posturing, myoclonic jerks, and chorea.

4.2. *SCN1A* variants and focal epilepsy

Seven patients manifested focal epilepsies. Two carried *de novo* *SCN1A* variants, whereas five were inherited. Focal epilepsies presented from early infancy to childhood with focal motor, focal to bilateral, focal with auditory aura, and reflex seizures. Only two patients have ongoing sporadic seizures. Cognitive functioning was normal in five, and two showed mild DD. Two siblings and

1
2
3 their father also presented **with** sporadic episodes of migraine without aura that **might** be related to
4 the *SCN1A* variant in this family.
5
6

7 **4.3 Literature review and **comparison of cohorts****

8
9
10 The literature review identified **44** patients, **42** with available clinical information and carrying
11 *SCN1A* pathogenic variants not associated with DS or GEFS+.^{7-10,38-42} Common findings with our
12 cohort include: a) early-onset seizure; b) frequent focal and focal to bilateral seizures; c) epileptic
13 spasms occurring only in a quarter of patients. Unlike our cohort, migrating seizures are more
14 prominent and observed in about 40%. Eleven patients with neonatal DEE had arthrogyriposis,
15 multiple contractures, and hip dysplasia and talipes, and two additional patients who died in utero
16 showed prenatal detection of arthrogyriposis. None of the patients included in our cohort manifested
17 arthrogyriposis. Status epilepticus presented in most, and premature death occurred in 21% of
18 reported patients. In our series, only one patient died of SUDEP at 27 years. **All patients but one**
19 had severe or profound DD with hyperkinetic MD and hypotonia in the majority of them.
20
21 One of the main objectives of our study was to investigate the response to ASMs, especially to
22 SCBs. Data analysis showed a striking, ‘unexpected’, response to SCBs, considering the genetic
23 etiology and previous reports.¹⁴⁻¹⁶ More than 70% of patients included in our cohort responded to
24 SCBs, comprising four with childhood FE who were seizure-free for long periods. The remaining
25 patients treated with SCBs were not aggravated. In the previously reported patients, **76%** received
26 more than one SCBs without seizure aggravation, and **about 50%** of them were responders.
27
28 **We performed** functional studies on three *SCN1A* variants representative of the phenotypic
29 spectrum of this cohort, **observing** a clear GoF for all of them. **The variant p.T1501A showed** the
30 largest effect, **inducing** GoF modifications in five biophysical parameters, whereas p.T162I and
31 p.R1892L modified a single parameter. The analysis of the overall effect of the variants on the
32 function of Na_v1.1 by means of action-potential-clamp experiments confirmed that p.T1501A
33 induces the largest GoF, followed by p.T162I and p.R1892L. Functional effects do not correlate
34 with phenotype severity since the most severe among the three tested variants is associated with
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 p.T162I, which causes neonatal DEE. The GoF induced by these variants is moderate compared to
4
5 those causing hemiplegic migraine,^{11,18,19,25,43} confirming that the amount of GoF is not directly
6
7 correlated to phenotype severity. All three patients carrying the GoF tested variants, associated with
8
9 different phenotypes with variable severity, shared a beneficial response to therapeutic doses of
10
11 SCBs, supporting our hypothesis that they could be effective medications counteracting channel
12
13 excess of function and possible neuronal hyperexcitability.
14
15

16
17 Two additional *SCN1A* pathogenic variants identified in our cohort (T226M and R1636Q) were
18
19 recently found to cause moderate GoF^{9,10,42}, similarly to the variants that we have characterized
20
21 here. Overall, five of the thirteen *SCN1A* variants detected in our cohort lead to a moderate GoF.
22
23

24 There is a striking genotype-phenotype correlation in patients carrying the recurrent p.R1636Q
25
26 variant, because they all showed early-onset DEE with variable seizure semiology and manifested a
27
28 favorable response to SCBs treatment.
29

30 **4.4 Loss-of-function vs gain-of-function *SCN1A* variants: is there a functional effect-** 31 32 **phenotype correlation?** 33

34
35 Numerous functional studies of *SCN1A* missense variants have demonstrated a predominant LoF,
36
37 which is expected to result in hyperexcitability of neuronal networks because Na_v1.1 channels are
38
39 particularly important for the function of inhibitory interneurons.^{1,44,45} Available data suggest that
40
41 complete LoF variants leading to haploinsufficiency are more likely to cause severe DS phenotype,
42
43 whereas phenotypes might be milder and fall within the GEFS+ spectrum when the LoF is partial,
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
27,45-47 but seizures may worsen the phenotype.⁴⁸ Although strong LoF is considered the main effect,
there are studies that have observed mixed GoF and LoF effects for a few DS missense variants.^{44,49}

51 The pharmacological implication of the discovery of pathogenic *SCN1A* haploinsufficiency in DS
52
53 has contributed to the clinical management of DS and GEFS+ patients. In general, SCBs like CBZ,
54
55 PHT, and LTG are not effective or can possibly worsen seizures; thus, all guidelines advise their
56
57 discontinuation in patients with DS and *SCN1A* variants.^{14,50,51}
58
59
60

1
2
3 Instead, most *SCN1A* GoF variants cause a spectrum of disorders beyond the DS/GEFS+
4
5 dichotomy, which extends from mild focal epilepsies to hemiplegic migraine and to severe
6
7 DEE.^{10,16,18,19,25,28} Few studies have described GEFS+ *SCN1A* variants causing Na_v1.1 GoF,^{52,53}
8
9 although this finding is controversial.² The pharmacological implication of this observation is that
10
11 GoF-associated phenotypes might benefit from the treatment with SCBs counteracting the increased
12
13 function of Na_v1.1 channels (¹⁰, present study).

14
15
16
17 More studies are needed to understand how the moderate GoF observed in Na_v1.1 variants can
18
19 cause such a wide phenotypic range, including very severe phenotypes. We might, however,
20
21 speculate that the larger GoF of hemiplegic migraine variants could block seizure onset because it
22
23 induces strong hyperexcitability of GABAergic neurons leading to extracellular K⁺ accumulation
24
25 and spreading depolarization of the entire neuronal network, as we have previously shown^{28,54}.
26
27 Notably, spreading depolarization has been proposed as a general antiseizure mechanism⁵⁵, and the
28
29 milder hyperexcitability induced by GoF Na_v1.1 epilepsy variants would not be sufficient to induce
30
31 it. Phenotypic variability in patients might also be conditioned by interactions with other genetic
32
33 variants, as recently observed in a family with large phenotypic variability⁵⁶. Additional source of
34
35 functional and phenotypic variability might be the recently observed dimer interaction between Na_v
36
37 alpha subunits⁵⁷ which might modify the functional properties of homomeric or heteromeric dimers,
38
39 possibly in a variant-specific manner.
40
41
42
43
44
45
46
47
48

49 5. CONCLUSION

50
51 Clinical analysis of the 16 plus 40 patients included in the study highlights that the boundaries of
52
53 GoF *SCN1A* disorders are wider than previously thought, although it is not clear yet how this
54
55 functional effect can cause these phenotypes. Seizures can arise from the first day of life, be very
56
57 severe from the onset and occur without triggers. On the other extreme, pathogenic variants are
58
59 associated with childhood-onset, mild epilepsies, and normal cognition. Overall, focal epilepsies
60

1
2
3 were diagnosed in 43.7% of probands, suggesting that focal seizures are a **major clinical feature** of
4
5 our cohort. This result supports the recommendation to include *SCN1A* testing in the diagnostic
6
7 workup of pediatric, familial and non-familial, focal epilepsies. This phenotypic evidence is in line
8
9 with a recent paper reporting an *SCN1A* gene overexpression in brain tissue of temporal lobe
10
11 epilepsy patients with a specific genotype (rs7587026)⁵⁸.
12
13

14 A relevant observation of our study is that it ‘unmasks the myth’ that SCBs should not be used in
15
16 patients with *SCN1A* disease-causing variants. There are patients in whom, due to the type of
17
18 variant causing a GoF, SCBs might be beneficial **because the variant they carry induces GoF**, and
19
20 functional analysis to evaluate the consequences of variants can be relevant in some cases to tailor
21
22 the appropriate ASM.
23
24
25
26
27


28 **Ethical Publication Statement**

29
30 We confirm that we have read the Journal’s position on issues involved in ethical publication and
31
32 affirm that this report is consistent with those guidelines.
33
34
35

36 **Data Availability**

37
38 Anonymized data not published within this article will be made available by request from any
39
40 qualified investigator.
41
42
43
44
45

46 **Study Funding**

47
48  S.C. and M.M. ~~and~~ work was supported by the Investissements d’Avenir-Laboratory of Excellence
49
50 “Ion Channels Science and Therapeutics” (LabEx ICST, ANR-11-LABX-0015-01 to MM), the
51
52 University Côte d’Azur-IDEX Jedi (ANR-15-IDEX-01 to MM) and by the Foundation Famiglie
53
54 Dravet ONLUS – Italy (FDO-2018).
55
56
57
58
59

60 **Acknowledgments**

1
2
3 We thank the Salesi Foundation Onlus – Children’s Hospital “G. Salesi”, Ancona, Italy, for the
4
5 support in the realization of this study which is included as a pilot-project of a more comprehensive
6
7 project aiming to evaluate the overall diagnostic genetic yield in pediatric epilepsies.
8
9

10 11 **Disclosure of Conflicts of Interest.**

12
13 None of the authors has any conflict of interest to disclose.
14
15

16 17 **Authors’ contribution:**

18 Sara Matricardi, Sandrine Cestèle, Massimo Mantegazza, and Carla Marini conceptualized and
19
20 designed the study. Sara Matricardi, Marina Trivisano, Benedetta Kassabian, Roberta Vittorini,
21
22 Margherita Nosadini, Elisabetta Cesaroni, Sabrina Siliquini, Cristina Marinaccio, Francesca
23
24 Longaretti, Barbara Podestà, Francesca Felicia Operto, Concetta Luisi, Stefano Sartori, Clementina
25
26 Boniver, Nicola Specchio, Federico Vigevano, and Carla Marini selected and enrolled patients,
27
28 critically reviewing all medical charts and records. Sandrine Cestèle, Nathalie Leroudier, and
29
30 Massimo Mantegazza performed the functional analysis. Sara Matricardi and Carla Marini
31
32 performed the literature review and were involved with the dataset and analysis. Sara Matricardi,
33
34 Sandrine Cestèle, Massimo Mantegazza, and Carla Marini drafted the manuscript. All the Authors
35
36 are responsible for each case data collection and diagnostic process. All Authors edited the
37
38 manuscript.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REFERENCES

1. Mantegazza M, Cestèle S, Catterall WA. Sodium channelopathies of skeletal muscle and brain. *Physiol Rev.* 2021;101(4):1633-1689.
2. Mantegazza M, Broccoli V. SCN1A/Na_v 1.1 channelopathies: Mechanisms in expression systems, animal models, and human iPSC models. *Epilepsia.* 2019;60 Suppl 3:S25-S38.
3. Meisler MH, Hill SF, Yu W. Sodium channelopathies in neurodevelopmental disorders. *Nat Rev Neurosci.* 2021;22:152-166.
4. Marini C, Mantegazza M. Na⁺ channelopathies and epilepsy: recent advances and new perspectives. *Expert Rev Clin Pharmacol.* 2010;3:371-84.
5. Scheffer IE, Nabbout R. SCN1A-related phenotypes: Epilepsy and beyond. *Epilepsia.* 2019;60 Suppl 3:S17-S24.
6. Zhang YH, Burgess R, Malone JP, Glubb GC, Helbig KL, Vadlamudi L, et al. Genetic epilepsy with febrile seizures plus: Refining the spectrum. *Neurology.* 2017;89(12):1210-1219.
7. Carranza Rojo D, Hamiwka L, McMahon JM, Dibbens LM, Arsov T, Suls A, et al. De novo SCN1A mutations in migrating partial seizures of infancy. *Neurology.* 2011;77(4):380-3.
8. Freilich ER, Jones JM, Gaillard WD, Conry JA, Tsuchida TN, Reyes C, et al. Novel SCN1A mutation in a proband with malignant migrating partial seizures of infancy. *Arch Neurol.* 2011;68(5):665-71.
9. Sadleir LG, Mountier EI, Gill D, Davis S, Joshi C, DeVile C, et al. Not all SCN1A epileptic encephalopathies are Dravet syndrome: Early profound Thr226Met phenotype. *Neurology.* 2017;89(10):1035-1042.
10. Brunklaus A, Brünger T, Feng T, Fons C, Lehtikoinen A, Panagiotakaki E, et al. The gain of function SCN1A disorder spectrum: novel epilepsy phenotypes and therapeutic implications. *Brain.* 2022 Jun 13:awac210. Epub ahead of print.

- 1
2
3 11. Mantegazza M, Cestèle S. Pathophysiological mechanisms of migraine and epilepsy:
4
5 Similarities and differences. *Neurosci Lett*. 2018;667:92-102.
6
- 7
8 12. Bechi G, Scalmani P, Schiavon E, Rusconi R, Franceschetti S, Mantegazza M. Pure
9
10 haploinsufficiency for Dravet syndrome Na(V)1.1 (SCN1A) sodium channel truncating
11
12 mutations. *Epilepsia*. 2012;53(1):87-100.
13
- 14
15 13. Brunklaus A, Ellis R, Reavey E, Forbes GH, Zuberi SM. Prognostic, clinical and
16
17 demographic features in SCN1A mutation-positive Dravet syndrome. *Brain*. 2012;135(Pt
18
19 8):2329-36.
20
- 21
22 14. Guerrini R, Dravet C, Genton P, Belmonte A, Kaminska A, Dulac O. Lamotrigine and
23
24 seizure aggravation in severe myoclonic epilepsy. *Epilepsia*. 1998;39(5):508-12.
25
- 26
27 15. Xu X, Zhang Y, Sun H, Liu X, Yang X, Xiong H, et al. Early clinical features and diagnosis
28
29 of Dravet syndrome in 138 Chinese patients with SCN1A mutations. *Brain Dev*.
30
31 2014;36(8):676-81.
32
- 33
34 16. Berecki G, Bryson A, Terhag J, Maljevic S, Gazina EV, Hill SL, et al. SCN1A gain of
35
36 function in early infantile encephalopathy. *Ann Neurol*. 2019;85(4):514-525.
37
- 38
39 17. Beck VC, Hull JM, Isom LL. Beyond Dravet Syndrome: Characterization of a Novel, More
40
41 Severe SCN1A-Linked Epileptic Encephalopathy. *Epilepsy Curr*. 2019;19(4):266-268.
42
- 43
44 18. Cestèle S, Schiavon E, Rusconi R, Franceschetti S, Mantegazza M. Nonfunctional NaV1.1
45
46 familial hemiplegic migraine mutant transformed into gain of function by partial rescue of
47
48 folding defects. *Proc Natl Acad Sci U S A*. 2013;110(43):17546-51.
49
- 50
51 19. Dhifallah S, Lancaster E, Merrill S, Leroudier N, Mantegazza M, Cestèle S. Gain of
52
53 Function for the *SCN1A/hNav1.1-L1670W* Mutation Responsible for Familial Hemiplegic
54
55 Migraine. *Front Mol Neurosci*. 2018;11:232.
56
- 57
58 20. Zuberi SM, Wirrell E, Yozawitz E, Wilmschurst JM, Specchio N, Riney K, et al. ILAE
59
60 classification and definition of epilepsy syndromes with onset in neonates and infants:

- 1
2
3 Position statement by the ILAE Task Force on Nosology and Definitions. *Epilepsia*
4
5 2022;63(6):1349-97.
6
7
8 21. Fisher RS, Cross JH, French JA, Higurashi N, Hirsch E, Jansen FE, et al. Operational
9
10 classification of seizure types by the International League Against Epilepsy: Position Paper
11
12 of the ILAE Commission for Classification and Terminology. *Epilepsia*. 2017;58(4):522-
13
14 530.
15
16
17 22. Antoniadis T, Buxton C, Dennis G, Forrester N, Smith D, Lunt P, et al. Application of
18
19 targeted multi-gene panel testing for the diagnosis of inherited peripheral neuropathy
20
21 provides a high diagnostic yield with unexpected phenotype-genotype variability. *BMC Med*
22
23 *Genet*. 2015;16:84.
24
25
26 23. Association of Clinical Genetics Science: Practice Guidelines for the Evaluation of
27
28 Pathogenicity and the Reporting of Sequence Variants in Clinical Molecular Genetics. 2013.
29
30 http://www.acgs.uk.com/media/774853/evaluation_and_reporting_of_sequence_variants_bp
31
32 [gs_june_2013_-_finalpdf.pdf](http://www.acgs.uk.com/media/774853/evaluation_and_reporting_of_sequence_variants_bp). Accessed 2 Apr 2015.
33
34
35 24. Mantegazza M, Yu FH, Powell AJ, Clare JJ, Catterall WA, Scheuer T. Molecular
36
37 determinants for modulation of persistent sodium current by G-protein betagamma subunits.
38
39 *J Neurosci*. 2005;25(13):3341-9.
40
41
42 25. Cestèle S, Scalmani P, Rusconi R, Terragni B, Franceschetti S, Mantegazza M. Self-limited
43
44 hyperexcitability: functional effect of a familial hemiplegic migraine mutation of the Nav1.1
45
46 (SCN1A) Na⁺ channel. *J Neurosci*. 2008;28(29):7273-83.
47
48
49 26. Cestèle S, Labate A, Rusconi R, Tarantino P, Mumoli L, Franceschetti S, et al. Divergent
50
51 effects of the T1174S SCN1A mutation associated with seizures and hemiplegic migraine.
52
53 *Epilepsia*. 2013;54(5):927-35.
54
55
56 27. Mantegazza M, Gambardella A, Rusconi R, Schiavon E, Annesi F, Cassulini RR, et al.
57
58 Identification of an Nav1.1 sodium channel (SCN1A) loss-of-function mutation associated
59
60 with familial simple febrile seizures. *Proc Natl Acad Sci U S A*. 2005;102(50):18177-82.

- 1
2
3 28. Chever O, Zerimech S, Scalmani P, Lemaire L, Pizzamiglio L, Loucif A, et al. Initiation of
4 migraine-related cortical spreading depolarization by hyperactivity of GABAergic neurons
5 and NaV1.1 channels. *J Clin Invest*. 2021;131(21):e142203.
6
7
8
9
10 29. Alame S, El-Houwayek E, Nava C, Sabbagh S, Fawaz A, Gillart AC, et al. Dravet
11 Syndrome in Lebanon: First Report on Cases with *SCN1A* Mutations. *Case Rep Med*.
12 2019;2019:5270503.
13
14
15
16 30. Nair P, Sabbagh S, Mansour H, Fawaz A, Hmaimess G, Noun P, et al. Contribution of next
17 generation sequencing in pediatric practice in Lebanon. A Study on 213 cases. *Mol Genet*
18 *Genomic Med*. 2018;6(6):1041-1052.
19
20
21
22 31. Harkin LA, McMahon JM, Iona X, Dibbens L, Pelekanos JT, Zuberi SM, et al. The
23 spectrum of *SCN1A*-related infantile epileptic encephalopathies. *Brain*. 2007;130(Pt 3):843-
24 52.
25
26
27
28
29
30 32. Butler KM, da Silva C, Alexander JJ, Hegde M, Escayg A. Diagnostic Yield From 339
31 Epilepsy Patients Screened on a Clinical Gene Panel. *Pediatr Neurol*. 2017;77:61-66.
32
33
34 33. Gokben S, Onay H, Yilmaz S, Atik T, Serdaroglu G, Tekin H, et al. Targeted next
35 generation sequencing: the diagnostic value in early-onset epileptic encephalopathy. *Acta*
36 *Neurol Belg*. 2017;117(1):131-138.
37
38
39
40 34. Cetica V, Chiari S, Mei D, Parrini E, Grisotto L, Marini C, et al. Clinical and genetic factors
41 predicting Dravet syndrome in infants with *SCN1A* mutations. *Neurology*.
42 2017;88(11):1037-1044.
43
44
45
46 35. Wang JW, Shi XY, Kurahashi H, Hwang SK, Ishii A, Higurashi N, et al. Prevalence of
47 *SCN1A* mutations in children with suspected Dravet syndrome and intractable childhood
48 epilepsy. *Epilepsy Res*. 2012;102(3):195-200.
49
50
51
52 36. Patino GA, Claes LR, Lopez-Santiago LF, Slat EA, Dondeti RS, Chen C, et al. A functional
53 null mutation of *SCN1B* in a patient with Dravet syndrome. *J Neurosci*. 2009;29(34):10764-
54 78.
55
56
57
58
59
60

- 1
2
3 37. Terragni B, Scalmani P, Franceschetti S, Cestè S, Mantegazza M. Post-translational
4
5 dysfunctions in channelopathies of the nervous system. *Neuropharmacology*. 2018;132:31-
6
7 42.
8
9
10 38. Ohashi T, Akasaka N, Kobayashi Y, Magara S, Kawashima H, Matsumoto N, et al. Infantile
11
12 epileptic encephalopathy with a hyperkinetic movement disorder and hand stereotypies
13
14 associated with a novel SCN1A mutation. *Epileptic Disord*. 2014;16(2):208-12.
15
16
17 39. Lim BC, Hwang H, Kim H, Chae JH, Choi J, Kim KJ, et al. Epilepsy phenotype associated
18
19 with a chromosome 2q24.3 deletion involving SCN1A: Migrating partial seizures of infancy
20
21 or atypical Dravet syndrome? *Epilepsy Res*. 2015;109:34-9.
22
23
24 40. Spagnoli C, Frattini D, Rizzi S, Salerno GG, Fusco C. Early infantile SCN1A epileptic
25
26 encephalopathy: Expanding the genotype-phenotype correlations. *Seizure*. 2019;65:62-64.
27
28
29 41. Jaber D, Gitiaux C, Blesson S, Marguet F, Buard D, Varela Salgado M, et al. De novo
30
31 mutations of *SCN1A* are responsible for arthrogryposis broadening the *SCN1A*-related
32
33 phenotypes. *J Med Genet*. 2021;58(11):737-742.
34
35 42. Clatot J, Parthasarathy S, Cohen S, McKee JL, Massey S, Somarowthu A, et al. SCN1A
36
37 gain-of-function mutation causing an early onset epileptic encephalopathy. *Epilepsia*. 2022
38
39 Oct 26. Epub ahead of print.
40
41
42 43. Bertelli S, Barbieri R, Pusch M, Gavazzo P. Gain of function of sporadic/familial
43
44 hemiplegic migraine-causing SCN1A mutations: Use of an optimized cDNA. *Cephalalgia*.
45
46 2019;39(4):477-488.
47
48
49 44. Ohmori I, Kahlig KM, Rhodes TH, Wang DW, George AL Jr. Nonfunctional SCN1A is
50
51 common in severe myoclonic epilepsy of infancy. *Epilepsia* 2006;47(10):1636-42.
52
53
54 45. Kluckova D, Kolnikova M, Lacinova L, Jurkovicova-Tarabova B, Foltan T, Demko V, et al.
55
56 A Study among the Genotype, Functional Alternations, and Phenotype of 9 SCN1A
57
58 Mutations in Epilepsy Patients. *Sci Rep*. 2020;10(1):10288.
59
60

- 1
2
3 46. Ragsdale DS. How do mutant Nav1.1 sodium channels cause epilepsy? *Brain Res Rev*
4
5 2008;58(1):149-59.
6
7
8 47. Escayg A, Goldin AL. Sodium channel SCN1A and epilepsy: mutations and mechanisms.
9
10 *Epilepsia* 2010;51(9):1650-8
11
12 48. Salgueiro-Pereira AR, Duprat F, Pousinha P, Loucif A, Douchamps V, Regondi C, et al. A
13
14 two-hit story: seizures and genetic mutation interaction sets phenotype severity in SCN1A
15
16 epilepsies. *Neurobiol Dis* 2019;125:31-44.
17
18
19 49. Rhodes TH, Lossin C, Vanoye CG, Wang DW, George ALJr. Noninactivating
20
21 voltage-gated sodium channels in severe myoclonic epilepsy of infancy. *Proc Natl*
22
23 *Acad Sci U. S. A.* 2004;101:11147-52.
24
25
26 50. Fang ZX, Hong SQ, Li TS, Wang J, Xie LL, Han W et al. Genetic and phenotypic
27
28 characteristics of SCN1A-related epilepsy in Chinese children. *Neuroreport.* 2019 Jun
29
30 12;30(9):671-680.
31
32
33 51. Ottman R, Hirose S, Jain S, Lerche H, Lopes-Cendes I, Noebels JL et al. Genetic testing in
34
35 the epilepsies--report of the ILAE Genetics Commission. *Epilepsia.* 2010;51(4):655-70
36
37
38 52. Lossin C, Rhodes TH, Desai RR, Vanoye CG, Wang D, Carniciu S, et al. Epilepsy-
39
40 associated dysfunction in the voltage-gated neuronal sodium channel SCN1A. *J Neurosci*
41
42 2003;23:11289-95.
43
44
45 53. Spampinato J, Kearney J A, de Haan G, McEwen D P, Escayg A, Aradi I, et al. A novel
46
47 epilepsy mutation in the sodium channel SCN1A identifies a cytoplasmic domain for beta
48
49 subunit interaction. *J Neurosci* 2004;24(44):10022-34.
50
51
52 54. Lemaire L, Desroches M, Krupa M, Pizzamiglio L, Scalmani P, Mantegazza M. Modeling
53
54 NaV1.1/SCN1A sodium channel mutations in a microcircuit with realistic ion concentration
55
56 dynamics suggests differential GABAergic mechanisms leading to hyperexcitability in
57
58 epilepsy and hemiplegic migraine. *PLoS Comput Biol.* 2021;17(7):e1009239.
59
60

- 1
2
3 55. Tamim I, Chung DY, de Morais AL, Loonen ICM, Qin T, Misra A, et al. Spreading
4 depression as an innate antiseizure mechanism. *Nat Commun.* 2021;12(1):2206.
5
6
7 56. Hammer MF, Pan Y, Cumbay M, Pendziwiat M, Afawi Z, Goldberg-Stern H, et al. Whole
8 exome sequencing and co-expression analysis identify an SCN1A variant that modifies
9 pathogenicity in a family with genetic epilepsy and febrile seizures plus. *Epilepsia.*
10 2022;63(8):1970-1980.
11
12
13 57. Clatot J, Hoshi M, Wan X, Liu H, Jain A, Shinlapawittayatorn K, et al. Voltage-gated
14 sodium channels assemble and gate as dimers. *Nat Commun.* 2017;8(1):2077.
15
16
17 58. Silvennoinen K, Gawel K, Tsortouktzidis D, Pitsch J, Alhusaini S, van Loo KMJ, et al.,
18 SCN1A overexpression, associated with a genomic region marked by a risk variant for a
19 common epilepsy, raises seizure susceptibility. *Acta Neuropathol.* 2022 May 12
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Figure 1.** Representative whole-cell sodium current families for hNa_v1.1-WT (**A**), hNa_v1.1-T162I
4 (**B**), hNa_v1.1-T1501A (**C**), hNa_v1.1-R1892L (**D**) recorded with 100-ms-long depolarizing voltage
5 steps from -80 to +60 mV in 5 mV increments from a holding potential of -100 mV. (**E**) Current
6 density calculated for cells transfected with the hNa_v1.1-WT or with the mutant channels. (**F**) Mean
7 voltage dependence of activation, the lines are mean Boltzmann fits. Comparison of the voltage of
8 half activation $V_{1/2}$ (**G**) and of the activation slope factor, k_a (**H**) obtained from the Boltzmann fits.
9
10 (**I**) Mean voltage dependence of inactivation, the lines are Boltzmann fits. Comparison of the voltage
11 of half inactivation (**J**) and of the inactivation slope factor, k_h , (**K**) obtained from the Boltzmann fits.
12
13 (**L**) Time constant (τ in ms) of the recovery from fast inactivation at -80 mV, obtained from the
14 exponential fits of recovery curves.

15
16
17
18
19
20
21
22
23
24
25
26
27
28 **Figure 2.** Comparison of mean normalized currents elicited with a 150-ms-long depolarizing step to
29 -10 mV from a holding potential of -100 mV. (**A**) hNa_v1.1-WT (black), hNa_v1.1-T162I (red), (**B**)
30 hNa_v1.1-WT (black), hNa_v1.1-T1501A (blue), (**C**) hNa_v1.1-R1892L (green). The left insets show a
31 9-ms window of the current traces in order to compare the activation kinetics and the current decay;
32 the right insets show the traces between 70 and 80 ms to compare the I_{NaP} . (**D**) Comparison of the
33 time constants (τ in ms) of the current decay at -10 mV (single exponential fits at the indicated
34 potential). (**E**) Comparison of the persistent current quantified as mean current between 70 and 80 ms
35 of the 150-ms-long depolarizing step (expressed as % of I_T max).

36
37
38
39
40
41
42
43
44
45
46
47
48
49 **Figure 3.** Action sodium currents recorded using as voltage stimulus an action potential discharge of
50 a GABAergic fast spiking neuron, shown in panel (**A**). Action Na^+ currents for hNa_v1.1-WT (**B**),
51 hNa_v1.1-T162I (**C**), hNa_v1.1-T1501A (**D**), hNa_v1.1-R1892L (**E**); traces are presented as mean current
52 densities (CD), for clarity error bars are not shown. (**F**) Comparison of the peak current density of the
53 first action current. (**G**) Comparison of the peak current density at the end of the discharge (mean of
54 the 3 last action currents).

1
2
3 **Figure S1 in Supplemental Material.** The pedigrees of inherited variants.
4
5
6
7

8 **Figure S2 in Supplemental Material.** Current density(CD)-voltage curves of peak sodium current
9 for hNa_v1.1-WT, hNa_v1.1-T162I, hNa_v1.1-T1501A and hNa_v1.1-R1892L. Data points are displayed
10 as mean±SEM.
11
12
13
14
15
16

17 **Figure S3 in Supplemental Material.** Action sodium currents recorded using as voltage stimulus an
18 action potential discharge of a GABAergic fast spiking neuron, shown in panel (A), in tsA-201 cells
19 co-expressing hβ1 and hβ2 subunits. Action Na⁺ currents for hNa_v1.1-WT+ hβ1+hβ2 (B), hNa_v1.1-
20 T162I+ hβ1+hβ2 (C), hNa_v1.1-T1501A+ hβ1+hβ2 (D), hNa_v1.1-R1892L+ hβ1+hβ2 (E); traces are
21 displayed as mean current densities (CD), for clarity error bars are not shown. (F) Comparison of the
22 peak current density of the first action current. (G) Comparison of the peak current density at the end
23 of the discharge (mean of the 3 last action currents).
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Clinical and genetic features of the 16 novel, unpublished, patients

Patient ID	SCN1A variant	Current age & Gender	Age at sz onset	Follow-up	Sz type	Epilepsy Syndrome	ASMs prescribed & other treatments	Effective treatments	Sz off-set	Development, other features
1 *	c.469G>A; p.V157I; familial	19m M	1 m	14m	F motor	Early Infantile FE	PB	PB	2m; sz freedom no ASM	Mild DD
2	c.485C>T; p.T162I	13y 8m F	1 d	13y 8m	F motor; F to bilateral; Tonic; Myoclonic; Spasms	Neonatal DEE	PB, CBZ, CLB, VGB, VPA, LEV, ESM, PER	CBZ	Sz reduction	Severe ID, Spastic quadriplegia
3	c.677C>T; p.T226M	8m M	2.5 m	5.5m	F motor; F to bilateral; Tonic; Myoclonic; F dysautonomic; spasms; SE	Early Infantile DEE	PB, Pyridoxine, CBZ, NZP, PHT, LEV, VPA, CLB, STP	VPA+CLB+STP	Sz reduction	Severe DD, Hypotonia, Hyperkinetic MD
4	c.1172C>T; p.T391I	10y 1m F	36 m	7y	F motor; F to bilateral;	Childhood FE	VPA, LEV, CBZ, ACZ, CLB, ESM, TPM	CBZ, TPM	Sz freedom	Mild ID
5	c.1259C>A; p.A420D	12m F	2 m	10m	F motor; F to bilateral; F dysautonomic; Migrating; Tonic; Myoclonic, SE	Early Infantile DEE	Pyridoxine, CBZ, LEV, LCM, MDZ, PHT, VGB, VPA, ACTH, KD, PB, STR, CLB	CLB, PB, VPA	Sz reduction (last sz 3/12/20 due to VPA discontinuation)	Severe DD, Hyperkinetic MD
6 *	c.4012A>C; p.N1338H; familial	39y F	11 y	15y	F motor; F to bilateral; reflex sz	Childhood FE	CLB, CNZ, CBZ, OXC, VGB, ZNS, PER, LCS, PB, PHT, LEV, ESM, LTG, VPA, TPM; FLB, VGB, KD, VNS, Epilepsy Surgery	CBZ, VPA, KD	Sz reduction	Normal Cognitive functioning
7	c.4501 A>G; p.T1501A	24y M	9 m	23y 3m	F motor; F to bilateral	Infantile onset FE	VPA, LEV, LCM, CBZ	LCM, VPA	Transitory sz freedom	Borderline cognitive functioning
8	c.4907 G>A; p.R1636Q	1y 8m F	1 m	1y 7m	F motor; Tonic; Myoclonic, F dysautonomic	Early Infantile DEE	Pyridoxine; PB, VPA, CBZ, CLB, ACTH	CBZ, ACTH	Prolonged Sz freedom	Moderate DD, Hyperkinetic MD, Hypotonia
9	c.4907 G>C; p.R1636P + c.3386C>T; p.T129M	1y 3m F	2 d	1y 3m	F motor; Tonic; Myoclonic; F dysautonomic; Migrating; SE	Neonatal DEE	Pyridoxine; PLP, PB, LEV, PHT, MDZ, Thiopental, Ketamine, LCM, VPA, CLB	LCM, VPA	Sz reduction	Severe DD, Hypotonia
10	c.4907 G>A; p.R1636Q	14y 3m M	2 d	14y 1m	F motor; F to bilateral; Tonic; Myoclonic	Neonatal DEE	PB, CBZ, CLB, VPA, CNZ, HC, LEV, ESM, LTG	CBZ, VPA	Sz reduction	Severe ID, Pyramidal signs
11	c.5006C>A; p.A1669E	10y 3m M	5 m	9y 10m	F motor; F to bilateral; Tonic; Myoclonic, Spasms	Early infantile DEE	Pyridoxine, LEV, TPM, VGB, VPA, CBZ, ACTH, HC, PB, ESM, CLB	None	Ongoing sz	Severe ID, Spastic quadriplegia, Hyperkinetic MD
12	c.5020G>C; p.G1674C	Deceased at 27y; M	2 m	26y10m	F motor; F to bilateral; Tonic; Myoclonic, SE	Early infantile DEE	Pyridoxine, LEV, PB, LTG, VPA, CNZ	None	Ongoing sz	Severe ID, Spastic quadriplegia
13*		10y 11m M	5 y	5y 11m	F motor; F to bilateral;	Childhood FE	CBZ	CBZ	Sz freedom	Normal Cognitive functioning; Migraine
14*	c.5675G>T p.R1892L; familial	8y 6m F	4 y	4y 6m	F motor	Childhood FE	CBZ	CBZ	Sz freedom	Normal Cognitive functioning; Migraine
15*		43y 1m M	10 y	33y 1m	F motor; F to bilateral;	Childhood FE	PB, CBZ	CBZ	Sz freedom	Normal Cognitive functioning; Migraine
16	c.5732 A>G; p.K1911R	8y 9m F	3 m	8y 6m	F motor; F to bilateral; spasms	Early Infantile DEE	VGB, ACTH, VPA, NTZ, HC, TPM, CLB, PDN, PER, CNZ, LZP, LTG, KD, LCM	ACTH, CBZ, LCM	Sz reduction	Moderate ID, ADHD

Abbreviations: ACTH: adrenocorticotrophic hormone; ACZ: acetazolamide; ADHD: attention deficit hyperactivity disorder; ASM: antiseizure medication; CBZ: carbamazepine; CLB: clobazam; CNZ: clonazepam; d: days; DD: developmental delay; DEE: developmental epileptic encephalopathy; ESM: ethosuximide; F: focal; FE: focal epilepsy; FLB:

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

felabamate; HC: hydrocortisone; ID: intellectual disability; KD: ketogenic diet; m: months; MD: movement disorder; MDZ: midazolam; LCM: lacosamide; LEV: levetiracetam; LTG: lamotrigine; LZP: lorazepam; NTZ: nitrazepam; PB: phenobarbital; PDN: prednisone; PER: perampanel; PLP: pyridoxal phosphate; Sz: seizures; TPM: topiramate; VPA: valproate; VGB: vigabatrin; VNS: vagus nerve stimulation; y: years. *= familial pathogenic SCN1A variants

For Review Only

	WT	T162I	T1501A	R1892L
Current Density (pA/pF)	153.6±20.3 (n=16)	147.0±11.7 (n=11)	115.0±10.5 (n=13)	111.1±15.6 (n=9)
V_a (mV)	-27.6±1.2 (n=16)	-34.2±0.9 (n=11) p = 3*10 ⁻⁴	-30.3±0.8 (n=13)	-24.9±1.2 (n=9)
K_a (mV)	6.9±0.4 (n=16)	6.7±0.3 (n=11)	7.2±0.4 (n=13)	7.6±0.5 (n=9)
V_h (mV)	-63.3±0.9 (n=13)	-61.7±1.1 (n=11)	-53.8±1.1 (n=13) p = 3*10 ⁻⁷	-55.6±1.3 (n=9) p = 9*10 ⁻⁵
K_h (mV)	6.3±0.3 (n=13)	4.9±0.2 (n=11) p = 0.008	5.5±0.3 (n=13)	6.8±0.4 (n=9)
τ Recovery from Fast Inactivation at -80mV (ms)	3.1±0.3 (n=6)	2.1±0.2 (n=6)	0.93±0.04 (n=9) p = 3*10 ⁻⁶	2.7±0.4 (n=6)
τ Decay at -10mV (ms)	0.53±0.02 (n=16)	0.48±0.02 (n=11)	0.81±0.04 (n=13) p < 3*10 ⁻¹⁰	0.53±0.02 (n=9)
I_{NaP} (%I_{NaTmax})	1.6±0.3 (n=16)	2.2±0.4 (n=11)	6.3±0.3 (n=13) p < 10 ⁻¹⁰	1.5±0.2 (n=9)
CD First peak AP discharge (pA/pF)	-30.9±6.7 (n=9)	-50.5±10.3 (n=8)	-53.4±5.2 (n=9)	-41.2±5.3 (n=9)

			<i>(p=0.054)</i>	
Mean CD last 3 peaks AP discharge (pA/pF)	-4.9±0.6 (n=9)	-11.5±1.7 (n=9) p = 0.027	-12.7±1.8 (n=9) p = 0.004	-10.2±1.7 (n=9) <i>(p=0.09)</i>
CD First peak AP discharge with β1 & β2 (pA/pF)	-32.6±7.3 (n=7)	-50.9±5.9 (n=6)	-56.1±5.6 (n=7) <i>(p=0.062)</i>	-43.4±6.2 (n=7)
Mean CD last 3 peaks AP discharge with β1 & β2 (pA/pF)	-5.2±0.9 (n=7)	-12.1±1.7 (n=6) p = 0.029	-15.2±2.1 (n=7) p = 0.0007	-11.2±1.4 (n=7) p=0.048

Table 2. Functional properties of sodium currents recorded from tsA-201 cells. In bold the statistically significant differences in comparison with the WT (see Figures for all the statistical comparisons).

Table 3 Clinical finding of 42 probands reported in literature

	Males	24
	Females	18
	Age at seizure onset	Median: 5 weeks (range: 1 day–16 weeks)
	Age at the time of the study	15 months (range: 14 days–19 years)
	Length of the follow-up after seizure onset	13 months (range 21 days–18.8 years)
Epilepsy type	<i>Neonatal DEE</i>	17: 40.4%
	<i>Early infantile DEE</i>	25: 59.5%
Seizure Types	<i>Focal motor</i>	15: 35.7%
	<i>Focal to bilateral</i>	15: 35.7%
	<i>Generalized Tonic-Clonic</i>	14: 33.3%
	<i>Tonic</i>	32: 76.1%
	<i>Myoclonic</i>	20: 47.6%
	<i>Focal with dysautonomic features</i>	8: 19%
	<i>Spasms</i>	5: 11.9%
	<i>Migrating</i>	11: 26.2%
	Seizures triggered by fever	13: 30.9%
	Status Epilepticus	28: 66.6%
	SUDEP	3: 7.1%
EEGs	Focal epileptiform activity	17: 40.4%
	Multifocal epileptiform activity	31: 73.8%
	Migrating pattern	7: 16.6%
	Diffuse epileptiform activity	12: 28.5%
Brain MRI	Normal	27: 64.2%
	Nonspecific abnormalities: dysmorphic corpus callosum, small hippocampi, arachnoid cyst, pineal cyst	10: 23.8%
	Progressive brain atrophy	11: 26.2%
Cognitive functioning	Severe/Profound global impairment	41: 97.6%
Additional features	Hyperkinetic MD	22: 52.3%
	Hypotonia	20: 47.6%
	Microcephaly	11: 26.2%
	Arthrogyriposis	11: 26.2%
	Stiffness, clubfeet, hip dysplasia, multiple contractures, and talipes	1: 2.3%




Figure 1. Representative whole-cell sodium current families for hNav1.1-WT (A), hNav1.1-T162I (B), hNav1.1-T1501A (C), hNav1.1-R1892L (D) recorded with 100-ms-long depolarizing voltage steps from -80 to +60 mV in 5 mV increments from a holding potential of -100 mV. (E) Current density calculated for cells transfected with the hNav1.1-WT or with the mutant channels. (F) Mean voltage dependence of activation, the lines are mean Boltzmann fits. Comparison of the voltage of half activation $V_{1/2}$ (G) and of the activation slope factor, k_a (H) obtained from the Boltzmann fits. (I) Mean voltage dependence of inactivation, the lines are Boltzmann fits. Comparison of the voltage of half inactivation (J) and of the inactivation slope factor, k_h , (K) obtained from the Boltzmann fits. (L) Time constant (τ in ms) of the recovery from fast inactivation at -80 mV, obtained from the exponential fits of recovery curves.

272x208mm (300 x 300 DPI)




Figure 2. Comparison of mean normalized currents elicited with a 150-ms-long depolarizing step to -10 mV from a holding potential of -100 mV. (A) hNav1.1-WT (black), hNav1.1-T162I (red), (B) hNav1.1-WT (black), hNav1.1-T1501A (blue), (C), hNav1.1-R1892L (green). The left insets show a 9-ms window of the current traces in order to compare the activation kinetics and the current decay; the right insets show the traces between 70 and 80 ms to compare the I_{NaP} . (D) Comparison of the time constants (τ in ms) of the current decay at -10 mV (single exponential fits at the indicated potential). (E) Comparison of the persistent current quantified as mean current between 70 and 80 ms of the 150-ms-long depolarizing step (expressed as % of I_r max).

272x208mm (300 x 300 DPI)




Figure 3. Action sodium currents recorded using as voltage stimulus an action potential discharge of a GABAergic fast spiking neuron, shown in panel (A). Action Na^+ currents for hNav1.1-WT (B), hNav1.1-T162I (C), hNav1.1-T1501A (D), hNav1.1-R1892L (E); traces are presented as mean current densities (CD), for clarity error bars are not shown. (F) Comparison of the peak current density of the first action current. (G) Comparison of the peak current density at the end of the discharge (mean of the 3 last action currents).

120x85mm (300 x 300 DPI)