



Ferrocifen loaded Nanoparticles: Targeting MDR tumors via the Thioredoxin Reductase Pathway

Milad Baroud, Elise Lepeltier, Anne Vessieres, Nicolas Clere, Gérard Jaouen,
Catherine Passirani

► To cite this version:

Milad Baroud, Elise Lepeltier, Anne Vessieres, Nicolas Clere, Gérard Jaouen, et al.. Ferrocifen loaded Nanoparticles: Targeting MDR tumors via the Thioredoxin Reductase Pathway. SFNano 2022, Dec 2022, Strasbourg, France. 130, pp.146 - 153, 2008, 10.1016/j.jconrel.2008.05.027 . hal-04127365

HAL Id: hal-04127365

<https://hal.science/hal-04127365>

Submitted on 14 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Ferrocifen loaded Nanoparticles: Targeting MDR tumors via the Thioredoxin Reductase Pathway

Milad BAROUD¹, Elise LEPELTIER¹, Anne VESSIERES², Nicolas CLERE¹, Gérard JAOUEN², Catherine PASSIRANI^{1,*}

¹ Micro et Nanomédecines Translationnelles, MINT, UNIV Angers, UMR INSERM 1066, UMR CNRS 6021, 4 Rue Larrey, 49333, Angers, France

² Sorbonne Université, CNRS, IPCM, 4 Place Jussieu, 75005, Paris, France


*corresponding author : catherine.passirani@univ-angers.fr

Introduction

Regardless of the leaps being achieved in the field of cancer treatment, 90% of the failures of chemotherapy arise from the multidrug resistance aspect gained by the tumors¹. Aiming to overcome these limitations, ferrocifens, a family of molecules obtained chemically by combination of one ferrocene with hydroxytamoxifen derivatives², encapsulated in lipid nanocapsules, could offer an innovative and promising approach to combat this resistance. Moreover, the functionalization of the obtained nanoparticles with polyethylene glycol would imbue them with stealth properties³, increasing their half-life and blood circulation time, while utilizing the enhanced permeability and retention effect (EPR) to improve the internalization into cancer cells⁴.

Ferrocifen has been hypothesized to act via targeting the thioredoxin reductase (TrxR), a system responsible for thiol redox homeostasis and overexpressed in cancer cells, leading to an increased resistance^{5,6}. This action might stem from TrxR displaying a selenocysteine residue at its C-terminal active site, which can be targeted by metal complexes. This project, along-side producing ferrocifen loaded nanoparticles, aims to verify and establish the mode of action of this active substance on TrxR, using different genetic and molecular experimental techniques.

1. Formulation and characterization of the Lipid Nanocapsules (LNCs)



	Diameter (nm)	PDI	Zeta Potential (mV)
Blank LNC	53.2 ± 0.3	0.09 ± 0.01	-2.8 ± 0.5
Blank LNC-PEG	57.4 ± 0.3	0.09 ± 0.02	-20.3 ± 2.8
P15 LNC	51.0 ± 0.2	0.05 ± 0.01	16.2 ± 1.5
P15 LNC-PEG	55.5 ± 0.2	0.07 ± 0.01	-5.5 ± 0.7
P53 LNC	50.0 ± 0.3	0.02 ± 0.01	-0.5 ± 0.7
P53 LNC-PEG	56.0 ± 0.7	0.02 ± 0.01	-26.3 ± 0.5
P722 LNC	49.0 ± 0.1	0.06 ± 0.01	-2.5 ± 0.7
P722 LNC-PEG	52.4 ± 0.2	0.07 ± 0.09	-17.0 ± 0.8

2. Cytotoxicity studies via MTT

A549		IC ₅₀ in μM							Hep G2		IC ₅₀ in μM								
Auranofin	P15	P53	P722	Blank LNC-PEG	P15 LNC-PEG	P53 LNC-PEG	P722 LNC-PEG	A549	TXNRD2	TXNRD1	GPX4	Auranofin	P15	P53	P722	Blank LNC-PEG	P15 LNC-PEG	P53 LNC-PEG	P722 LNC-PEG
4h	18	17	106	187	75	0.03	509	49	9	1452	134	8.8	18	97	89	85	126	1188	94
24h	7.3	6.7	10	184	10	47	17	6.2	9	1452	134	3.18	1.86	9.1	72	12	14	13	20
48h	2.9	4.3	7.7	485	4.8	0.75	9	0.79	1.7	1452	134	1.7	1.6	12	9.1	10	10	15	28
72h	3.6	1.69	1.55	189	6.95	0.76	4.3	0.3	6.4	1.8	9	4.8	10.7	6.3	6.9	28	28	28	28
SK-HEP-1		IC ₅₀ in μM							Calu-3		IC ₅₀ in μM								
Auranofin	P15	P53	P722	Blank LNC-PEG	P15 LNC-PEG	P53 LNC-PEG	P722 LNC-PEG	SK-HEP-1	TXNRD2	TXNRD1	GPX4	Auranofin	P15	P53	P722	Blank LNC-PEG	P15 LNC-PEG	P53 LNC-PEG	P722 LNC-PEG
4h	4	21	182	93	19	97	108	103	35	733	192	12	19	93	93	122	183	79.86	14
24h	1.59	6.17	17	19	7.63	11	22	28	2.4	20	6.6	12.4	70	68.83	92	0.2	155	155	155
48h	1.54	13	14	19	6.34	11	22	29	1.7	6.6	0.7	3.8	20	11	14	6	101	101	101
72h	1.55	12	1.23	15	2.2	7.5	31	17	2.3	5.9	6.2	3.6	9.8	9.5	8.7	3.6	3.6	3.6	3.6

Conclusion

- P722 LNC-PEG is the best acting ferrocifen formulation with the lowest IC₅₀ at 72 hours.
- Encapsulation of P722 in LNCs enhances its action starting from earlier time points.
- P722 LNC-PEG has a 630-fold increased activity after encapsulation into LNC.
- P722 LNC-PEG has a 55-fold increased activity on HIGH compared to MEDIUM TrxR1 cell lines (A549 vs SKHEP-1).
- P722 LNC-PEG have an 63-fold increases activity on HIGH compared to LOW TrxR1 cell lines (A549 vs HepG2).

Perspectives

- Annexin V / PI apoptosis studies via FACS
- Ferroptosis studies via MTT
- Lipid peroxidation studies via FACS
- Western blot to determine the expression levels of keys proteins
- QRT-PCR to determine the expression levels of keys genes

- In vivo studies on PDX murine models
- Ex vivo studies on tissues / organs extracted from PDX models

References:

- B. Mansoori et al. Adv. Pharm. Bull. 2017. <https://doi.org/10.15171/apb.2017.041>
- G. Jaouen et al. Chem. Soc. Rev. 2015. <https://doi.org/10.1039/C5CS00486A>
- E. Allard et al. J. Control. Release. 2008. <https://doi.org/10.1016/j.jconrel.2008.05.027>
- E. Lepeitier et al. Drug Resist. Updat. 2020. <https://doi.org/10.1016/j.drup.2020.100704>
- A. Saei et al. Redox Biol. 2020. <https://doi.org/10.1016/j.redox.2020.101491>