



**HAL**  
open science

## Pulsating bubbles with nanoparticle shells

Nicole Anderton, Craig Stuart Carlson, Michiel Postema

► **To cite this version:**

Nicole Anderton, Craig Stuart Carlson, Michiel Postema. Pulsating bubbles with nanoparticle shells. Workshop on Bionanomaterial and Nanomaterial Engineering, Jun 2023, Łódź, Poland. hal-04130586

**HAL Id: hal-04130586**

**<https://hal.science/hal-04130586v1>**

Submitted on 16 Jun 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Pulsating bubbles with nanoparticle shells




Nicole Anderton, Craig S. Carlson, Michiel Postema


Antibubbles are microscopic gas bubbles comprising one or multiple liquid or solid cores, typically surrounded by stabilising shells. Hydrophobised silica nanoparticles are used to create this stabilised shell, a process called Pickering-stabilising. Pickering-stabilised antibubbles possess acoustic properties required to act as both diagnostic and therapeutic agents. This research aimed to determine the dynamic response of Pickering-stabilised antibubbles to pulsed ultrasound and identify and quantify the contribution of the shell and core to that behaviour.

## Aims

- 1 Quantify oscillation asymmetry of antibubbles.
- 2 Determine the influence of endoskeletal and shell friction on antibubble pulsation phase.
- 3 Predict the fragmentation threshold of antibubbles



Schematic representation of an endoskeletal antibubble.




Scanning electron micrograph of an antibubble

## Methods

### Experiments

Radial excursions of antibubbles during ultrasound exposure were extracted from high-speed footage. Medically safe conditions were maintained throughout the experiments; with an ultrasonic frequency of 1MHz and peak negative pressure of either 200kPa or 1MPa.



### Simulations

#### Fundamental pulsation

$$R\ddot{R} + \frac{3}{2}\dot{R}^2 = \frac{1}{\rho} \left[ \left( p_0 - p_v + \frac{2\sigma R_0}{R_0^2 - R_c^2} \right) \left( \frac{R_0^3 - R_c^3}{R^3 - R_c^3} \right)^\gamma + p_v - \frac{2\sigma R}{R^2 - R_c^2} - 2\chi \left( \frac{1}{R_0} - \frac{1}{R} \right) - \frac{4\eta\dot{R}}{R} - p_0 - P(t) \right]$$

#### Pulsation phase


$$\delta = \delta_v + \delta_r + \delta_{\text{sh}} + \delta_{\text{sk}} + \delta_{\text{g}}$$

$$\alpha = f(\delta, R_0, \omega)$$


#### Point of fragmentation

$$E_k \approx 2\pi\rho\dot{R}^2R^3 > \Delta E_s \approx (N^{\frac{1}{3}} - 1) \cdot 4\pi R^2\sigma$$



## Results



Scatter plot of initial resting radius ( $R_0$ ) versus oscillation asymmetry ( $\xi^+ - \xi^-$ ) as a result of a single cycle of ultrasound at pulse pressure amplitudes of (a) 200 kPa and (b) 1 MPa for antibubbles with cores (●) and without (○). The dotted purple line (---) represents perfectly symmetrical excursion, where  $(\xi^+ - \xi^-) = 0$ .



Pulsation phase with respect to incident sound field as a function of  $R_0$  for a free gas bubble (---), an antibubble with no core (—), and antibubbles with core content having endoskeletal damping coefficients equal to 0.3 (—), 0.6 (—), 0.9 (—) and 2 (—).



Fragmentation threshold as a function of driving frequency for two different core volume ratios (a,b). Shell stiffnesses are represented by different line colours:  $\chi = 0 \text{ N m}^{-1}$  (—),  $\chi = 1.1 \text{ N m}^{-1}$  (—),  $\chi = 7.6 \text{ N m}^{-1}$  (—), and  $\chi = 15 \text{ N m}^{-1}$  (—). High-speed video frames are shown before (c) and after (d) fragmentation.

## Conclusions

1. Antibubbles oscillate stably and symmetrically at low amplitudes, and with unique asymmetry at higher amplitudes.
2. The damping properties of the shell and skeletal materials of minute antibubbles may alter the degree to which a particle oscillates in phase with the sound field.
3. As the rigidity of the antibubble is increased, either through the addition of core content or a stiffer shell, the wall velocity during excursion decreases, reducing the kinetic energy of the shell, ultimately increasing the fragmentation threshold.

