



HAL
open science

Robust and non asymptotic estimation of probability weighted moments with application to extreme value analysis

Anna Ben-Hamou, Philippe Naveau, Maud Thomas

► **To cite this version:**

Anna Ben-Hamou, Philippe Naveau, Maud Thomas. Robust and non asymptotic estimation of probability weighted moments with application to extreme value analysis. 2023. hal-04131622

HAL Id: hal-04131622

<https://hal.science/hal-04131622>

Preprint submitted on 16 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Robust and non asymptotic estimation of probability weighted moments with application to extreme value analysis

Anna BEN-HAMOU¹, Philippe NAVEAU² and Maud THOMAS¹

¹ Sorbonne Université, CNRS, Laboratoire de Probabilités, Statistique et Modélisation,
LPSM, 4 place Jussieu, F-75005 Paris, France,
² Laboratoire des Sciences du Climat et de l'Environnement, UMR8212
CEA-CNRS-UVSQ, IPSL & Université Paris-Saclay, 91191 Gif-sur-Yvettes,
France
E-mails: anna.ben_hamou@sorbonne-universite.fr,
philippe.naveau@lsce.ipsl.fr,
maud.thomas@sorbonne-universite.fr

Abstract

In extreme value theory and other related risk analysis fields, probability weighted moments (PWM) have been frequently used to estimate the parameters of classical extreme value distributions. This method-of-moment technique can be applied when second moments are finite, a reasonable assumption in many environmental domains like climatological and hydrological studies. Three advantages of PWM estimators can be put forward: their simple interpretations, their rapid numerical implementation and their close connection to the well-studied class of U -statistics. Concerning the later, this connection leads to precise asymptotic properties, but non asymptotic bounds have been lacking when off-the-shelf techniques (Chernoff method) cannot be applied, as exponential moment assumptions become unrealistic in many extreme value settings. In addition, large values analysis is not immune to the undesirable effect of outliers, for example, defective readings in satellite measurements or possible anomalies in climate model runs. Recently, the treatment of outliers has sparked some interest in extreme value theory, but results about finite sample bounds in a robust extreme value theory context are yet to be found, in particular for PWMs or tail index estimators. In this work, we propose a new class of robust PWM estimators, inspired by the median-of-means framework of Devroye et al. [2016]. This class of robust estimators is shown to satisfy a sub-Gaussian inequality when the assumption of finite second moments holds. Such non asymptotic bounds are also derived under the general contamination model. Our main proposition confirms theoretically the trade-off between efficiency and robustness pointed out by Brazauskas

and Serfling [2006]. Our simulation study indicates that, while classical estimators of PWMs can be highly sensitive to outliers, our new approach remains weakly affected by the degree contamination.

Key words: Probability weighted moments; Concentration inequalities; Robustness; Extreme Value Analysis.

1 Introduction

Let X be an integrable real-valued random variable with cumulative distribution function F . The probability weighted moments (PWMs) of X are defined as

$$E(XF(X)^r\bar{F}(X)^s)$$

where r and s are non-negative integers, and $\bar{F} = 1 - F$ denotes the survival function associated with F . The use of these moments have been motivated by hydrologists and applied statisticians [see, e.g. Hosking and Wallis, 1987, Landwehr et al., 1979, Greenwood et al., 1979]. They also appear naturally in the expression of the parameters of several distributions used in extreme value theory [see, e.g. de Haan and Ferreira, 2006]. For example, if F corresponds to a generalized extreme value distribution with shape parameter $\xi \in \mathbb{R}$, then

$$\frac{3E(XF^2(X)) - E(X)}{2E(XF(X)) - E(X)} = \frac{3^\xi - 1}{2^\xi - 1},$$

and a similar formula is available for the generalized Pareto distribution.

Such moment equalities provide simple building blocks to quickly and efficiently implement a method-of-moment to estimate both generalized extreme value distribution or generalized Pareto parameters. Two main approaches have been used to infer PWMs. The first one consists in replacing the function F by its empirical version and taking the mean over the sample. The second one takes advantage of the link between PWMs and order statistics. More precisely, if (X_1, \dots, X_m) is an independent and identically distributed (i.i.d.) sample with common distribution function F , and if $X_{(1:m)} \leq \dots \leq X_{(m:m)}$ is the ordered sample, then a simple calculation shows that, for all $1 \leq k \leq m$,

$$\theta_{k:m} = E(X_{(k:m)}) = k \binom{m}{k} E(XF(X)^{k-1}\bar{F}(X)^{m-k}).$$

This indicates that, in the i.i.d. setting, the estimation of PWMs can be deduced from the order statistics. A natural choice for estimating $\theta_{k:m}$, with $1 \leq k \leq m \leq n$, is thus to use the unbiased estimator

$$\frac{1}{\binom{n}{m}} \sum_{1 \leq i_1 < \dots < i_m \leq n} \Psi_k(X_{i_1}, \dots, X_{i_m}) \quad (1.1)$$

where $\Psi_k(X_{i_1}, \dots, X_{i_m})$ corresponds to the k -th order statistic in the subsample $(X_{i_1}, \dots, X_{i_m})$. For instance, Landwehr et al. [1979] considered the special case of $k = 1$. Those two approaches are closely related and their asymptotic properties have been studied in detail [see, e.g. Hosking et al., 1985, Ferreira and de Haan, 2015, Diebolt et al., 2008, Guillou et al., 2009, Diebolt et al., 2003, 2004, 2007].

The literature on non asymptotic properties of PWM estimators is, to our knowledge, sparse. Furrer and Naveau [2007] derived explicit variance expressions for finite samples, but only in the case where the sample distribution is a generalized Pareto distribution. Estimators such as (1.1) have at least two drawbacks. First, in heavy-tailed scenarios where the underlying distribution has only low-order moments, estimate properties are not established for finite samples. In particular, classical concentration inequalities based on exponential decay of the tail cannot be directly applied to quantities like (1.1). Second, they may be extremely sensitive to the presence of outliers in the sample. The main motivation of this work is thus to design estimators of $\theta_{k:m}$ with good concentration properties under a second moment assumption only, and that would be robust to the presence of outliers.

Let us mention that the treatment of outliers for PWM estimation has rarely been covered within the extreme value theory community [see, e.g. Hubert et al., 2008, Dupuis and Victoria-Feser, 2006]. Reducing the negative impact of outliers, i.e. large corrupted anomalies, on the estimation of extreme value parameters demands a careful statistical analysis. Therefore, inference tools based on robust statistics [see e.g. Minsker and Wei, 2020, Hubert et al., 2008, Lecu e and Lerasle, 2019, Devroye et al., 2016] need to be adapted to extreme value theory. For example, Brazauskas and Serfling [2006] leveraged the concept of generalized quantiles to obtain favourable trade-offs between efficiency and robustness in the estimation of the parameters of a generalized Pareto distribution. Recently, Bhattacharya et al. [2019] studied a trimmed version of the Hill estimator to infer positive ξ and they proposed a methodology to identify extreme outliers in heavy-tailed data. Bhattacharya and Beirlant [2019] extended their work to light tail distributions and built a tail-adjusted boxplot. Still, all these studies focused on developing asymptotic distributions for their estimators, but non asymptotic bounds were not obtained.

To derive concentration bounds without exponential moment assumption and to achieve robustness, we propose, in Section 2, to adapt the so-called median-of-means concentration technique [see e.g. Devroye et al., 2016, Joly and Lugosi, 2016, Lecu e and Lerasle, 2019, 2020] to the estimation of $\theta_{k:m}$. Our estimator is actually defined in the much more general context of estimating the mean of symmetric multivariate kernels, when usual U -statistics may not give reliable estimates. In Section 3, we establish non asymptotic performance bounds, for degenerate and non-degenerate kernels, with sharp variance proxys. In addition, we show that our estimator is strongly robust

to the presence of outliers in the sample, under a very generic contamination scheme introduced by Lecué and Lerasle [2019]. Section 4 combines the problem of tail index estimation with our robust median-of-means inference scheme. In Section 5, numerical experiments are used to compare the classical PWM approach with our method. All proofs can be found in Section 6. In the appendix, the bounds derived in Section 3 are generalized beyond the i.i.d. setting by considering exchangeable sequences satisfying a negative dependence condition known as conditional negative association.

2 Median-of-means estimators

In this section, we recall the median-of-means techniques and the construction of the associated estimators. As its name suggests, a median-of-means estimator is obtained as the median of means, the latter being, in this work, computed as U -statistics on independent blocks of the original given sample.

In the sequel, we assume that samples are all independent and identically distributed, unless otherwise specified. Let X_1, \dots, X_n be a sample with values in some measurable set \mathcal{X} . We are interested in the robust estimation of quantities of the form

$$\theta = E(\Psi(X_1, \dots, X_m)) \in \mathbb{R} \quad (2.1)$$

where, for an integer $m \geq 1$, $\Psi : \mathcal{X}^m \rightarrow \mathbb{R}$ is a symmetric function, called kernel. Let $v_k = \text{var}(E[\Psi(X_1, \dots, X_m) | X_1, \dots, X_k])$, the kernel Ψ is said to be q -degenerate, for $q \in \{1, \dots, m\}$, if $v_1 = \dots = v_{q-1} = 0$ and $v_q > 0$. If $v_1 > 0$, Ψ is said to be non-degenerate.

Assuming $n \geq m$, a natural estimator for θ is given by the following U -statistics

$$U_n = \frac{1}{\binom{n}{m}} \sum_{\substack{A \subset [n] \\ |A|=m}} \Psi(X_A),$$

where $[n] = \{1, \dots, n\}$ and $X_A = (X_i)_{i \in A}$. In the case of PWM, $\theta = \theta_{k:m} = E(X_{(k:m)})$ and $\Psi = \Psi_k$ corresponds to the k -th order statistics in a sample of size m .

As shown by Joly and Lugosi [2016], robust estimation of the parameter θ , defined in (2.1), can be obtained using median-of-means techniques. We present here a new estimator of θ based on these techniques. The main idea is to divide the sample (X_1, \dots, X_n) into K blocks, that is disjoint subsets. To fix this number K of blocks, the practitioner first needs to choose an error level $\delta \in [e^{-\lfloor n/m \rfloor}, 1[$ and then the integer K can be defined as $K = \lceil \log(1/\delta) \rceil$. By construction, $1 \leq K \leq \lfloor n/m \rfloor$ and the set $[n] = \{1, \dots, n\}$ is then divided into K disjoint blocks B_1, \dots, B_K , each of size $|B_j| \geq \lfloor n/K \rfloor \geq m$.

Within each block j , we construct the U-statistic estimator

$$\hat{\theta}^{(j)} = \frac{1}{\binom{|B_j|}{m}} \sum_{\substack{A \subset B_j \\ |A|=m}} \Psi(X_A),$$

and then compute the median among blocks, i.e.

$$\hat{\theta}_n = \text{median} \left(\hat{\theta}^{(1)}, \dots, \hat{\theta}^{(K)} \right), \quad (2.2)$$

where $\text{median}(z_1, \dots, z_K)$ corresponds to the smallest value $z \in \{z_1, \dots, z_K\}$ such that

$$|\{j \in [K], z_j \leq z\}| \geq K/2 \quad \text{and} \quad |\{j \in [K], z_j \geq z\}| \geq K/2.$$

Obtaining non asymptotic concentration inequalities for U_n with the correct variance factor turns out to be a difficult problem, even in the non-degenerate case. Hoeffding or Bernstein-type inequalities have been obtained, but they require the kernel Ψ to be bounded or to have sufficiently light tails. For instance, if $|\Psi| \leq c$, Maurer [2019] that, for all $t \geq 0$,

$$\text{pr}(|U_n - E(U_n)| \geq t) \leq 2 \exp \left(- \frac{nt^2}{2m^2v_1 + \frac{4m^4v_m}{n} + \frac{16m^2ct}{3}} \right)$$

[see also Arcones, 1995]. Previously, Hoeffding [1963] had shown a sub-Gaussian inequality for U_n .

However, when Ψ is heavy-tailed, exponential inequalities might not hold anymore. Since, PWM estimators are classically used to construct estimators in extreme value analysis. We are interested in deriving non asymptotic bounds with minimum moments conditions. In the next section, we show that our median-of-means estimator exhibits exponential concentration with the correct variance factor, see Proposition 3.1. We show also that this estimator is robust to the presence of outliers in the sample, in a very generic contamination scheme, see Proposition 3.3.

3 Sub-Gaussian and Bernstein-type bounds for median-of-means estimators

In this section, we derive sub-Gaussian and Bernstein-type bounds for general median-of-means estimators with non-degenerate or degenerate kernel. Then, the issue of the robustness to the presence of outliers in the sample is also addressed. The results are stated in a general context and PWM estimators simply correspond to a particular case.

Proposition 3.1. *Let X_1, \dots, X_n be an i.i.d. sample with values in \mathcal{X} , and, for a positive integer $m \in [n]$, let $\Psi : \mathcal{X}^m \rightarrow \mathbb{R}$ be a symmetric q -degenerate kernel, $q \in [m]$, with $E[\Psi(X_1, \dots, X_m)] = \theta$ and $v_m < \infty$. Then, for all $\delta \in [e^{-\lceil n/m \rceil}, 1)$, the median-of means estimator $\hat{\theta}_n$ defined by (2.2) with $K = \lceil \log(1/\delta) \rceil$ satisfies*

$$\text{pr} \left(|\hat{\theta}_n - \theta| > 2e \sqrt{\frac{\binom{m-1}{q-1} (2m)^q v_m \lceil \log(1/\delta) \rceil^q}{n^q}} \right) \leq \delta, \quad (3.1)$$

and

$$\text{pr} \left(|\hat{\theta}_n - \theta| > 2e \sqrt{\frac{\binom{m}{q} (2m)^q v_q \lceil \log(1/\delta) \rceil^q}{n^q} + \frac{\binom{m-1}{q} (2m)^{q+1} v_m \lceil \log(1/\delta) \rceil^{q+1}}{n^{q+1}}} \right) \leq \delta. \quad (3.2)$$

Bound (3.1) is similar to the one obtained by Joly and Lugosi [2016] for their estimator. Bound (3.2) is new and gives a significant improvement over (3.1), especially in the regime where $\log(1/\delta)/n = o(1)$. In that case, the second term under the square root can be neglected, and the variance factor in the first term is improved to

$$\frac{\binom{m}{q} (2m)^q v_q}{n^q},$$

which is close to the asymptotic variance obtained by Hoeffding [1948] which states that

$$\text{var}(U_n) \underset{n \rightarrow \infty}{\sim} q! \binom{m}{q}^2 \frac{v_q}{n^q},$$

for non-degenerate kernels, $\text{var}(U_n) \underset{n \rightarrow \infty}{\sim} m^2 v_1/n$, and U_n is asymptotically normal:

$$\sqrt{n}(U_n - \theta) \underset{n \rightarrow \infty}{\rightsquigarrow} \mathcal{N}(0, m^2 v_1).$$

In the case $q = 1$, i.e. in the non-degenerate case, then Equation (3.1) states that $\hat{\theta}_n - \theta$ is sub-Gaussian on both tails with variance factor proportional to mv_m/n :

$$\text{pr} \left(|\hat{\theta}_n - \theta| > 2e \sqrt{\frac{2mv_m \lceil \log(1/\delta) \rceil}{n}} \right) \leq \delta, \quad (3.3)$$

while Equation (3.2) states that it is sub-gamma on both tails with variance factor proportional to $m^2 v_1/n$ and scale factor proportional to $\sqrt{m^3 v_m/n}$:

$$\text{pr} \left(|\hat{\theta}_n - \theta| > 2e \sqrt{\frac{2m^2 v_1 \lceil \log(1/\delta) \rceil}{n} + \frac{4m^3 v_m \lceil \log(1/\delta) \rceil^2}{n^2}} \right) \leq \delta. \quad (3.4)$$

Since $mv_1 \leq v_m$, the variance factor in (3.4) is always smaller than the one in (3.3). Due to the scale factor in (3.4), either inequality might be better, depending on the value of δ .

Remark 3.2. *The estimator $\hat{\theta}_n$ depends on the pre-chosen confidence threshold δ . As shown by Devroye et al. [2016, Theorem 4.2], when an upper-bound v_m^* on v_m is available, an estimator $\tilde{\theta}_n$ independent of δ may be constructed. The notation $\hat{\theta}_{n,\delta}$ below highlights this connection. From Proposition 3.1, for all $k \in \{1, \dots, \lfloor n/m \rfloor\}$, the interval*

$$\hat{I}_k = \left[\hat{\theta}_{n,e^{-k}} \pm 2e \sqrt{\frac{2mv_m^* k}{n}} \right]$$

is a confidence interval with level $1 - e^{-k}$, where $\hat{\theta}_{n,e^{-k}}$ denote the estimator (2.2) obtained with $\delta = e^{-k}$. Now, let

$$\hat{k} = \min \left\{ 1 \leq k \leq \lfloor n/m \rfloor, \bigcap_{j=k}^{\lfloor n/m \rfloor} \hat{I}_j \neq \emptyset \right\},$$

and define the estimator $\tilde{\theta}_n$ as the midpoint of the interval $\bigcap_{j=\hat{k}}^{\lfloor n/m \rfloor} \hat{I}_j$. Then, the estimator $\tilde{\theta}_n$ satisfies, for all $\delta \in [e^{-\lfloor n/m \rfloor}/(1 - e^{-1}), 1[$,

$$\text{pr} \left(|\tilde{\theta}_n - \theta| > 4e \sqrt{\frac{2mv_m^* \left(1 + \log \left(\frac{1}{1-e^{-1}} \right) + \log \left(\frac{1}{\delta} \right) \right)}{n}} \right) \leq \delta. \quad (3.5)$$

Recall that our main goal was two-fold. First, we aim at proposing a family of estimators of θ for which sharp concentration bounds are available under minimal moment conditions. Then, we also wish to address the issue of robustness of these estimators. This is the purpose of the next result which concerns the robustness of the estimator $\hat{\theta}_n$ to the presence of outliers in the original sample. To translate the presence of outliers, we consider the contamination scheme introduced by Lecué and Lerasle [2019]: the index set $[n]$ is divided into two disjoint subsets, the subset \mathcal{I} of inliers, and the subset \mathcal{O} of outliers. The sequence $(X_i)_{i \in \mathcal{I}}$ is i.i.d. while no assumption is made on the variables $(X_i)_{i \in \mathcal{O}}$. In what follows, $\text{pr}_{\mathcal{O} \cup \mathcal{I}}$ corresponds to the distribution of such a contaminated sample.

Proposition 3.3. *Let (X_1, \dots, X_n) be a contaminated sample under the $\mathcal{O} \cup \mathcal{I}$ model. For all $\delta \in [e^{-\lfloor n/m \rfloor}, 1)$, let $\hat{\theta}_n$ be defined as in (2.2) with $K = \lceil \log(1/\delta) \rceil$. If $|\mathcal{O}| \leq K/4$, then*

$$\text{pr}_{\mathcal{O} \cup \mathcal{I}} \left(|\hat{\theta}_n - \theta| > \frac{16e^2}{3\sqrt{3}} \sqrt{\frac{2mv_m \lceil \log(1/\delta) \rceil}{n}} \right) \leq \delta,$$

and

$$\Pr_{\mathcal{O} \cup \mathcal{I}} \left(|\hat{\theta}_n - \theta| > \frac{16e^2}{3\sqrt{3}} \sqrt{\frac{2m^2 v_1 \lceil \log(1/\delta) \rceil}{n} + \frac{4m^3 v_m \lceil \log(1/\delta) \rceil^2}{n^2}} \right) \leq \delta.$$

To obtain non asymptotic bounds for $\theta_{k:m}$, it suffices to take $\Psi = \Psi_k$, that is the k -th order statistic in a sample of size m , which is symmetric and non-degenerate. This introduces our new class of robust estimators $\hat{\theta}_{k:m}$ of $\theta_{k:m} = E(X_{(k:m)})$ as defined in (2.2) with $K = \lceil \log(1/\delta) \rceil$ for some $\delta \in [e^{-\lfloor n/m \rfloor}, 1[$. Thus, $\hat{\theta}_{k:m}$ satisfies sub-Gaussian and Bernstein-type inequalities for uncontaminated or contaminated samples, that is $\hat{\theta}_{k:m}$ satisfies Propositions 3.1 and 3.3 with $v_m = \text{var}(X_{(k:m)})$ and $v_1 = \text{var}(E(X_{(k:m)}|X_1))$.

4 Sub-Gaussian and Bernstein-type bounds for the median-of-means PWM estimator of the tail index

Considering a sample X_1, \dots, X_n distributed according to a generalized extreme value distribution G_ξ . As mentioned in the introduction, the parameter ξ can be linked to PWM. Here, we consider the following explicit expression proposed by Ferreira and de Haan [2015, Remark 2.2]:

$$\xi = \frac{1}{\log 2} \log \left(\frac{4E(XG_\xi^3(X)) - 2E(XG_\xi(X))}{2E(XG_\xi(X)) - E(X)} \right) = \frac{1}{\log 2} \log \left(\frac{\theta_4 - \theta_2}{\theta_2 - \theta_1} \right), \quad (4.1)$$

where $\theta_j = \theta_{j:j} = E(\max(X_1, \dots, X_j))$. Therefore, a natural estimator of ξ is obtained by simply estimating the θ_j , $j = 1, 2, 4$ in (4.1) and a median-of-means PWM estimator of ξ is thus given by

$$\hat{\xi}_n = \frac{1}{\log 2} \log \left(\frac{\hat{\theta}_4 - \hat{\theta}_2}{\hat{\theta}_2 - \hat{\theta}_1} \right), \quad (4.2)$$

where $\hat{\theta}_j$ be the median-of-means estimate θ_j for $j \in \{1, 2, 4\}$ constructed with $K = \lceil \log(1/\delta) \rceil$ blocks for $\delta \in [e^{-\lfloor n/4 \rfloor}, 1[$. The next result shows that ξ_n is sub-Gaussian on both tails for uncontaminated or contaminated samples.

Proposition 4.1. *Let X_1, \dots, X_n be a sample with distribution function F . Then,*

$$\Pr \left(|\hat{\xi}_n - \xi| \geq \frac{2e(2^\xi + 1)(2^{-\xi} + 2^{-\xi})}{\log(2)(\hat{\theta}_2 - \hat{\theta}_1)} \sqrt{\frac{8 \max(v_1, v_2, v_4) \lceil \log(1/\delta) \rceil}{n}} \right) \leq 3\delta,$$

where $v_j = \text{var}(\max(X_1, \dots, X_j))$.

Moreover, if the sample X_1, \dots, X_n is contaminated by outliers with the same scheme of Proposition 3.3, then

$$\text{pr}_{\mathcal{O} \cup \mathcal{I}} \left(|\hat{\xi}_n - \xi| \geq \frac{16e^2(2^\xi + 1)(2^{-\xi} + 2^{-\xi})}{3 \log(2)\sqrt{3}(\hat{\theta}_2 - \hat{\theta}_1)} \sqrt{\frac{8 \max(v_1, v_2, v_4) \lceil \log(1/\delta) \rceil}{n}} \right) \leq 3\delta.$$

In extreme value theory, statisticians are interested in the estimation of this parameter ξ which reflects the heaviness of the tail of the distribution. Our median-of-mean estimator thus provide a robust and with good finite sample properties estimation of such quantity.

5 Numerical experiments

This section illustrates the robustness of our median-of-means PWM estimators $\hat{\theta}_{k:m}$ and $\hat{\xi}$. We consider the following setting. We simulate 1 000 generalized extreme value samples contaminated by outliers. For that purpose, we first draw samples of inliers of size $n_{\mathcal{I}}$ distributed according a generalized extreme value distribution with 3 different values for the tail index $\xi = -0.4, 0$ and 0.4 . These samples are then contaminated by $n_{\mathcal{O}}$ outliers, such that $n = n_{\mathcal{I}} + n_{\mathcal{O}}$ constant equal to 200. To simulate the outliers, we distinguish two cases: $\xi < 0$ and $\xi \geq 0$. When $\xi < 0$, the probability that a generalized extreme value variable with parameter $\xi < 0$ exceeds any value $-1/\xi$ is equal to 0. The outliers $X_i, i = 1, \dots, n_{\mathcal{O}}$ are then obtained as

$$X_i \sim \text{Unif}[0, 20 - \frac{1}{\xi}], \quad i = 1, \dots, n_{\mathcal{O}}.$$

When $\xi \geq 0$, The outliers $X_i, i = 1, \dots, n_{\mathcal{O}}$ are obtained

$$X_i \sim \mathcal{N}(q_\xi(1 - 10^{-4}), 1), \quad i = 1, \dots, n_{\mathcal{O}}.$$

where $q_\xi(1 - 10^{-4})$ denotes the $(1 - 10^{-4})$ -quantile of a generalized extreme value of parameter ξ .

Let us highlight that when the kernel $\Psi = \Psi_k$, the procedure to build $\hat{\theta}_{k:m}$ does not require to solve an optimisation problem (in contrast to maximum likelihood techniques) and is computationally straightforward to implement. In this special case, computing the U -statistics $\hat{\theta}^{(j)}$ within bloc B_j does not require going through all the m subsets of B_j . Instead, we may notice that

$$\frac{1}{\binom{n}{m}} \sum_{\substack{A \subset [n] \\ |A|=m}} \Psi_k(X_A) = \frac{1}{\binom{n}{m}} \sum_{i=1}^n \binom{n-i}{m-k} \binom{i-1}{k-1} X_{(i:n)} := T_{k:m}. \quad (5.1)$$

For comparison, we also consider the estimator $T_{k:m}$ of $\theta_{k:m}$ which corresponds to the classical U -statistics estimate over the whole sample (taking




Figure 1: Estimates of a) $\theta_{3:4}$, b) $\theta_{4:4}$, c) ξ and d) $q_\xi(0.95)$ obtained with the median-of-mean estimator (MM, in blue, see Equation (4.2)), and the estimator computed as a linear combination (LC, in red, see Equation (5.1)), when the shape parameter ξ is equal to -0.4, 0 and 0.4, in columns, and the number of outliers n_O is equal to 0, 5, 15 and 20, in rows. In each graph, the blue dotted line corresponds to the true value of the quantity to be estimated.

$K = 1$ block). As a weighted mean of the order statistics, it is very sensitive to the presence of outliers.

In both cases, we estimate $\theta_{k:m}$, for $m = 1, \dots, 4$ and $k \leq m$, the parameter ξ and the 0.95-quantile, denoted $q_\xi(0.95)$, with the linear combination estimator $T_{k:m}$ and the median-of-means estimator $\hat{\theta}_{k:m}$. Figure 1 displays the boxplots of both estimators as the number of outliers $n_{\mathcal{O}}$ varies for $\theta_{3:4}$, $\theta_{4:4}$, ξ and $q_\xi(0.95)$ (the other graphs can be found along with the code online [git@github.com:maudmthomas/PWM.git](https://github.com:maudmthomas/PWM.git)). It can be seen that the classical estimator is highly sensitive to outliers. Our median-of-means estimator inherits the robustness properties of the median.

6 Proofs

To prove Proposition 3.1, we first establish the following lemma, which gives non asymptotic bounds on the variance of U -statistics.

Lemma 6.1. *Let $\Psi : \mathcal{X}^m \rightarrow \mathbb{R}$ be a symmetric q -degenerate kernel, with $q \in [m]$, such that $v_m < \infty$. For $n \geq m$, let X_1, \dots, X_n be an i.i.d. sample, and define*

$$U_n = \frac{1}{\binom{n}{m}} \sum_{\substack{A \subset [n] \\ |A|=m}} \Psi(X_A).$$

Then,

$$\text{var}(U_n) \leq \frac{\binom{m-1}{q-1} m^q v_m}{n^q}, \quad (6.1)$$

and

$$\text{var}(U_n) \leq \frac{\binom{m}{q} m^q v_q}{n^q} + \frac{\binom{m-1}{q} m^{q+1} v_m}{n^{q+1}}. \quad (6.2)$$

of Lemma 6.1. Recall the identity

$$\text{var}(U_n) = \frac{1}{\binom{n}{m}} \sum_{k=1}^m \binom{m}{k} \binom{n-m}{m-k} v_k. \quad (6.3)$$

Now if Ψ is q -degenerate, then $v_1 = \dots = v_{q-1} = 0$, and the sum above may be started at $k = q$. Hoeffding [1948, Theorem 5.1] showed that, for all $1 \leq k \leq \ell \leq m$,

$$\frac{v_k}{k} \leq \frac{v_\ell}{\ell}. \quad (6.4)$$

Hence, noting that $v_k \leq kv_m/m$,

$$\begin{aligned} \text{var}(U_n) &\leq \frac{1}{\binom{n}{m}} \sum_{k=q}^m \binom{m}{k} \binom{n-m}{m-k} \frac{k}{m} v_m \\ &= \frac{v_m}{\binom{n}{m}} \sum_{k=q-1}^{m-1} \binom{m-1}{k} \binom{n-m}{m-1-k} \\ &\leq \frac{v_m \binom{m-1}{q-1} \binom{n-q}{m-q}}{\binom{n}{m}}, \end{aligned}$$

where the last inequality comes from the fact that the number of ways to choose $m-1$ elements in a set of size $n-1$ with at least $q-1$ elements taken from a given subset of size $m-1$ is less than the number of ways to first choose $q-1$ elements in that subset, and then $m-q$ elements in what remains. Then, since

$$\frac{\binom{n-q}{m-q}}{\binom{n}{m}} \leq \left(\frac{m}{n}\right)^q,$$

we obtain

$$\text{var}(U_n) \leq \frac{\binom{m-1}{q-1} m^q v_m}{n^q},$$

establishing (6.1). The second bound (6.2) is obtained by singling out the term corresponding to $k=q$ in (6.3) and using $v_k \leq kv_m/m$ only for $k \geq q+1$. This yields

$$\text{var}(U_n) \leq \frac{\binom{m}{q} \binom{n-m}{m-q}}{\binom{n}{m}} v_q + \frac{1}{\binom{n}{m}} \sum_{k=q+1}^m \binom{m}{k} \binom{n-m}{m-k} \frac{k}{m} v_m.$$

Now, on the one hand,

$$\frac{\binom{m}{q} \binom{n-m}{m-q}}{\binom{n}{m}} \leq \binom{m}{q} \frac{\binom{n-q}{m-q}}{\binom{n}{m}} \leq \binom{m}{q} \left(\frac{m}{n}\right)^q.$$

On the other hand,

$$\begin{aligned} \frac{1}{\binom{n}{m}} \sum_{k=q+1}^m \binom{m}{k} \binom{n-m}{m-k} \frac{k}{m} &= \frac{1}{\binom{n}{m}} \sum_{k=q+1}^m \binom{m-1}{k-1} \binom{n-m}{m-k} \\ &= \frac{1}{\binom{n}{m}} \sum_{k=q}^{m-1} \binom{m-1}{k} \binom{n-m}{m-1-k} \\ &\leq \frac{\binom{m-1}{q} \binom{n-q-1}{m-q-1}}{\binom{n}{m}} \\ &\leq \binom{m-1}{q} \left(\frac{m}{n}\right)^{q+1}. \end{aligned}$$

Finally,

$$\text{var}(U_n) \leq \frac{\binom{m}{q} m^q v_q}{n^q} + \frac{\binom{m-1}{q} m^{q+1} v_m}{n^{q+1}}$$

establishing (6.2). \square

Before proving Proposition 3.1, we first state a well-known but useful fact. We include the proof for completeness.

Lemma 6.2. *Let (Y_1, \dots, Y_K) be independent Bernoulli random variables such that for all $j \in [K]$, we have $E(Y_j) \leq p$, for some $p \in (0, 1)$. Then, for all $a \in (0, 1)$,*

$$\text{pr} \left(\sum_{j=1}^K Y_j \geq aK \right) \leq \left(\frac{p}{a} \right)^{aK} \left(\frac{1-p}{1-a} \right)^{(1-a)K}.$$

In particular, for $a = 1/2$,

$$\text{pr} \left(\sum_{j=1}^K Y_j \geq K/2 \right) \leq \left(2\sqrt{p(1-p)} \right)^K \leq (2\sqrt{p})^K.$$

of Lemma 6.2. Let $\lambda \geq 0$, then

$$\log E \left(e^{\lambda \sum_{j=1}^K Y_j} \right) = \sum_{j=1}^K \log E \left(e^{\lambda Y_j} \right) = \sum_{j=1}^K \log \left(E(Y_j) e^{\lambda} + 1 - E(Y_j) \right).$$

Since $E(Y_j) \leq p$,

$$\log E \left(e^{\lambda \sum_{j=1}^K Y_j} \right) \leq K \log \left(p e^{\lambda} + 1 - p \right).$$

Next, using a Chernoff bound,

$$\text{pr} \left(\sum_{j=1}^K Y_j \geq aK \right) \leq \exp \left\{ -K \sup_{\lambda \geq 0} \left(\lambda a - \log(p e^{\lambda} + 1 - p) \right) \right\}.$$

Observing that the supremum is attained for $\lambda = \log(a(1-p)/(p(1-a)))$,

$$\text{pr} \left(\sum_{j=1}^K Y_j \geq aK \right) \leq \left(\frac{p}{a} \right)^{aK} \left(\frac{1-p}{1-a} \right)^{(1-a)K}.$$

\square

of Proposition 3.1. Let $t = \min\{t_1, t_2\}$, with

$$t_1 = 2e\sqrt{\frac{\binom{m-1}{q-1}(2m)^q v_m [\log(1/\delta)]^q}{n^q}}, \quad (6.5)$$

and

$$t_2 = 2e\sqrt{\frac{\binom{m}{q}(2m)^q v_q [\log(1/\delta)]^q}{n^q} + \frac{\binom{m-1}{q}(2m)^{q+1} v_m [\log(1/\delta)]^{q+1}}{n^{q+1}}}. \quad (6.6)$$

By definition of the median, both the number of j such that $\hat{\theta}^{(j)} \geq \hat{\theta}_n$ and the number of j such that $\hat{\theta}^{(j)} \leq \hat{\theta}_n$ are at least $K/2$. This leads to write

$$\text{pr}\left(|\hat{\theta}_n - \theta| > t\right) \leq \text{pr}\left(\sum_{j=1}^K Y_j \geq K/2\right),$$

where $Y_j = \mathbf{1}_{\{|\hat{\theta}^{(j)} - \theta| > t\}}$. We now look for an upper bound on $E(Y_j)$ so as to apply Lemma 6.2. By Chebyshev Inequality, for all $j \in \{1, \dots, m\}$,

$$E(Y_j) = \text{pr}\left(|\hat{\theta}^{(j)} - \theta| > t\right) \leq \frac{\text{var}\left(\hat{\theta}^{(j)}\right)}{t^2}.$$

When $t = t_1$, one may use the bound (6.1) in Lemma 6.1 with $|B_j|$ instead of n . Since $|B_j| \geq \lfloor n/K \rfloor \geq n/(2K)$,

$$E(Y_j) \leq \frac{\binom{m-1}{q-1} m^q v_m}{|B_j|^q t_1^2} \leq \frac{\binom{m-1}{q-1} (2mK)^q v_m}{n^q t_1^2} = \frac{1}{4e^2}.$$

When $t = t_2$, the bound (6.2) in Lemma 6.1 entails

$$E(Y_j) \leq \frac{\frac{\binom{m}{q} m^q v_q}{|B_j|^q} + \frac{\binom{m-1}{q} m^{q+1} v_m}{|B_j|^{q+1}}}{t_2^2} \leq \frac{\frac{\binom{m}{q} (2mK)^q v_q}{n^q} + \frac{\binom{m-1}{q} (2mK)^{q+1} v_m}{n^{q+1}}}{t_2^2} = \frac{1}{4e^2},$$

where as above, we used that $|B_j| \geq \lfloor n/K \rfloor \geq n/(2K)$. In both cases, Lemma 6.2 can be applied with $a = 1/2$ and $p = e^{-2}/4$ to obtain

$$\text{pr}\left(\sum_{j=1}^K Y_j \geq K/2\right) \leq e^{-K} \leq \delta.$$

□

Concerning the proof of Equation (3.5) in Remark 3.2. Let $\delta \in [e^{-\lfloor n/m \rfloor}/(1 - e^{-1}), 1[$ and let k_δ be the smallest integer $k \in \{1, \dots, \lfloor n/m \rfloor\}$ such that $\delta \geq e^{-k}/(1 - e^{-1})$. By a union bound,

$$\text{pr}\left(\bigcap_{j=k_\delta}^{\lfloor n/m \rfloor} \{\theta \in \hat{I}_j\}\right) \geq 1 - \sum_{j=k_\delta}^{\lfloor n/m \rfloor} e^{-j} \geq 1 - \frac{e^{-k_\delta}}{1 - e^{-1}} \geq 1 - \delta.$$

Now, on the event $\bigcap_{j=k_\delta}^{\lfloor n/m \rfloor} \{\theta \in \hat{I}_j\}$, one has $\bigcap_{j=k_\delta}^{\lfloor n/m \rfloor} \hat{I}_j \neq \emptyset$, hence $\hat{k} \leq k_\delta$. But if $\hat{k} \leq k_\delta$, then $\tilde{\theta}_n$ also belongs to $\bigcap_{j=k_\delta}^{\lfloor n/m \rfloor} \hat{I}_j$ and

$$|\tilde{\theta}_n - \theta| \leq 4e \sqrt{\frac{2mv_m^* k_\delta}{n}} \leq 4e \sqrt{\frac{2mv_m^* \left(1 + \log\left(\frac{1}{1-e^{-1}}\right) + \log\left(\frac{1}{\delta}\right)\right)}{n}},$$

which proves Equation (3.5).

of Proposition 3.3. Let $t = 8e \min\{t_1, t_2\}/(3\sqrt{3})$, with t_1 and t_2 as defined in (6.5) and (6.6). Then,

$$\text{pr}_{\mathcal{O} \cup \mathcal{I}} \left(|\hat{\theta}_n - \theta| > t \right) \leq \text{pr}_{\mathcal{O} \cup \mathcal{I}} \left(\sum_{j=1}^K Y_j \geq K/2 \right),$$

with $Y_j = \mathbf{1}_{\{|\hat{\theta}^{(j)} - \theta| > t\}}$. Letting \mathcal{B} be the set of blocks that do not intersect \mathcal{O} ,

$$\text{pr}_{\mathcal{O} \cup \mathcal{I}} \left(\sum_{j=1}^K Y_j \geq K/2 \right) \leq \text{pr}_{\mathcal{O} \cup \mathcal{I}} \left(|\mathcal{B}^c| + \sum_{j \in \mathcal{B}} Y_j \geq K/2 \right).$$

Since $|\mathcal{B}^c| \leq |\mathcal{O}| \leq K/4$ by assumption, we get

$$\text{pr}_{\mathcal{O} \cup \mathcal{I}} \left(|\hat{\theta}_n - \theta| > t \right) \leq \text{pr} \left(\sum_{j=1}^K Y_j \geq K/4 \right),$$

where pr , here, corresponds to the probability measure for an i.i.d. sample. By Chebyshev Inequality, we have

$$E(Y_j) \leq \frac{\text{var}(\hat{\theta}^{(j)})}{t^2}.$$

In the proof of Proposition 3.1, we have shown that

$$\frac{\text{var}(\hat{\theta}^{(j)})}{\min\{t_1, t_2\}^2} \leq \frac{1}{4e^2},$$

leading to $E(Y_j) \leq 3^3 2^{-8} e^{-4}$. We may thus apply Lemma 6.2 with $a = 1/4$ and $p = 3^3 2^{-8} e^{-4}$ to obtain

$$\text{pr} \left(\sum_{j=1}^K Y_j \geq K/4 \right) \leq e^{-K} \leq \delta.$$

□

of Proposition 4.1. Let

$$c_n(\delta) = \mathbf{c} \sqrt{\frac{8 \max(v_1, v_2, v_4) \lceil \log(1/\delta) \rceil}{n}}$$

with \mathbf{c} denoting either $2e$ or $16e^2/(3\sqrt{3})$ depending on whether the sample is contaminated or not. For $j = 1, 2, 4$, consider the event

$$E_j = \left\{ |\hat{\theta}_j - \theta_j| \leq c_n(\delta) \right\}.$$

On the one hand, from Propositions (3.1) and 3.3, $\Pr(E_1 \cap E_2 \cap E_4) \geq 1 - 3\delta$ and $\Pr_{\mathcal{O} \cup \mathcal{I}}(E_1 \cap E_2 \cap E_4) \geq 1 - 3\delta$ depending on the value of \mathbf{c} .

On the other hand,

$$\begin{aligned} 2^{\hat{\xi}} - 2^{\xi} &= \frac{\hat{\theta}_4 - \hat{\theta}_2}{\hat{\theta}_2 - \hat{\theta}_1} - \frac{\theta_4 - \theta_2}{\theta_2 - \theta_1} \\ &= \frac{(\hat{\theta}_4 - \hat{\theta}_2)(\theta_2 - \theta_1) - (\theta_4 - \theta_2)(\hat{\theta}_2 - \hat{\theta}_1)}{(\hat{\theta}_2 - \hat{\theta}_1)(\theta_2 - \theta_1)} \\ &= \frac{(\hat{\theta}_4 - \theta_4)(\theta_2 - \theta_1) - (\hat{\theta}_2 - \theta_2)(\theta_4 - \theta_1) + (\hat{\theta}_1 - \theta_1)(\theta_4 - \theta_2)}{(\hat{\theta}_2 - \hat{\theta}_1)(\theta_2 - \theta_1)}. \end{aligned}$$

On the event $E_1 \cap E_2 \cap E_4$,

$$|2^{\hat{\xi}} - 2^{\xi}| \leq \frac{2c_n(\delta)(\theta_4 - \theta_1)}{(\hat{\theta}_2 - \hat{\theta}_1)(\theta_2 - \theta_1)} = \frac{2c_n(\delta)(2^{\xi} + 1)}{\hat{\theta}_2 - \hat{\theta}_1}.$$

Using that for $a, b > 0$,

$$|\log(a) - \log(b)| \leq \frac{1}{2}|a - b| \left(\frac{1}{a} + \frac{1}{b} \right),$$

we obtain that, with probability larger than $1 - 3\delta$,

$$|\hat{\xi} - \xi| \leq \frac{c_n(\delta)}{\log(2)(\hat{\theta}_2 - \hat{\theta}_1)} (2^{\xi} + 1)(2^{-\hat{\xi}} + 2^{-\xi}).$$

□

A Concentration bounds under exchangeability and negative association

Definition A.1. A sequence of real-valued random variables (X_1, \dots, X_n) is said to be negatively associated if for all subset $A \subset [n]$, and for all (coordinate-wise) non-decreasing functions $f : \mathbb{R}^{|A|} \rightarrow \mathbb{R}$ and $g : \mathbb{R}^{n-|A|} \rightarrow \mathbb{R}$ such that the expectations below are well-defined, one has

$$E[f(X_A)g(X_{[n] \setminus A})] \leq E[f(X_A)]E[g(X_{[n] \setminus A})].$$

Definition A.2. A sequence of real-valued random variables (X_1, \dots, X_n) is said to be conditionally negatively associated (CNA) if for all $S \subset [n]$ and $A \subset S$, and for all non-decreasing functions $f : \mathbb{R}^{|A|} \rightarrow \mathbb{R}$ and $g : \mathbb{R}^{|S \setminus A|} \rightarrow \mathbb{R}$ such that the expectations below are well-defined, one has

$$E [f(X_A) g(X_{S \setminus A}) \mid X_S] \leq E [f(X_A) \mid X_S] E [g(X_{S \setminus A}) \mid X_S].$$

In other words, the sequence is CNA if it is negatively associated and all conditionalizations are negatively associated.

A immediate consequence is that sums of negatively associated random variables concentrate at least as well as sums of independent random variables with the same marginals. More precisely, the Laplace transform can be bounded by the product of marginal transforms: if (X_1, \dots, X_n) is NA, then for all $\lambda \in \mathbb{R}$,

$$E \left[e^{\lambda \sum_{i=1}^n X_i} \right] \leq \prod_{i=1}^n E \left[e^{\lambda X_i} \right].$$

Let us also mention an important proposition of NA.

Proposition A.3. If (X_1, \dots, X_n) is NA, if A_1, \dots, A_k are disjoint subsets of $[n]$, and if h_1, \dots, h_k are non-decreasing functions defined on $\mathbb{R}^{|A_1|}, \dots, \mathbb{R}^{|A_k|}$ respectively, then the sequence $(h_1(X_{A_1}), \dots, h_k(X_{A_k}))$ is NA.

The following result generalizes Proposition 3.1 to the case where the sample (X_1, \dots, X_n) is exchangeable and CNA. For simplicity, we only prove it in the non-degenerate case ($q = 1$). Notice that the symmetric function Ψ needs to be non-decreasing. Under this additional assumptions, the same bounds hold, with probability $1 - 2\delta$ instead of $1 - \delta$.

Proposition A.4. Let (X_1, \dots, X_n) be an exchangeable CNA sample, and, for $m \in [n]$, let $\Psi : \mathbb{R}^m \rightarrow \mathbb{R}$ be a non-decreasing symmetric function such that $E[\Psi(X_1, \dots, X_m)] = \theta$ and $\text{var}(\Psi(X_1, \dots, X_m)) < \infty$. For all $1 \leq k \leq m$, let

$$v_k = \text{var}(E[\Psi(X_1, \dots, X_m) \mid X_1, \dots, X_k]).$$

Then, for all $\delta \in [e^{-n/m}, 1)$, the median-of-means estimator $\hat{\theta}_n$ defined by (2.2) satisfies

$$\text{pr} \left(|\hat{\theta}_n - \theta| > 2e \sqrt{\frac{2mv_m \lceil \log(1/\delta) \rceil}{n}} \right) \leq 2\delta, \quad (\text{A.1})$$

and

$$\text{pr} \left(|\hat{\theta}_n - \theta| > 2e \sqrt{\frac{2m^2v_1 \lceil \log(1/\delta) \rceil}{n} + \frac{4m^3v_m \lceil \log(1/\delta) \rceil^2}{n^2}} \right) \leq 2\delta. \quad (\text{A.2})$$

Proof of Proposition A.4. Let $t = \min\{t_1, t_2\}$, with

$$t_1 = 2e\sqrt{\frac{2mv_m \lceil \log(1/\delta) \rceil}{n}}$$

and

$$t_2 = 2e\sqrt{\frac{2m^2v_1 \lceil \log(1/\delta) \rceil}{n} + \frac{4m^3v_m \lceil \log(1/\delta) \rceil^2}{n^2}}.$$

To deal with monotone events, we first have to decompose the absolute value:

$$\text{pr}\left(|\hat{\theta}_n - \theta| > t\right) = \text{pr}\left(\hat{\theta}_n - \theta > t\right) + \text{pr}\left(\hat{\theta}_n - \theta < -t\right).$$

Let us show that both terms on the right-hand side above are less than δ . We only detail the argument for $\text{pr}\left(\hat{\theta}_n - \theta > t\right)$. The term $\text{pr}\left(\hat{\theta}_n - \theta < -t\right)$ can be treated similarly. We have

$$\text{pr}\left(\hat{\theta}_n - \theta > t\right) \leq \text{pr}\left(\sum_{j=1}^K Y_j \geq K/2\right),$$

with $Y_j = \mathbf{1}_{\{\hat{\theta}^{(j)} - \theta > t\}}$. Since Ψ is non-decreasing, the sequence (Y_1, \dots, Y_K) is NA thanks to Property A.3. The bounds of Lemma 6.2 (which come from bounds on the Laplace transform) thus apply here as well, and it remains to verify that $E(Y_j) \leq \frac{1}{4e^2}$, as in the proof of Proposition 3.1. To that aim, it suffices to show that the variance bounds of Lemma 6.1 also holds in the exchangeable CNA setting, under the assumption that Ψ is non-decreasing. We have

$$\begin{aligned} \text{var}(U) &= \frac{1}{\binom{n}{m}^2} E \left[\left(\sum_{\substack{A \subset [n] \\ |A|=m}} (\Psi(X_A) - \theta) \right)^2 \right] \\ &= \frac{1}{\binom{n}{m}^2} \sum_{k=0}^m \sum_{\substack{A, B \subset [n] \\ |A|=|B|=m, |A \cap B|=k}} E [(\Psi(X_A) - \theta)(\Psi(X_B) - \theta)]. \end{aligned}$$

By exchangeability and CNA, and since Ψ is non-decreasing and symmetric, we have, for all subsets A and B of size m such that $|A \cap B| = k$,

$$\begin{aligned} E [(\Psi(X_A) - \theta)(\Psi(X_B) - \theta)] &= E \left[E [(\Psi(X_A) - \theta)(\Psi(X_B) - \theta) \mid X_{A \cap B}] \right] \\ &\leq E \left[(E [\Psi(X_1, \dots, X_m) \mid X_1, \dots, X_k] - \theta)^2 \right]. \end{aligned}$$

Since the number of subsets A and B of size m such that $|A \cap B| = k$ is equal to $\binom{n}{m} \binom{m}{k} \binom{n-m}{m-k}$, we arrive at

$$\text{var}(U) \leq \frac{1}{\binom{n}{m}} \sum_{k=0}^m \binom{m}{k} \binom{n-m}{m-k} v_k,$$

where $v_k = E \left[\left(E [\Psi(X_1, \dots, X_m) | X_1, \dots, X_k] - \theta \right)^2 \right]$. At this point, all the proof of Lemma 6.1 can be repeated, after checking that the proof of Inequality (6.4) in [Hoeffding, 1948] only requires exchangeability. \square

References

- M. A. Arcones. A Bernstein-type inequality for u-statistics and u-processes. *Statistics & probability letters*, 22(3):239–247, 1995.
- S. Bhattacharya and J. Beirlant. Outlier detection and a tail-adjusted boxplot based on extreme value theory, 2019.
- S. Bhattacharya, M. Kallitsis, and S. Stoev. Data-adaptive trimming of the Hill estimator and detection of outliers in the extremes of heavy-tailed data. *Electronic Journal of Statistics*, 13(1):1872 – 1925, 2019.
- V. Brazauskas and R. Serfling. Robust estimation of tail parameters for two-parameter Pareto and exponential models via generalized quantile statistics. *Extremes*, 3:231–249, 2006.
- L. de Haan and A. Ferreira. *Extreme value theory*. Springer-Verlag, 2006.
- L. Devroye, M. Lerasle, G. Lugosi, and R. I. Oliveira. Sub-gaussian mean estimators. *The Annals of Statistics*, 44(6):2695–2725, 2016.
- J. Diebolt, , A. Guillo, and R. Worms. Asymptotic behaviour of the probability-weighted moments and penultimate approximation. *E.S.A.I.M. Probab. Statist.*, pages 219–238, 2003.
- J. Diebolt, , A. Guillo, , and I. Rached. A new look at probability-weighted moments estimators. *C. R. Acad. Sci. de Paris, Sér. 7*, pages 629–634, 2004.
- J. Diebolt, A. Guillo, and I. Rached. Approximation of the distribution of excesses through a generalized probability-weighted moments method. *Journal of Statistical Planning and Inference*, 137:841 – 857, 2007.
- J. Diebolt, A. Guillo, P. Naveau, and P. Ribereau. Improving probability-weighted moment methods for the generalized extreme value distribution. *REVSTAT - Statistical Journal*, 6(1):33–50, 2008.

- D. J. Dupuis and M.-P. Victoria-Feser. A robust prediction error criterion for pareto modelling of upper tails. *Canadian Journal of Statistics*, 34(4): 639–658, 2006.
- A. Ferreira and L. de Haan. On the block maxima method in extreme value theory: PWM estimators. *The Annals of Statistics*, 43(1):276–298, feb 2015. doi: 10.1214/14-aos1280. URL <https://doi.org/10.1214/2014-aos1280>.
- R. Furrer and P. Naveau. Probability weighted moments properties for small samples. *Statistics and Probability Letters*, 77:190–195, 2007.
- J. A. Greenwood, J. M. Landwehr, N. C. Matalas, and J. R. Wallis. Probability weighted moments: definition and relation to parameters of several distributions expressible in inverse form. *Water resources research*, 15(5): 1049–1054, 1979.
- A. Guillou, P. Naveau, D. Diebolt, and P. Ribereau. Return level bounds for discrete and continuous random variables. *Test*, 18:584–604, 2009.
- W. Hoeffding. A class of statistics with asymptotically normal distribution. *Ann. Math. Statist.*, 19(3):293–325, 1948.
- W. Hoeffding. Probability inequalities for sums of bounded random variables. *Journal of the American Statistical Association*, 58(13-30), 1963.
- J. R. M. Hosking and J. R. Wallis. Parameter and quantile estimation for the generalized Pareto distribution. *Technometrics*, 29:339–349, 1987.
- J. R. M. Hosking, J. R. Wallis, and E. F. Wood. Estimation of the generalized extreme-value distribution by the method of probability-weighted moments. *Technometrics*, 27:251–261, 1985.
- M. Hubert, P. J. Rousseeuw, and S. V. Aelst. High-Breakdown Robust Multivariate Methods. *Statistical Science*, 23(1):92 – 119, 2008.
- E. Joly and G. Lugosi. Robust estimation of u-statistics. *Stochastic Processes and their Applications*, 126(12):3760–3773, 2016.
- J. Landwehr, N. Matalas, and J. Wallis. Probability weighted moments compared with some traditional techniques in estimating Gumbel parameters and quantiles. *Water Resour. Res.*, 15:1055–1064, 1979.
- G. Lecué and M. Lerasle. Learning from mom’s principles: Le cam’s approach. *Stochastic Processes and their applications*, 129(11):4385–4410, 2019.
- G. Lecué and M. Lerasle. Robust machine learning by median-of-means: theory and practice. *The Annals of Statistics*, 48(2):906–931, 2020.

- A. Maurer. A bernstein-type inequality for functions of bounded interaction. *Bernoulli*, 25(2):1451–1471, 2019.
- S. Minsker and X. Wei. Robust modifications of U-statistics and applications to covariance estimation problems. *Bernoulli*, 26(1):694 – 727, 2020.