

Multi-decadal satellite monitoring of vegetation cover (MOD44B) in Fennoscandia: application to Saami' reindeer husbandry landscapes

Romain Courault, Marianne Cohen

▶ To cite this version:

Romain Courault, Marianne Cohen. Multi-decadal satellite monitoring of vegetation cover (MOD44B) in Fennoscandia: application to Saami' reindeer husbandry landscapes. XIXème édition des Journées scientifiques du Comité National Français des Recherches Arctique et Antarctique (CNFRAA), May 2023, Paris, France. hal-04132296

HAL Id: hal-04132296 https://hal.science/hal-04132296v1

Submitted on 19 Jun2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Multi-decadal satellite monitoring of vegetation cover (MOD44B) in Fennoscandia: application to Saami' reindeer





husbandry landscapes

R. Courault^{*}, M. Cohen^{**}





Mann Kendall S= ŽŽ SGN(3-7)

TYTE) if

V(S)=n(n-1)(2n+5), Z=

* Université Paris 1 Panthéon-Sorbonne, UMR 7533 Ladyss, GDR AREES, Labex Dynamite ****** Sorbonne-Université, UR Médiations INTRODUCTION

i. In subarctic & montane areas, large-scale surveys of vegetal cover are important to assess consequences of the increase in temperature & intensification of human activities

- ii. In Lapland for saami reindeer' herders, pastoral systems are unprecedently affected by global change, having regional to local consequences (be it biological, ecological, economical or cultural): herds' metabolism, pasture quality & availability, insect harassment, predators etc.
- iii. We propose a monitoring of the Gabna & Gällivare reindeer husbandry territories-landscapes- using MODIS 44B Vegetation continuous fields time series. Gabna' reindeer husbandry system is considered by Swedish and Saami authorities as a montane system, whereas Gällivare has forest features

MATERIAL & METHODS

- a) Vegetation cover time series (MOD44B Yearly Vegetation Cover Fields, 2000-2022; NASA-USGS)
- b) Declared territories of reindeer husbandry (Rengis 2.0)
- c) Computing spatiotemporal trend tests of Mann-Kendall (MK): Revegetation ? Devegetation? How much areas of territories are affected by such processes? Drivers?



d) Clipping and selecting significant trend tests (pvalue<0.05; MK>+0.4 & MK<-0.4)

HIGHLIGHTS & DISCUSSION

Forest cover : 26% of Gabna is significantly affected by forest changes, mostly de-vegetation (average - \bar{x} - of MK= -0.45). For Gällivare: 84%, $\bar{x} = -0.3$. Forestry is the main driver, clear-cuttings & pine recruitments, besides images' preprocessing refinements in detection of trees (coniferous & broadleaved) Non-forest: For this layer, 22 and



19% of Gabna and Gällivare surface areas are revegetating ($\bar{x} = +0.47$ & **Tundra'** +0.44,respectively). shrubification/extension of montane shrubs (Gabna) + pine' recruitment in stands for forested areas (Gabna & Gällivare) REFERENCES

Significant spatio-temporal trends for non-vegetated pixels, Gabna' community Swedish Saami Parliement, MODIS Continuous Vegetation Fields, 2000-2022; R. Courault, 2023

Non-vegetated: Differences between Gabna (montane reindeer herding, see map above) and Gällivare (forest): for the latter most of significant pixels shows a temporal increase of non-vegetated pixels $(\bar{x} = +0.4)$. Urban and mining areas are extending. Meanwhile, evolution of non-vegetated areas within Gabna shows heterogeneous processes, mentioned previously.

DIMICELI, Charlene, TOWNSHEND, John, CARROLL, Mark, et al. Evolution of the representation of global vegetation by vegetation by vegetation of global vegetation of global vegetation of global vegetation of global vegetation by vegetation by vegetation of global renbruksplaner 2005-2010. Skogsstyrelsen, 2011.; MAJASALMI, Titta et RAUTIAINEN, Miina. Representation of tree cover in global land cover products: Finland as a case study area. Environmental Monitoring and Assessment, 2021, vol. 193, no 3, p. 121.; TOWNSEND, J. et DIMICELI, study area. C. MOD44B MODIS/Terra vegetation continuous fields yearly L3 global 500m SIN grid. University of Maryland and MODAPS SIPS-NASA DOI, 2015, vol. 10.; Sàmediggi / Swedish Saami Parliement : https://www.sametinget.se/8382

Contact : Romain.Courault@univ-paris1.fr