



HAL
open science

Multi-decadal satellite monitoring of vegetation cover (MOD44B) in Fennoscandia: application to Saami' reindeer husbandry landscapes

Romain Courault, Marianne Cohen

► To cite this version:

Romain Courault, Marianne Cohen. Multi-decadal satellite monitoring of vegetation cover (MOD44B) in Fennoscandia: application to Saami' reindeer husbandry landscapes. XIXème édition des Journées scientifiques du Comité National Français des Recherches Arctique et Antarctique (CNFRAA), May 2023, Paris, France. hal-04132296

HAL Id: hal-04132296

<https://hal.science/hal-04132296v1>

Submitted on 19 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Multi-decadal satellite monitoring of vegetation cover (MOD44B) in Fennoscandia: application to Saami' reindeer husbandry landscapes



husbandry landscapes



R. Courault*, M. Cohen**

* Université Paris 1 Panthéon-Sorbonne, UMR 7533 Ladyss, GDR AREES, Labex Dynamite

** Sorbonne-Université, UR Médiations

INTRODUCTION

- i. In subarctic & montane areas, large-scale surveys of vegetal cover are important to assess consequences of the increase in temperature & intensification of human activities
- ii. In Lapland for saami reindeer' herders, pastoral systems are unprecedentedly affected by global change, having regional to local consequences (be it biological, ecological, economical or cultural): herds' metabolism, pasture quality & availability, insect harassment, predators etc.
- iii. We propose a monitoring of the Gabna & Gällivare reindeer husbandry territories-landscapes- using MODIS 44B Vegetation continuous fields time series. Gabna' reindeer husbandry system is considered by Swedish and Saami authorities as a montane system, whereas Gällivare has forest features

MATERIAL & METHODS


- a) Vegetation cover time series (MOD44B Yearly Vegetation Cover Fields, 2000-2022; NASA-USGS)
- b) Declared territories of reindeer husbandry (Rengis 2.0)
- c) Computing spatiotemporal trend tests of Mann-Kendall (MK): Revegetation ? Devegetation? How much areas of territories are affected by such processes? Drivers?
- d) Clipping and selecting significant trend tests (p-value < 0.05; MK > +0.4 & MK < -0.4)



HIGHLIGHTS & DISCUSSION

Forest cover : 26% of Gabna is significantly affected by forest changes, mostly de-vegetation (average \bar{x} of MK = -0.45). For Gällivare: 84%, $\bar{x} = -0.3$. Forestry is the main driver, clear-cuttings & pine recruitments, besides images' preprocessing refinements in detection of trees (coniferous & broadleaved)

Non-forest: For this layer, 22 and 19% of Gabna and Gällivare surface areas are revegetating ($\bar{x} = +0.47$ & $+0.44$, respectively). Tundra' shrubification/extension of montane shrubs (Gabna) + pine' recruitment in stands for forested areas (Gabna & Gällivare)



Non-vegetated: Differences between Gabna (montane reindeer herding, see map above) and Gällivare (forest): for the latter most of significant pixels shows a temporal increase of non-vegetated pixels ($\bar{x} = +0.4$). Urban and mining areas are extending. Meanwhile, evolution of non-vegetated areas within Gabna shows heterogeneous processes, mentioned previously.

REFERENCES