



**HAL**  
open science

# Subdivisions of four blocks cycles in digraphs with large chromatic number

Darine Al-Mniny

► **To cite this version:**

Darine Al-Mniny. Subdivisions of four blocks cycles in digraphs with large chromatic number. *Discrete Applied Mathematics*, 2021, 305, pp.71-75. 10.1016/j.dam.2021.08.005 . hal-04134238

**HAL Id: hal-04134238**

**<https://hal.science/hal-04134238>**

Submitted on 22 Jul 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

# Subdivisions of Four Blocks Cycles in Digraphs with Large Chromatic Number

Darine AL MNINY\*

*Baalbeck, Lebanon*

*KALMA Laboratory<sup>a,\*</sup>, Camille Jordan Institute<sup>b</sup>*

<sup>a</sup>*Department of Mathematics, Faculty of Sciences I, Lebanese University, Hadath, Lebanon*

<sup>b</sup>*Department of Mathematics, Claude Bernard University-Lyon 1, Villeurbanne, France*

---

## Abstract

A cycle with four blocks  $C(k_1, k_2, k_3, k_4)$  is an oriented cycle formed of four blocks of lengths  $k_1, k_2, k_3$  and  $k_4$  respectively. We conjecture that for every positive integers  $k_1, k_2, k_3, k_4$ , there is an integer  $g(k_1, k_2, k_3, k_4)$  such that every strongly connected digraph with chromatic number greater than  $g(k_1, k_2, k_3, k_4)$  contains a subdivision of  $C(k_1, k_2, k_3, k_4)$ . As evidence, we prove this conjecture for  $k_2 = k_3 = k_4 = 1$ .

*Key words:* Chromatic number, four blocks cycle, subdivision

*2010 MSC:* 00-01, 99-00

---

## 1. Introduction

Throughout this paper, all graphs are considered to be simple, that is, there are no loops and no multiple edges. By giving an orientation to each edge of a graph  $G$ , the obtained oriented graph is called a digraph. Reciprocally, the graph obtained from a digraph  $D$  by ignoring the directions of its arcs is called the underlying graph of  $D$ , and denoted by  $G(D)$  (a circuit of length 2 in  $D$  corresponds to one edge in  $G(D)$ ). The chromatic number of a digraph  $D$ , denoted by  $\chi(D)$ , is the chromatic number of its underlying graph. A digraph  $D$  is said

---

\*Corresponding author

*Email address:* [darine.mniny@liu.edu.lb](mailto:darine.mniny@liu.edu.lb) (Darine AL MNINY)

2  
3  
4  
5  
6  
7  
8  
9 to be  $k$ -chromatic if  $\chi(D) \geq k$ . If  $x$  is a vertex of  $D$ , we denote by  $N^+(x)$  (resp.  
10  $N^-(x)$ ) the set of vertices  $y$  such that  $(x, y)$  (resp.  $(y, x)$ ) is an arc of  $D$ . The  
11 out-degree (resp. in-degree) of  $x$  is the cardinality of  $N^+(x)$  (resp.  $N^-(x)$ ).  
12

13  
14 An oriented path (resp. oriented cycle) is an orientation of a path (resp. cy-  
15 cle). The length of a path (resp. cycle) is the number of its arcs. An oriented  
16 path (resp. oriented cycle) is said to be directed if all its arcs have the same  
17 orientation. Given an oriented path  $P$  (resp. oriented cycle  $C$ ), a block is a  
18 maximal directed subpath of  $P$  (resp. of  $C$ ). We denote by  $P(k_1, k_2, \dots, k_n)$   
19 (resp.  $C(k_1, k_2, \dots, k_n)$ ) the oriented path (resp. oriented cycle) formed of  $n$   
20 blocks of lengths  $k_1, k_2, \dots, k_{n-1}$  and  $k_n$  respectively.  
21  
22

23  
24 A digraph  $D$  is said to be acyclic if it contains no directed cycles.  $D$  is said to  
25 be strongly connected if for any two vertices  $x$  and  $y$  there is a directed path  
26 from  $x$  to  $y$ . Recall that a subdivision of a digraph  $D$  is a digraph  $D'$  obtained  
27 from  $D$  by replacing each arc  $(x, y)$  by an  $xy$ -dipath of length at least 1, all new  
28 paths being internally disjoint.  
29  
30  
31  
32

33  
34  
35 An important question to be asked is the following:  
36

37 **Problem 1.** *Which are the graphs  $G$  such that every graph with sufficiently*  
38 *high chromatic number contains  $G$  as a subgraph?*  
39  
40

41 In this context, Erdős and Hajnal [7] proved that every graph with chromatic  
42 number at least  $k$  contains an odd cycle of length at least  $k$ . A counterpart of  
43 this theorem for even length was settled by Mihók and Schiermeyer [13]: every  
44 graph with chromatic number at least  $k$  contains an even cycle of length at  
45 least  $k$ . Further results on graphs with prescribed lengths of cycles have been  
46 obtained [9, 10, 12, 13, 15].  
47  
48  
49  
50  
51

52 In their article, Cohen et al. [5] investigated a generalization of Problem 1  
53 by considering the analogous problem for directed graphs:  
54  
55

56 **Problem 2.** *Which are the digraphs  $D$  such that every  $k$ -chromatic digraph*  
57  
58

2  
3  
4  
5  
6  
7  
8  
9 contains  $D$  as a subdigraph?  
10

11 A famous theorem by Erdős [6] states that there exist digraphs with arbitrar-  
12 ily large chromatic number and arbitrarily high girth. This implies that if  $D$  is a  
13 40 digraph containing an oriented cycle, there exist digraphs with arbitrarily high  
14 chromatic number with no subdigraph isomorphic to  $D$ . Thus the only possible  
15 candidates to answer Problem 2 are the oriented trees. Burr [4] conjectured that  
16 every  $(2k - 2)$ -chromatic digraph contains every oriented tree  $T$  of order  $k$ , and  
17 he was able to prove that every  $(k - 1)^2$ -chromatic digraph contains a copy of  
18 45 any oriented tree  $T$  of order  $k$ . The best known bound, due to Addario-Berry et  
19 al. [2], is in  $(k/2)^2$ . For special oriented trees, better bounds on the chromatic  
20 number are known. The most famous one, known as Gallai-Roy theorem, deals  
21 with directed paths:  
22  
23  
24  
25  
26  
27

28 **Theorem 3.** (Gallai [8], Roy [14]) Every  $k$ -chromatic digraph contains a di-  
29 50 rected path of length  $k - 1$ .  
30  
31

32 However, for paths with two blocks, the best possible upper bound has been  
33 determined by Addario-Berry et al. as follows:  
34  
35

36 **Theorem 4.** (Addario-Berry et al. [1]) Let  $k_1$  and  $k_2$  be positive integers such  
37 55 that  $k_1 + k_2 \geq 3$ . Every  $(k_1 + k_2 + 1)$ -chromatic digraph  $D$  contains any two  
38 blocks path  $P(k_1, k_2)$ .  
39  
40

41 The following famous theorem of Bondy shows that the story does not stop  
42 here:  
43  
44

45 **Theorem 5.** (Bondy [3]) Every strong digraph  $D$  contains a directed cycle of  
46 60 length at least  $\chi(D)$ .  
47

48 The strong connectivity assumption is indeed necessary, because there exists  
49 acyclic digraphs (transitive tournaments) with large chromatic number and no  
50 directed cycle.  
51  
52

53 Since any directed cycle of length at least  $k$  can be seen as a subdivision of the  
54 65 directed cycle  $C_k$  of length  $k$ , Cohen et al. [5] conjectured that Bondy's theorem  
55 can be extended to all oriented cycles:  
56  
57  
58

2  
3  
4  
5  
6  
7  
8  
9 **Conjecture 6.** (Cohen et al. [5]) For every oriented cycle  $C$ , there exists a  
10 constant  $f(C)$  such that every strong digraph with chromatic number at least  
11  $f(C)$  contains a subdivision of  $C$ .  
12  
13

14  
15 70 Cohen et al. [5] noticed that the strongly connected connectivity assumption  
16 is also necessary in Conjecture 6. This follows from proving the existence of  
17 acyclic digraphs with large chromatic number and no subdivisions of  $C$  for any  
18 oriented cycle  $C$ :  
19  
20

21 **Theorem 7.** (Cohen et al. [5]) For any positive integers  $b, c$ , there exists an  
22  
23 75 acyclic digraph  $D$  with  $\chi(D) \geq c$  in which all oriented cycles have more than  $b$   
24 blocks.  
25

26  
27 In fact, Cohen et al. [5] proved Conjecture 6 in their article for cycles with  
28 two blocks. In particular, they showed that the chromatic number of strong  
29 digraphs with no subdivisions of two blocks cycles  $C(k_1, k_2)$  is bounded from  
30 above by  $O((k_1 + k_2)^4)$ . More recently, Kim et al. [11] improved this upper  
31 80 bound to  $O((k_1 + k_2)^2)$ , and asked whether this upper bound can be improved  
32 to  $O(k_1 + k_2)$ , which is remained to be open.  
33  
34  
35  
36  
37

38  
39 Furthermore, Cohen et al. confirmed the above conjecture for the case of  
40 85 cycles with four blocks as follows:  
41

42 **Theorem 8.** (Cohen et al. [5]) Let  $D$  be a strongly connected digraph with no  
43 subdivisions of four blocks cycles  $C(1, 1, 1, 1)$ , then  $\chi(D) \leq 24$ .  
44

45  
46 In this paper, we are interested in the chromatic number of digraphs con-  
47 taining no subdivisions of cycles with four blocks  $C(k_1, k_2, k_3, k_4)$ . In particular,  
48 90 we conjecture the following:  
49

50  
51 **Conjecture 9.** For every positive integers  $k_1, k_2, k_3$  and  $k_4$ , there is an integer  
52  $g(k_1, k_2, k_3, k_4)$  such that every strongly connected digraph with no subdivisions  
53 of  $C(k_1, k_2, k_3, k_4)$  has chromatic number at most  $g(k_1, k_2, k_3, k_4)$ .  
54  
55

56 We confirm this conjecture when  $k_2 = k_3 = k_4 = 1$  (for arbitrary  $k_1$ ) using  
57  
58

2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

95 the simple notion of a maximal tree and the technique of digraphs decomposing.  
Precisely, we prove that  $g(k_1, 1, 1, 1) = 8^3 \cdot k_1$ .

## 2. Main Result

We introduce some basic definitions and terminologies that will be elementary to state our main theorem.

100  
A tree is a connected graph containing no cycles. An oriented tree is an orientation of a tree. An out-tree is an oriented tree in which all vertices have in-degree at most 1. This implies that an out-tree has exactly one vertex of in-degree 0, called the source. Given a digraph  $D$  having a spanning out-tree  
105  $T$  with source  $r$ , the level of a vertex  $x$  with respect to  $T$ , denoted by  $l_T(x)$ , is the length of the unique  $(r, x)$ -directed path in  $T$ . For a nonnegative integer  $i$ , we define  $L_i(T) := \{x \in V(T) \mid l_T(x) = i\}$ . For a vertex  $x$  of  $D$ , the ancestors of  $x$  are the vertices that belong to  $T[r, x]$ . If  $y$  is an ancestor of  $x$  with respect to  $T$ , we write  $y \leq_T x$ . In this case, the vertices  $x$  and  $y$  are said to be comparable  
110 with respect to the order  $(V(T), \leq_T)$  induced by  $T$ . Conversely, we denote by  $S(x)$  the set of vertices  $v$  of  $G$  such that  $x$  is an ancestor of  $v$ . Furthermore,  $T_x$  denotes the subtree of  $T$  rooted at  $x$  and induced by  $S(x)$ . For two vertices  $x_1$  and  $x_2$  of  $D$ , the least common ancestor  $z$  of  $x_1$  and  $x_2$  is the common ancestor of  $x_1$  and  $x_2$  having the highest level in  $T$ . Note that the latter notion  
115 is well-defined since  $r$  is a common ancestor of all vertices. An arc  $(x, y)$  of  $D$  is said to be forward with respect to  $T$  if  $l_T(x) < l_T(y)$ , otherwise it is called a backward arc. We say that  $T$  is a maximal out-tree of  $D$  if for every backward arc  $(x, y)$  of  $D$ ,  $y$  is an ancestor of  $x$ .

120 The next proposition shows an interesting structural property on digraphs having a spanning out-tree:

**Proposition 10.** *Given a digraph  $D$  having a spanning out-tree  $T$ , then  $D$  contains a maximal out-tree.*

2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

*Proof.* Initially, set  $T_0 := T$ . If  $T_0$  is maximal, there is nothing to do. Otherwise  
125 there is an arc  $(x, y)$  of  $D$  which is backward with respect to  $T_0$  such that  $y$  is  
not an ancestor of  $x$ . Let  $T_1$  be the out-tree obtained from  $T_0$  by adding  $(x, y)$   
to  $T_0$ , and deleting the arc of head  $y$  in  $T_0$ . We can easily see that the level of  
each vertex in  $T_1$  is at least its level in  $T_0$ , and there exists a vertex ( $y$ ) whose  
level has strictly increased. Since the level of a vertex cannot increase infinitely,  
130 we can see that after a finite number of repeating the above process we reach  
an out-tree which is maximal.  $\square$

An oriented cycle  $C$  is said to be antirected if every vertex of  $C$  has either  
out-degree 2 or in-degree 2 in  $C$ . That is, it is a cycle in which all blocks have  
length equal 1. For a positive integer  $k$ , an antirected cycle of length at least  
135  $k$  is denoted by  $\mathcal{A}_{\geq k}$ . In their article, Cohen et al. [5] proved the following:

**Theorem 11.** (Cohen et al.[5]) *Let  $D$  be an oriented graph and  $k$  be an integer  
greater than 1. If  $\chi(D) \geq 8k - 7$ , then  $D$  contains an antirected cycle  $\mathcal{A}_{\geq 2k}$   
of length at least  $2k$ .*

Given two digraphs  $D_1$  and  $D_2$ ,  $D_1 \cup D_2$  is defined as the digraph whose  
140 vertex-set is  $V(D_1) \cup V(D_2)$  and whose arc-set is  $A(D_1) \cup A(D_2)$ . The next  
lemma will be useful for the coming proofs:

**Lemma 12.**  $\chi(D_1 \cup D_2) \leq \chi(D_1) \times \chi(D_2)$  for any two digraphs  $D_1$  and  $D_2$ .

*Proof.* For  $i \in \{1, 2\}$ , let  $\phi_i : V(D_i) \rightarrow \{1, 2, \dots, \chi(D_i)\}$  be a proper  $\chi(D_i)$ -  
coloring of  $D_i$ . Define  $\psi$ , the coloring of  $V(D_1 \cup D_2)$ , as follows:

$$\psi(x) = \begin{cases} (\phi_1(x), 1) & x \in V(D_1) \setminus V(D_2); \\ (\phi_1(x), \phi_2(x)) & x \in V(D_1) \cap V(D_2); \\ (1, \phi_2(x)) & x \in V(D_2) \setminus V(D_1). \end{cases}$$

We may easily verify that  $\psi$  is a proper coloring of  $D_1 \cup D_2$  with color-set  
 $\{1, 2, \dots, \chi(D_1)\} \times \{1, 2, \dots, \chi(D_2)\}$ . Consequently, it follows that  $\chi(D_1 \cup D_2) \leq$   
145  $\chi(D_1) \times \chi(D_2)$ .  $\square$

Now we are ready to state our main theorem:

**Theorem 13.** *If  $D$  is a digraph having a spanning out-tree  $T$  with no subdivisions of  $C(k, 1, 1, 1)$ , then the chromatic number of  $D$  is at most  $8^3.k$ .*

*Proof.* Due to Proposition 10, we may assume without loss of generality that  $T$  is a maximal out-tree in  $D$ . For  $0 \leq i \leq k-1$ , let  $V_i := \cup_{\alpha \geq 0} L_{i+\alpha k}(T)$ . Define  $D_i$  to be the subdigraph of  $D$  induced by  $V_i$ , and then partition the arcs of  $D_i$  as follows:

$$A_1 := \{(x, y) | l_T(x) < l_T(y) \text{ and } x \leq_T y\};$$

$$A_2 := \{(x, y) | l_T(x) > l_T(y) \text{ and } y \leq_T x\};$$

$$A_3 := A(D_i) \setminus (A_1 \cup A_2).$$

For  $0 \leq i \leq k-1$  and  $j = 1, 2, 3$ , let  $D_i^j$  be the spanning subdigraph of  $D_i$  whose arc-set is  $A_j$ .

**Claim 14.**  $\chi(D_i^1) \leq 8$  for all  $i \in \{0, 1, \dots, k-1\}$ .

*Proof of Claim 14.* Assume to the contrary that  $\chi(D_i^1) \geq 9$ . Then Theorem 11 implies that  $D_i^1$  contains an antidirected cycle of length at least 4, say  $\mathcal{A}_{2k}$  with  $k \geq 2$ . Set  $\mathcal{A}_{2k} := x_1 x_2 \dots x_k x_{k+1} x_{k+2} \dots x_{2k} = S_1 \cup S_2$ , where  $S_1 = \{x_i \in V(\mathcal{A}_{2k}) | |N^+(x_i) \cap V(\mathcal{A}_{2k})| = 2\}$  and  $S_2 = \{x_i \in V(\mathcal{A}_{2k}) | |N^-(x_i) \cap V(\mathcal{A}_{2k})| = 2\}$ . Let  $x_i \in S_1$  such that  $l_T(x_i)$  is maximal. Assume that  $N^+(x_i) \cap V(\mathcal{A}_{2k}) = \{x_\alpha, x_\beta\}$ . Clearly,  $x_\alpha$  and  $x_\beta$  belong to  $S_2$ . By the definition of  $S_2$ , there exists  $1 \leq \alpha', \beta' \leq 2k$  other than  $i$  such that  $(x_{\alpha'}, x_\alpha)$  and  $(x_{\beta'}, x_\beta)$  belong to  $A(D_i^1)$ . Note that probably  $\alpha' = \beta'$ . By the definition of  $D_i^1$ ,  $x_{\alpha'}, x_{\beta'}$  and  $x_i$  are all comparable with respect to the order induced by  $T$ . Without loss of generality, assume that  $l_T(x_{\beta'}) < l_T(x_{\alpha'})$  and  $l_T(x_\alpha) \leq l_T(x_\beta)$ . Thus by the maximality of  $l_T(x_i)$  in  $S_1$ , we get  $l_T(x_{\beta'}) < l_T(x_{\alpha'}) < l_T(x_i)$ . Therefore,  $T[x_i, x_\alpha] \cup (x_i, x_\beta) \cup T[x_{\beta'}, x_{\alpha'}] \cup (x_{\alpha'}, x_\alpha) \cup (x_{\beta'}, x_\beta)$  is a subdivision of  $C(k, 1, 1, 1)$ , a contradiction.  $\square$

**Claim 15.**  $\chi(D_i^2) \leq 16$  for all  $i \in \{0, 1, \dots, k-1\}$ .



2  
 3  
 4  
 5  
 6  
 7  
 8  
 9 *Proof of Claim 15.* Assume to the contrary that  $\chi(D_i^2) \geq 17$ . Then Theorem  
 10 11 implies that  $D_i^2$  contains an antirected cycle of length at least 6, say  $\mathcal{A}_{2k}$   
 12 with  $k \geq 3$ . Set  $\mathcal{A}_{2k} := x_1x_2\dots x_kx_{k+1}x_{k+2}\dots x_{2k} = S_1 \cup S_2$ , where  $S_1 = \{x_i \in$   
 13  $V(\mathcal{A}_{2k}) \mid |N^+(x_i) \cap V(\mathcal{A}_{2k})| = 2\}$  and  $S_2 = \{x_i \in V(\mathcal{A}_{2k}) \mid |N^-(x_i) \cap V(\mathcal{A}_{2k})| =$   
 14 170  $2\}$ . Let  $x_i \in S_2$  such that  $l_T(x_i)$  is maximal. Assume that  $N^-(x_i) \cap V(\mathcal{A}_{2k}) =$   
 15  $\{x_\alpha, x_\beta\}$ . Clearly,  $x_\alpha$  and  $x_\beta$  belong to  $S_1$ . By the definition of  $S_1$ , there exists  
 16  $1 \leq \alpha', \beta' \leq 2k$  other than  $i$  such that  $(x_\alpha, x_{\alpha'})$  and  $(x_\beta, x_{\beta'})$  belong to  $A(D_i^2)$ .  
 17 Note that probably  $\alpha' \neq \beta'$ , since otherwise we get  $k = 2$ , a contradiction to  
 18 our assumption. By the definition of  $A_2$ ,  $x_{\alpha'}, x_{\beta'}$  and  $x_i$  are all comparable  
 19 with respect to the order induced by  $T$ . By the maximality of  $l_T(x_i)$  in  $S_2$ , we  
 20 get  $l_T(x_{\alpha'}) < l_T(x_i)$  and  $l_T(x_{\beta'}) < l_T(x_i)$ . Without loss of generality, we may  
 21 assume that  $l_T(x_{\alpha'}) > l_T(x_{\beta'})$ . By the definition of  $D_i^2$ ,  $l_T(x_{\alpha'}) - l_T(x_{\beta'}) \geq k$ .  
 22 Thus the union of  $(x_\beta, x_{\beta'}) \cup T[x_{\beta'}, x_{\alpha'}], (x_\beta, x_i), (x_\alpha, x_{\alpha'})$  and  $(x_\alpha, x_i)$  is a sub-  
 23 175 division of  $C(k, 1, 1, 1)$ , a contradiction.  $\square$   
 24  
 25  
 26  
 27  
 28  
 29  
 30  
 31  
 32  
 33  
 34  
 35  
 36  
 37  
 38  
 39  
 40  
 41  
 42  
 43  
 44  
 45  
 46  
 47  
 48  
 49  
 50  
 51  
 52  
 53  
 54  
 55  
 56  
 57  
 58  
 59  
 60  
 61  
 62  
 63  
 64  
 65

**Claim 16.**  $\chi(D_i^3) \leq 4$  for all  $i \in \{0, 1, \dots, k-1\}$ .

35 *Proof of Claim 16.* Assume to the contrary that  $\chi(D_i^3) \geq 5$ . Due to Theorem  
 36 4,  $D_i^3$  contains a copy  $Q$  of  $P(2, 2)$ , which is the union of two directed paths  $Q_1$   
 37 and  $Q_2$  which are disjoint except in their initial vertex  $y_0$ , say  $Q_1 = (y_0, y_1, y_2)$   
 38 and  $Q_2 = (y_0, z_1, z_2)$ .  
 39  
 40  
 41

42 Assume that  $y_2$  and  $z_2$  are not comparable with respect to the tree-order, and  
 43 let  $x$  be the least common ancestor of  $y_2$  and  $z_2$ . By the definition of  $D_i^3$ ,  $x$  is  
 44 distinct from  $y_1$  and  $z_1$ .  
 45

46 Assume that  $(T[x, y_2] \cup T[x, z_2]) \cap Q = \{y_2, z_2\}$ . Observe that  $l(T[x, y_2]) < k$   
 47 and  $l(T[x, z_2]) < k$ , since otherwise  $T[x, y_2] \cup T[x, z_2] \cup Q$  is a subdivision of  
 48  $C(k, 1, 1, 1)$ , a contradiction. Consider  $v$  to be the least common ancestor of  $y_0$   
 49 and  $y_1$ . Note that  $(T[v, y_0] \cup T[v, y_1]) \cap (T[x, y_2] \cup T[x, z_2]) = \emptyset$ , since otherwise we  
 50 get  $l(T[x, y_2]) \geq k$  or  $l(T[x, z_2]) \geq k$ , a contradiction. In what follows, if  $y_1$  and  
 51  $z_1$  are comparable with respect to the tree-order, we will assume without loss of  
 52 195 generality that  $y_1 \leq_T z_1$ . This implies that  $T[v, y_1] \cap Q = \{y_1\}$ . Thus the union  
 53  
 54  
 55  
 56  
 57  
 58  
 59  
 60  
 61  
 62  
 63  
 64  
 65

2  
 3  
 4  
 5  
 6  
 7  
 8  
 9 of  $T[v, y_1] \cup (y_1, y_2)$ ,  $T[v, y_0] \cup Q_2$ ,  $T[x, y_2]$  and  $T[x, z_2]$  would be a subdivision of  
 10  $C(k, 1, 1, 1)$ , a contradiction. This implies that  $(T[x, y_2] \cup T[x, z_2]) \cap Q \neq \{y_2, z_2\}$ .  
 11 Without loss of generality, we will assume that  $T[x, y_2] \cap Q - y_2 \neq \emptyset$ . Let  $v$  be  
 12 the last vertex of  $T[x, y_2]$  in  $Q - y_2$ . By our assumption,  $z_2$  is not an ancestor of  
 13  $y_2$ , so  $v$  is different from  $z_2$ . Also by the definition of  $A_3$ ,  $y_1$  is not an ancestor  
 14 of  $y_2$ , so  $v$  is different from  $y_1$ . Thus  $v$  is either  $z_1$  or  $y_0$ . Assume that  $v = z_1$ .  
 15 Let  $t$  be the least common ancestor of  $y_0$  and  $z_2$ . Due to our assumption and  
 16 to the definition of  $A_3$ , it is easy to see that  $y_2$  and  $z_1$  are not on  $T[t, z_2]$ .  
 17 Observe that  $T[t, z_2] \cap Q_1 - y_0 = \emptyset$ , since otherwise  $y_1$  belongs to  $T[t, z_2]$  and  
 18 hence  $T[y_1, z_2] \cup (y_1, y_2) \cup (z_1, z_2) \cup T[z_1, y_2]$  is a subdivision of  $C(k, 1, 1, 1)$ , a  
 19 contradiction. Thus the union of  $T[t, z_2]$ ,  $T[t, y_0] \cup Q_1$ ,  $(z_1, z_2)$  and  $T[z_1, y_2]$  is  
 20 a subdivision of  $C(k, 1, 1, 1)$ , a contradiction. Therefore,  $v = y_0$ .

21 Assume that  $z_1$  and  $y_1$  are not comparable with respect to the tree-order, and  
 22 let  $t$  be the least common ancestor of  $z_1$  and  $y_1$ . By the definition of  $A_3$ ,  $y_0$  is not  
 23 an ancestor neither of  $y_1$  nor of  $z_1$ , and so  $(T[t, y_1] \cup T[t, z_1]) \cap (Q \cup T[y_0, y_2]) =$   
 24  $\{y_1, z_1\}$ . Thus the union of  $T[y_0, y_2]$ ,  $(y_0, z_1)$ ,  $T[t, y_1] \cup (y_1, y_2)$  and  $T[t, z_1]$   
 25 is a subdivision of  $C(k, 1, 1, 1)$ , a contradiction. This implies that  $y_1 \leq_T z_1$   
 26 or vice versa. If  $y_1 \leq_T z_1$ , then  $T[y_0, y_2] \cup (y_0, z_1) \cup (y_1, y_2) \cup T[y_1, z_1]$  is a  
 27 subdivision of  $C(k, 1, 1, 1)$ , a contradiction. Thus  $z_1 \leq_T y_1$  and so the union of  
 28  $T[x, z_2]$ ,  $T[x, y_0] \cup T[y_0, y_2]$ ,  $(z_1, z_2)$  and  $T[z_1, y_1] \cup (y_1, y_2)$  is a subdivision of  
 29  $C(k, 1, 1, 1)$ , a contradiction. Therefore,  $y_2$  and  $z_2$  are comparable with respect  
 30 to the tree-order.

31 Without loss of generality, we will assume that  $y_2 \leq_T z_2$ . By the definition of  
 32  $A_3$ ,  $T[y_2, z_2] \cap Q = \{y_2, z_2\}$ . We suppose that  $y_0$  is an ancestor of  $y_2$ . Again by  
 33 the definition of  $A_3$ ,  $T[y_0, y_2] \cap Q = \{y_0, y_2\}$ . Now we assume that  $z_1$  and  $y_1$  are  
 34 not ancestors, and we consider  $t$  to be the least common ancestor of  $y_1$  and  $z_1$ .  
 35 It is clear that  $(T[t, y_1] \cup T[t, z_1]) \cap (Q \cup T[y_0, y_2]) = \{y_1, z_1\}$ . Hence the union of  
 36  $T[y_0, y_2]$ ,  $(y_0, z_1)$ ,  $T[x, y_1] \cup (y_1, y_2)$  and  $T[x, z_1]$  is a subdivision of  $C(k, 1, 1, 1)$ .  
 37 This means that  $y_1$  and  $z_1$  are comparable with respect to the tree-order. If  
 38  $y_1 \leq_T z_1$ , then we may easily check that  $T[y_1, z_1] \cup (y_1, y_2) \cup T[y_0, y_2] \cup (y_0, z_1)$  is a  
 39 subdivision of  $C(k, 1, 1, 1)$ , a contradiction. Thus  $z_1 \leq_T y_1$  and hence we obtain

2  
3  
4  
5  
6  
7  
8  
9 a subdivision of  $C(k, 1, 1, 1)$  which is formed by the union of  $T[y_0, y_2] \cup T[y_2, z_2]$ ,  
10  $(y_0, y_1)$ ,  $(z_1, z_2)$  and  $T[z_1, y_1]$ , a contradiction.

11  
12 230 It follows that  $y_0$  is not an ancestor of  $y_2$ . By the definition of  $A_3$ ,  $y_1$  is not an  
13 ancestor of  $y_2$ , so we consider  $x$  to be the least common ancestor of  $y_1$  and  $y_2$ . It  
14 is clear that  $T[x, y_2] \cap Q = \{y_2\}$ , since otherwise we get either  $y_0 \leq_T y_2$ , or  $z_1 \leq_T$   
15  $z_2$ , a contradiction. Moreover, we may easily verify that  $T[x, y_1] \cap Q - y_1 \subseteq \{z_1\}$ .  
16 Assume that  $z_1$  does not belong to  $T[x, y_1]$ , then the union of  $T[x, y_2] \cup T[y_2, z_2]$ ,  
17  $T[x, y_1]$ ,  $Q_2$  and  $(y_0, y_1)$  is a subdivision of  $C(k, 1, 1, 1)$ , a contradiction. Thus  
18 235  $z_1$  belongs to  $T[x, y_1]$ , and so  $T[x, y_2] \cup T[x, z_1] \cup Q_1 \cup (y_0, z_1)$  forms a subdivision  
19 of  $C(k, 1, 1, 1)$ , a contradiction.  $\square$   
20  
21  
22  
23  
24

25 Therefore, Claims 14, 15, 16 together with Lemma 12 imply that  $\chi(D_i) \leq 8^3$ .  
26 As  $V(D)$  is partitioned into  $V(D_i)$  for  $i = 0, 1, \dots, k-1$ , then  $D$  can be colored  
27 properly using  $8^3 \cdot k$  colors by assigning  $8^3$  distinct colors to each  $D_i$ . This  
28 240 completes the proof.  $\square$   
29  
30  
31

32 Note that the upper bound in the theorem above can easily be lowered.  
33 However, we made no attempt to it here for two reasons: Firstly, we would like  
34 to keep the proof as simple as possible; secondly using our method, there is no  
35 hope to get an optimal value for  $\chi(D)$  and all what we look for is to prove that  
36 245 it is bounded.  
37  
38  
39

40  
41 Due to the fact that every strongly connected digraph contains a spanning  
42 out-tree, we can immediately deduce the following:  
43  
44

45 250 **Corollary 17.** *Let  $D$  be a strong digraph with no subdivisions of  $C(k, 1, 1, 1)$ ,*  
46 *then the chromatic number of  $D$  is at most  $8^3 \cdot k$ .*  
47  
48  
49

## 50 References

- 51  
52 [1] L. Addario-Berry, F. Havet, and S. Thomassé, Paths with two blocks in  
53  $n$ -chromatic digraphs, *J. of Combinatorial Theory, Series B*, 97 (4), pages  
54 255 620-626, 2007.  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

[2] L. Addario-Berry, F. Havet, C. L. Sales, B. A. Reed, and S. Thomassé, Oriented trees in digraphs, *Discrete Math*, 313 (8), pages 967-974, 2013.

[3] J. A. Bondy, Disconnected orientations and a conjecture of Las Vergnas, *J. London Math. Soc. (2)*, 14 (2), pages 277-282, 1976.

260 [4] S. A. Burr, Subtrees of directed graphs and hypergraphs, *Proceedings of the Eleventh Southeastern Conference on Combinatorics, Graph Theory and Computing*, Boca Raton, Congr. Numer, 28, pages 227-239, 1980.

[5] N. Cohen, F. Havet, W. Lochet, and N. Nisse, Subdivisions of oriented cycles in digraphs with large chromatic number, *J. Graph Theory*, 89, pages 265 439-456, 2018.

[6] P. Erdős, Graph Theory and probability, *Canad. J. Math.*, 11, pages 34-38, 1959.

[7] P. Erdős and A. Hajnal, On chromatic number of graphs and set-systems, *Acta Math. Acad. Sci. Hungar*, 17, pages 61-99, 1966.

270 [8] T. Gallai, On directed paths and circuits, *In Theory of Graphs (Proc. Colloq. Titany, 1966)*, pages 115-118, Academic Press, New York, 1968.

[9] A. Gyárfás, Graphs with  $k$  odd cycle lengths, *Discrete Math*, 103, pages 41-48, 1992.

[10] T. Kaiser, O. Rucký, and R. Skrekovski, Graphs with odd cycle lengths 5 and 7 are 3-colorable, *SIAM J. Discrete Math*, 25, pages 1069-1088, 2011.

275 [11] R. Kim, SJ. Kim, J. Ma, and B. Park, Cycles with two blocks in  $k$ -chromatic digraphs, *J. Graph Theory*, 00, pages 1-14, 2017.

[12] C. Löwenstein, D. Rautenbach, and I. Schiermeyer, Cycle length parities and the chromate number, *J. Graph Theory*, 64, pages 210-218, 2010.

280 [13] P. Mihók and I. Schiermeyer, Cycle lengths and chromatic number of graphs, *Discrete Math*, 286, pages 147-149, 2004.

2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

[14] B. Roy, Nombre chromatique et plus longs chemins d'un graphe, *Rev. Française Informat. Recherche Opérationnelle*, 1 (5), pages 129-132, 1967.

[15] S. S. Wang, Structure and coloring of graphs with only small odd cycles, *SIAM J. Discrete Math*, 22, pages 1040-1072, 2008.