



**HAL**  
open science

# Stability of optimal shapes and convergence of thresholding algorithms in linear and spectral optimal control problems: Supplementary material

Antonin Chambolle, Idriss Mazari-Fouquer, Yannick Privat

## ► To cite this version:

Antonin Chambolle, Idriss Mazari-Fouquer, Yannick Privat. Stability of optimal shapes and convergence of thresholding algorithms in linear and spectral optimal control problems: Supplementary material. 2023. hal-04140334

**HAL Id: hal-04140334**

**<https://hal.science/hal-04140334>**

Preprint submitted on 25 Jun 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# Stability of optimal shapes and convergence of thresholding algorithms in linear and spectral optimal control problems: Supplementary material

Antonin Chambolle\*    Idriss Mazari-Fouquer†    Yannick Privat‡§

June 25, 2023

## Abstract

We address the convergence of thresholding schemes for spectral optimisation problems and linear control problems.

**Keywords:** Convergence analysis, Numerical algorithms in optimal control, PDE constrained optimisation, Shape optimisation, Quantitative inequalities, Thresholding scheme.

**AMS classification:** 49M05, 49M41, 49N05, 49Q10.

## A Proof of Theorem II

As mentioned in the introduction, when the proofs are identical to those of Theorem I we omit them. We introduce the analog of Definition 7.

**Definition 1** (Critical points). *For any  $V_0 \in (0; 1)$ , for any  $f \in \mathcal{F}(V_0)$ , we say  $f$  is a **critical point** of  $\lambda$  if:*

1. *There exists a unique  $\mu_f = \mu_{u_f, V_0}$  such that*

$$\text{Vol}(\{\eta_f > \mu_f\}) = V_0 \text{Vol}(\Omega) = \text{Vol}(\{\eta_f \geq \mu_f\}),$$

2. *and  $f = \mathbb{1}_{\{\eta_f > \mu_f\}}$ .*

A **critical set** is a subset  $E$  of  $\Omega$  such that  $\mathbb{1}_E \in \mathcal{F}(V_0)$  and such that  $\mathbb{1}_E$  is a critical point of  $\lambda$ .

**Definition 2.** *An admissible control  $f^* \in \mathcal{F}(V_0)$  is called a **stable local minimiser** of  $\lambda$  in  $\mathcal{F}(V_0)$  if there exist  $C = C(f^*)$ ,  $\delta = \delta(f^*) > 0$  such that*

$$\forall f \in \mathcal{F}(V_0), \|f - f^*\|_{L^1(\Omega)} \leq \delta(f^*) \Rightarrow \lambda(f) \geq \lambda(f^*) + C(f^*) \|f - f^*\|_{L^1(\Omega)}^2.$$

---

\*CEREMADE, UMR 7534, CNRS & Université Paris Dauphine PSL, 75775 Paris Cedex 16, France ([chambolle@ceremade.dauphine.fr](mailto:chambolle@ceremade.dauphine.fr))

†CEREMADE, UMR 7534, Université Paris Dauphine PSL, 75775 Paris Cedex 16, France ([mazari@ceremade.dauphine.fr](mailto:mazari@ceremade.dauphine.fr))

‡IRMA, Université de Strasbourg, CNRS UMR 7501, Inria, 7 rue René Descartes, 67084 Strasbourg, France ([yannick.privat@unistra.fr](mailto:yannick.privat@unistra.fr)).

§Institut Universitaire de France (IUF).

## A.1 First consequences of a large volume constraint

We show regularity properties similar to those of Lemma 13. We let, for any  $f \in \mathcal{F}(V_0)$ ,  $\mu_{f,V_0}$  be the unique real number such that

$$\frac{\text{Vol}(\{\eta_f > \mu_{f,V_0}\})}{\text{Vol}(\Omega)} \leq V_0 \leq \frac{\text{Vol}(\{\eta_f \geq \mu_{f,V_0}\})}{\text{Vol}(\Omega)}.$$

We also define

$$\omega_{f,V_0} := \{\eta_f > \mu_{f,V_0}\}.$$

**Lemma 3.** *There exists  $\varepsilon_0 > 0$  and  $\delta_0 > 0$  such that, for any  $V_0 \in (1 - \varepsilon_0; 1)$ , for any  $f \in \mathcal{F}(V_0)$ , we have on the one hand*

$$\text{Vol}(\{\eta_f = \mu_{f,V_0}\}) = 0$$

and, on the other hand, the following regularity properties:

1.  $\omega_{f,V_0}$  has a  $\mathcal{C}^{1,\alpha}$  boundary (for any  $\alpha \in (0; 1)$ ),

2.

$$\min_{\partial\omega_{f,V_0}} \left| \frac{\partial\eta_f}{\partial\nu} \right| \geq \delta_0,$$

3.  $\partial\omega_{f,V_0}$  is locally a graph over  $\partial\Omega$ .

Finally, for any  $\alpha \in (0; 1)$ ,  $\partial\omega_{f,V_0}$  converges to  $\partial\Omega$  in  $\mathcal{C}^{1,\alpha}$  as  $V_0 \rightarrow 1$ , uniformly in  $f \in \mathcal{F}(V_0)$ .

*Lemma 3.* The proof is identical, except that the analog of (2.4) deals with the first Dirichlet eigenfunction  $\psi_1$  of  $\Omega$ . Let us explain in more details: let  $(\lambda^D(\Omega), \psi_1)$  be the first Dirichlet eigenpair of the domain  $\Omega$ :

$$\begin{cases} -\Delta\psi_1 = \lambda^D(\Omega)\psi_1 & \text{in } \Omega, \\ \psi_1 \in W_0^{1,2}(\Omega), \\ \psi_1 > 0 & \text{in } \Omega, \\ \int_{\Omega} \psi_1^2 = 1. \end{cases} \quad (\text{A.1})$$

From the variational formulation (1.3) of  $\lambda(f)$  and the fact that  $f \in \mathcal{F}(V_0)$  we have the bound

$$\lambda^D(\Omega) - 1 \leq \lambda(f) \leq \lambda^D(\Omega)$$

whence, from a standard bootstrapping argument there holds, for any  $p \in [1; +\infty)$ ,

$$\sup_{V_0 \in (0;1)} \sup_{f \in \mathcal{F}(V_0)} \|\eta_f\|_{W^{2,p}(\Omega)} < \infty. \quad (\text{A.2})$$

It is expected that as  $V_0 \rightarrow 1$  we should have  $\lambda(f) + 1 \approx \lambda^D$ . Let us prove this. From the variational formulation (1.3) we have

$$-\int_{\Omega} \eta_f^2 (1-f) \leq \lambda(f) - (\lambda^D - 1) \leq \int_{\Omega} \psi_1^2 (1-f).$$

From (A.2) we deduce that

$$\lim_{V_0 \rightarrow 1} \sup_{f \in \mathcal{F}(V_0)} |\lambda(f) - (\lambda^D(\Omega) - 1)| = 0.$$

Let us now consider a sequence  $\{V_k\}_{k \in \mathbb{N}}$  converging to 1 and, for any  $k \in \mathbb{N}$ , a function  $f_k \in \mathcal{F}(V_k)$ . Passing to the limit in the weak formulation of the equation (1.4) we deduce that any closure point  $\eta_\infty$  of  $\{\eta_{f_k}\}_{k \in \mathbb{N}}$  solves the equation

$$\begin{cases} -\Delta \eta_\infty = \lambda^D(\Omega) \eta_\infty & \text{in } \Omega, \\ \eta_\infty \in W_0^{1,2}(\Omega), \\ \eta_\infty > 0 & \text{in } \Omega, \\ \int_\Omega \eta_\infty^2 = 1. \end{cases} \quad (\text{A.3})$$

From the simplicity of the first eigenvalue  $\lambda^D(\Omega)$  it follows that  $\eta_\infty = \psi_1$ . From this fact it is then easy to show, as in Lemma 13, that for any  $\alpha \in (0; 1)$ ,

$$\lim_{V_0 \rightarrow 1} \sup_{f \in \mathcal{V}_0} \|\psi_1 - \eta_f\|_{\mathcal{C}^{1,\alpha}(\Omega)} = 0. \quad (\text{A.4})$$

We now use the maximum principle of Hopf: there exists a constant  $\delta_\Omega > 0$  such that

$$\inf_{\partial\Omega} |\nabla \psi_1| \geq \delta_\Omega > 0$$

and we conclude the proof of Lemma 3 by the same arguments.  $\square$

In particular we also obtain the existence of a constant  $M$  such that

$$\forall V_0 \in (1 - \varepsilon_0; 1), \forall f \in \mathcal{F}(V_0), \text{Lip}(\partial\omega_{f,V_0}) + \text{Per}(\omega_{f,V_0}) \leq M. \quad (\text{A.5})$$

We also have in a similar manner the following topological information

**Lemma 4.** *There exists  $\varepsilon'_0 > 0$  such that, for any  $V_0 \in (1 - \varepsilon'_0; 1)$ , for any  $f \in \mathcal{F}(V_0)$  the boundary  $\partial\omega_{f,V_0}$  is connected.*

We may, analogously to Lemma 14, choose  $\varepsilon_0 = \varepsilon'_0$  where  $\varepsilon_0$  is given in Lemma 13.

## A.2 Applications of the quantitative bathtub principle

We apply the quantitative bathtub principle ([39, Proposition 26]) to the thresholding algorithm (3) and, as in Lemma 16 we obtain the following result:

**Lemma 5.** *Let  $\varepsilon_0, \delta_0$  be in the conditions of Lemma 3. For any  $V_0 \in (1 - \varepsilon_0; 1)$ , for any  $f_0 \in \mathcal{F}(V_0)$ , the sequence  $\{f_k\}_{k \in \mathbb{N}}$  is uniquely defined (in the sense that for any  $k \in \mathbb{N}$  there holds  $\text{Vol}(\{\eta_{f_k} = \mu_{f_k, V_0}\}) = 0$ ) and there holds*

$$\sum_{k=0}^{\infty} \|f_{k+1} - f_k\|_{L^1(\Omega)}^2 < \infty. \quad (\text{A.6})$$

The following result is adapted from Lemma 17.

**Lemma 6.** *Let  $\varepsilon_0$  be in the conditions of Lemma 13. For any  $V_0 \in (1 - \varepsilon_0; 1)$ , for any initialisation  $f_0 \in \mathcal{F}(V_0)$ , any  $L^\infty - *$  closure point  $f_\infty$  of the sequence  $\{f_k\}_{k \in \mathbb{N}}$  generated by the thresholding algorithm is a bang-bang function:*

$$\exists E_\infty \subset \Omega, f_\infty = \mathbf{1}_{E_\infty}. \quad (\text{A.7})$$

Furthermore  $E_\infty = \{\eta_{f_\infty} > \mu_{u_{f_\infty}, V_0}\}$ , which is uniquely defined. In other words,  $E_\infty$  is a critical set in the sense of Definition 1.

Using this result we get the following dichotomy for the sequence  $\{f_k\}_{k \in \mathbb{N}}$  (see Lemma 18 for a proof).

**Lemma 7.** *Let  $\varepsilon_0$  be chosen as in Lemma 13. For any  $V_0 \in (1 - \varepsilon_0; 1)$ , for any initialisation  $f_0 \in \mathcal{F}(V_0)$ , let  $\{f_k\}_{k \in \mathbb{N}}$  be the sequence generated by the thresholding algorithm. Then:*

1. *Either  $\{f_k\}_{k \in \mathbb{N}}$  has a unique closure point  $f_\infty$ , in which case  $f_k \xrightarrow[k \rightarrow \infty]{} f_\infty$  strongly in  $L^1(\Omega)$ ,*
2. *Or  $\{f_k\}_{k \in \mathbb{N}}$  has an infinite number of closure points.*

**The goal of the next sections** In the following section we study the shape Hessians at critical shape. This is where the main differences with the proof of Theorem I occur.

### A.3 Qualitative study of critical points

In this section we analyse the critical sets (in the sense of Definition 1). For the second order shape derivatives of the Lagrangians to be meaningful we need the critical sets to have an (at least)  $\mathcal{C}^3$  boundary. Here we are exactly in the setting for which [11, Lemma 3] was introduced and, similar to Lemma 19, we have the following regularity result:

**Lemma 8.** *Let  $\varepsilon_0 > 0$  be as in Lemma 3. Any critical set  $E$  (in the sense of Definition 7) has a compact analytic boundary and, furthermore, is uniformly bounded in the  $\mathcal{C}^2$  topology.*

**Shape derivative formalism** We first identify the functional  $\lambda$  with a shape functional  $\Lambda$ , by defining

$$\Lambda : \Omega \supset E \mapsto \lambda(\mathbf{1}_E).$$

For any compact  $\mathcal{C}^3$  subset  $E$  of  $\Omega$  and for any compactly supported vector field  $\Phi \in W^{2,\infty}(\Omega; \mathbb{R}^d)$ , we can define, for any  $t \in (-1; 1)$  small enough, the function

$$e_{E,\Phi}(t) := \Lambda\left((\text{Id} + t\Phi)E\right).$$

It was proved in [39] (using the aforementioned implicit function method of [41]) that  $\Lambda$  is twice shape differentiable, and we define:

$$\Lambda'(E)[\Phi] = e'_{E,\Phi}(0), \text{ resp. } \Lambda''(E)[\Phi, \Phi] = e''_{E,\Phi}(0).$$

**First order shape derivative and definition of the Lagrange multiplier** As for the Dirichlet energy, we work with a Lagrangian and we shall use the expressions for the derivative of the volume functional given in (2.24).

We introduce the associated notion of shape criticality:

**Definition 9** (Critical shape). *A  $\mathcal{C}^3$  shape  $E$  is a **critical shape** for  $\Lambda$  if, for any compactly supported  $\Phi \in W^{2,\infty}(\Omega)$ ,*

$$\int_{\partial E} \langle \Phi, \nu_E \rangle = 0 \Rightarrow \Lambda'(E)[\Phi] = 0.$$

We now compute the Lagrange multiplier at a given critical shape. If a shape  $E$  is critical, then the condition given in Definition 22 rewrites as: there exists a constant  $\mu_E$  such that, for any compactly supported vector field  $\Phi \in W^{2,\infty}(\Omega; \mathbb{R}^d)$ ,

$$(\Lambda + \mu_E \text{Vol})'(E)[\Phi] = 0.$$

To compute  $\mu_E$  we rely on the following lemma:

**Lemma 10.** For any  $\mathcal{C}^3$  shape, for any compactly supported vector field  $\Phi \in W^{2,\infty}(\Omega; \mathbb{R}^d)$ ,

$$\Lambda'(E)[\Phi] = - \int_{\partial E} \eta_{\mathbf{1}_E}^2 \langle \Phi, \nu_E \rangle.$$

Furthermore, the shape derivative of the map  $E \mapsto \eta_{\mathbf{1}_E}$  at  $E$  in the direction  $\Phi$  is the unique solution of

$$\begin{cases} -\Delta \eta'_\Phi - \lambda(\mathbf{1}_E) \eta'_\Phi - \mathbf{1}_E \eta'_\Phi = \Lambda'(E)[\Phi] \eta_{\mathbf{1}_E} & \text{in } \Omega, \\ \llbracket \partial_\nu \eta'_\Phi \rrbracket = -\eta_{\mathbf{1}_E} \langle \Phi, \nu_E \rangle & \text{on } \partial E, \\ \int_\Omega \eta_E \eta'_\Phi = 0 \end{cases} \quad (\text{A.8})$$

where  $\llbracket \cdot \rrbracket$  denotes the jump of a function across a hypersurface.

This lemma is proved in [35, Appendix (B.2)]. It allows to prove, exactly in the same way as Lemma 24, the following statement:

**Lemma 11.** A shape  $E \subset \Omega$  is a critical set in the sense of Definition 1 if, and only if it is a critical shape in the sense of Definition 9.

To alleviate notations we introduce a notation for the shape Lagrangian:

**Definition 12** (Shape Lagrangian). Let  $E$  be a critical shape for  $\mathcal{F}$ . The **associated shape Lagrangian** is

$$\Theta_E := \Lambda + \mu_E \text{Vol}$$

where  $E = \{\eta_{\mathbf{1}_E} > \mu_E\}$ .

With the above ingredients at hand, we now move to second-order shape derivatives.

**First computations and elements for shape Hessians** In the subsequent paragraphs we use indifferently the expressions "shape Hessians" and "second order shape derivatives". We begin with an expression of the shape hessian at a critical shape  $E$  in the direction  $\Phi$ .

**Lemma 13.** Let  $E$  be a critical shape in the sense of Definition 9 and let  $\Phi \in W^{2,\infty}(\Omega; \mathbb{R}^d)$  be a compactly supported vector field. The second-order shape derivative of the shape Lagrangian  $\Theta_E$  (defined in Definition 12) is given by the expression

$$\begin{aligned} \frac{1}{2} \Theta_E''(E)[\Phi, \Phi] &= - \int_{\partial E} \eta_E \eta'_\Phi \langle \Phi, \nu_E \rangle - \int_{\partial E} \eta_E \frac{\partial \eta_E}{\partial \nu} \langle \Phi, \nu_E \rangle^2 \\ &= \mu_E \left( - \int_{\partial E} \eta'_\Phi \langle \Phi, \nu_E \rangle - \int_{\partial E} \frac{\partial \eta_E}{\partial \nu} \langle \Phi, \nu_E \rangle^2 \right) \end{aligned} \quad (\text{A.9})$$

where  $\eta'_\Phi$  solves (A.8) and where we used the fact that  $\eta_E \equiv \mu_E$  on  $\partial E$ .

We want to prove that critical shapes are stable in the following sense:

**Definition 14** ( $L^2$ -stability). A critical shape  $E$  is said to be  $L^2$ -**stable** if there exists a constant  $c_E > 0$  such that for any compactly supported vector field  $\Phi \in W^{2,\infty}(\Omega; \mathbb{R}^d)$ ,

$$\Theta_E''(E)[\Phi, \Phi] \geq c_E \|\langle \Phi, \nu \rangle\|_{L^2(\partial E)}^2.$$

Our main result is the following proposition:

**Proposition 15.** *There exists  $\varepsilon_1 > 0, \varepsilon_1 \leq \varepsilon_0, \underline{c} > 0$  such that, for any  $V_0 \in (1 - \varepsilon_1; 1)$ , for any critical set  $E$  (in the sense of Definition 9), for any compactly supported vector field  $\Phi \in W^{2,\infty}(\Omega; \mathbb{R}^d)$ ,*

$$\Theta''_E(E)[\Phi, \Phi] \geq \underline{c}\mu_E \|\langle \Phi, \nu \rangle\|_{L^2(\partial E)}^2.$$

Before we prove this proposition we need to introduce the diagonalisation basis. We consider a fixed critical shape  $E \subset \Omega$  and work with  $V_0 \in (1 - \varepsilon_0; 1)$ , where  $\varepsilon_0$  is given by Lemma 3. Heuristically, given that  $\eta'_\Phi$  satisfies (A.8) and that  $\Lambda'(E)[\Phi] = 0$  as we are working at a critical shape, it is natural to solve the following eigenvalue problem: find  $(\lambda, \phi) \in \mathbb{R} \times W^{1,2}(\Omega)$  such that

$$\begin{cases} -\Delta\phi - \Lambda(E)\phi - \mathbf{1}_E\phi = 0 & \text{in } E \cup (E)^c \\ \llbracket \partial_\nu \phi \rrbracket_E = -\lambda\rho\phi & \text{on } \partial E, \\ \int_{\partial E} \phi^2 > 0, \\ \phi \in W_0^{1,2}(\Omega), \\ \int_\Omega \phi \eta_E = 0. \end{cases} \quad (\text{A.10})$$

where  $\rho \in L^\infty(\partial E; [\delta; 1/\delta])$  ( $\delta > 0$ ) is a weight to be determined. Let us simply consider the eigenvalue problem (A.10) with a weight  $\rho \in L^\infty(\partial E; [\delta; 1/\delta])$  for some small  $\delta > 0$ . Note that as we work with a critical shape  $E$ ,

$$\eta_E \equiv \mu_E \text{ on } \partial E.$$

We consider the weighted space

$$L_\rho^2(\partial E) := \left\{ f \in L^2(\partial E), \int_{\partial E} \rho f^2 < \infty \right\} = L^2(\partial E).$$

It is endowed with the scalar product

$$\langle \cdot, \cdot \rangle_\rho : (f, g) \mapsto \int_{\partial E} \rho fg.$$

As we need to take care of the orthogonality condition  $\int_\Omega \eta_E \eta'_\Phi = 0$  we introduce the space

$$X := \left\{ u \in W_0^{1,2}(\Omega) : \int_\Omega \eta_E u = 0 \right\} = \langle \eta_E \rangle^\perp. \quad (\text{A.11})$$

To use the spectral theorem we need to ensure that the solution mapping (analogous to  $T_\rho$ ) we consider is well-defined. This relies on the simplicity of the first eigenvalue  $\Lambda(E) = \lambda(\mathbf{1}_E)$ .

**Lemma 16.** *Let  $E$  be a critical shape in the sense of Definition 9. For any  $f \in L^2(\partial E)$ , for any  $\rho \in L^\infty(\partial E; [\delta; 1/\delta])$  ( $\delta > 0$ ) there exists a unique solution  $v_{f,\rho}$  to the equation*

$$\begin{cases} -\Delta v_{f,\rho} - \Lambda(E)v_{f,\rho} - \mathbf{1}_E v_{f,\rho} = 0 & \text{in } \Omega, \\ \llbracket \partial_\nu v_{f,\rho} \rrbracket = -\rho f & \text{on } \partial E, \\ v_{f,\rho} \in X \text{ defined in (A.11): } v_{f,\rho} \in W_0^{1,2}(\Omega) \text{ and } \int_\Omega v_{f,\rho} \eta_E = 0. \end{cases} \quad (\text{A.12})$$

*Lemma 16.* The first eigenvalue  $\lambda(\mathbf{1}_E) = \Lambda(E)$  is simple. Consequently, there exists a parameter  $\underline{m} > 0$  such that,  $X$  being the space introduced in (A.11),

$$\forall u \in X, \int_\Omega |\nabla u|^2 - \int_\Omega \mathbf{1}_E u^2 - \Lambda(E) \int_\Omega u^2 \geq \underline{m} \int_\Omega u^2. \quad (\text{A.13})$$

As a consequence, for any  $f \in L^2(\partial E)$ , the energy functional

$$W_{\rho,f} : X \ni u \mapsto \frac{1}{2} \left( \int_{\Omega} |\nabla u|^2 - \int_{\Omega} \mathbf{1}_E u^2 - \Lambda(E) \int_{\Omega} u^2 \right) - \int_{\partial E} \rho f u$$

is coercive. Consequently,  $W_{\rho,f}$  admits a minimiser, which is a solution of (A.12). Furthermore, once again by simplicity of  $\Lambda(E)$ , there is at most one solution of (A.12). The proof is concluded.  $\square$

Define the operator  $T_{\rho} : L^2_{\rho}(\partial E) \rightarrow L^2(\partial E)$  as follows: for any  $f \in L^2_{\rho}(\partial E)$  let  $v_{f,\rho} \in W^{1,2}(\Omega)$  be the unique solution of (A.12). Let  $\text{Tr}_E : W^{1,2}(\Omega) \rightarrow L^2(\partial E)$  be the trace operator.  $E$  is analytic (Lemma 8) so that this operator is well-defined. Introduce

$$T_{\rho}(f) := \text{Tr}_E(v_{f,\rho}). \quad (\text{A.14})$$

While the symmetry and compactness of the operator  $T_{\rho}$  are immediate, the positivity uses once more the simplicity of  $\Lambda(E)$ . Indeed, for any  $f \in L^2_{\rho}(\partial E)$  we have

$$\begin{aligned} \int_{\partial E} \rho f T_{\rho}(f) &= \int_{\partial E} \rho f v_{f,\rho} \\ &= \int_{\Omega} |\nabla v_{f,\rho}|^2 - \int_{\Omega} \mathbf{1}_E v_{f,\rho}^2 - \Lambda(E) \int_{\Omega} v_{f,\rho}^2 \\ &\geq \underline{m} \int_{\Omega} v_{f,\rho}^2 \geq 0 \text{ from (A.13),} \end{aligned}$$

and the last inequality is strict unless  $f \equiv 0$ .

We may thus apply the spectral theorem: there exists a sequence of eigenpairs  $\{(\sigma_{k,\rho}, \psi_{k,\rho})\}_{k \in \mathbb{N}} \in (\mathbb{R}_+^* \times L^2_{\rho}(\partial E))^{\mathbb{N}}$  such that

$$\sigma_{k,\rho} \text{ is non-increasing in } k \text{ and } \sigma_{k,\rho} \xrightarrow[k \rightarrow \infty]{} 0,$$

$$\{\psi_{k,\rho}\}_{k \in \mathbb{N}} \text{ is an orthonormal basis of } L^2_{\rho}(\partial E)$$

and

$$\forall k \in \mathbb{N}, \quad T_{\rho}(\psi_{k,\rho}) = \sigma_{k,\rho} \psi_{k,\rho}.$$

We set  $\phi_{k,\rho} := v_{\psi_{k,\rho}}$  to extend it to a function on  $\Omega$ . Hence,

$$\begin{cases} -\Delta \phi_{k,\rho} - \Lambda(E) \phi_{k,\rho} - \mathbf{1}_E \phi_{k,\rho} = 0 & \text{in } \Omega, \\ \llbracket \partial_{\nu} \phi_{k,\rho} \rrbracket = -\rho \psi_{k,\rho} = -\frac{1}{\sigma_{k,\rho}} \rho \phi_{k,\rho} & \text{on } \partial E, \\ \phi_{k,\rho} \in X : \phi_{k,\rho} \in W_0^{1,2}(\Omega), \int_{\Omega} \phi_{k,\rho} \eta_E = 0. \end{cases} \quad (\text{A.15})$$

We thus obtain, defining

$$\lambda_{k,\rho} := \frac{1}{\sigma_{k,\rho}}$$

the system

$$\forall k \in \mathbb{N}^*, \quad \begin{cases} -\Delta \phi_{k,\rho} - \Lambda(E) \phi_{k,\rho} - \mathbf{1}_E \phi_{k,\rho} = 0 & \text{in } \Omega, \\ \llbracket \partial_{\nu} \phi_{k,\rho} \rrbracket = -\lambda_{k,\rho} \rho \phi_{k,\rho} & \text{on } \partial E, \\ \phi_{k,\rho} \in X : \phi_{k,\rho} \in W_0^{1,2}(\Omega), \int_{\Omega} \phi_{k,\rho} \eta_E = 0 \end{cases}$$



$$\text{and } \forall k, k' \in \mathbb{N}^*, \int_{\partial E} \rho \phi_{k,\rho} \phi_{k',\rho} = \delta_{k,k'}. \quad (\text{A.16})$$

Thus the family  $\{\phi_{0,\rho}\} \cup \{\phi_{k,\rho}\}_{k \in \mathbb{N}^*} = \{\phi_{k,\rho}\}_{k \in \mathbb{N}}$  is an orthonormal basis of  $L^2_\rho(\partial E)$  for the scalar product  $\langle \cdot, \cdot \rangle_\rho$ .

From the Courant-Fisher principle we have, furthermore, the following characterisation of the first eigenvalue:

$$\lambda_{0,\rho} = \min_{u \in X, \int_{\partial E} u^2 > 0} \frac{\int_\Omega |\nabla u|^2 - \int_\Omega \mathbf{1}_E u^2 - \Lambda(E) \int_\Omega u^2}{\int_{\partial E} \rho u^2}. \quad (\text{A.17})$$

**Diagonalisation of  $\Theta''_E(E)$**  We start from (B.7) and we seek to determine the weight  $\rho$ . We use the following notational convention: for any  $\Phi \in W^{2,\infty}(\Omega; \mathbb{R}^d)$ ,

$$\varphi := \langle \Phi, \nu \rangle, \tilde{\varphi} := \frac{1}{\rho} \varphi.$$

The first-order shape derivative  $\eta'_\Phi$  satisfies

$$\begin{cases} -\Delta \eta'_\Phi - \Lambda(E) \eta'_\Phi - \mathbf{1}_E \eta'_\Phi = 0 & \text{in } \Omega, \\ \left[ \frac{\partial \eta'_\Phi}{\partial \nu} \right] = -\rho \tilde{\varphi} & \text{on } \partial E, \\ \eta'_\Phi \in X : \eta'_\Phi \in W_0^{1,2}(\Omega) \text{ and } \int_\Omega \eta'_\Phi \eta_E = 0. \end{cases} \quad (\text{A.18})$$

We decompose  $\tilde{\varphi}$  in the basis  $\{\phi_{k,\rho}\}_{k \in \mathbb{N}}$  as

$$\tilde{\varphi} = \sum_{k=0}^{\infty} \alpha_{k,\rho} \phi_{k,\rho}. \quad (\text{A.19})$$

Thus,

$$\eta'_\Phi = \sum_{k=0}^{\infty} \frac{\alpha_{k,\rho}}{\lambda_{k,\rho}} \phi_{k,\rho}.$$

As a conclusion

$$\begin{aligned} \frac{1}{2\mu_E} \Theta''_E(E)[\Phi, \Phi] &= - \int_{\partial E} \eta'_\Phi \langle \Phi, \nu_E \rangle - \int_{\partial E} \frac{\partial \eta_E}{\partial \nu} \langle \Phi, \nu_E \rangle^2 \\ &= - \int_{\partial E} \rho \tilde{\varphi} \eta'_\Phi - \int_{\partial E} \rho^2 \frac{\partial \eta_E}{\partial \nu} (\tilde{\varphi})^2. \end{aligned}$$

Let us choose

$$\rho_E = - \frac{1}{\partial_\nu \eta_E}.$$

From Lemma 3, we have  $\rho_E \in L^\infty([\delta; 1/\delta])$  for  $\delta$  small enough. Finally, this yields

$$\begin{aligned} \frac{1}{2\mu_E} \Theta''_E(E)[\Phi, \Phi] &= - \int_{\partial E} \eta'_\Phi \langle \Phi, \nu_E \rangle - \int_{\partial E} \frac{\partial \eta_E}{\partial \nu} \langle \Phi, \nu_E \rangle^2 \\ &= - \int_{\partial E} \rho_E \tilde{\varphi} \eta'_\Phi - \int_{\partial E} \rho_E^2 \frac{\partial \eta_E}{\partial \nu} (\tilde{\varphi})^2 \\ &= \sum_{k=0}^{\infty} \alpha_{k,\rho_E}^2 \left( 1 - \frac{1}{\lambda_{k,\rho_E}} \right). \end{aligned}$$

In particular, given that the sequence  $\{\lambda_{k,\rho_E}\}_{k \in \mathbb{N}}$  is non-decreasing, we have the estimate from below:

$$\frac{1}{2\mu_E} \Theta_E''(E)[\Phi, \Phi] \geq \left(1 - \frac{1}{\lambda_{0,\rho_E}}\right) \sum_{k=0}^{\infty} \alpha_{k,\rho_E}^2 = \left(1 - \frac{1}{\lambda_{0,\rho_E}}\right) \|\langle \Phi, \nu_E \rangle\|_{L^2(\partial E)}^2. \quad (\text{A.20})$$

Hence we have the following sufficient condition for the stability of a critical shape:

**Lemma 17.** *If*

$$\lambda_{0,\rho_E} > 1$$

*then  $E$  is stable in the sense of Definition 14.*

To prove Proposition 15 we study the asymptotic behaviour of  $\lambda_{0,\rho_E}$  as  $V_0 \rightarrow 1$ .

**Asymptotic behaviour of  $\lambda_{0,\rho_E}$  as  $V_0 \rightarrow 1$**  We prove here the following lemma:

**Lemma 18.** *There holds*

$$\min_{E \text{ critical shape}} \lambda_{0,E} \xrightarrow{V_0 \rightarrow 1} \infty.$$

*Proof of Lemma 18.* From Lemma 3 and the Rayleigh quotient formulation (A.17) of  $\lambda_{0,\rho_E}$  we have, for  $V_0 \in (1 - \varepsilon_0; \varepsilon)$ ,

$$\lambda_{0,\rho_E} \geq \delta_0 \min_{u \in X, \int_{\partial E} u^2 > 0} \frac{\int_{\Omega} |\nabla u|^2 - \int_{\Omega} \mathbf{1}_E u^2 - \Lambda(E) \int_{\Omega} u^2}{\int_{\partial E} u^2}.$$

As in Lemma 31 we argue by contradiction: assume that there exists a sequence  $\varepsilon \rightarrow 0$ , a constant  $M_0 > 0$ , and, for any  $\varepsilon > 0$ , a critical set  $E_\varepsilon$  such that  $\mathbf{1}_{E_\varepsilon} \in \mathcal{F}(V_0)$  with  $V_0 \in (1 - \varepsilon; 1)$  that satisfies

$$\lambda_{0,\rho_{E_\varepsilon}} \leq M_0.$$

Define, for any  $\varepsilon > 0$ ,

$$\lambda(\varepsilon) := \min_{u \in X, \int_{\partial E_\varepsilon} u^2 \neq 0} \frac{\int_{\Omega} |\nabla u|^2 - \int_{\Omega} \mathbf{1}_{E_\varepsilon} u^2 - \Lambda(E_\varepsilon) \int_{\Omega} u^2}{\int_{\partial E_\varepsilon} u^2}.$$

By assumption the sequence  $\{\lambda(\varepsilon)\}_{\varepsilon \rightarrow 0}$  is uniformly bounded. In other words we have

$$0 < \lambda(\varepsilon) \leq M_1 < \infty \quad (\text{A.21})$$

for some suitable constant  $M_1$ . We define, for any  $\varepsilon > 0$ ,  $\varphi_\varepsilon$  as the first eigenfunction associated with  $\lambda(\varepsilon)$ ; in other words,  $\varphi_\varepsilon$  solves the equation

$$\begin{cases} -\Delta \varphi_\varepsilon - \Lambda(E_\varepsilon) \varphi_\varepsilon - \mathbf{1}_{E_\varepsilon} \varphi_\varepsilon = 0 & \text{in } \Omega, \\ \left[ \frac{\partial \varphi_\varepsilon}{\partial \nu} \right] = -\lambda(\varepsilon) \varphi_\varepsilon & \text{on } \partial E_\varepsilon, \\ \varphi_\varepsilon \in X, \\ \int_{\partial E_\varepsilon} \varphi_\varepsilon^2 = 1. \end{cases} \quad (\text{A.22})$$

In the same way we proved that (A.21) implied (2.41), it is easy to show that if (A.21) holds, then:

$$\forall \beta \in (0; 1), \exists c_\beta, \sup_{\varepsilon > 0} \|\varphi_\varepsilon\|_{\mathcal{C}^{0,\beta}} \leq c_\beta. \quad (\text{A.23})$$

The only difficulty here is to initialise the bootstrap to obtain higher integrability of  $\varphi_\varepsilon$  but this relies once again on the spectral gap estimate (A.13), as we immediately obtain

$$\int_{\Omega} |\nabla \varphi_\varepsilon|^2 \leq M \left( \lambda(\varepsilon) + \int_{\Omega} \varphi_\varepsilon^2 \right) \leq M''$$

for some  $M''$  under the assumption (A.21). The rest of the proof is identical.  $\square$

We then prove the coercivity result (Proposition 15) with the same proof as that of Proposition 29.

**Local quantitative inequalities around critical shapes** As in [39] we have the following results:

**Lemma 19.** *Let  $\varepsilon_1 > 0$  be given by Proposition 15. For any  $V_0 \in (1 - \varepsilon_1; 1)$  the critical shapes are isolated in the sense that*

$$\inf_{E, E' \text{ critical}, E' \neq E} |E' \Delta E| = \delta(V_0) > 0.$$

**Lemma 20.** *Let  $\varepsilon_1$  be given by Proposition 15. For any  $V_0 \in (1 - \varepsilon_1; 1)$ , any critical shape  $E$  is a local  $L^1$  minimiser: there exist  $c_E > 0$  and  $r_E > 0$  such that, for any  $f \in \mathcal{F}(V_0)$  with  $\|f - \mathbb{1}_E\|_{L^1(\Omega)} \leq r_E$ , there holds*

$$\lambda(f) - \lambda(\mathbb{1}_E) \geq c_E \|f - \mathbb{1}_E\|_{L^1(\Omega)}^2.$$

## A.4 Conclusion of the proof of Theorem II

*Proof of Theorem II.* Finally, the proof of Theorem II is identical to that of Theorem I.  $\square$

## B Proof of Theorem III

Similar to the proof of Theorem II, when the proofs are identical to those of Theorem I we omit them. We introduce the analog of Definitions 7-8.

**Definition 21** (Critical points). *For any  $V_0 \in (0; 1)$ , for any  $f \in \mathcal{F}(V_0)$ , we say  $f$  is a **critical point** of  $J$  if:*

1. *There exists a unique  $\mu_f = \mu_{u_f, V_0}$  such that*

$$\text{Vol}(\{p_f > \mu_f\}) = V_0 \text{Vol}(\Omega) = \text{Vol}(\{p_f \geq \mu_f\}),$$

2. *and  $f = \mathbb{1}_{\{p_f > \mu_f\}}$ .*

A **critical set** is a subset  $E$  of  $\Omega$  such that  $\mathbb{1}_E \in \mathcal{F}(V_0)$  and such that  $\mathbb{1}_E$  is a critical point of  $J$ .

**Definition 22.** *An admissible control  $f^* \in \mathcal{F}(V_0)$  is called a **stable local minimiser** of  $J$  in  $\mathcal{F}(V_0)$  if there exist  $C = C(f^*)$ ,  $\delta = \delta(f^*) > 0$  such that*

$$\forall f \in \mathcal{F}(V_0), \|f - f^*\|_{L^1(\Omega)} \leq \delta(f^*) \Rightarrow J(f) \geq J(f^*) + C(f^*) \|f - f^*\|_{L^1(\Omega)}^2.$$

## B.1 First consequences of a large volume constraint

We show regularity properties similar to those of Lemma 13. We let, for any  $f \in \mathcal{F}(V_0)$ ,  $\mu_{f,V_0}$  be the unique real number such that

$$\frac{\text{Vol}(\{p_f > \mu_{f,V_0}\})}{\text{Vol}(\Omega)} \leq V_0 \leq \frac{\text{Vol}(\{p_f \geq \mu_{f,V_0}\})}{\text{Vol}(\Omega)}.$$

We also define

$$\omega_{f,V_0} := \{p_f > \mu_{f,V_0}\}.$$

**Lemma 23.** *There exists  $\varepsilon_0 > 0$  and  $\delta_0 > 0$  such that, for any  $V_0 \in (1 - \varepsilon_0; 1)$ , for any  $f \in \mathcal{F}(V_0)$ , we have on the one hand*

$$\text{Vol}(\{p_f = \mu_{f,V_0}\}) = 0$$

and, on the other hand, the following regularity properties:

1.  $\omega_{f,V_0}$  has a  $\mathcal{C}^{3,\alpha}$  boundary (for any  $\alpha \in (0; 1)$ ),
- 2.

$$\min_{\partial\omega_{f,V_0}} \left| \frac{\partial p_f}{\partial \nu} \right| \geq \delta_0,$$

3.  $\partial\omega_{f,V_0}$  is locally a graph over  $\partial\Omega$ .

Finally, for any  $\alpha \in (0; 1)$ ,  $\partial\omega_{f,V_0}$  converges to  $\partial\Omega$  in  $\mathcal{C}^{3,\alpha}$  as  $V_0 \rightarrow 1$ , uniformly in  $f \in \mathcal{F}(V_0)$ .

*Lemma 23.* Here the proof is slightly different; by using an adjoint state, we get more regularity on the free boundary of the set  $\omega_{f,V_0}$ . We already observed in the proof of Lemma 13 that for any  $\alpha \in (0; 1)$  there held  $\sup_{V_0 \in (0;1), f \in \mathcal{F}(V_0)} \|u_f\|_{\mathcal{C}^{1,\alpha}(\bar{\Omega})} < \infty$ . By standard Schauder estimates applied to equation (1.5) we get

$$\sup_{V_0 \in (0;1), f \in \mathcal{F}(V_0)} \|p_f\|_{\mathcal{C}^{3,\alpha}(\bar{\Omega})} < \infty.$$

From (2.4) is also clear that

$$\forall \alpha \in (0; 1), \lim_{V_0 \rightarrow 1} \sup_{f \in \mathcal{F}(V_0)} \|p_f - p_\Omega\|_{\mathcal{C}^{3,\alpha}(\Omega)} = 0 \quad (\text{B.1})$$

where  $p_\Omega$  is the unique solution to

$$\begin{cases} -\Delta p_\Omega = \frac{\partial j}{\partial u}(x, w_\Omega) & \text{in } \Omega, \\ p_\Omega = 0 & \text{on } \partial\Omega. \end{cases} \quad (\text{B.2})$$

In this equation  $w_\Omega$  is the torsion function (2.1). As  $w_\Omega > 0$  and as  $\partial_u j(x, \cdot) > 0$  the maximum principle of Hopf entails that  $\inf_{\Omega(r_0)} |\nabla p_\Omega| > 0$  in the tubular neighbourhood  $\Omega(r_0)$  defined in (2.8) for  $r_0 > 0$  small enough. The rest of the proof is identical and the implicit function theorem implies that for  $V_0$  close enough to 1  $\partial\omega_{f,V_0}$  is  $\mathcal{C}^{3,\alpha}$ . □

Similarly we obtain the existence of a constant  $M$  such that

$$\forall V_0 \in (1 - \varepsilon_0; 1), \forall f \in \mathcal{F}(V_0), \text{Lip}(\partial\omega_{f,V_0}) + \text{Per}(\omega_{f,V_0}) \leq M. \quad (\text{B.3})$$

We also give the following information:

**Lemma 24.** *There exists  $\varepsilon'_0 > 0$  such that, for any  $V_0 \in (1 - \varepsilon'_0; 1)$ , for any  $f \in \mathcal{F}(V_0)$  the boundary  $\partial\omega_{f,V_0}$  is connected.*

We may, analogously to Lemma 14, take  $\varepsilon_0 = \varepsilon'_0$  where  $\varepsilon_0$  is given in Lemma 23.

## B.2 Applications of the quantitative bathtub principle

We apply the quantitative bathtub principle ([39, Proposition 26]) and obtain the following results for the thresholding algorithm (4).

**Lemma 25.** *Let  $\varepsilon_0, \delta_0$  be in the conditions of Lemma 23. For any  $V_0 \in (1 - \varepsilon_0; 1)$ , for any  $f_0 \in \mathcal{F}(V_0)$ , the sequence  $\{f_k\}_{k \in \mathbb{N}}$  is uniquely defined (in the sense that for any  $k \in \mathbb{N}$  there holds  $\text{Vol}(\{p_{f_k} = \mu_{f_k, V_0}\}) = 0$ ) and there holds*

$$\sum_{k=0}^{\infty} \|f_{k+1} - f_k\|_{L^1(\Omega)}^2 < \infty. \quad (\text{B.4})$$

**Lemma 26.** *Let  $\varepsilon_0$  be in the conditions of Lemma 13. For any  $V_0 \in (1 - \varepsilon_0; 1)$ , for any initialisation  $f_0 \in \mathcal{F}(V_0)$ , any  $L^\infty - *$  closure point  $f_\infty$  of the sequence  $\{f_k\}_{k \in \mathbb{N}}$  generated by the thresholding algorithm is a bang-bang function:*

$$\exists E_\infty \subset \Omega, f_\infty = \mathbb{1}_{E_\infty}. \quad (\text{B.5})$$

Furthermore  $E_\infty = \{p_{f_\infty} > \mu_{u_{f_\infty}, V_0}\}$ , which is uniquely defined. In other words,  $E_\infty$  is a critical set in the sense of Definition 21.

We obtain the following dichotomy (see Lemma 18).

**Lemma 27.** *Let  $\varepsilon_0$  be chosen as in Lemma 23. For any  $V_0 \in (1 - \varepsilon_0; 1)$ , for any initialisation  $f_0 \in \mathcal{F}(V_0)$ , let  $\{f_k\}_{k \in \mathbb{N}}$  be the sequence generated by the thresholding algorithm. Then:*

1. *Either  $\{f_k\}_{k \in \mathbb{N}}$  has a unique closure point  $f_\infty$ , in which case  $f_k \xrightarrow[k \rightarrow \infty]{} f_\infty$  strongly in  $L^1(\Omega)$ ,*
2. *Or  $\{f_k\}_{k \in \mathbb{N}}$  has an infinite number of closure points.*

**The goal of the next sections** In the following section we study the shape Hessians at critical shape. This is where the main differences with the proofs of Theorems I-II occur.

## B.3 Qualitative study of critical points

In this section we analyse the critical sets (in the sense of Definition 21). For the second order shape derivatives of the Lagrangians to be meaningful we need the critical sets to have an (at least)  $\mathcal{C}^3$  boundary, which is guaranteed by Lemma 23.

**Shape derivative formalism** We associate to the functional  $J$  a shape functional  $\mathcal{J}$ , by defining

$$\mathcal{J} : \Omega \supset E \mapsto J(\mathbb{1}_E).$$

For any compact  $\mathcal{C}^3$  subset  $E$  of  $\Omega$  and for any compactly supported vector field  $\Phi \in W^{2,\infty}(\Omega; \mathbb{R}^d)$ , we can define, for any  $t \in (-1; 1)$  small enough, the function

$$e_{E,\Phi}(t) := \mathcal{J}((\text{Id} + t\Phi)E).$$

From the same arguments that established the differentiability of  $E \mapsto u_{\mathbb{1}_E}$  (Lemma 23) that  $\mathcal{J}$  is twice shape differentiable, and we define:

$$\mathcal{J}'(E)[\Phi] = e'_{E,\Phi}(0), \text{ resp. } \mathcal{J}''(E)[\Phi, \Phi] = e''_{E,\Phi}(0).$$

**First order shape derivative and definition of the Lagrange multiplier** We use the formulas given in (2.24).

**Definition 28** (Critical shape). *A  $\mathcal{C}^3$  shape  $E$  is a **critical shape** for  $\mathcal{J}$  : if, for any compactly supported  $\Phi \in W^{2,\infty}(\Omega)$ ,*

$$\int_{\partial E} \langle \Phi, \nu_E \rangle = 0 \Rightarrow \mathcal{J}'(E)[\Phi] = 0.$$

We can now compute the Lagrange multiplier. Let  $E$  be a critical shape; then there exists  $\mu_E$  such that, for any compactly supported vector field  $\Phi \in W^{2,\infty}(\Omega; \mathbb{R}^d)$ ,

$$(\mathcal{J} + \mu_E \text{Vol})'(E)[\Phi] = 0.$$

We need, as in Lemma 23, the expression of the first-order shape derivative; we give it in the following statement:

**Lemma 29.** *For any  $\mathcal{C}^3$  shape, for any compactly supported vector field  $\Phi \in W^{2,\infty}(\Omega; \mathbb{R}^d)$ ,*

$$\mathcal{J}'(E)[\Phi] = \int_{\partial E} p_{\mathbb{1}_E} \langle \Phi, \nu_E \rangle.$$

Furthermore, the shape derivative of the map  $E \mapsto u_{\mathbb{1}_E}$  at  $E$  in the direction  $\Phi$  is the solution of (2.25).

Similar to Lemma 24 we have the following statement:

**Lemma 30.** *A shape  $E \subset \Omega$  is a critical set in the sense of Definition 21 if, and only if it is a critical shape in the sense of Definition 28.*

We now define the shape Lagrangian

**Definition 31** (Shape Lagrangian). *Let  $E$  be a critical shape for  $\mathcal{F}$ . The **associated shape Lagrangian** is*

$$\mathcal{K}_E := \mathcal{J} + \mu_E \text{Vol}$$

where  $E = \{p_{\mathbb{1}_E} > \mu_E\}$ .

With the above ingredients at hand, we now move to second-order shape derivatives.

**First computations and elements for shape Hessians** We first express the shape hessian of the Lagrangian. To express it we need to compute the derivative of  $p_E$ ; through the same argument as in [41] we obtain that the map  $E \mapsto p_E$  is differentiable and that, for any compactly supported vector field  $\Phi \in W^{2,\infty}(\Omega; \mathbb{R}^d)$ , its shape derivative  $p'_\Phi$  satisfies

$$\begin{cases} -\Delta p'_\Phi = u'_\Phi \partial_{uu}^2 j(x, u_\Phi) & \text{in } \Omega, \\ p'_\Phi \in W_0^{1,2}(\Omega). \end{cases} \quad (\text{B.6})$$

**Lemma 32.** *Let  $E$  be a critical shape in the sense of Definition 28 and let  $\Phi \in W^{2,\infty}(\Omega; \mathbb{R}^d)$  be a compactly supported vector field. The second-order shape derivative of the shape Lagrangian  $\mathcal{K}_E$  (defined in Definition 31) is given by the expression*

$$\mathcal{K}''_E(E)[\Phi, \Phi] = \int_{\partial E} p'_\Phi \langle \Phi, \nu_E \rangle + \int_{\partial E} \frac{\partial p_E}{\partial \nu} \langle \Phi, \nu_E \rangle^2 \quad (\text{B.7})$$

where  $p'_\Phi$  solves (B.6).

We want to prove that critical shapes are stable in the following sense:

**Definition 33** ( $L^2$ -stability). *A critical shape  $E$  is said to be  $L^2$ -stable if there exists a constant  $c_E > 0$  such that for any compactly supported vector field  $\Phi \in W^{2,\infty}(\Omega; \mathbb{R}^d)$ ,*

$$\mathcal{K}_E''(E)[\Phi, \Phi] \geq c_E \|\langle \Phi, \nu \rangle\|_{L^2(\partial E)}^2.$$

Our main result is the following proposition:

**Proposition 34.** *There exists  $\varepsilon_1 > 0, \varepsilon_1 \leq \varepsilon_0, \underline{c} > 0$  such that, for any  $V_0 \in (1 - \varepsilon_1; 1)$ , for any critical set  $E$  (in the sense of Definition 28), for any compactly supported vector field  $\Phi \in W^{2,\infty}(\Omega; \mathbb{R}^d)$ ,*

$$\mathcal{K}_E''(E)[\Phi, \Phi] \geq \underline{c} \|\langle \Phi, \nu \rangle\|_{L^2(\partial E)}^2.$$

The diagonalisation basis is defined in the same way it was introduced when studying ( $\mathbf{P}_{\text{Dir}}$ )-( $\mathbf{P}_{\text{Eigen}}$ ). We consider a fixed critical shape  $E \subset \Omega$  and work with  $V_0 \in (1 - \varepsilon_0; 1)$ , where  $\varepsilon_0$  is given by Lemma 23. As  $p'_\Phi$  satisfies (B.6) and as  $\mathcal{K}'(E)[\Phi] = 0$ , the operator the eigenvalues of which we should be considering is the following: consider a weight  $\rho \in L^\infty(\partial E, [\delta; 1/\delta])$ ,  $\delta > 0$  and for any function  $f \in L^2(\partial E)$ , let  $v_f$  and  $q_f$  be, respectively, the unique solutions of

$$\begin{cases} -\Delta v_f = 0 & \text{in } \Omega, \\ \llbracket \partial_\nu v_f \rrbracket = -\rho f & \text{on } \partial E, \\ v_f \in W_0^{1,2}(\Omega) \end{cases} \quad (\text{B.8})$$

and

$$\begin{cases} -\Delta q_f = v_f \partial_{uu}^2 j(x, u_E) & \text{in } \Omega \\ q_f \in W_0^{1,2}(\Omega). \end{cases} \quad (\text{B.9})$$

Finally, define the operator

$$T_\rho : f \mapsto \text{Tr}_{\partial E}(q_f).$$

It is easily checked that  $T_\rho$  is symmetric, compact and positive. We can thus use the spectral decomposition theorem and obtain the existence of an increasing, diverging sequence of positive eigenvalues  $\{\lambda_{k,\rho}\}_{k \in \mathbb{N}}$  and a sequence of couples  $\{\phi_{k,\rho}, \psi_{k,\rho}\}_{k \in \mathbb{N}}$  such that, for any  $k \in \mathbb{N}$ , there holds

$$\begin{cases} -\Delta \phi_{k,\rho} = 0 & \text{in } \Omega, \\ -\Delta \psi_{k,\rho} = \partial_{uu}^2 j(x, u_{\mathbb{1}_E}) \phi_{k,\rho} & \text{in } \Omega, \\ \llbracket \partial_\nu \phi_{k,\rho} \rrbracket = -\lambda_{k,\rho} \rho \psi_{k,\rho} & \text{on } \partial E, \\ \psi_{k,\rho}, \phi_{k,\rho} \in W_0^{1,2}(\Omega), \\ \int_{\partial E} \rho \psi_{k,\rho}^2 = 1. \end{cases} \quad (\text{B.10})$$

But now observe that the eigen-system (B.10) may be recase in a simpler form (that involves a fourth-order operator). Indeed, by noticing that the strict convexity of  $j$  in  $u$  allows us to write

$$\phi_{k,\rho} = -\frac{1}{j''(x, u_{\mathbb{1}_E})} \Delta \psi_{k,\rho}$$

(B.10) rewrites

$$\begin{cases} \Delta \left( \frac{1}{j''(x, \mathbb{1}_E)} \Delta \psi_{k,\rho} \right) = 0 & \text{in } \Omega, \\ \llbracket \partial_\nu \frac{1}{\partial_{uu}^2 j(x, u_{\mathbb{1}_E})} \Delta \psi_{k,\rho} \rrbracket = \lambda_{k,\rho} \rho \psi_{k,\rho} & \text{on } \partial E, \\ \psi_{k,\rho} \in W_0^{1,2}(\Omega), \Delta \psi_{k,\rho} = 0 \text{ on } \partial \Omega. \end{cases} \quad (\text{B.11})$$

The form (B.11) has one advantage, which is that it makes the associated Rayleigh quotient more easily read; it is indeed simply given by

$$R_\rho : W^{2,2}(\Omega) \cap W_0^{1,2}(\Omega) \ni \psi \mapsto \frac{\int_\Omega \frac{1}{\partial_{uu}^2 j(x, u_{1_E})} (\Delta \psi)^2}{\int_{\partial E} \rho \psi^2}.$$

Now, as before, choosing

$$\rho_E := \frac{1}{\left| \frac{\partial p_E}{\partial \nu} \right|}$$

and decomposing  $\tilde{\phi} := \frac{1}{\rho_E} \langle \Phi, \nu \rangle$  as

$$\tilde{\phi} = \sum_{k=0}^{\infty} a_k \psi_k$$

it appears that we have the diagonalised expression

$$\mathcal{K}_E''(E)[\Phi, \Phi] \leq \left( \frac{1}{\lambda_{0, \rho_E}} - 1 \right) \sum_{k=0}^{\infty} a_k^2$$

whence the question of the coercivity of  $\mathcal{K}_E''$  boils down to studying the asymptotic behaviour of  $\lambda_{0, \rho}$  as  $V_0 \rightarrow 1$ .

In other words, we have the following stability criterion:

**Lemma 35.** *If*

$$\lambda_{0, \rho_E} > 1$$

*then  $E$  is stable in the sense of Definition 33.*

**Asymptotic behaviour of  $\lambda_{0, \rho_E}$  as  $V_0 \rightarrow 1$**  The following lemma is then almost the conclusion of this section:

**Lemma 36.** *There holds*

$$\min_{E \text{ critical shape}} \lambda_{0, \rho_E} \xrightarrow{V_0 \rightarrow 1} \infty.$$

*Proof of Lemma 36.* Notice that  $\lambda_{0, \rho_E}$  has the variational formulation

$$\lambda_{0, \rho_E} = \min_{\psi \in W^{2,2}(\Omega) \cap W_0^{1,2}(\Omega), \int_{\partial E} \psi^2 > 0} \frac{\int_\Omega \frac{1}{\partial_{uu}^2 j(x, u_{1_E})} (\Delta \psi)^2}{\int_{\partial E} \rho_E \psi^2}. \quad (\text{B.12})$$

From Lemma 23, the strict convexity of  $j$  and the fact that  $j''(x, \cdot)$  is bounded, and, finally, from the Rayleigh quotient formulation (B.12) of  $\lambda_{0, \rho_E}$  there exists a constant  $c_0 > 0$  such that for  $V_0 \in (1 - \varepsilon_0; \varepsilon)$ ,

$$\lambda_{0, \rho_E} \geq c_0 \min_{\psi \in W^{2,2}(\Omega) \cap W_0^{1,2}(\Omega), \int_{\partial E} \psi^2 > 0} \frac{\int_\Omega (\Delta \psi)^2}{\int_{\partial E} \psi^2}.$$

Following the strategy we introduced in Lemma 31 we argue by contradiction: assume that there exists a sequence  $\varepsilon \rightarrow 0$ , a constant  $M_0 > 0$ , and, for any  $\varepsilon > 0$ , a critical set  $E_\varepsilon$  such that  $\mathbb{1}_{E_\varepsilon} \in \mathcal{F}(V_0)$  with  $V_0 \in (1 - \varepsilon; 1)$  that satisfies

$$\lambda_{0, \rho_{E_\varepsilon}} \leq M_0.$$



Let for any  $\varepsilon > 0$

$$\xi(\varepsilon) := \min_{\psi \in W^{2,2}(\Omega) \cap W_0^{1,2}(\Omega), \int_{\partial E_\varepsilon} \psi^2 > 0} \frac{\int_\Omega (\Delta \psi)^2}{\int_{\partial E_\varepsilon} \psi^2}.$$

However notice now that by the Poincaré inequality, for any  $\psi \in W^{2,2}(\Omega) \cap W_0^{1,2}(\Omega)$  there holds

$$\int_\Omega (\Delta \psi)^2 \geq c_1 \int_\Omega |\nabla \psi|^2$$

for some constant  $c_1 > 0$ , so that

$$\xi(\varepsilon) \geq c_1 \lambda(\varepsilon)$$

where  $\lambda(\varepsilon)$  was defined in the proof of Theorem I. From Lemma 31 we deduce that  $\xi(\varepsilon) \xrightarrow{\varepsilon \rightarrow 0} \infty$ , the desired contradiction.  $\square$

We then prove the coercivity result (Proposition 34) with the same proof as that of Proposition 29.

**Local quantitative inequalities around critical shapes** Similarly, we have:

**Lemma 37.** *Let  $\varepsilon_1 > 0$  be given by Proposition 34. For any  $V_0 \in (1 - \varepsilon_1; 1)$  the critical shapes are isolated in the sense that*

$$\inf_{E, E' \text{ critical}, E' \neq E} |E' \Delta E| = \delta(V_0) > 0.$$

**Lemma 38.** *Let  $\varepsilon_1$  be given by Proposition 34. For any  $V_0 \in (1 - \varepsilon_1; 1)$ , any critical shape  $E$  is a local  $L^1$  minimiser: there exist  $c_E > 0$  and  $r_E > 0$  such that, for any  $f \in \mathcal{F}(V_0)$  with  $\|f - \mathbb{1}_E\|_{L^1(\Omega)} \leq r_E$ , there holds*

$$J(f) - J(\mathbb{1}_E) \geq c_E \|f - \mathbb{1}_E\|_{L^1(\Omega)}^2.$$

## B.4 Conclusion of the proof of Theorem III

The conclusion of the proof is identical to the conclusion of the proof of Theorem I.

## References

- [1] C. Amrouche, C. Conca, A. Ghosh, and T. Ghosh. Uniform  $W^{1,p}$ - estimates for an elliptic operator with Robin boundary condition in a  $C^1$  domain. *Calculus of Variations and Partial Differential Equations*, 59(2), Mar. 2020.
- [2] S. Amstutz. Analysis of a level set method for topology optimization. *Optim. Methods Softw.*, 26(4-5):555–573, 2011.
- [3] S. Amstutz and H. Andrä. A new algorithm for topology optimization using a level-set method. *J. Comput. Phys.*, 216(2):573–588, 2006.
- [4] G. Barles and C. Georgelin. A simple proof of convergence for an approximation scheme for computing motions by mean curvature. *SIAM Journal on Numerical Analysis*, 32(2):484–500, apr 1995.

- [5] E. Bednarczuk, M. Pierre, E. Rouy, and J. Sokolowski. Calculating tangent sets to certain sets in functional spaces. Research Report RR-3190, INRIA, 1997.
- [6] J. Bintz and S. Lenhart. Optimal resource allocation for a diffusive population model. *Journal of Biological Systems*, 28(04):945–976, dec 2020.
- [7] B. Braida, J. Dalphin, C. Dapogny, P. Frey, and Y. Privat. Shape and topology optimization for maximum probability domains in quantum chemistry. *Numerische Math.*, To appear.
- [8] J. Céa, A. Gioan, and J. Michel. Quelques resultats sur l'identification de domaines. *Calcolo*, 10(3-4):207–232, sep 1973.
- [9] A. Chambolle, I. Mazari-Fouquer, and Y. Privat. Stability of optimal shapes and convergence of thresholding algorithms in linear and spectral optimal control problems. Supplementary material.
- [10] S. Chanillo, D. Grieser, M. Imai, K. Kurata, and I. Ohnishi. Symmetry breaking and other phenomena in the optimization of eigenvalues for composite membranes. *Communications in Mathematical Physics*, 214(2):315–337, Nov. 2000.
- [11] S. Chanillo, D. Grieser, and K. Kurata. The free boundary problem in the optimization of composite membranes, 2000.
- [12] S. Chanillo and C. Kenig. Weak uniqueness and partial regularity for the composite membrane problem. *Journal of the European Mathematical Society*, pages 705–737, 2008.
- [13] S. Chanillo, C. E. Kenig, and T. To. Regularity of the minimizers in the composite membrane problem in  $\mathbb{R}^2$ . *Journal of Functional Analysis*, 255(9):2299–2320, Nov. 2008.
- [14] A. Cianchi and A. Ferone. A strengthened version of the Hardy-Littlewood inequality. *Journal of the London Mathematical Society*, 77(3):581–592, Feb. 2008.
- [15] R. Cominetti and J.-P. Penot. Tangent sets of order one and two to the positive cones of some functional spaces. *Applied Mathematics and Optimization*, 36(3):291–312, nov 1997.
- [16] M. Dambrine and J. Lamboley. Stability in shape optimization with second variation. *Journal of Differential Equations*, 267(5):3009–3045, Aug. 2019.
- [17] M. Dambrine and M. Pierre. About stability of equilibrium shapes. *ESAIM: Mathematical Modelling and Numerical Analysis*, 34(4):811–834, July 2000.
- [18] K. Eppler, H. Harbrecht, and R. Schneider. On convergence in elliptic shape optimization. *SIAM Journal on Control and Optimization*, 46(1):61–83, jan 2007.
- [19] L. C. Evans. Convergence of an algorithm for mean curvature motion. *Indiana University mathematics journal*, pages 533–557, 1993.
- [20] D. Gilbarg and N. Trudinger. *Elliptic Partial Differential Equations of Second Order*. Springer Berlin Heidelberg, 1983.
- [21] A. Henrot. *Extremum problems for eigenvalues of elliptic operators*. Frontiers in Mathematics. Birkhäuser Verlag, Basel, 2006.
- [22] A. Henrot and M. Pierre. *Shape Variation and Optimization*. European Mathematical Society Publishing House, Feb. 2018.

- [23] A. Henrot, M. Pierre, and M. Rihani. Positivity of the shape Hessian and instability of some equilibrium shapes. *Mediterr. J. Math.*, 1(2):195–214, 2004.
- [24] M. Hintermüller, C.-Y. Kao, and A. Laurain. Principal eigenvalue minimization for an elliptic problem with indefinite weight and Robin boundary conditions. *Applied Mathematics & Optimization*, 65(1):111–146, dec 2011.
- [25] H. Ishii. A generalization of the Bence, Merriman and Osher algorithm for motion by mean curvature. *Curvature flows and related topics*, pages 111–127, 1995.
- [26] H. Ishii, G. E. Pires, and P. E. Souganidis. Threshold dynamics type approximation schemes for propagating fronts. *Journal of the Mathematical Society of Japan*, 51(2), apr 1999.
- [27] C.-Y. Kao, Y. Lou, and E. Yanagida. Principal eigenvalue for an elliptic problem with indefinite weight on cylindrical domains. *Math. Biosci. Eng.*, 5(2):315–335, 2008.
- [28] C.-Y. Kao and S. A. Mohammadi. Extremal rearrangement problems involving Poisson’s equation with Robin boundary conditions. *Journal of Scientific Computing*, 86(3), feb 2021.
- [29] C.-Y. Kao, S. A. Mohammadi, and B. Osting. Linear convergence of a rearrangement method for the one-dimensional poisson equation. *Journal of Scientific Computing*, 86(1), jan 2021.
- [30] J. Lamboley, A. Laurain, G. Nadin, and Y. Privat. Properties of optimizers of the principal eigenvalue with indefinite weight and Robin conditions. *Calculus of Variations and Partial Differential Equations*, 55(6), Dec. 2016.
- [31] T. Laux and F. Otto. Convergence of the thresholding scheme for multi-phase mean-curvature flow. *Calculus of Variations and Partial Differential Equations*, 55(5):1–74, 2016.
- [32] T. Laux and D. Swartz. Convergence of thresholding schemes incorporating bulk effects. *Interfaces and Free Boundaries*, 19(2):273–304, 2017.
- [33] G. Leoni. *A First Course in Sobolev Spaces*. Graduate studies in mathematics. American Mathematical Society, 2nd. edition, 2017.
- [34] E. Lieb and M. Loss. *Analysis*. American Mathematical Society, Providence, Rhode Island, 2001.
- [35] I. Mazari. Quantitative inequality for the eigenvalue of a Schrödinger operator in the ball. *Journal of Differential Equations*, 269(11):10181–10238, Nov. 2020.
- [36] I. Mazari. *Shape optimization and spatial heterogeneity in reaction-diffusion equations*. Theses, Sorbonne Université, July 2020.
- [37] I. Mazari. Quantitative estimates for parabolic optimal control problems under  $L^\infty$  and  $L^1$  constraints in the ball: Quantifying parabolic isoperimetric inequalities. *Nonlinear Analysis*, 215:112649, Feb. 2022.
- [38] I. Mazari, G. Nadin, and Y. Privat. Optimisation of the total population size for logistic diffusive equations: bang-bang property and fragmentation rate. *Communications in Partial Differential Equations*, 47(4):797–828, dec 2021.
- [39] I. Mazari and D. Ruiz-Balet. Quantitative stability for eigenvalues of Schrödinger operator, quantitative bathtub principle, and application to the turnpike property for a bilinear optimal control problem. *SIAM Journal on Mathematical Analysis*, 54(3):3848–3883, jun 2022.

- [40] B. Merriman, J. Bence, and S. Osher. Diffusion generated motion by mean curvature. In L. A. Department of Mathematics, University of California, editor, *CAM Report 92-33*, 1992.
- [41] F. Mignot, J. Puel, and F. Murat. Variation d'un point de retournement par rapport au domaine. *Communications in Partial Differential Equations*, 4(11):1263–1297, 1979.
- [42] G. Nadin and A. I. T. Marrero. On the maximization problem for solutions of reaction–diffusion equations with respect to their initial data. *Mathematical Modelling of Natural Phenomena*, 15:71, 2020.
- [43] O. Pironneau. *Optimal Shape Design for Elliptic Systems*. Springer Berlin Heidelberg, 1984.
- [44] S. J. Ruuth, B. Merriman, J. Xin, and S. Osher. Diffusion-generated motion by mean curvature for filaments. *Journal of Nonlinear Science*, 11(6):473–493, 2001.
- [45] D. Swartz and N. K. Yip. Convergence of diffusion generated motion to motion by mean curvature. *Communications in Partial Differential Equations*, 42(10):1598–1643, sep 2017.
- [46] M. Yousefnezhad, C.-Y. Kao, and S. A. Mohammadi. Optimal chemotherapy for brain tumor growth in a reaction-diffusion model. *SIAM Journal on Applied Mathematics*, 81(3):1077–1097, jan 2021.