

PK/PD modelling of the CAMHB nutrients concentrations effect on polymyxin B antimicrobial activity against Acinetobacter baumannii

Mathilde Lacroix, Jérémy Moreau, Claudia Zampaloni, Caterina Bissantz, Hamasseh Shirvani, Sandrine Marchand, William Couet, Alexia Chauzy

▶ To cite this version:

Mathilde Lacroix, Jérémy Moreau, Claudia Zampaloni, Caterina Bissantz, Hamasseh Shirvani, et al.. PK/PD modelling of the CAMHB nutrients concentrations effect on polymyxin B antimicrobial activity against Acinetobacter baumannii. 33rd ECCMID: European Congress of Clinical Microbiology & Infectious Diseases, Apr 2023, Copenaghen, Denmark. hal-04141371

HAL Id: hal-04141371 https://hal.science/hal-04141371

Submitted on 26 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. Inserm







100%

TK tubes with 10° CFU/mL. PMB



PK/PD modelling of the CAMHB nutrients concentrations effect on polymyxin B antimicrobial activity against Acinetobacter baumannii

Mathilde Lacroix ^{1,2}, Jérémy Moreau ², Claudia Zampaloni ^{3*}, Caterina Bissantz ^{3**}, Hamasseh Shirvani ¹, Sandrine Marchand ^{2,4}, William Couet ^{2,4}, Alexia Chauzy ²

- ¹ Institut Roche Boulogne-Billancourt (France)
- ² INSERM U1070, University of Poitiers Poitiers (France)
- ³ Roche Pharma Research and Early Development, Roche Innovation Center Basel, F. Hoffmann-La Roche Ltd, Grenzacherstrasse 124 Basel (Switzerland)
 - ³* Immunology, Infectious Diseases and Opthalmology, ³** Pharmaceutical Science
- ⁴ Department of Toxicology and Pharmacokinetics, University Hospital of Poitiers Poitiers (France)

Background

Artificial culture media used for in vitro PK/PD experiments are intended to promote bacterial growth, not to mimic in vivo conditions, which complicates in vitro / in vivo extrapolations.

The aim of this study was to assess the effect of Cation Adjusted Mueller-Hinton Broth (CAMHB) dilutions on polymyxin B (PMB) activity against Acinetobacter baumannii in vitro, using a PK/PD modelling approach.

Materials and methods

Poster P2198

Time-kill experiments (TK) were performed using cation adjusted CAMHB at 100% or diluted at 50%, 25% and 10%, with 3 A.baumannii strains: ATCC 19606 (MIC_{PMB} = 1mg/L), and two clinical isolates¹ AB121 (MIC_{PMB}=0.5-1mg/L), and AB122 (MIC_{PMB}=32-64mg/L). The starting inoculum for TK was 10^6 CFU/mL. PMB concentrations were ranging between 0.125 and 256mg/L. Pre-culture unti 10⁸ CFU/mL in culture in Bacterial counts were determined (3 replicates) at various times up to 30h, CAMHB 100% **CAMHB 100%** Figure 1 : Time-kill experiment's design and data were used to develop a PK/PD model with NONMEM[®]. Data for each strain were analysed separately.

Results

CAMHB dilution had no effect on bacterial growth (controls: Figure 3, 1rst column for each strain), nor on the initial rapid CFU decay in the presence of PMB. However the regrowth observed for concentrations up to 8xMIC at CAMHB 100%, was progressively reduced with dilutions (Figure 3).

A PK/PD model including two bacterial subpopulations with different susceptibilities (S/R)² was used to describe TK data (Figure 2). The medium effect was characterized by an increase of PMB bactericidal effect with nutrients depletion, modeled by an increase of the Emax_R and EC50_R for ATCC 19606 and AB121 but by a decrease of EC_{50} (common to both S/R subpopulations) for AB122 (Figure 4).

Because cations interfere with PMB activity³ complementary experiments with adjusted Ca²⁺ and Mg²⁺ concentrations were conducted and showed that cations depletion was not responsible for the observed medium effect (data not shown).

ATCC 19606 & AB121

AB122

Figure 2 : PK/PD model representation Knet: apparent growth rate constant (h⁻¹)



S: susceptible bacterial subpopulation R: resistant bacterial subpopulation PMB: polymyxin B concentration (mg/L) Emax: maximum rate of PMB effect (h⁻¹) EC₅₀: PMB concentration needed to reach 50% of Emax (mg/L) y: Hill coefficient

Sampling at T0h, T1h

[3h, T6h, T10h, T24h and T30h

Serial dilutions and plating on Mueller

Hinton agar + 1% activated charcoal







Figure 4 : Drug effect (bacterial killing rate) as a function of PMB concentration and CAMHB dilution for susceptible and resistant bacterial subpopulations.

Figure 3 : Visual predictive checks (VPC) for the final model. Circles represent experimental data, solid lines depict the mean of simulated data and colored areas depict the 90% prediction interval for 1,000 simulated profiles.

References:

1. Jaidane et al., 2018 International Journal of Antimicrobial Agents 2. Mouton JW et al., 1997 Antimicrob Agents Chemotherapy 3. Bulitta et *al.,* 2010 Antimicrob Agents Chemotherapy

Conclusion

PK/PD modelling demonstrated an impaired apparent regrowth of A.baumannii with nutrients depletion, presumably due to an increased PMB bactericidal effect.

128 mg/L

128 mg/L

128 mg/L

Further experiments would be necessary to understand this effect which might change the mutation frequency or impact the triggering of adaptation mechanism(s) responsible of the apparent bacterial regrowth.

