



HAL
open science

Gravity-driven large-scale deformation system in the Tumbes-Guayaquil forearc basin, Northern Andes (Northern Peru-Southern Ecuador)

Andréa Peuzin, Marianne Saillard, Nicolas Espurt, François Michaud, Cédric Bulois, Daniel Praeg, Marc Régnier, Ysabel Calderon

► **To cite this version:**

Andréa Peuzin, Marianne Saillard, Nicolas Espurt, François Michaud, Cédric Bulois, et al.. Gravity-driven large-scale deformation system in the Tumbes-Guayaquil forearc basin, Northern Andes (Northern Peru-Southern Ecuador). *Journal of Structural Geology*, 2023, 173, pp.104909. 10.1016/j.jsg.2023.104909 . hal-04143145


HAL Id: hal-04143145

<https://hal.science/hal-04143145v1>

Submitted on 27 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Gravity-driven large-scale deformation system in the Tumbes-Guayaquil forearc basin, Northern Andes (Northern Peru-Southern Ecuador)

Andréa Peuzin^{a,*}, Marianne Saillard^a, Nicolas Espurt^a, François Michaud^{a,b}, Cédric Bulois^{a,1}, Daniel Praeg^a, Marc Régnier^a, Ysabel Calderon^c

^a Université Côte d'Azur, IRD, CNRS, Observatoire de la Côte d'Azur, Géoazur, Valbonne, France

^b Sorbonne Université, Faculté des Sciences et Ingénierie, F-75252, Paris, France

^c PERUPETRO S.A., Lima, Luis Aldana 320, San Borja, 15034, Lima, Peru

ARTICLE INFO

Keywords:

Gravity tectonics
High sedimentation
Overpressured shales
Crustal deformation
North Peru-South Ecuador forearc

ABSTRACT

The offshore Tumbes-Guayaquil forearc basin in the accretionary prism of Northern Peru-Southern Ecuador shows evidence of gravity-driven large-scale deformation systems active during the Late Neogene-Quaternary period. Subsurface data and the construction of eight structural cross-sections show that the ~8 km-thick Oligocene-Quaternary sedimentary infill is detached seaward and completely decoupled from the underlying inner accretionary prism systems. The Corvina décollement in the Tumbes basin and the Posorja décollement in the Guayaquil basin constitute two thin-skinned gravity tectonic systems associated with kilometer-scale, updip "raft" extensional structures paired with downdip fold-thrust systems (Barracuda and Domito thrust systems). Although many previous studies have described the structural and stratigraphic architecture of the Tumbes-Guayaquil forearc basin, no model explicitly accounts for this anomalous large-scale gravity tectonics. We propose that this gravity tectonic style, more commonly observed in passive continental margins, is primarily controlled by the combination of tectonostratigraphic features, including crustal-scale transtensional deformation related to oblique convergence along the Northern Andean margin, basal décollement slope tilting, strong sediment accumulation, and the presence of overpressured shales.

1. Introduction

Because of their ability to record tectonic events, forearc basins are a key structural feature of oceanic subduction zones, although their dynamics still include some grey areas (e.g., Dickinson, 1995; Heuret et al., 2012; Tsuji et al., 2015; Noda, 2016; Vannucchi et al., 2016; Noda and Miyakawa, 2017). The offshore Tumbes-Gulf of Guayaquil forearc basin is part of the Northern Peruvian-Southern Ecuadorian accretionary prism developed over the Nazca (Farallon)-South American plate convergence system through the Late Cretaceous to Cenozoic period (Daly, 1989; Espurt et al., 2018; Aizprua et al., 2019; Jaillard, 2022, Fig. 1). The complex structural architecture and petroleum systems of this frontier forearc region have been strongly studied by academic and petroleum industry through numerous seismic reflection and exploration well data (e.g., Benitez, 1995; Deniaud et al., 1999; Deniaud, 2000;

Collot et al., 2002; Higley, 2004a, b; Calahoranno, 2005; Fernández et al., 2005; Witt et al., 2006; Cobos, 2010; Espurt et al., 2018; Reynaud et al., 2018; Aizprua et al., 2019; Lemgruber-Traby et al., 2020; Guzmán et al., 2022; Márquez et al., 2022). These studies show that the Tumbes-Gulf of Guayaquil forearc basin is characterized by: (1) a massive Cenozoic prograding sedimentary infill containing ductile basal shale layers (Higley, 2004a,b; Espurt et al., 2018); (2) a large crustal-scale strike-slip fault zone, the Puná fault zone that crosses obliquely the forearc basin, leading to significant transtensional deformation favoring available space and sedimentary trapping in the forearc basin (Benitez, 1995; Deniaud et al., 1999; Deniaud, 2000; Cobos, 2010). The combination of these sedimentary and tectonic features might have exerted a strong control on the kinematics of structural growth of the forearc basin.

This paper aims to present an original contribution with eight

* Corresponding author.

E-mail addresses: peuzin@geoazur.unice.fr (A. Peuzin), saillard@geoazur.unice.fr (M. Saillard), espurt@geoazur.unice.fr (N. Espurt), michaud@geoazur.unice.fr (F. Michaud), bulois@geologie.ens.fr (C. Bulois), praeg@geoazur.unice.fr (D. Praeg), regnier@geoazur.unice.fr (M. Régnier), ycalderon@perupetro.com.pe (Y. Calderon).

¹ Now at Laboratoire de Géologie, CNRS/UMR 8538, Ecole Normale Supérieure, PSL Research University, 24 Rue Lhomond, 75005 Paris, France.

structural cross-sections based on seismic reflection profiles and well data to understand the three-dimensional structure and dynamics of the Tumbes-Guayaquil forearc basin (Fig. 1). Here we present a new tectono-stratigraphic interpretation for the Tumbes-Guayaquil forearc basin, involving a tectonic style reported for the first time in the dynamics of the region. Results reveal a new view of the gravity tectonics that controlled the structural growth of the Tumbes-Guayaquil forearc depocenter. This large-scale gravity tectonics is active during the Late Neogene and the Quaternary in a context of ongoing oceanic subduction, rather classically observed in passive continental margins. We discuss the regional driving tectono-stratigraphic features controlling such atypical gravity tectonic style and its relationship with the geodynamic processes affecting the Northern Andean active margin.

2. Geological setting

2.1. Tectonic context

Currently, along the Northern Peruvian-Southern Ecuadorian active margin, the Nazca Plate is subducting N83°E-trending beneath the Northern Andes at a velocity of 60 mm/a (Villegas-Lanza et al., 2016, Fig. 1). The accretionary prism structure is characterized by the accretion of successive forearc depocenters, separated by thrust wedges involving continental, oceanic and sedimentary rocks (Espurt et al., 2018; Aizprua et al., 2019). The Tumbes and Gulf of Guayaquil offshore

basins form an approximately 16000 km² large, thick Cenozoic forearc depocenter lying between coastal basement wedges (Carpitas, Zorritos, Pallatanga and Santa Elena) and outer wedges (Banco Peru and Domito) made of off-scraped sediments (Fig. 1). The Tumbes and Gulf of Guayaquil basins are separated by the broad Barracuda antiform in the forearc basin center (Fig. 1; Fernández et al., 2005; Vega, 2009). The Gulf of Guayaquil basin is cut obliquely by the SW-trending Puná dextral strike-slip fault zone, which accommodates the relative motion of two crustal slivers: the North Andean Sliver, moving NE-ward and the Inca Sliver to the south, moving SE-ward (Nocquet et al., 2014; Alvarado et al., 2016; Villegas-Lanza et al., 2016, Fig. 1). The fault zone participates to the opening of the Gulf of Guayaquil basin (Deniaud, 2000; Witt et al., 2006). The collision of the Carnegie Ridge with the Ecuadorian margin could have controlled the North Andean Sliver NE-ward escape and favored the opening of the Gulf of Guayaquil during at least the Quaternary (Witt et al., 2006; Michaud et al., 2009, 2018, Fig. 1).

The historical seismicity of the Tumbes-Guayaquil forearc zone does not show large magnitude subduction earthquakes (only three ~ Mw 7–7.5 events in 1901, 1933 and 1953 were possibly tsunamigenic; Lockridge, 1984; Espinoza, 1992; Beauval et al., 2013; Ioualalen et al., 2014; Yepes et al., 2016; Chunga et al., 2018; Vaca et al., 2019). Low interseismic coupling and creeping on the subduction interface contrast with strong interseismic coupling regions to the north characterized by large earthquake occurrence (Chlieh et al., 2014; Nocquet et al., 2014; Villegas-Lanza et al., 2016). Crustal faults in the onshore and offshore




Fig. 1. Geodynamic and structural settings of the Tumbes-Guayaquil forearc basin. (a) Geodynamic map of the Central-North Andean active margin related to the Nazca-South American convergence plate system. The dotted red square indicates the location of the Tumbes-Guayaquil forearc basin. The black arrows show the relative convergence rate (mm/a) and the oblique convergence trend between the Nazca (NAZ) and South American (SAM) Plates and the relative motions of the North Andean Sliver (NAS) and the Inca Sliver (INS) in mm/a (from Villegas-Lanza et al., 2016). (b) Structural map of the Tumbes-Guayaquil forearc basin. The thick lines with numbers indicate the locations of the eight cross-sections (labelled 1 to 8) constructed in this study along seismic reflection profiles. The white and black circles correspond to exploration wells. M: Marina-1X well, C: Corvina-40-X-1 well, B: Barracuda-15-X-1 well, E: Esperanza-1 well, D: Domito-1 well, T: Tiburon-1 well. Major rivers are indicated in blue. Coordinate system is UTM zone 17S. IW: inner wedge; OW: outer wedge. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

region of the Gulf of Guayaquil basin are seismically active (e.g., Puná fault zone), as revealed by strong to moderate present-day seismicity (Alvarado et al., 2018; Vaca et al., 2019) and morpho-tectonics analysis (Dumont et al., 2005).

2.2. Lithostratigraphy

Many previous works have described the Cenozoic sedimentary infill across the Tumbes and Gulf of Guayaquil basins and surrounding areas based on seismic profile and well data, correlated with field observations (e.g., Séranne, 1987; Benitez, 1995; Jaillard et al., 1995; Deniaud, 2000; Higley, 2004a, b; Fernández et al., 2005; Witt et al., 2006; Fildani et al., 2008; Vega, 2009; Cobos, 2010; Espurt et al., 2018; Aizprua et al., 2019; Reynaud et al., 2018; Jaillard, 2022; Aizprua et al., 2022). The lithostratigraphy is presented using the Corvina-40-X-1 well in Peru and the Esperanza-1 well in Ecuador (Fig. 2) and described hereafter.

In the study area, basement rocks are either exposed along the coast or revealed by exploration wells (Fernández et al., 2005; Vega, 2009;

Riel et al., 2014; Espurt et al., 2018; Aizprua et al., 2019; Lajo-Yáñez et al., 2022; Jaillard, 2022, Fig. 1). In North Peru, the Amotape-Tahuin basement is formed by Paleozoic to Triassic metamorphic and granitic rocks, unconformably covered by terrestrial to marine Eocene or lower Oligocene strata forming the Carpitás and Zorritos inner wedges along the coast (Fernández et al., 2005; Vega, 2009; Espurt et al., 2018). In South Ecuador, the Pallatanga-Piñon basement shows upper Cretaceous oceanic terranes unconformably covered by Paleocene to Eocene, terrestrial and deep marine strata forming the Pallatanga and Santa Elena inner wedges (Riel et al., 2014; Aizprua et al., 2019; Jaillard, 2022).

The sedimentary infill of the Tumbes-Guayaquil forearc depocenter is underlined by the upper Oligocene ductile deep marine shales (Heath Formation in Peru/Playa Rica Formation in Ecuador; Fig. 2). These shales form a potential source rock and are expected to have generated hydrocarbons in the deeper parts of the depocenters (Fildani et al., 2005; Lemgruber-Traby et al., 2020). The upper Oligocene shales are overlain by lower Miocene coarse-grained fluviodeltaic to deep-marine strata (Zorritos Formation in Peru/Subibaja Formation in Ecuador). These siliciclastic strata are unconformably overlain by middle Miocene-Pliocene marginal marine and deltaic strata (Cardalitos, Tumbes, and Mal Pelo Formations in Peru/Progreso and Lower Puná Formations in Ecuador), and finally by Quaternary claystones, sands, and dolomitized limestones (La Cruz Formation in Peru/Upper Puná Formation in Ecuador; Benitez, 1995; Deniaud, 2000; Fildani et al., 2005; Cobos, 2010; Reynaud et al., 2018; Fig. 2). Distal equivalents of the Tumbes-Guayaquil forearc depocenter sedimentary sequences are involved in the Banco Peru and Domito outer wedges (Fig. 1).

3. Subsurface structural data and interpretations

To illustrate the structural architecture and kinematics of structural growth of the offshore Tumbes-Guayaquil forearc basin, we show five, E- to ESE-trending serial cross-sections (labelled 1 to 5; Fig. 4) perpendicular to the forearc basin axis (approximately parallel to the thrust transport direction) and three ~ N- to NE-trending cross-sections (labelled 6 to 8; Fig. 5) parallel to the forearc basin axis (approximately parallel to the extensional trend in Guayaquil basin) (Figs. 1 and 3). Cross-sections 1, 2 and 6 are located in the Tumbes depocenter, and cross-sections 3, 4, 5, 7 and 8 in the Guayaquil depocenter (Figs. 1 and 3). The geology of these cross-sections has been interpreted using fifteen seismic reflection profiles (AIP92-19, Z1-3D extracted from Z1 seismic cube, VMX09-23, PC99-01, AIP92-61, VMX09-71, OXY98-114, AIP92-60, g83-s23e, g83-s23, g83-s23w, gt83-w23, s-9, g83-s09, g83-01, g83-n16, g83-n29; Supplementary Material Figs. S1 and S2) in second two-way travel time (sTWT) and six exploration wells (Marina-1X, Corvina-40-X-1, Barracuda-15-X-1, Esperanza-1, Domito-1, Tiburon-1) provided by Perupetro S.A. and Petroecuador (Figs. 4 and 5). The cross-sections range from 44 km to 58 km in length and reach 6–8 sTWT (second two-way travel time) in depth. Well data indicate that the entire sedimentary pile is relatively homogeneous across the basin. The seismic velocity values increases with depth from ~1800 m/s to ~3000 m/s through the Quaternary-Oligocene sedimentary section (Cobos, 2010; Espurt et al., 2018). We also used all available seismic reflection profile data and previously published data (Benitez, 1995; Deniaud, 2000; Collot et al., 2002; Fernández et al., 2005; Witt et al., 2006; Cobos, 2010; Espurt et al., 2018; Aizprua et al., 2019, 2022) to construct an isopach map of the Quaternary infill in sTWT of the Tumbes-Guayaquil forearc depocenter (Fig. 3).

3.1. Structures

The structural architecture of the Tumbes-Guayaquil forearc basin, off the coast of northern Peru and southern Ecuador, is described from south to north, from the Tumbes forearc depocenter to the Guayaquil forearc depocenter (Figs. 4–6).




Fig. 2. Stratigraphy of the Tumbes-Guayaquil forearc basin and lateral correlations in between based on the Corvina-40-X-1 (total depth ~4321 m Peru) and Esperanza-1 (total depth ~ 4000 m Ecuador) wells (For location, See Fig. 1). We have established a lateral correlation between the lithological nature and the ages of the formations. The upper Oligocene shales of the Heath Playa-Rica source rock correspond to top of overpressure conditions indicated by a blue circle with arrow (Fildani et al., 2005; Lemgruber-Traby et al., 2020; Perupetro S.A data). This blue circle indicates overpressured zone in the Corvina-40-X-1 well (Perupetro S.A data). Fm: Formation. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)




Fig. 3. Isopach map in sTWT of the Quaternary infill (La Cruz and Upper Puná Formations; as described in Fig. 2). The map reveals the position of the Guayaquil and Tumbes depocenters. The dotted contouring lines are 250 ms and intervals is 50 m. Location of available subsurface data are indicated: thin black lines are seismic profiles and red circles are exploration wells. The lines with numbers labelled and wells annotated with a letter are those used in this study. M: Marina-1X well, C: Corvina-40-X-1 well, B: Barracuda-15-X-1 well, E: Esperanza-1 well, D: Domito-1 well, T: Tiburon-1 well. Bathymetric data are extracted from the GEBCO site. Coordinate system is UTM zone 17S. FD: Forearc depocenter. IW: inner wedge; OW: outer wedge. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

3.1.1. The Tumbes forearc depocenter

The Tumbes forearc depocenter develops ahead of the Carpitas-Zorritos inner thrust wedge system involving metamorphic/granitic rocks and Cenozoic sediment (Fernández et al., 2005; Vega, 2009; Espurt et al., 2018). Cross-sections 1 and 2 (Fig. 4a and b and 6a), perpendicular to the trend of the forearc basin, show that the Tumbes forearc depocenter consists of a seaward-thickening sedimentary wedge of about 40–45 km-long. It contains a sedimentary section thicker than ~6 sTWT (~7.2 km) in thickness composed mainly by Oligocene-Neogene strata and thin Quaternary section of ~0.6 sTWT (~400m) (Fig. 3). The sedimentary pile is deformed by NE-trending listric normal faults of the Corvina fault system (Fig. 4a and b). Normal faults dip regionally basinward (with rare counter-regional faults) and branch downward onto a NW-dipping décollement level, the Corvina décollement, developed in the Oligocene ductile shales of the Heath Formation. The seaward sliding of the sedimentary cover of the Tumbes depocenter along the Corvina décollement is associated with typical rollover folds (Dula, 1991; Xiao and Suppe, 1992, Fig. 4a and b). Neogene and Quaternary strata in the hanging walls of listric normal faults are characterized by reflectors that exhibit fan-shaped geometries (Fig. 6a).

Seaward, the sedimentary pile is deformed by the 15 km-long Barracuda antiform (Fernández et al., 2005; Vega, 2009; Brusset et al., 2018) mainly developed on cross-section 2 (Fig. 4b and 6a). This structure is formed by thin-skinned imbrications of NW-verging thrusts connected at depth into the Oligocene ductile shales of the Heath Formation. The thrust branches upward into several blind back-thrusts that define a triangle zone (Fig. 6a). The Barracuda antiform can be

interpreted as a compressional structure accommodating the downdip gravitational sliding of the Corvina fault system as proposed by Vega (2009). The interpretation of the seismic profiles suggests that the Corvina décollement connects downdip to the Banco Peru outer wedge (Fig. 4b and 6a). This wedge is made of off-scraped distal Cenozoic sediments including oceanic mafic bodies (intrusive and volcanic) accreted above the subducting oceanic Nazca crust (Fig. 4a and b; Shepherd and Moberly, 1981; Fernández et al., 2005; Espurt et al., 2018). The southwestern side of the Banco Peru wedge is cut by the NNE-trending SE-dipping Banco Peru normal/strike-slip fault zone (Figs. 3 and 4a). This latter is only developed of the Tumbes depocenter and intersects the sea floor.

The NE-trending cross-section 6 (Fig. 5a), parallel to the trend of the depocenter, reveals that the NE-trending geometry of the basin remains similar to those revealed by the NW-trending cross-sections. However, the cross-section 6 shows NW-trending fault system including NE- and SW-dipping listric normal faults involves the Oligocene to Quaternary strata (Witt et al., 2006; Auguy et al., 2017; Brusset et al., 2018). This fault system and these several thin-skinned imbricates are connected at depth into the flat-lying Oligocene shale level of the Heath Formation (Fig. 4a and b and 5a).

3.1.2. The Guayaquil forearc depocenter

The Guayaquil forearc depocenter develops above the Pallatanga-Santa Elena inner thrust wedge system involving upper Cretaceous oceanic terranes and Paleocene-Eocene sediments (Aizprua et al., 2019). The wedge system is cut by the Puná crustal fault zone (Fig. 3). This fault zone runs SW-ward into the Guayaquil forearc depocenter through the




Fig. 4. Along-dip interpreted serial cross-sections of the Tumbes-Guayaquil forearc basin. For location, please refer to Figs. 1 and 3. The seismic profiles AIP92-19, Z1-3D, PC99-01, AIP92-61, VMX09-71, g83-s23e, g83-s23, g83-s23w, gt83-w23, s-9, g83-s09, g83-01 have been calibrated with the wells Domito-1, Esperanza-1, Corvina-40-X-1 wells. PIW: Pallatanga inner wedge. pjt.: projected.




Fig. 5. Along-strike interpreted cross-sections of the Tumbes-Guayaquil forearc basin. For locations, please refer to Figs. 1 and 3. The seismic profiles g83-n16, g83-n29 and OXY98-114 have been calibrated with the Tiburon-1, Domito-1, Barracuda 15-X-1 and Marina-1X wells. pjt.: projected.

Amistad antiform as suggested by seismic reflection and seismicity data (Witt et al., 2006; Vaca et al., 2019; Aizprua et al., 2019, 2022, Figs. 3 and 4c,d,e). It probably extends farther SW beneath the Barracuda antiform, although seismic reflection profiles do not show a fault zone beneath the Corvina décollement (Fig. 4b). Cross-sections 4, 5 and 6 (Fig. 4d and e), perpendicular to the trend of the Guayaquil forearc depocenter, show that the depocenter consists of a seaward-thickening sedimentary wedge of a maximum length of 50 km (decreasing to 30 km northward) and has an up to ~ 7.3 sTWT (~ 9 km-thick) Oligocene to Quaternary sedimentary section in its center. The Guayaquil forearc depocenter is marked by a much thicker Quaternary section than in the Tumbes depocenter, in excess of ~ 5 sTWT (~ 6 km) (Figs. 3 and 4d). Cross-sections show that the sedimentary pile is deformed by NW-trending listric normal faults of the Posorja fault system along the southern edge of the uplifted Santa Elena High (Fig. 3). In cross-sections 3, 4 and 5, the structural style consists of a broad deep rollover fold along the hanging wall of the Posorja décollement developed into the Oligocene ductile shales of the Playa Rica Formation (Fig. 4c,d,e). Upward, the décollement connects to inter-sequence lithologic contrast zones (e.g., erosional unconformities (Benitez, 1995; Deniaud, 2000); intra Neogene or Neogene-Quaternary interface along cross-sections 3 and 4) or reach the surface (northern cross-section 5). Like in the Tumbes forearc depocenter, the rollover folding is characterized by reflectors depicting spectacular fan-shaped geometries in the upper Neogene and especially in the Quaternary sediments (Fig. 4c,d,e and 6b).

Along-dip cross-sections 3, 4 and 5 (Fig. 4c,d,e) show that the Posorja décollement connects downdip with a ~ 25 km-large zone dominated by thin-skinned compressional structures deforming the Oligocene-Quaternary sequences. Seismic profiles show that some thrusts deform the rollover fold developed along the hanging wall of the Posorja décollement, while other thrusts, farther downdip, are associated with the Domito outer wedge (Fig. 4c,d,e and 6b). Cross-sections 7 and 8, parallel to the basin trend (Fig. 5b and c) show that the Posorja and Domito fault systems are associated with secondary NW-trending listric normal/strike-slip faults connected downdip into the Oligocene flat-lying shale layer.

3.2. Kinematics and timing of deformations

The subsurface data presented in the previous sections clearly show that the structural architecture of the Tumbes-Guayaquil forearc basin is characterized by two thin-skinned deformation domains: an updip

extensional zone defined by raft structures and a downdip contractional zone defined by fold-thrust systems (Figs. 4 and 5). The two domains are connected by narrow translational zones (~ 7 km) through décollements located in the Oligocene ductile shales (Heath and Playa Rica Formations) that lie directly above the seaward slope of the inner wedges (Fig. 6). The downdip fold-thrust structural systems are complex and comprise interference between structures related to frontal accretion-subduction and gravitational tectonics. In the Tumbes forearc depocenter, we interpret that the downdip Barracuda compressional structure balances part of the extension accommodated higher on the slope by the Corvina fault system (Fig. 4a and b). Thus, the growth of the Barracuda structure was mostly controlled by the downdip gravitational sliding of the Corvina raft system (Vega, 2009). Like the Tumbes forearc depocenter, part of the sedimentary infill of the Guayaquil depocenter is gravitationally rafted along the Posorja décollement toward the sea (Fig. 4d and e). Although the structures of the Banco Peru and Domito wedges are more compatible with frontal accretion related to subduction, we interpret that part of the thrusting and folding in these outer wedges could be therefore associated to gravitational tectonics. For instance, we interpret the current form of the rollover fold in cross-section 5 (Fig. 6a) as significantly deformed by a series of three thrusts connected onto the seaward-dipping Posorja décollement.

The sedimentary infill of the Tumbes-Guayaquil forearc basin contains syntectonic horizons that constrain the timing of deformation. Growth strata and submarine unconformities reveal that the Corvina-Barracuda structure of the Tumbes depocenter was active in the Middle(?)–Late Neogene (Fig. 4a and b; Brusset et al., 2018; Espurt et al., 2018). The onlap of the Quaternary strata on the Barracuda antiform and the truncation of its crest domain at the seafloor suggest ongoing deformation and uplift through the Quaternary until the present-day (Fig. 4b and 5). Seaward, the deformation of the Banco Peru outer accretionary wedge is attested by the pinch out of the Neogene-Quaternary strata (Fig. 4a and b; Brusset et al., 2018; Espurt et al., 2018). In the Guayaquil forearc depocenter, the activity of the Posorja raft system is recorded by spectacular Late Neogene and Quaternary growth strata fanning (Witt et al., 2006, Fig. 4c,d,e). The strong Quaternary sedimentation in the Guayaquil basin is related to large extensional displacement along the Posorja raft system. Synchronous and somewhat younger thrusting and folding in the Domito outer thrust wedge is locally recorded by upper Neogene and Quaternary growth strata (Fig. 4c,d,e). Deformation in this outer thrust wedge might therefore result from the combined effect of frontal accretion and gravitational sliding of the Posorja raft system.




Fig. 6. Line drawing of seismic profiles along the cross-sections 2 and 5 (See Fig. 4 for locations). The seismic close ups show the detail of the down-dip fold-thrust belts balancing part of the extensional displacement of the up-dip Corvina and Posorja raft systems.

The Quaternary to present-day emplacement of a regional transtensional stress regime in the forearc basin is partly accommodated by potential recent submarine strike-slip structures. For instance, the activity of the Puná fault-related Amistad antiform is recorded by Quaternary growth strata on the northwestern limb of the antiform (Fig. 4c; Witt et al., 2006). The recent activity of some fault segments (e. g., Puná and Banco Peru transtensional faults) is attested by fault scarps offsetting the seafloor (Fig. 4a; Witt et al., 2006).

4. Discussion: gravity tectonics in a forearc basin

The large-scale gravity-driven deformation system observed in the Tumbes-Guayaquil forearc basin is similar to that classically observed in deltaic systems along passive continental margins (e.g., Morley and Guerin, 1996; Stewart, 1999; Hooper et al., 2002; Rowan et al., 2004; Bilotti and Shaw, 2005; Ahmed et al., 2022). In the following, we discuss some of the key tectono-stratigraphic features responsible for the establishment of the gravity tectonic style through an active accretionary prism (Fig. 7).

4.1. Oblique convergence and crustal-scale transtensional deformation

The Northern Peru-Southern Ecuador accretionary prism system exhibits a complex three-dimensional structural architecture resulting from successive stages of deformation through the Mesozoic to Cenozoic period, including the Farallon-Nazca Plate subduction dynamics, crustal block accretion, ridge collision, and motion of forearc sliver (Benitez, 1995; Deniaud, 2000; Gutscher et al., 2002; Jaillard et al., 2004; Witt and Bourgois, 2010; Riel et al., 2014; Espurt et al., 2018; Hernández et al., 2020; Aizprua et al., 2022; Jaillard, 2022). Post-Oligocene clockwise rotations in the coastal areas of Ecuador and Northern Peru (Kissel and Laj, 1989; Mitouard et al., 1990; Siravo et al., 2021) are related to oblique convergence and partitioning of the deformation in the Northern Andes subduction zone (Fig. 7; Alvarado et al., 2016). This induced change and vertical rotation of the maximal stress in the forearc zone and the oblique reactivation of inherited fault and suture zones between crustal blocks (Aizprua, 2021). For instance, the development of NE-trending strike-slip fault (e.g., NE-trending Banco Peru and Puná strike-slip faults) and NW-trending normal faults could be the consequence of the NE-ward tectonic escape of the continental North Andean




Fig. 7. Three-dimensional structural model illustrating the tectono-stratigraphic features leading to the gravity-driven large-scale deformation system in the Tumbes-Guayaquil forearc basin in relation to the geodynamic setting of the Northern Peruvian-Southern Ecuadorian subduction zone. See text for further explanations.

Sliver during at least the Quaternary (Deniaud, 2000; Witt et al., 2006; Bourgois, 2013; Nocquet et al., 2014; Villegas-Lanza et al., 2016; Espurt et al., 2018). This modern crustal-scale transtensional regime in the accretionary prism (Noda and Miyakawa, 2017) might have favored the creation of accommodation space for gravity-driven deformation in the Tumbes-Guayaquil forearc basin (Fig. 7).

4.2. Basal décollement slope tilting

Gravity tectonics can be influenced by the dip of the basal décollement level (Mauduit et al., 1997; Rowan et al., 2004). For instance, Mauduit et al. (1997) demonstrated that an increase in basal slope dip increases noticeably the gravity sliding and the amount of extensional deformation. This increase in basal slope dip can be caused by a margin tilt induced by differential uplift across the margin. Indeed, the coastal domain of the Tumbes-Guayaquil forearc basin shows evidences of uplift characterized by the presence of large Plio-Pleistocene perched marine terraces sequences (called Tablazos), located at more than 300 m above sea level (Séranne, 1987; DeVries, 1988; Pedoja et al., 2006). These morpho-tectonic markers recorded the uplift of the Carpitás-Zorritos, Pallatanga and Santa Elena inner wedges during at least the Quaternary period (Pedoja et al., 2006; Regard et al., 2012; Bourgois, 2013; Espurt et al., 2018). We propose that the gravitational failure in the offshore Tumbes-Guayaquil forearc basin is partially controlled by the seaward tilting margin related to the uplift of the inner wedges. This is comparable to the continental passive margin of Namibia where the uplift event controlled the gravity sliding in the Orange basin (De Vera et al., 2010). The Quaternary uplift of the Northern Peruvian and Southern Ecuadorian active margins largely coincides with the major sedimentary and gravity tectonic activity in the Tumbes-Guayaquil depocenter (Fig. 7).

4.3. Sediment accumulation

The Tumbes-Guayaquil forearc basin is characterized by a 7 to 9 km-thick upper Neogene-Quaternary coarse-grained fluvio-deltaic to deep-marine sedimentary pile (Deniaud et al., 1999; Deniaud, 2000; Witt et al., 2006; Cobos, 2010, Figs. 4 and 5). The Quaternary sedimentary pile alone can exceed 5 km-thick in the Gulf of Guayaquil basin and records sedimentation rate with a maximum rate of up to 8600 m/Ma (Deniaud, 2000). Today, ongoing basin filling is sustained by more than twenty rivers (the Guayas river in Ecuador and the Tumbes river in Peru are the major ones) draining the Andean reliefs towards the Pacific margin (Salomons et al., 2005), associated with a sedimentary particle discharge estimated about 30 Mt/a (Milliman and Farnsworth, 2013). Thus, the massive sedimentary load could have strongly controlled gravitational raft tectonic instabilities in the Tumbes-Gulf of Guayaquil forearc basin. For instance, Mauduit et al. (1997) demonstrated that the sedimentation increases the rate of displacement and overall extension along a décollement. This implies that the shear stress parallel to the décollement is a function of the overburden of the sedimentary pile for a given angle of the basal slope. Furthermore, this massive sedimentary load would cause an increase in the shear stress within the décollement which, deforms and controls the geometry of the décollements. Further quantitative analyses of the bed-parallel slip along décollements are required to better illustrate the development of these shales décollement-related gravitational raft tectonics (e.g., Chapman and Williams, 1984; Delogkos et al., 2017; Alsop et al., 2020). In addition, restored cross-sections are needed to quantify the amounts of extensional and compressional deformations and to evaluate lateral compaction in the downdip fold-thrust systems (Butler and Paton, 2010; De Vera et al., 2010; Scarselli et al., 2016). Meanwhile, the geometry of the tectonic system is conformable with gravitational raft tectonics type described in passive continental margins (e.g., Rowan et al., 2004; De Vera et al., 2010; Scarselli et al., 2016). This indicates that massive sedimentary load strongly controlled gravitational raft tectonic

instabilities in the Tumbes-Gulf of Guayaquil forearc basin.

4.4. Overpressured shales

Shales and fluid pressures are very important in gravity tectonics (Hesthammer and Fossen, 1999; Oldenziel et al., 2002; Rowan et al., 2004). Shale can behave as a viscous-plastic solid involving brittle and ductile fractures if the deviatoric stress approaches the strength of the shale (Rowan et al., 2004; Wood, 2010; Soto et al., 2021). Interestingly, fluid pressures are observed in the Corvina-40-X-1 well, located at the southeastern border of the Tumbes depocenter (Figs. 1 and 2). They can be found below fluid retention depth (zRFD) of ~3.1–3.3 km where extensional stress exists based on the deviated borehole failure analysis (data provided by Perupetro S.A.). The sonic data acquired in the Corvina 40-X-1 well show that there is a high sonic slowdown at 3.7 km-depth, above Oligocene Heath Formation sequences. This implies a different compaction and burial history and an overpressure stress state (Fig. 2; data provided by Perupetro S.A.; Fildani et al., 2005; Lemgruber-Traby et al., 2020). The top of the overpressure zone corresponds to the top of the ductile shales of the Oligocene Heath Formation (Fig. 2) and results in a reversal of the Vp (2500 m/s to 3000 m/s; data provided by Perupetro S.A) in the ductile deep-marine shales of the Oligocene Heath/Playa Rica Formation compared to the overlying coarse-grained fluviodeltaic to deep-marine strata. According to our structural interpretations, the sedimentary cover of the Tumbes-Guayaquil forearc basin is detached below the overpressure zone in the ductile shales of the Oligocene Heath/Playa Rica Formation (Figs. 2, 4, 5 and 7; Corvina and Posorja décollements). As in many continental passive margins, the pore fluid pressures would promote the development of the Corvina and Posorja normal fault systems which are connected at depth in the décollement layer (Figs. 2, 4, 5 and 7; e.g., Mourgues et al., 2009; Ahmed et al., 2022). Moreover, in the context of a prograding deltaic system like in the study area, fluid pressures can decrease near the coast and increase in the seaward side of the basin, where the sedimentary thickness increases (Mourgues et al., 2009). This is coherent with the location of Posorja décollement that develops in the deepest depocenter of the Guayaquil forearc depocenter (Figs. 2, 4, 5 and 7). Consequently, friction at the level of the overpressured shales is reduced throughout the Tumbes-Guayaquil forearc depocenter. Thus, the sedimentary load of the Tumbes-Guayaquil forearc basin and the presence of overpressured shales promote low frictional properties at depth, which might control the trenchward displacement of the Banco Peru and Domito outer thrust wedges.

4.5. Seismicity

The study area is characterized by low interseismic coupling on the subduction interface (Chlieh et al., 2014; Nocquet et al., 2014; Villegas-Lanza et al., 2016). Few historical subduction earthquakes with intermediate magnitudes are located in the region (e.g., Beauval et al., 2013). However crustal faults are seismically active in the onshore and offshore region of the Gulf of Guayaquil basin (e.g., Puná dextral fault zone; Fig. 5), as revealed by national earthquakes catalogue (Alvarado et al., 2018). The gravitational sliding on the basal décollement levels is slow and appears aseismic, due to overpressured shales, sedimentary loading and tectonic strains. This raises the question of whether the basal décollement levels could accommodate stresses such as fault. It follows that we cannot exclude that aseismic slip on the basal décollements may be triggered by either dynamic or static stress changes (Du et al., 2003). One is dynamic stress change or transient deformation generated by the passage of seismic waves and the other is the static stress change associated with a nearby faulting process. Both possibilities are likely under the local seismo-tectonic conditions. Concerning the gravity sliding at the sea-bottom, the absence of slope (since we are on the continental shelf), of overhanging sedimentary mass (the sea-bottom is rather flat) and of an above significant water layer (only

several tens of meters to some hundreds of meters), make that potential intra-basin tsunamis are unlikely and the associated tsunami hazard is therefore almost null.

In addition, a potential tsunami that could be triggered by an earthquake on the Puná dextral strike-slip fault is also unlikely because of the dominant strike-slip component of the event and its short-length segments that imply moderate magnitudes along the fault. Instead, in the study area and nearby coastal areas, the tsunami hazard is related to a subduction earthquake occurring near the trench on the subduction interface (Ioualalen et al., 2014).

5. Conclusion

We propose a new structural interpretation for the Tumbes-Guayaquil forearc basin involving a tectonic style reported for the first time in the dynamics of the area. Our results provide a model for the evolution of a forearc basin, belonging to an accretionary prism, which exhibits widespread gravitational tectonic instabilities in the Late Neogene-Quaternary interval. Subsurface data and the construction of structural cross-sections show that the forearc basin is detached seaward from the underlying accretionary prism and deformed by large-scale, updip listric normal fault systems (Corvina and Posorja raft systems) associated with down-dip fold-thrust systems (Barracuda and part of the Domito thrust systems). The décollement layer is located within the Oligocene ductile shales of the Heath/Playa Rica Formation. We propose that this gravity tectonic style, rather observed in passive continental margins, is primarily controlled by the combination of tectonostratigraphic features including crustal-scale transtensional deformation related to oblique convergence along the Northern Andean margin and opening of the basin along the Puná fault zone, basal décollement slope tilting related to the Quaternary coastal uplift, the massive sediment accumulation, and the presence of overpressured shales.

Author statement

Andréa Peuzin - Conceptualization, Formal analysis, Writing - original draft preparation.

Marianne Saillard - Conceptualization, Formal analysis, Funding acquisition, Project administration, Writing - original draft preparation.

Nicolas Espurt - Conceptualization, Formal analysis, Funding acquisition, Supervision, Writing - original draft preparation.

François Michaud - Conceptualization, Formal analysis, Funding acquisition, Supervision, Writing - original draft preparation.

Cédric Bulois - Conceptualization, Formal analysis, original draft preparation.

Daniel Praeg - Conceptualization, Formal analysis, original draft preparation.

Marc Régnier - Formal analysis, original draft preparation.

Ysabel Calderon - Formal analysis, Funding acquisition.

Declaration of competing interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: Andrea Peuzin reports was provided by University of Côte d'Azur. Andrea Peuzin reports a relationship with University of Côte d'Azur that includes: employment.

Data availability

Data will be made available on request.

Acknowledgments

This work is part of the A. Peuzin's PhD thesis (EDSFA 284) and supported by the ANR MARACAS (ANR-18-CE31-0022; PI: M. Saillard).

The interpretation of the seismic profiles and the construction of the cross-sections were performed with the MOVE Software Suite donated by Petroleum Experts Limited. We thank C. Matonti for the interesting discussions on the reservoir and fluid overpressure. We also thank J. I. Soto and an anonymous reviewer for their constructive comments that allowed us to improve the manuscript and the figures in many points.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jsg.2023.104909>.

References

- Ahmed, B., McClay, K., Scarselli, N., Bilal, A., 2022. New insights on the gravity-driven deformation of late Albian–early Turonian stacked delta collapse systems in the Ceduna sub-basin, Bight Basin, southern margin of Australia. *Tectonophysics* 823, 229184.
- Aizprua, C., 2021. Forearc Crustal Structure and Controlling Factors on Basin Formation across the Southernmost Northern Andes. PhD thesis. Université de Lille; Norwegian University of Science and Technology, Trondheim, Norvège, p. 220 (English).
- Aizprua, C., Witt, C., Johansen, S.E., Barba, D., 2019. Cenozoic stages of forearc evolution following the accretion of a sliver from the late cretaceous-caribbean large igneous province: SW Ecuador-nw Peru. *Tectonics* 38, 1441–1465. <https://doi.org/10.1029/2018TC005235>.
- Aizprua, C., Witt, C., Barba, D., 2022. An appraisal of basin development at the trailing edge of the Northern Andes, SW Ecuador–NW Peru. *Andean Structural Styles* 237–247. <https://doi.org/10.1016/B978-0-323-85175-6.00018-3>. Elsevier.
- Alsop, G.L., Weinberger, R., Marco, S., Levi, T., 2020. Bed-parallel slip: identifying missing displacement in mass transport deposits. *J. Struct. Geol.* 131, 103952.
- Alvarado, A., Audin, L., Nocquet, J.M., Jaillard, E., Mothes, P., Jarrin, P., Segovia, M., Rolandone, F., Cisneros, D., 2016. Partitioning of oblique convergence in the Northern Andes subduction zone: migration history and the present-day boundary of the North Andean Sliver in Ecuador. *Tectonics* 35, 1048–1065. <https://doi.org/10.1002/2016TC004117>.
- Alvarado, A., Ruiz, M., Mothes, P., Yepes, H., Segovia, M., Vaca, M., Córdova, A., 2018. Seismic, volcanic, and geodetic networks in Ecuador: building capacity for monitoring and research. *Seismol. Res. Lett.* 89 (2A), 432–439.
- Auguy, C., Calvès, G., Calderon, Y., Brusset, S., 2017. Seismic evidence of gas hydrates, multiple BSRs and fluid flow offshore Tumbes Basin, Peru. *Mar. Geophys. Res.* 38, 409–423. <https://doi.org/10.1007/s11001-017-9319-2>.
- Beauval, C., Yepes, H., Palacios, P., Segovia, M., Alvarado, A., Font, Y., Aguilar, J., Troncoso, L., Vaca, S., 2013. An earthquake catalog for seismic hazard assessment in Ecuador. *Bull. Seismol. Soc. Am.* 103, 773–786. <https://doi.org/10.1785/0120120270>.
- Benitez, S., 1995. Évolution géodynamique de la province côtière sud-équatorienne au Crétacé supérieur-Tertiaire. Géologie appliquée. Université Joseph-Fourier - Grenoble I. Français, p. 156.
- Bilotti, F., Shaw, J.H., 2005. Deep-water Niger Delta fold and thrust belt modeled as a critical-taper wedge: the influence of elevated basal fluid pressure on structural styles. *AAPG Bull.* 89 (11), 1475–1491.
- Bourgeois, J., 2013. A review on tectonic record of strain buildup and stress release across the andean forearc along the Gulf of Guayaquil-tumbes basin (GGTB) near Ecuador-Peru border. *Int. J. Geosci.* 4, 618–635. <https://doi.org/10.4236/ijg.2013.43057>.
- Brusset, S., Soula, J.C., Roddaz, M., Calvès, G., Espurt, N., Vega, M., Baby, P., Calderon, Y., 2018. Reappraisal of the tectonic style of the talara-tumbes forearc basin: regional insights for hydrocarbon exploration. *Petroleum Basins and Hydrocarbon Potential of the Andes of Peru and Bolivia* 323–338. <https://doi.org/10.1306/13622126M1173772>.
- Butler, R.W.H., Paton, D.A., 2010. Evaluating lateral compaction in deepwater fold and thrust belts: how much are we missing from “nature’s sandbox”. *GSA Today (Geol. Soc. Am.)* 20 (3), 4–10.
- Calahorran, A., 2005. Structure de la marge du Golfe de Guayaquil (Équateur) et propriétés physiques du chenal de subduction à partir de données de sismique marine réflexion et réfraction. Thèse de doctorat. Université Pierre et Marie Curie, Paris, p. 227 (Français).
- Chapman, T.J., Williams, G.D., 1984. Displacement-distance methods in the analysis of fold-thrust structures and linked-fault systems. *J. Geol. Soc.* 141 (1), 121–128.
- Chlieh, M., Mothes, P.A., Nocquet, J.-M., Jarrin, P., Charvis, P., Cisneros, D., Font, Y., Collot, J.-Y., Villegas-Lanza, J.-C., Rolandone, F., Vallée, M., Regnier, M., Segovia, M., Martin, X., Yepes, H., 2014. Distribution of discrete seismic asperities and aseismic slip along the Ecuadorian megathrust. *Earth Planet. Sci. Lett.* 400, 292–301. <https://doi.org/10.1016/j.epsl.2014.05.027>.
- Chunga, K., Ochoa-Cornejo, F., Mulas, M., Toulkeridis, T., Menéndez, E., 2018. Characterization of seismogenic crustal faults in the Gulf of Guayaquil, Ecuador. *Andean Geol.* 46, 66. <https://doi.org/10.5027/andgeoV46n1-2991>.
- Cobos, L., 2010. Estudio Ontegrado del Golfo de Guayaquil del Mioceno al Reciente. Facultad de Ingeniería en Ciencias de la Tierra, Guayaquil - Ecuador.
- Collot, J.-Y., Charvis, P., Gutscher, M.-A., Operto, E., 2002. Exploring the Ecuador-Colombia active margin and inter-plate seismogenic zone. *Eos, Transactions American Geophysical Union* 83 (17), 185–190. <https://doi.org/10.1029/2002EO000120>.
- Daly, M.C., 1989. Correlations between Nazca/Farallon plate kinematics and forearc basin evolution in Ecuador. *Tectonics* 8 (4), 769–790.
- De Vera, J., Granado, P., McClay, K., 2010. Structural evolution of the Orange Basin gravity-driven system, offshore Namibia. *Mar. Petrol. Geol.* 27 (1), 223–237.
- Delogkos, E., Childs, C., Manzocchi, T., Walsh, J.J., Pavlides, S., 2017. The role of bed-parallel slip in the development of complex normal fault zones. *J. Struct. Geol.* 97, 199–211.
- Deniaud, Y., 2000. Enregistrements sédimentaire et structural de l'évolution géodynamique des Andes équatoriennes au cours du Néogène : étude des bassins d'avant-arc et bilans de masse. Thèse de doctorat en Tectonique. Université Joseph Fourier, Grenoble, France, p. 153.
- Deniaud, Y., Baby, P., Basile, C., Ordoñez, M., Montenegro, G., Mascle, G., 1999. Ouverture et évolution tectono-sédimentaire du Golfe de Guayaquil: bassin d'avant-arc néogène et quaternaire du Sud des Andes équatoriennes. *C. R. Acad. Sci., Paris* 328, 181–187.
- DeVries, T.J., 1988. The geology of late Cenozoic marine terraces (tablazos) in northwestern Peru. *J. S. Am. Earth Sci.* 1, 121–136. [https://doi.org/10.1016/0895-9811\(88\)90030-2](https://doi.org/10.1016/0895-9811(88)90030-2).
- Dickinson, W.R., 1995. Forearc basins. In: Busby, C.J., Ingersoll, R.V. (Eds.), *Tectonics of Sedimentary Basins*. Blackwell Science, Oxford, UK, pp. 221–261.
- Du, W.-X., Sykes, L.R., Shaw, B.E., Scholz, C.H., 2003. Triggered aseismic fault slip from nearby earthquakes, static or dynamic effect? *J. Geophys. Res.* 108 (B2), 2131. <https://doi.org/10.1029/2002JB002008>.
- Dula Jr., W.F., 1991. Geometric models of listric normal faults and rollover folds. *AAPG Bull.* 75 (10), 1609–1625.
- Dumont, J.F., Santana, E., Vilema, W., 2005. Morphologic evidence of active motion of the zambapala fault, Gulf of Guayaquil (Ecuador). *Geomorphology* 65 (3–4), 223–239. <https://doi.org/10.1016/j.geomorph.2004.09.003>. ISSN 0169-555X.
- Espinoza, J., 1992. Terremotos tsunamigénicos en el Ecuador. *Acta Oceanográfica del Pacífico. INOCAR* 25, 66–82. *Geologica del Peru* 32 (2), 225–238. <http://hdl.handle.net/1834/2181>.
- Espurt, N., Brusset, S., Baby, P., Henry, P., Vega, M., Calderon, Y., Saillard, M., 2018. Deciphering the Late Cretaceous-Cenozoic structural evolution of the North Peruvian forearc system. *Tectonics* 37, 251–282. <https://doi.org/10.1002/2017TC004536>.
- Fernández, J., Martínez, E., Calderón, Y., Hermoza, W., Galdos, C., 2005. Tumbes and Talara Basins Hydrocarbon Evaluation. *Perupetro S.A. Basin Evaluations Group Exploration Department*, p. 130 internal report. Retrieved from www.perupetro.com.pe.
- Fildani, A., Hanson, A.D., Chen, Z., Moldovan, J.M., Graham, S.A., Arriola, P.R., 2005. Geochemical characteristics of oil and source rocks and implications for petroleum systems, Talara basin, northwest Peru. *AAPG (Am. Assoc. Pet. Geol.) Bull.* 89, 1519–1545. <https://doi.org/10.1306/06300504094>.
- Fildani, A., Hessler, A.M., Graham, S.A., 2008. Trench-forearc interactions reflected in the sedimentary fill of Talara basin, northwest Peru. *Basin Res.* 20 (3), 305–331.
- Gutscher, M.-A., Malod, J., Rehault, J.-P., Contrucci, I., Klingelhoefer, F., Mendes-Victor, L., Spakman, W., 2002. Evidence for active subduction beneath Gibraltar. *Geology* 30, 1071. [https://doi.org/10.1130/0091-7613\(2002\)030<1071:EFASBG>2.0.CO;2](https://doi.org/10.1130/0091-7613(2002)030<1071:EFASBG>2.0.CO;2).
- Guzmán, M.A., Márquez, G., Boente, C., Witt, C., Morato, A., Tocco, R., 2022. Geochemical evaluation of sedimentary rocks and hydrocarbon fluids in the Amistad offshore field, Progreso Basin Province, Southwest Ecuador. *J. Petrol. Sci. Eng.* 213, 110410. <https://doi.org/10.1016/j.petrol.2022.110410>.
- Hernández, M.J., Michaud, F., Collot, J.-Y., Proust, J.-N., d'Acremont, E., 2020. Evolution of the Ecuador offshore nonaccretionary-type forearc basin and margin segmentation. *Tectonophysics* 781, 228374. <https://doi.org/10.1016/j.tecto.2020.228374>.
- Hesthammer, J., Fossen, H., 1999. Evolution and geometries of gravitational collapse structures with examples from the Statfjord Field, northern North Sea. *Mar. Petrol. Geol.* 16, 259–281. [https://doi.org/10.1016/S0264-8172\(98\)00071-3](https://doi.org/10.1016/S0264-8172(98)00071-3).
- Heuret, A., Conrad, C.P., Funicello, F., Lallemand, S., Sandri, L., 2012. Relation between subduction megathrust earthquakes, trench sediment thickness and upper plate strain. *Geophys. Res. Lett.* 39. <https://doi.org/10.1029/2011GL050712>.
- Higley, D.K., 2004a. The progreso basin province of northwestern Peru and southwestern Ecuador: Neogene and cretaceous-paleogene. *Total Petroleum Systems* 31.
- Higley, D.K., 2004b. The Talara Basin Province of Northwestern Peru, Cretaceous-Tertiary Total Petroleum System. US Department of the Interior, US Geological Survey.
- Hooper, R.J., Fitzsimmons, R.J., Grant, N., Vendeville, B.C., 2002. The role of deformation in controlling depositional patterns in the south-central Niger Delta, West Africa. *J. Struct. Geol.* 24 (4), 847–859.
- Ioualalen, M., Larroque, C., Scotti, O., 2014. Tsunami mapping related to local earthquakes on the French-Italian riviera (western mediterranean). *Pure Appl. Geophys.* 171, 1423–1443. <https://doi.org/10.1007/s00024-013-0699-1>.
- Jaillard, E., 2022. Late cretaceous-paleogene orogenic build-up of the Ecuadorian Andes: review and discussion. *Earth Sci. Rev.* 230, 104033. <https://doi.org/10.1016/j.earscirev.2022.104033>.
- Jaillard, E., Ordoñez, M., Benitez, S., Berrones, G., Jiménez, N., Montenegro, G., Zambrano, I., 1995. Basin Development in an Accretionary, Oceanic-Floored Fore-Arc Setting: Southern Coastal Ecuador during Late Cretaceous-Late Eocene Time.
- Jaillard, E., Ordoñez, M., Suárez, J., Toro, J., Iza, D., Lugo, W., 2004. Stratigraphy of the late Cretaceous-Paleogene deposits of the cordillera Occidental of central Ecuador: geodynamic implications. *J. S. Am. Earth Sci.* 17, 49–58. <https://doi.org/10.1016/j.jsames.2004.05.003>.
- Kissel, C., Laj, C. (Eds.), 1989. *Paleomagnetic Rotations and Continental Deformation*. Springer Netherlands, Dordrecht. <https://doi.org/10.1007/978-94-009-0869-7>.

- Lajo-Yáñez, J.A., Flint, S.S., Brunt, R.L., Huuse, M., Searle, S.R.A., Sheppard, J.M., 2022. Disentangling tectonic and eustatic controls on forearc basin stratigraphy, Talara Basin, Peru. *Sediment. Geol.* 442, 106277 <https://doi.org/10.1016/j.sedgeo.2022.106277>.
- Lemgruber-Traby, A., Espurt, N., Souque, C., Henry, P., Calderon, Y., Baby, P., Brusset, S., 2020. Thermal structure and source rock maturity of the North Peruvian forearc system: insights from a subduction-sedimentation integrated petroleum system modeling. *Mar. Petrol. Geol.* 122, 104664 <https://doi.org/10.1016/j.marpetgeo.2020.104664>.
- Lockridge, P.A., 1984. The criteria for Identification of tsunami events likely to produce far-field damage, A data application. In: Brennan, A.M., Lander, J.F. (Eds.), *Second UJNR Tsunami Workshop*, pp. 43–50. Honolulu, Hawaii, November 1990.
- Márquez, G., Morato, A., Guzmán, M.A., Boente, C., Witt, C., 2022. Geochemical characterization of crude oils and potential source rocks from southern coastal Ecuador: hypothetical petroleum systems. *Mar. Petrol. Geol.* 138, 105545 <https://doi.org/10.1016/j.marpetgeo.2022.105545>.
- Mauduit, T., Guerin, G., Brun, J.-P., Lecanu, H., 1997. Raft tectonics: the effects of basal slope angle and sedimentation rate on progressive extension. *J. Struct. Geol.* 19, 1219–1230. [https://doi.org/10.1016/S0191-8141\(97\)00037-0](https://doi.org/10.1016/S0191-8141(97)00037-0).
- Michaud, F., Witt, C., Royer, J.-Y., 2009. Influence of the subduction of the Carnegie volcanic ridge on Ecuadorian geology: reality and fiction. *Backbone of the Americas: Shallow Subduction, Plateau Uplift, and Ridge and Terrane Collision*. [https://doi.org/10.1130/2009.1204\(10\)](https://doi.org/10.1130/2009.1204(10)). Geological Society of America.
- Michaud, F., Collot, J.Y., Ratzov, G., Proust, J.N., Dano, A., Lebrun, J.F., De Min, L., 2018. A honeycomb seafloor morphology in carbonate sediment of the Carnegie Ridge (offshore Ecuador): formation and potential geodynamic significance. *Geology* 46 (11), 979–982.
- Milliman, J.D., Farnsworth, K.L., 2013. *River Discharge to the Coastal Ocean: a Global Synthesis*. Cambridge University Press.
- Mitouard, P., Kissel, C., Laj, C., 1990. Post-Oligocene rotations in southern Ecuador and northern Peru and the formation of the Huancabamba deflection in the Andean Cordillera. *Earth Planet Sci. Lett.* 98, 329–339. [https://doi.org/10.1016/0012-821X\(90\)90035-V](https://doi.org/10.1016/0012-821X(90)90035-V).
- Morley, C.K., Guerin, G., 1996. Comparison of gravity-driven deformation styles and behavior associated with mobile shales and salt. *Tectonics* 15 (6), 1154–1170.
- Mourgues, R., Lecomte, E., Vendeville, B., Raillard, S., 2009. An experimental investigation of gravity-driven shale tectonics in progradational delta. *Tectonophysics* 474, 643–656. <https://doi.org/10.1016/j.tecto.2009.05.003>.
- Nocquet, J.-M., Villegas-Lanza, J.C., Chlieh, M., Mothes, P.A., Rolandone, F., Jarrin, P., Cisneros, D., Alvarado, A., Audin, L., Bondoux, F., Martin, X., Font, Y., Rognier, M., Vallée, M., Tran, T., 2014. Motion of continental slivers and creeping subduction in the northern Andes. *Nat. Geosci.* 7 (4), 287–291.
- Noda, A., 2016. Forearc basins: types, geometries, and relationships to subduction zone dynamics. *Geol. Soc. Am. Bull.* 128, 879–895. <https://doi.org/10.1130/B31345.1>.
- Noda, A., Miyakawa, A., 2017. *Deposition and Deformation of Modern Accretionary-type Forearc Basins: Linking Basin Formation and Accretionary Wedge Growth*, vol. 26.
- Oldenziel, T., van Dithuijzen, R., van Kruijsdijk, C., 2002. Geologic parameterization of reservoir model enhances history-match procedure. *Lead. Edge* 21, 544–551. <https://doi.org/10.1190/1.1490648>.
- Pedoja, K., Ortlieb, L., Dumont, J.F., Lamothe, M., Ghaleb, B., Auclair, M., Labrousse, B., 2006. Quaternary coastal uplift along the Talara Arc (Ecuador, Northern Peru) from new marine terrace data. *Mar. Geol.* 228, 73–91. <https://doi.org/10.1016/j.margeo.2006.01.004>.
- Regard, V., Pedoja, K., Saillard, M., 2012. *Les terrasses marines, marqueurs de l'interaction entre soulèvement et oscillation du niveau de la mer*, vol. 4.
- Reynaud, J.Y., Witt, C., Pazmiño, A., Gilces, S., 2018. Tide-dominated deltas in active margin basins: insights from the Guayas estuary, Gulf of Guayaquil, Ecuador. *Mar. Geol.* 403, 165–178.
- Riel, N., Martelat, J.-E., Guillot, S., Jaillard, E., Monié, P., Yuquilema, J., Duclaux, G., Mercier, J., 2014. Fore arc tectonothermal evolution of the El Oro metamorphic province (Ecuador) during the Mesozoic. *Tectonics* 33, 1989–2012. <https://doi.org/10.1002/2014TC003618>.
- Rowan, M.G., Peel, F.J., Vendeville, B.C., 2004. Gravity-driven foldbelts on passive margins. *AAPG (Am. Assoc. Pet. Geol.) Bull.* 157–182.
- Salomons, W., Kremer, H.H., Turner, R.K., Andreeva, E.N., Arthurton, R.S., Behrendt, H., Burbridge, P., Chen, C.-T.A., Crossland, C.J., Gandrass, J., Gordeev, V.V., Harvey, N., Hong, G.H., Kjerfve, B., de Lacerda, L.D., Marshall Crossland, J.I., Morcom, N., Odada, E., Pacyna, J., Rabalais, N.N., Swaney, D., Wiebe, W.J., 2005. The catchment to coast continuum. In: Crossland, C.J., Kremer, H.H., Lindeboom, H.J., Marshall Crossland, J.I., Le Tissier, M.D.A. (Eds.), *Coastal Fluxes in the Anthropocene, Global Change — the IGBP Series*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 145–200. https://doi.org/10.1007/3-540-27851-6_4.
- Scarselli, N., McClay, K., Elders, C., 2016. Seismic geomorphology of Cretaceous megaslices offshore Namibia (Orange Basin): insights into segmentation and degradation of gravity-driven linked systems. *Mar. Petrol. Geol.* 75, 151–180.
- Séranne, M., 1987. *Évolution tectono-sédimentaire du bassin de Talara (nord-ouest du Pérou)*.
- Shepherd, G.L., Moberly, R., 1981. *Coastal Structure of the Continental Margin, Northwest Peru and Southwest Ecuador*.
- Siravo, G., Speranza, F., Mulas, M., Costanzo-Alvarez, V., 2021. Significance of northern Andes terrane extrusion and genesis of the interandean valley: paleomagnetic evidence from the “Ecuadorian orocline.”. *Tectonics* 40. <https://doi.org/10.1029/2020TC006684>.
- Soto, J.I., Heidari, M., Hudec, M.R., 2021. Proposal for a mechanical model of mobile shales. *Sci. Rep.* 11, 23785 <https://doi.org/10.1038/s41598-021-02868-x>.
- Stewart, S.A., 1999. Geometry of thin-skinned tectonic systems in relation to detachment layer thickness in sedimentary basins. *Tectonics* 18 (4), 719–732.
- Tsuji, T., Ashi, M., Strasser, M., Kimura, G., 2015. Identification of the static backstop and its influence on the evolution of the accretionary prism in the Nankai Trough. *Earth Planet Sci. Lett.* 431, 15–25. <https://doi.org/10.1016/j.epsl.2015.09.011>. ISSN 0012-821X.
- Vaca, S., Vallée, M., Nocquet, J.-M., Alvarado, A., 2019. Active deformation in Ecuador enlightened by a new waveform-based catalog of earthquake focal mechanisms. *J. S. Am. Earth Sci.* 93, 449–461. <https://doi.org/10.1016/j.jsames.2019.05.017>.
- Vannucchi, P., Morgan, J.-P., Balestrieri, M.-L., 2016. Subduction erosion, and the deconstruction of continental crust: the Central America case and its global implications. *Gondwana Res.* 40, 184–198. <https://doi.org/10.1016/j.gr.2016.10.001>. ISSN 1342-937X.
- Vega, M., 2009. *Architecture tectonique et stratigraphique du bassin d'avant-arc de Tumbes, (Nord Pérou) : Implications pour l'exploration des hydrocarbures*. Ph.d. thesis. University of Toulouse III-Paul Sabatier, p. 191 (Français).
- Villegas-Lanza, J.C., Chlieh, M., Cavalié, O., Tavera, H., Baby, P., Chire-Chira, J., Nocquet, J.-M., 2016. Active tectonics of Peru: heterogeneous interseismic coupling along the Nazca megathrust, rigid motion of the Peruvian Sliver, and Subandean shortening accommodation: active Tectonics of Peru. *J. Geophys. Res. Solid Earth* 121, 7371–7394. <https://doi.org/10.1002/2016JB013080>.
- Witt, C., Bourgeois, J., 2010. Forearc basin formation in the tectonic wake of a collision-driven, coastwise migrating crustal block: the example of the North Andean block and the extensional Gulf of Guayaquil-Tumbes Basin (Ecuador-Peru border area). *Geol. Soc. Am. Bull.* 122, 89–108. <https://doi.org/10.1130/B26386.1>.
- Witt, C., Bourgeois, J., Michaud, F., Ordoñez, M., Jiménez, N., Sosson, M., 2006. Development of the Gulf of Guayaquil (Ecuador) during the quaternary as an effect of the North andean block tectonic escape. *Tectonics* 25. <https://doi.org/10.1029/2004TC001723> n/a-n/a.
- Wood, L.J., 2010. *Shale Tectonics: A Preface*, vol. 93. American Association of Petroleum Geologists Memoir. <https://doi.org/10.1306/13231305M93730> (Chapter 1), 1–4.
- Xiao, H., Suppe, J., 1992. Origin of rollover (1). *AAPG Bull.* 76 (4), 509–529.
- Yepes, H., Audin, L., Alvarado, A., Beauval, C., Aguilar, J., Font, Y., Cotton, F., 2016. A new view for the geodynamics of Ecuador: implication in seismogenic source definition and seismic hazard assessment. *Tectonics* 35, 1249–1279. <https://doi.org/10.1002/2015TC003941>.