



**HAL**  
open science

# Zooarchaeological investigation of the Hoabinhian exploitation of reptiles and amphibians in Thailand and Cambodia with a focus on the Yellow-Headed tortoise (*Indotestudo elongata* (Blyth, 1854))

Corentin Bochaton, Sirikanya Chantasri, Melada Maneechote, Julien Claude, Christophe Griggo, Wilailuck Naksri, Hubert Forestier, Heng Sophady, Prasit Auertrakulvit, Jutinach Bowonsachoti, et al.

## ► To cite this version:

Corentin Bochaton, Sirikanya Chantasri, Melada Maneechote, Julien Claude, Christophe Griggo, et al.. Zooarchaeological investigation of the Hoabinhian exploitation of reptiles and amphibians in Thailand and Cambodia with a focus on the Yellow-Headed tortoise (*Indotestudo elongata* (Blyth, 1854)). BioRxiv, 2023, 10.1101/2023.04.27.538552 . hal-04145045

**HAL Id: hal-04145045**

**<https://hal.science/hal-04145045v1>**

Submitted on 28 Jun 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - ShareAlike 4.0 International License

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28

# Zooarchaeological investigation of the Hoabinhian exploitation of reptiles and amphibians in Thailand and Cambodia with a focus on the Yellow-Headed tortoise (*Indotestudo elongata* (Blyth, 1854))

Corentin Bochaton<sup>1,2\*</sup>, Sirikanya Chantasri<sup>3</sup>, Melada Maneechote<sup>4</sup>, Julien Claude<sup>2,5</sup>, Christophe Griggo<sup>6</sup>, Wilailuck Naksri<sup>7</sup>, Hubert Forestier<sup>8</sup>, Heng Sophady<sup>9</sup>, Prasit Auertrakulvit<sup>4</sup>, Jutinach Bowonsachoti<sup>3</sup>, and Valéry Zeitoun<sup>1</sup>

<sup>1</sup>UMR 7207. CR2P-CNRS-MNHN-SU, Sorbonne Université, campus Jussieu, T. 46-56, 5ème étage, case 104, 4, place Jussieu, 75 252 Paris Cedex 05, France.

<sup>2</sup>Institut des sciences de l'évolution de Montpellier, UMR 5554 CNRS/UM/IRD/EPHE, Université de Montpellier, Montpellier, France.

<sup>3</sup>12th Regional Office of Fine Art Department, Nakhon Si Thammarat, Thailand.

<sup>4</sup>Department of Archaeology, Faculty of Archaeology, Silpakorn University, Bangkok, Thailand.

<sup>5</sup>Faculty of Science, University of Chulalongkorn, Bangkok, Thailand.

<sup>6</sup>Laboratoire Environnements, DYnamiques et Territoires de Montagne, UMR 5204 – Université Grenoble Alpes, Le Bourget-du-Lac, France.

<sup>7</sup>Northeastern Research Institute of Petrified Wood and Mineral Resources, Nakhon Ratchasima Rajabhat University, Nakhon Ratchasima, Thailand.

<sup>8</sup>Muséum national d'Histoire naturelle, Musée de l'Homme, UMR 7194 HNHP, Paris, France.

<sup>9</sup>Ministry of Culture and Fine Arts of Cambodia, Phnom Penh, Cambodia.

\*Corresponding author: [corentin.bochaton@cnrs.fr](mailto:corentin.bochaton@cnrs.fr)

## ABSTRACT

Although non-marine turtles are nearly ubiquitous in the Southeast Asian archaeological record, their zooarchaeological study has been very poorly undertaken in that tropical region of the world. This lack of study makes the understanding of past human subsistence strategies very complex especially regarding the prehistoric hunter gatherer populations which may have massively exploited inland chelonian taxa. In order to try to start a new dynamic regarding the study of the past human-turtle interactions in Southeast Asia we propose here an in-depth zooarchaeological analysis of the turtle bone remains recovered

37 from four Hoabinhian Hunter-gatherer archaeological assemblages located in Thailand and  
38 Cambodia, and dated from the Late Pleistocene to the first half of the Holocene. Our study  
39 is focused on the bone remains attributed to the Yellow-Headed Tortoise (*Indotestudo*  
40 *elongata*) as they account for the majority of the turtle archaeological assemblages  
41 identified in the target area. For this species, we developed osteo-metric equations enabling  
42 the estimation of the carapace size of the archaeological individuals of this species. This  
43 allowed us to study the size structure of the archaeological populations in the different sites  
44 and to reveal the human exploitation strategies of these animals. We found a strong  
45 taphonomic homogeneity between the studied assemblages suggesting similarities of the  
46 subsistence behaviors in the different sites despite their very different environmental  
47 settings. We thus hypothesize putative cultural similarities across time and space. In  
48 addition, we also provide a baseline for future zooarchaeological studies as well as a  
49 methodological frame for the detailed studies of archaeological turtle bones in continental  
50 Southeast Asia.

51

52

53 **Keywords:** Reptiles, Hunter-gatherer, Southeast Asia, Turtle, Zooarchaeology

54

55

## Introduction

56           The Hoabinhian has been a major topic in prehistoric research in Mainland Southeast Asia for  
57 nearly 90 years. Since its first definition by the French archaeologist Madeleine Colani in the early 1930s  
58 (Collectif 1932), the Hoabinhian has no doubt been one of the most debated topics of the field. A wide  
59 range of subjects pertaining to the Hoabinhian populations have been discussed, such as their  
60 distribution in space and time, their definition, the technological and functional characteristics of their  
61 lithic industries, their economic organization, and their environmental context (Forestier *et al.* 2021). The  
62 chronology of the Hoabinhian has been largely extended since the earliest known Hoabinhian sites.  
63 Indeed, the occupation of the Xiaodong Rock shelter in Southwest China has been dated to about 43, 000  
64 BP (Ji *et al.* 2016) and is considered to be the earliest occurrence of the Hoabinhian culture while the site  
65 of Huai Hin in Northwest Thailand, dated to about 3700 BP is considered to be the last occurrence with a  
66 lithic production associated with ceramic sherds (Zeitoun *et al.* 2008; Forestier *et al.* 2013). More than a  
67 hundred Hoabinhian sites have been reported in Southeast Asia (Moser 2001; Chung 2008; Zeitoun *et al.*  
68 2008; White 2011; Forestier *et al.* 2017) leading inevitably to the question of the variability of Hoabinhian  
69 lithic assemblages with unexpected operational sequences on pebble matrix (Forestier *et al.* 2022, 2023).  
70 Despite this, the Hoabinhian people remain, however, quite poorly understood from cultural and  
71 material point of views. In fact, the homogeneity and lack of diversity of their lithic material culture,  
72 probably related to their putative heavy use of objects made of perishable vegetal material (Forestier  
73 2003) does not allow to characterize the precise uses of the sites by past populations, and it is difficult to  
74 expect a cultural stasis over 30,000 years among different hunter-gatherer groups in such an extensive  
75 region presenting diverse environmental, ecological and geographic settings.

76           White (2011) proposed that the emergence of cultural diversity in mainland Southeast Asia  
77 began precisely in the late Late Pleistocene which is also suggested by burial practices (Imdirakphol *et al.*  
78 2017). Forestier *et al.* (2013) argued that more analyses were still needed to evaluate the whole corpus of  
79 Southeast Asian lithic industries to describe putative “cultural variations” and since then some patterns  
80 started to emerge (Forestier *et al.* 2017, 2021). However, while a technological definition of the  
81 Hoabinhian culture is in progress, the lack of detailed zooarchaeological analyses makes our knowledge  
82 of the Hoabinhian paleoecology and subsistence strategies variability blurry. This is also an issue for the  
83 characterization of the archaeological deposits as faunal data are of major importance to characterize the  
84 use and occupation periods of the archaeological sites. The economic aspects of the Hoabinhian culture  
85 were previously addressed by several authors (Gorman 1969, 1970, 1971; Glover 1977; Yen 1977; Vu  
86 1994). Yet, the pioneering prehistoric zooarchaeological studies relying on occurrence data (Gorman  
87 1971) offer only a poor understanding of the choices made by the hunters by focusing on qualitative data  
88 and the diversity of the exploited animals rather than on quantitative information reflecting the intensity  
89 of the exploitation of each species. These first works also lack the detailed taphonomic and taxonomic  
90 analyses needed to describe past human behaviors and bone accumulation processes in more depth.  
91 Because of these lacks, we currently have only a vague idea of the spatial and chronological variability of  
92 the subsistence strategies of the Hoabinhian people. These problems impacting southeast Asian  
93 zooarchaeology have previously been reviewed in depth by Conrad (2015). Among the identified issues is  
94 the lack of detailed study of each animal group but especially of the non-mammal taxa including reptiles  
95 and non-marine turtles that are often accounting for a large part of the animal bone assemblages found  
96 in the archaeological record.

97           This limit is not proper to continental Southeast Asia but is more impactful here than in many  
98 other areas. Zooarchaeological studies fully focused on non-marine turtles have been conducted in non-  
99 tropical areas as Europe (Blasco 2008; Nabais & Zilhão 2019; Nabais *et al.* 2019), Near East (Speth &  
100 Tchernov 2002; Blasco *et al.* 2016; Biton *et al.* 2017), South Africa (Avery *et al.* 2004; Thompson &

101 Henshilwood 2014), and Northern America (Rhodin 1992). Such studies remained, however, limited in  
102 tropical areas including in South-East Asia despite the fact that turtle bones are way more common in  
103 tropical settings than in the temperate regions where there is a strong tradition of prehistoric studies. In  
104 Southeast Asia, this problem is partly rooted in the facts that there is a general lack of detailed  
105 anatomical data allowing for the identification of the taxa on the basis of isolated plate remains, and that  
106 appropriate methodological framework were never developed to analyze this material. Several works  
107 have been conducted regarding the osteology of Southeast Asian turtles most of which focusing on the  
108 Geoemydidae family in order to address questions related to the phylogeny and the paleo-biodiversity of  
109 the group (Naksri 2007, 2013; Naksri *et al.* 2013; Garbin *et al.* 2018). However, only few works have been  
110 interested in the study of the isolated elements found in the archaeological record (Pritchard *et al.* 2009;  
111 Claude *et al.* 2019). Despite these limitations several zooarchaeological studies of Hoabinhian  
112 archaeological deposits in mainland Southeast Asia have started to characterize the exploitation of the  
113 non-marine turtles by these prehistoric populations. In the few existing studies, turtle remains are often  
114 left unidentified: Ban Rai Rockshelter (Treerayapiwat 2005); Banyan Valley Cave (Higham 1977); Gua  
115 Gunung Runtuh (Zuraina 1994); Gua Harimau (Bulbeck 2003); Gue Kechil (Dunn 1964; Medway 1969);  
116 Gua Ngaum (Bulbeck 2003); Gua Peraling (Adi 2000); Gua Teluk Kelawar (Bulbeck 2003); Tham Lod  
117 Rockshelter (Amphansri 2011); Lang Kamnan Cave (Shoocongdej 1996); Khao Toh Chong Rockshelter (Van  
118 Vlack 2014); Moh Khiew II Rockshelter (Auetrakulvit 2004); Spirit Cave (Higham 1977); Tham Phaa Can  
119 (Higham 2002); Thung Nong Nien Rockshelter (Auetrakulvit 2004). In some studies they are identified but  
120 not quantified by species for instance in the Lang Rongrien Rock Shelter assemblage (Anderson 1990;  
121 Mudar & Anderson 2007). In the rare studies in which turtle bones are identified and quantified, the most  
122 abundant species is often by far the Yellow-Headed tortoise (*Indotestudo elongata* (Blyth, 1854)) -Doi Pha  
123 Kan Rockshelter (Frère *et al.* 2018); Laang Spean Cave (Frère *et al.* 2018); Spirit Cave (Conrad *et al.* 2016)-  
124 at least in its modern distribution area as Geoemydidae turtles are the best represented in the Malaysian  
125 sites Gua Sagu (Rabett 2012), and Gua Tenggek (Rabett 2012). There is also a site in Vietnam (Hiem Cave)  
126 where the tortoise *Manouria* is the most abundant turtle taxa but the small size of the faunal assemblage  
127 does not allow drawing conclusions from this observation (Masojc *et al.* 2023). However, even in the  
128 quantified above mentioned studies, the study of the turtle bone remains is still superficial. For instance,  
129 a detailed analysis of the taphonomy of the bone assemblages is not conducted and the population of  
130 turtle exploited is not characterized further than from the aspect of its species composition.

131 To start proposing solutions to these issues and also to provide the first detailed data regarding  
132 the prehistoric exploitation of Southeast Asian turtles we conducted an in-depth zooarchaeological  
133 analysis of the turtle bone remains recovered from four hunter-gatherer archaeological assemblages.  
134 These sites are located in Thailand and Cambodia and dated from the Late Pleistocene to the first half of  
135 the Holocene. They are the Doi Pha Kan rockshelter, the Moh Khiew Cave, and the Khao Tha Phlai Cave  
136 located in Thailand, and the Laang Spean Cave located in Cambodia. In addition, in order to be able to  
137 characterize with more precision the exploitation strategies of the non-marine turtles by the  
138 archaeological human populations, we developed osteo-metric equations enabling to estimate the  
139 carapace size of the archaeological individuals of *Indotestudo elongata*. We choose to focus our  
140 methodological approach on this species as most of the rich assemblages of turtle bones collected in the  
141 four considered sites correspond to this species (Frère *et al.* 2018; C. B., J. C. preliminary observations).  
142 This analytical tool allows for the study of the size structures of the archaeological populations in the  
143 different sites, and the characterization the choices made by the hunters. Altogether these data provide  
144 the first characterization of the exploitation of non-marine turtle by Pleistocene and Holocene hunter-  
145 gatherer populations of continental Southeast Asia.

146

147

## Material and Methods

## 148 **Main characteristics of *Indotestudo elongata*, the Yellow-Headed tortoise**

149 The genus *Indotestudo* currently includes three species: the Forsten's Tortoise *Indotestudo*  
150 *forstenii* (Schlegel & Müller, 1845), the Travancore Tortoise *Indotestudo travancorica* (Boulenger, 1907),  
151 and the Yellow-Headed tortoise *Indotestudo elongata* (Blyth, 1854), the two later being sister taxa  
152 (Iverson *et al.* 2001). These species are currently distributed in India (*I. travancorica*), Sulawesi (*I.*  
153 *forstenii*), and northern India and continental Southeast Asia (*I. elongata*) (Rhodin *et al.* 2021). This last  
154 species is the only *Indotestudo* species present in continental Southeast Asia. It is nowadays present in  
155 most areas of Thailand, Cambodia, Vietnam, and Laos (Ihlow *et al.* 2016; Rhodin *et al.* 2021). It is also  
156 occurring in northwestern Malaysia but is absent from most of this country and from the insular Sunda  
157 (Ihlow *et al.*, 2016). *I. elongata* is a medium size tortoise whose adult Straight Carapace Length (SCL)  
158 reaches about 300 mm (Taylor 1970). The largest specimen ever recorded is a male of about 380 mm SCL  
159 (Rhodin *et al.* 2021). It is possible that the sexual dimorphism in this species vary from one population to  
160 another although males are generally larger than females (Ihlow *et al.* 2016). The size of the hatchlings  
161 ranges from 50 to 55 mm SCL (Ihlow & Handschuh 2011). The specie is present in a wide variety of  
162 environments including many forest types (Ihlow *et al.*, 2016). *I. elongata* is active during daytime mostly  
163 in the early morning and late afternoon. It present seasonal activity patterns in order to avoid the highest  
164 temperatures during the dry season during which it aestivate spending most of its time hiding in former  
165 burrows of other animal species including porcupines (Van Dijk 1998; Som & Cottet 2016; Ihlow *et al.*  
166 2016). In their active period the individuals spend most of their time in open areas during the rainy  
167 season then move in more closed environment (mostly semi-evergreen and pine forests) when the  
168 climate becomes dryer (Van Dijk 1998; Som & Cottet 2016). Depending of their sex, the individuals reach  
169 sexual maturity between 175 mm and 240 mm SCL at an age of 6-8 years old (Van Dijk 1998; Eberling  
170 2011; Sriprateep *et al.* 2013). Reproduction takes place during the rainy season.

## 171 **Presentation of the studied sites and assemblages**

### 172 *The Doi Pha Kan rockshelter*

173 The site of Doi Pha Kan is a rockshelter located in northern Thailand (E 99° 46' 37.2" ; N 18° 26'  
174 57.0"). The site is studied since 2011 by P. Auetrakulvit and V. Zeitoun and its excavation is still in  
175 progress. The site is mostly known for its three Hoabinhian sepultures dated between 11,170 ± 40 and  
176 12,930 ± 50 BP (Imdirakphol *et al.* 2017; Zeitoun *et al.* 2019). Two of these sepultures contained turtle  
177 shells elements in anatomical connection interpreted as funeral offerings. The site also provided a rich  
178 archaeological assemblage corresponding to a Hoabinhian occupation older than the sepultures. The  
179 stratigraphy of the site being homogeneous from a sedimentary point of view, its archaeological material  
180 has been so far considered as a single assemblage.

181 Samples of the lithic material and of the animal bone assemblages collected on the site have  
182 already been the object of studies (Celiberti *et al.* 2018; Frère *et al.* 2018) but most of the material is still  
183 under study. In this paper we will present zooarchaeological data collected on the herpetofaunal taxa  
184 bone remains recovered on the site until 2019. To collect this dataset the whole archaeofaunal material  
185 has been observed to extract and then study the reptile and amphibian bone remains. The material  
186 screened this way corresponds to the material previously studied by S. Frère (Frère *et al.* 2018) with the  
187 addition of the material collected following this first study. In the first study, S. Frère analyzed 4256  
188 animal remains among which 2541 were attributed to vertebrate species. So far, no complete study of  
189 this faunal assemblage has been completed and no Minimal Number of Individual (MNI) data have ever  
190 been published. In these conditions, the overall weight of the herpetofauna we collected in respect to the  
191 full sample cannot be assessed with precision. However, the existing data indicates it could account for a  
192 very significant part of the full assemblage as it corresponds to 17.1% of the total bone weight and 51% of



193 the vertebrate total Number of Identified Skeletal Parts (NISP) analyzed in the previous study (Frère *et*  
194 *al.* 2018).

#### 195 *The Moh Khiew Cave*

196 The site of Moh Khiew Cave is a 30m long archaeological rock-shelter located in southern  
197 Thailand in the Krabi Province (E 98° 55' 49.27''; N 08° 09' 36.32''). It was first excavated by S. Pookajorn  
198 between 1991 and 1994 (Pookajorn 2001) before being the object of a later excavation in 2008 by P.  
199 Auetrakulvit in order to clarify its stratigraphy (Auetrakulvit *et al.* 2012). The stratigraphy of the site is  
200 composed of several archaeological layers dated from the Holocene to the Late Pleistocene, most of  
201 which corresponding to Hoabinhian occupations. Five sepultures were also discovered, one during the  
202 first excavations, and four in 2008.

203 Regarding the zooarchaeological data, the complete assemblages of the first excavations was  
204 studied by P. Auetrakulvit (Auetrakulvit 2004). In this assemblage, MNI data indicate that herpetofaunal  
205 species account for 24.9% of the assemblage. This proportion dramatically increases to more than 70% of  
206 the material if the NISP is considered with nearly 60% for the non-marine turtles bone alone.  
207 Unfortunately, the turtle remains were not identified further at the moment of this first study and we  
208 were not able to locate this material in order to study it again in the framework of the present analysis.  
209 We however had access to the material collected during the 2008 excavation but only to previously  
210 studied herpetofaunal bone samples that were extracted from the complete bone samples by several  
211 master students. These bones were recovered from the four different layers identified during the 2008  
212 excavation of Moh Khiew Cave (Auetrakulvit *et al.*, 2012). The first layer (Layer 1) is composed of the  
213 upper first 90 cm of the sequence. It is a disturbed layer that has not been dated and which corresponds  
214 to the levels 1 and 2 identified by S. Pookajorn in the first excavations. The second layers include the  
215 sediment collected between 90 and 170 cm of depth. It has been the object of three radiocarbon dates  
216 between 7520+-420 BP. and 8660+-480 BP. This layer corresponds to the level 3 identified in the previous  
217 excavations. The third layer correspond to a depth between 170 and 210 cm and has been dated with  
218 two radiocarbon dates of 8730+-480 BP. and 9270+-510 BP. It was also identified as corresponding to the  
219 level 3 previously described by S. Pookajorn. The last layer corresponds to the remaining of the  
220 stratigraphy. It is a layer of scree and was not dated but associated to the level 4 described in the  
221 previous excavations.

#### 222 *The Khao Tha Phlai cave*

223 The site of Khao Tha Phlai is a cave located in southern Thailand in the province of Chumphon (E  
224 10°36'12.39"; N 99° 5'49.08"). It has been excavated by the 12<sup>th</sup> Regional Office of Fine Arts Department,  
225 Nakhon Si Thammarat since 2014, and has so far been the object of two excavation campaigns with two  
226 test-pits conducted in the deposit. A first test-pit of 9 m<sup>2</sup> (TP1) in 2014, and a second of 20 m<sup>2</sup> (TP2) in  
227 2021-2022.

228 The archaeofaunal material collected in the TP1 has been the object of a preliminary study by S.  
229 Jeawkok in the framework of a Bachelor thesis that remained unpublished. This first study conducted on  
230 6945 bone remains indicated that the reptiles represent around 25% of the NISP of the complete  
231 assemblage, the remaining bones being nearly all attributed to large mammals. In the present study we  
232 considered the herpetofaunal material previously extracted by S. Jeawkok in the faunal samples collected  
233 in the TP1. We also consulted the full archaeofaunal sample recovered from the TP2 in order to extract  
234 the reptile and amphibian bones from it. We considered separately the samples collected in the two test-  
235 pits and subdivided the samples in two assemblages corresponding to Metal Ages (between 75 and 130  
236 cm of depth in the TP2 and between 65 and 180 cm in TP1), and Neolithic periods (between 130 and 320  
237 cm of depth in the TP2 and between 180 and 320 in TP1). These layers have been dated on the basis of  
238 the typology of the archaeological artifacts they have provided.

239 *The Laang Spean cave*

240 The site of Laang Spean is a large cave of more than 1000 m<sup>2</sup> located in northwest Cambodia, in  
241 the Battambang province (E 102° 51' 00.0"; N 12°51' 00.0"). The site has been the object of a first  
242 excavation between 1965 and 1968 by R. Mourer and C. Mourer-Chauviré (Mourer-Chauviré *et al.* 1970;  
243 Mourer-Chauviré & Mourer 1970). The archaeofaunal material collected during these excavations has  
244 been the object of a preliminary study but has never been studied in depth at the exception of the  
245 rhinoceros remains (Guerin & Mourer-Chauviré 1969), and no zooarchaeological study have been  
246 conducted. Following this first exploration, the site has been the object of a new detailed archaeological  
247 excavation by H. Forestier between 2009 and 2019 aiming to document in more detail the Hoabinhian  
248 occupation previously identified (Forestier *et al.* 2015; Sophady *et al.* 2016). These excavations conducted  
249 on a surface of over 40 m<sup>2</sup> led to the discovery of an important undisturbed Hoabinhian layer dated  
250 between 5018 ± 29 cal. BP and 10 042 ± 43 cal. BP as well as several Neolithic burials dug in the  
251 Hoabinhian level (Zeitoun *et al.* 2012, 2021). These sepultures have been dated from 3335 ± 30 to 2960 ±  
252 30 BP (Sophady 2016). Regarding the subdivision of the material, the archaeological remains collected in  
253 the squares lacking traces of Neolithic perturbations (see Forestier *et al.*, 2015) have been associated to  
254 the Hoabinhian occupation. There is no stratigraphic evidence suggesting a subdivision of this Hoabinhian  
255 assemblage obviously representing several occupations over a time span of more than 5000 years.  
256 Regarding the other squares, the squares in which sepultures were found are grouped together under the  
257 term "Sepulture layer" and the first 120 cm of the disturbed squares are considered as being a "Neolithic  
258 layer". These subdivisions are however very artificial and probably correspond to a mix of Neolithic and  
259 Hoabinhian material as it has been evidenced on the lithic material these contexts provided (H.  
260 Forestier, *com. pers.*).

261 A sample of the complete faunal assemblage collected in the Hoabinhian squares has been the  
262 object of a first zooarchaeological study by S. Frère (Frère *et al.* 2018). In this first study, among the 5885  
263 vertebrate remains identified, turtle account for 44% of the NISP, monitor lizards for 1%, and large  
264 mammals for 37%. Unfortunately no MNI data were reported and the fact that most of the small  
265 fragments were not attributed to a least a size class of animal make these results difficult to interpret.  
266 The study presented in this paper corresponds to the herpetofaunal material collected in the complete  
267 assemblage of bones recovered during all the excavations since 2009. This material has been extracted by  
268 C. B. upon the consultation of all the bone samples collected on the site. The zooarchaeological study of  
269 the other groups of vertebrate for the complete Hoabinhian assemblage of Laang Spean is currently in  
270 progress and the final results are not available at the moment.

271 **Quantification of the zooarchaeological data**

272 The basic units of quantification considered are the Number of Identified Skeletal Parts (NISP)  
273 and the Weight of the Remains (WR). The fragmentation of each bone has been recorded by describing  
274 the Percentage of Completion (PC) of the anatomical elements represented by the fragment. The  
275 laterality of the bones has been registered when possible for the best represented and easiest to identify  
276 anatomical elements (i. e. peripheral plates, and all paired elements of the plastron). A Minimal Number  
277 of Elements (MNE) has been compute for each anatomical part in order to study the anatomical  
278 distribution of the remains in the different archaeological contexts. To do so, we have added the PCs of a  
279 given element and divided the result by 100. The results were rounded to the superior unit to obtain the  
280 MNEs. The Minimal Number of Individual (MNI) is defined using the anatomical element with the highest  
281 MNE. The anatomical distributions are represented by the Percentage of Representation (PR) of Dodson  
282 and Wexlar (1979) using the MNE of each anatomical elements and the MNI of the considered  
283 assemblage. All the archaeological bones have been weighted individually. In order to avoid a potential  
284 impact of taxonomic identification bias on the anatomical distribution of the remains we considered all



285 the turtle/tortoises taxa in these analyses and not only the bone fragments attributed to *Indotestudo*. As  
286 tortoises are by far the best represent taxa in the different assemblage most of the unidentified turtle  
287 bones likely represents *Indotestudo*. The positions of the peripheral plates have been identified but only  
288 for *Indotestudo* remains. The peripheral plates for which it was not possible to give a position have been  
289 posteriorly assigned to the different ranks following the distribution of those for which a position was  
290 determined. Regarding the size estimations (see below), the mean of the obtained size estimation is  
291 considered in the case several measurements were recorded on a single bone. Chi<sup>2</sup> tests were performed  
292 on the Microsoft Excel software 2007 version.

### 293 **Size estimation of archaeological *Indotestudo***

294 In order to reliably estimate the body size of the archaeological individuals of *Indotestudo* sp. we  
295 build size estimation equations on the models of what was previously done for Southeast Asian monitor  
296 lizards (Bochaton *et al.* 2019), and recently on the size and weight of species of tortoises (Esler *et al.*  
297 2019; Codron *et al.* 2022). These approaches are more powerful than considering isolated measurement  
298 (e. g. Klein and Cruz-Urbe 1983) to describe the size of subfossil animal populations because: 1) they  
299 enable to take account on several measurements from different anatomical parts to reconstruct the body  
300 size structure of a past population, and 2) it converts measurements taken on the skeleton into a variable  
301 used to describe the size of modern individuals which make easier the comparisons needed to address  
302 biological questions. To build the equations we defined a set of 86 measurements (See Supplementary  
303 Table 1; Appendix 1) that we recorded on a sample of 34 museum specimens of *Indotestudo* sp. from the  
304 Florida Museum of Natural History (UF), and the Comparative Anatomy collection of the Muséum  
305 national d'Histoire naturelle (MNHN-ZA-AC) (see all details in Supplementary Table 1). In order to have  
306 enough specimens to produce relevant and reliable predictive equations we pooled altogether the three  
307 species currently includes in the genus *Indotestudo*: *I. forstenii* (Schlegel & Müller, 1845): 8 specimens, *I.*  
308 *travancorica* (Boulenger, 1907): 14 specimens, and *I. elongata* (Blyth, 1854): 12 specimens. Differences of  
309 body proportions among species were controlled before applying this strategy to avoid including bias  
310 related to interspecific differences in the size estimations.


311 The measurements recorded were distributed on all the bones of the plastron and the carapace  
312 as well as on the long bones. Vertebrae and skull elements were not taken into account as their  
313 occurrences were too rare in the archaeological record. In addition to these measurements, in order to  
314 be able to choose a variable accounting for the “body size” of each individual, we took three  
315 measurements considered as “size variables” on the complete carapace of the modern specimens: the  
316 Carapace Straight Length (CSL), the Shell Height (SH), and the Plastron Length (PL). All the measurements  
317 collected on modern and archaeological specimens were recorded using a digital dial calliper [IP 67  
318 (Mitutoyo Corporation, Japan)]. All the measurements recorded are included in the Supplementary Table  
319 1. All statistical analyses were performed using the basic library Stats of the open-source software R (R  
320 Core Team 2020). Each size estimation equation produced is the result of a linear regression between a  
321 given log transformed measurement recorded on a bone/plate and the log transformed “size variable” of  
322 the specimens. The variables are log transformed in order to make linear the simple allometric  
323 relationship between the used variables (Huxley 1932; Gould 1966). Consequently, the obtained CSL  
324 estimation has to be log reversed using an exponential function to be obtained in the same unit as the  
325 used measurements. Obtained equations are of the form:

$$326 \quad \text{Log (“size variable”)} = (\text{Beta1}) * \text{log (osteological measurement in mm)} + (\text{Beta0})$$

327 From this initial set of equations we choose to discard all the equation that were not significant  
328 (p.value above 0.01) and/or with a low coefficient of determination (R<sup>2</sup>) (below 0.85) in order to keep  
329 only the best equations to estimate the size of the archaeological individuals.

330 **Specific identification of the *I. elongata* archaeological bone sample**

331 The archaeological bones attributed to *I. elongata* have been identified on the basis of a direct  
332 comparison with pictures of the skeletal specimens of this species used to build the SCL estimation  
333 equations. The fact that the family of the Testudinidae is only represented by very few species in  
334 Southeast Asia (*Geochelone platynota*, *Indotestudo elongata*, *Manouria emys*, and *Manouria impressa*)  
335 (Das 2010) presenting different sizes, morphologies, and distribution areas allowed to attribute with  
336 relative ease nearly all of the studied Testudinidae remains to *I. elongata*. Among the *Indotestudo* genus,  
337 as the only currently available qualitative diagnostic criteria for *I. elongata* is located on the nuchal plate  
338 (presence of a long and narrow cervical scute), we based most of our identification obtained from other  
339 plates/bones on the exact similarity between the archaeological bone remains and the morphologies  
340 present on the modern specimens of different ages we observed. An overview of the carapace  
341 morphologies of juvenile and adult *I. elongata* is provided here (Fig. 1). The remains attributed to this  
342 species also present the morphological traits common to most Testudinidae: a carapace lacking lateral  
343 keels, a costal pattern of odd costal plates with short distal end and long medial end, and even costal  
344 plates with long distal end and short medial end, octagonal and squared neural plates, peripheral plates  
345 without musk ducts, a costo-marginal sulcus superimposed to the costo-peripheral suture, a pygal plate  
346 not intersected by the posterior sulcus of the fifth vertebral scute, thickened epiplastra, and thin and  
347 vertical inguinal and axillary buttresses. Among Testudinidae the genus *Indotestudo* is characterized by  
348 the fact that the humeropectoral sulcus is crossing the entoplastron (Auffenberg 1974). The  
349 establishment of robust and quantified diagnostic criteria for the identification of isolated bones of  
350 Southeast Asian turtles has still to be performed. As a comment, we signal that the characteristic nuchal  
351 scute morphology of *I. elongata* was present on all the nuchal plate (N=109) attributed to this species at  
352 the exception of a single remain from Laang Spean cave. It has been previously signaled that the nuchal  
353 scute could be absent on some specimens (Ihlow *et al.* 2016) and our data indicate a frequency of such  
354 feature inferior to 1%.



355  
 356 **Figure 1** - A) Drawing of the carapace of an adult specimen of *Indotestudo elongata* (CUMZ-R-  
 357 TT181); B) Drawing of the plastron of an adult specimen of *Indotestudo elongata* (CUMZ-R-TT181);  
 358 C) Drawing of the carapace of a juvenile specimen of *Indotestudo elongata* lacking peripherals (UF-  
 359 34760); D) Drawing of the plastron of a juvenile specimen of *Indotestudo elongata* showing central  
 360 fontanel (UF-34760). **Abbreviations:** A. s.: Abdominal scute, An. s.: Anal scute, G. s.: Gular scute,  
 361 Co.: Costal plate, F. s.: Femoral scute, Ep.: Epiplastron, En.: Entoplastron, F.: plastral fontanel, H. s.:  
 362 Humeral scute, Hyo.: Hyoplastron, Hyp.: Hypoplastron, M. s.: Marginal scute, Ne.: Neural plate, Nu.  
 363 p.: Nuchal plate, C. s.: Cervical scute, P. s.: Pectoral scute, Pe.: Peripheral plate, Pl. s.: Pleural scute,  
 364 Py.: Pygal plate, S.p. 1: Supra-pygal 1, S.p. 2: Supra-pygal 2, S. s.: Supra-caudal scute, V. s.: Vertebral  
 365 scute, Xi.: Xiphiplastron.

366  
 367  
 368

## Results

369 **Size predictive equations**

370 In order to choose the variable we will use as our “size” variable we tested the correlations  
371 between the three “body size” measurements taken on the complete carapaces of our *Indotestudo* spp.  
372 modern specimens. Our results indicate that the Straight Carapace Length (SCL) is strongly correlated to  
373 the Plastron Length (PL) ( $R^2=0.97$ ), and that the Shell Height is more weakly correlated to the two other  
374 measurements ( $R^2=0.93$  and  $0.92$ ). This could be related to sex specific or interspecific differences  
375 regarding the height of the carapace among the considered specimens but these hypotheses are not  
376 possible to test considering the small size of our sample. As a result and because the carapace length is  
377 the most used variable to describe the size of a turtle we choose the SCL as our size scalar but the PL  
378 could have been equally considered.

379 From our complete modern sample, a set of 86 equations corresponding to the 86 initial  
380 recorded measurements has been produced. This set has been refined to keep only the equations  
381 providing the most reliable (significant linear relationships) and precise (high  $R^2$  linear relationships) SCL  
382 estimations. This final set of equations include 52 equations from 52 measurements distributed on  
383 epiplastron, entoplastron, hyoplastron, hypoplastron, xiphiplastron, nuchal plate, neural plates (ranks 1,  
384 2, 3, 4, 6, and 7), peripheral plates (ranks 1, 2, 3, 8, 9, and 10), 2nd supra-pygal plate, pygal plate,  
385 humerus, radius, ulna, femur, tibia, and fibula (Fig. 2; Table. 1).


386

387 **Table 1** - Equations retained for the prediction of the Straight Carapace Length (SCL) of the  
388 archaeological *Indotestudo* specimens. Are indicated: the measurement used (to be recorded on  
389 the archaeological specimen, the slope X (“Beta1” to integrate in the equation indicated in the  
390 Material and Method section), and the intercept Y (“Beta0” to integrate in the equation indicated in  
391 the Material and Method section). Is also indicated the coefficient of determination ( $R^2$ ) of each  
392 relation, the p.values, and the degree of freedom of each linear regression.

Measurement Used	Y	X	R <sup>2</sup>	P. value	Freedom degree
Greatest Length Nuchal plate (GL)	1.44	1.05	0.89	2.48E-14	27
Greatest Heigth Peripheral plate 1 (GH)	2.30	0.91	0.89	2.77E-14	27
Greatest Width Peripheral plate 1 (GW)	1.95	1.01	0.90	3.43E-15	27
Greatest Heigth Peripheral plate 2 (GH)	1.68	1.11	0.92	4.10E-16	26
Greatest Width Peripheral plate 3 (GW)	1.96	1.06	0.88	5.70E-14	27
Greatest Heigth Peripheral plate 3 (GH)	2.42	0.84	0.93	2.20E-16	27
Greatest Width Peripheral plate 8 (GW)	2.45	0.89	0.86	5.22E-13	27
Greatest Heigth Peripheral plate 8 (GH)	2.19	0.91	0.94	2.20E-16	27
Greatest Width Peripheral plate 9 (GW)	2.58	0.86	0.89	1.47E-14	27
Greatest Heigth Peripheral plate 9 (GH)	2.27	0.89	0.90	7.29E-15	27
Greatest Width Peripheral plate 10 (GW)	2.26	0.95	0.89	3.04E-14	27
Greatest Heigth Peripheral plate 10 (GH)	2.69	0.77	0.90	3.55E-15	27
Greatest Length Neural plate 2 (GL)	2.54	0.91	0.87	2.82E-13	27
Greatest Width Neural plate 2 (GW)	1.55	1.14	0.92	3.36E-16	27
Greatest Length Neural plate 3 (GL)	2.20	1.07	0.90	4.73E-15	27
Greatest Length Neural plate 4 (GL)	2.52	0.93	0.85	1.09E-12	27
Greatest Width Neural plate 4 (GW)	1.82	1.05	0.92	2.20E-16	27
Greatest Width Neural plate 6 (GW)	2.10	0.98	0.91	6.00E-16	27
Greatest Width Neural plate 7 (GW)	2.23	0.96	0.86	3.95E-13	27
Greatest Width Supra-pygal plate 2 (GW)	2.19	0.86	0.88	1.61E-13	26
Greatest Heigth Pygal plate (GH)	2.98	0.68	0.93	2.20E-16	26
Greatest Length Epiplastron (GL)	1.48	1.06	0.95	2.20E-16	27
Greatest Length Entoplastron (GL)	1.54	1.09	0.86	3.09E-13	27
Greatest Width Entoplastron (GW)	1.76	1.02	0.92	4.25E-16	27
Greatest medial Length Hyoplatron (GmL)	1.81	0.96	0.87	1.14E-13	27
Greatest Width Hyoplastron (GW)	0.02	1.29	0.94	2.20E-16	25
Greatest medial Length Hypoplastron (GmL)	0.44	1.25	0.90	7.32E-15	27
Greatest Width Hypoplastron (GW)	-0.09	1.32	0.95	2.20E-16	25
Greatest Length Xiphiplastron (GL)	1.53	1.02	0.88	6.69E-14	27
Greatest Width Xiphiplastron (GW)	1.09	1.15	0.92	2.20E-16	27
Greatest Length Humerus (GL)	1.16	1.08	0.95	2.20E-16	24
Greatest proximal antero-posterior Width Humerus (GpapW)	2.74	0.99	0.91	3.32E-14	24
Greatest proximal dorso-ventral Width Humerus (GpdvW)	2.59	1.00	0.94	8.45E-16	24
Greatest Length Ulna (GL)	1.52	1.12	0.96	2.20E-16	23
Greatest distal antero-posterior Width Ulna (GdapW)	3.60	0.93	0.90	4.69E-13	26
Greatest proximal latero-medial Width Ulna (GplmW)	3.13	1.02	0.87	1.22E-11	23
Greatest Length Radius (GL)	1.34	1.17	0.94	1.82E-15	23
Greatest proximal latero-medial Width Radius (GplmW)	3.18	1.10	0.92	3.64E-14	23
Greatest proximal antero-posterior Width Radius (GpapW)	3.96	0.89	0.89	1.20E-12	23
Greatest distal antero-posterior Width Radius (GdapW)	3.91	0.99	0.86	2.08E-11	23
Greatest Length Femur (GL)	1.41	1.03	0.96	2.20E-16	23
Greatest proximal antero-posterior Width Femur (GpapW)	3.15	0.86	0.93	1.89E-14	23
Greatest proximal dorso-ventral Width Femur (GpdvW)	2.91	0.90	0.97	2.20E-16	23
Greatest distal antero-posterior Width Femur (GdapW)	3.46	0.77	0.86	2.91E-11	23
Greatest Length Fibula (GL)	1.57	1.06	0.94	1.81E-14	21
Greatest distal latero-medial Width Fibula (GdlmW)	3.59	0.94	0.89	1.06E-11	21
Greatest distal antero-posterior Width Fibula (GdapW)	3.92	0.96	0.90	9.30E-12	21
Greatest Length Tibia (GL)	1.39	1.09	0.95	1.25E-15	22
Greatest proximal latero-medial Width Tibia (GplmW)	3.06	1.01	0.92	2.16E-13	22
Greatest proximal antero-posterior Width Tibia (GpapW)	3.20	1.02	0.94	3.03E-15	22
Greatest distal latero-medial Width Tibia (GdlmW)	3.45	1.00	0.92	1.99E-13	22
Greatest distal antero-posterior Width Tibia (GdapW)	3.60	0.97	0.91	9.04E-13	22







405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417

**Figure 3** - Examples of the turtle bone remains studied. A) Plastron of an adult *I. elongata* in anatomical connection from the Hoabinhian layer of Laang Spean Cave (ventral view); B) Plastron of a juvenile *I. elongata* in anatomical connection from the Hoabinhian layer of Laang Spean Cave (ventral view); C) Nuchal plate of *I. elongata* from the Hoabinhian layer of Laang Spean Cave (dorsal view); D) Pygal plate of a juvenile *I. elongata* from the Neolithic layer of Laang Spean Cave (posterior view); E) Pygal plate of an adult *I. elongata* from the Neolithic layer of Laang Spean Cave (posterior view); F) Left peripheral plates (1<sup>st</sup> to 3<sup>rd</sup>) of *I. elongata* in anatomical connection from the layer 2 of Moh Khiew Cave presenting burning traces limited to the internal side of the carapace; G) Left peripheral plates (8<sup>th</sup> to 11<sup>th</sup>) of a *Geoemydidae* in anatomical connection whose ventral part has been cut-down from the site of Doi Pha Kan.

418 *Doi Pha Kan Rockshelter*  
419 Composition of the herpetofaunal assemblages

420 The herpetofaunal assemblage of Doi Pha Kan consist of 8414 bone remains weighting a total of  
 421 6875 gr. and representing a least 115 individuals. Most of these bones correspond to non-marine turtles  
 422 (Fig. 3; Table. 2) in term of WR (74%), NISP (56%), and MNI (47%) with the second most represented  
 423 group being Monitor lizards (23% of the WR, 38% of the NISP, and 38% of the MNI) followed by rare  
 424 snakes, amphibians, and small lizards bone remains.

425 **Table 2** - Weight of the Remains (WR), Number of Identified Skeletal Parts (NISP), and Minimal  
 426 Number of Individual (MNI) corresponding to the different taxa identified in the complete Doi Pha  
 427 Kan Rock-shelter herpetofaunal assemblage.

	NISP	WR	NMI
Turtle/Tortoise	4762	5113.02	55
Monitor lizard	3203	1610.39	44
Snake	375	141	3
Amphibian	67	10.57	10
Small lizards	7	0.82	3
<b>Total</b>	<b>8414</b>	<b>6875.8</b>	<b>115</b>

428  
 429

430 Regarding the bones attributed to turtles and tortoises (Table. 3), 66.8% of them (50.8% of the  
 431 WR) could not be attributed to a given family. Among the fragments that were identified to at least the  
 432 family level, 71 % of them were attributed to *Indotestudo elongata* (71% of the WR and 76% of the MNI),  
 433 28% (28% of the WR and 18% of the MNI) to Geoemydidae, and less than 1% (0.6% of the WR and 5% of  
 434 the MNI) to Trionychidae. Note that for Trionychids remains from all localities, attribution can be done  
 435 for all the plates because of their ornamentation, they are therefore not underrepresented because they  
 436 cannot be easily identified.

437

438 **Table 3** - Number of Identified Skeletal Parts (NISP), Weight of the Remains (WR), and Minimal  
 439 Number of Individual (MNI) corresponding to the different turtle/tortoise taxa identified in the  
 440 complete Doi Pha Kan Rock-shelter assemblage.


	NISP	WR	NMI
<i>Indotestudo elongata</i>	1122	1787.17	42
Geoemydidae	447	711.34	10
Trionychidae	12	14.13	3
Turtle ind.	3181	2600.38	
<b>Total</b>	<b>4762</b>	<b>5113.02</b>	<b>55</b>

441  
 442

#### 443 Taphonomy of the turtle/tortoise bone assemblage

444 Among the 4762 bone fragments attributed to turtle/tortoises, 303 are complete elements  
 445 (6.3%), and 249 are nearly complete (at least 90% of the bone is preserved) while 1389 (29%) are small  
 446 fragments representing less than 5% of the complete anatomical part. The average percentage of  
 447 completion of the bones is 32%. The overall PR is 28%. The best represented bones (Fig. 4 – A) are the  
 448 stylopods (humerus and femur with PR>75%) followed by the tibia, the radius, the scapula, the coracoid,  
 449 and the epiplastron (PR>50%). Then follow the rest of the long bones, and most of the plates that are the  
 450 easiest to identify. The peripheral plates of the bridge are less represented (PR=10%) than all the other  
 451 peripheral plates (PR=39%). The skull, vertebrae and all small elements of the hands and foot are nearly  
 452 absent. The largest plates (hyoplastron and hypoplastron) are the most fragmented with completion  
 453 means of less than 26%. The peripheral plates corresponding to the bridge are also more heavily


454 fragmented (completion mean of 59%) compare to the other peripheral plates (completions means 67-  
455 89%). Burning (black) and carbonization (grey/white) traces were observed on 690 bones (14% of the  
456 total NISP). Such traces were present indiscriminately on every anatomical element of the carapace and  
457 skeleton. Cut marks were observed on only five bones, six peripheral plates, and one hypoplastron. A  
458 series of peripheral plates attributed to a Geoemydidae turtle bears clear traces of a clean cut aiming to  
459 cut the ventral part of the carapace (Fig. 3-G). Lastly, 120 bones have been observed as corresponding to  
460 preserved anatomical connections on the field between unfused plates. This indicates that at least a part  
461 of the assemblage was undisturbed prior to the excavation.  
462



463  
464 **Figure 4** - Anatomical distributions of the turtle/tortoise remains collected in the sites of Doi Pha  
465 Kan rockshelter (A), and Moh Khiew Cave (B-E). The percentage of representation (PR) is considered  
466 here to provide a graphical visualization of the different values observed for the different  
467 anatomical elements.

468 Size of *Indotestudo elongata* archaeological individuals

469 The measurements recorded on the *I. elongata* archaeological material of Doi Pha Kan enabled  
470 for the reconstruction of 201 SCL estimations included between 64 and 292 mm with a mean of 182 mm  
471 (Figure. 5-A) and corresponding to at least 42 individuals. The distribution of these sizes was not  
472 unimodal (Hartigans' dip test,  $p.val>0.05$ ) but could be bi-modal with a population of small individuals  
473 around 120 mm, and a second one between 140 and 260 mm.  
474



475 **Figure 5** - Histograms of the *Indotestudo elongata* size reconstructions (Standard Carapace Length)  
476 obtained from the different herpetofaunal bone assemblages studied: A) Doi Pha Kan Rock-shelter  
477 (NMI=42); B) Moh Khiew Cave (NMI=59); C) Laang Spean Cave (NMI=75); D) Khao Ta Phlai Cave  
478 (NMI=26). The black bars represent the minimal size of the sexually mature specimens based on  
479 modern data collected on modern *I. elongata* populations.  
480

481 Moh Khiew cave

482 Composition of the herpetofaunal assemblages

483 The herpetofaunal assemblage of Moh Khiew consists in 9 108 bone remains weighting 8351 gr.  
484 Those bones are mostly distributed in the layers 2 (51% of the NISP, and 52% of the WR), 1 (26% of the  
485 NISP, and 24% of the WR), and 3 (17% of the NISP, and 18% of the WR) (Table. 4). The complete  
486 assemblage includes bone fragment from a least 152 individuals.  
487 The full sample mostly correspond to non-marine turtle remains (63% of the NISP, 74% of the WR, and  
488 52% of the MNI), followed by Monitor lizards (25% of the NISP, 16% of the WR, and 24% of the MNI), and  
489 snakes (8% of the NISP, 8% of the WR, and 4% of the MNI). The small size lizards (excluding snakes),  
490 amphibians and crocodile remains altogether represent less than 5% of the assemblage in term of NISP  
491 and WR. The distribution of these groups among the layers is however far from homogenous. Indeed,



492 turtle bone remains represent more than 75% of the NISP, 85% of the WR, and 60% of the MNI in the  
 493 layers 2 to 4 but only 20% of the NISP, 35% of the WR, and 21% of the MNI in the layer 1. At the opposite,  
 494 Monitor lizards represent 46% of the NISP, 31% of the WR, and 35% of the MNI in the layer 1 but less  
 495 than 20% of NISP, 13% of the WR, and 25% of the MNI in the other layers. Snakes are also better  
 496 represented in the layer 1 (23% of the NISP) than in the subsequent levels (less than 2.6% of the NISP).  
 497 Chi<sup>2</sup> tests performed on the NISP indicate that the faunal composition of the layer 1 significantly differs  
 498 (P.val< 0.01) from the layers 2 and 3, the effective of the layer 4 being too low to conduct a statistical  
 499 test.

500  
 501  
 502  
 503

**Table 4** - Number of Identified Skeletal Parts (NISP), Weight of the remains (WR), and Minimal Number of Individuals (MNI) identified in the complete herpetofaunal assemblage of the different layers of the 2008 excavation of Moh Khiew Cave.

	Layer 1			Layer 2			Layer 3			Layer 4			Total		
	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI
Turtle/Tortoise	503	702.8	6	3614	3789	53	1184	1288	15	439	481.3	5	<b>5740</b>	<b>6261</b>	<b>79</b>
Monitor lizard	1111	641.4	17	872	463.8	12	298	190.1	5	43	25.4	2	<b>2324</b>	<b>1321</b>	<b>36</b>
Snake	560	612.2	3	117	52.8	1	37	29.15	1	7	6.4	1	<b>721</b>	<b>700.6</b>	<b>6</b>
Amphibian	144	38.45	14	34	6.5	4	22	3.92	3	0	0	0	<b>200</b>	<b>48.87</b>	<b>21</b>
Small lizards	89	8.27	7	4	0.3	1	28	3.1	1	<b>0</b>	<b>0</b>	<b>0</b>	<b>121</b>	<b>11.67</b>	<b>9</b>
Crocodile	2	8.5	1	0	0	0	0	0	0	0	0	0	<b>2</b>	<b>8.5</b>	<b>1</b>
<b>Total</b>	<b>2409</b>	<b>2012</b>	<b>48</b>	<b>4641</b>	<b>4312</b>	<b>71</b>	<b>1569</b>	<b>1514</b>	<b>25</b>	<b>489</b>	<b>513.1</b>	<b>8</b>	<b>9108</b>	<b>8351</b>	<b>152</b>

504  
 505

506 Regarding the identification of non-marine turtle taxa (Table 5), between 66 and 79% of the bone  
 507 fragment have not been associated with at least a family rank identification. Regarding the identified  
 508 families, Testudinidae (*Indotestudo elongata*) represent a stable major part of the identified bone  
 509 remains (between 68 and 63% of the NISP, 70-56% of the WR, and 86-50% of the MNI depending of the  
 510 layer). The other fragments have been attributed to Geoemydidae turtles (36-29% of the NISP, 36-28% of  
 511 the WR, and 33-12% of the MNI), and Trionychidae (0-2% of the NR and WR). The composition of the  
 512 turtle assemblages in term of families seems then to be fairly stable across the layers as indicated by  
 513 results of Chi<sup>2</sup> tests that did not indicated significant differences in the distribution of the NISP from the  
 514 different testable layers (P.val> 0.01).

515  
 516  
 517  
 518

**Table 5** - Number of Identified Skeletal Parts (NISP), Weight of the remains (WR), and Minimal Number of Individuals (MNI) in the non-marine turtle assemblage from the different layers of the 2008 excavation of Moh Khiew Cave.

	Layer 1			Layer 2			Layer 3			Layer 4			Total		
	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI
<i>Indotestudo elongata</i>	109	239.9	4	853	1274	43	193	323.9	10	60	105	2	<b>1215</b>	<b>1943</b>	<b>59</b>
Geoemydidae	61	107.3	2	386	512.9	6	111	253.9	3	26	62.2	1	<b>584</b>	<b>936</b>	<b>12</b>
Trionychidae	1	0.7	0(1)	16	37.8	1	0	0	1	2	2.5	1	<b>19</b>	<b>41</b>	<b>3</b>
Turtle ind.	332	354.9	0	2359	1964	3	880	709.9	1	351	311.6	1	<b>3922</b>	<b>3340</b>	<b>5</b>
<b>Total</b>	<b>503</b>	<b>703</b>	<b>6</b>	<b>3614</b>	<b>3789</b>	<b>53</b>	<b>1184</b>	<b>1288</b>	<b>15</b>	<b>439</b>	<b>481</b>	<b>5</b>	<b>5740</b>	<b>6261</b>	<b>79</b>

519  
 520

#### 521 Taphonomy of the turtle/tortoise bone assemblage

522 The fragmentation of the 5740 bone fragments analyzed increase with the depth. In the layer 1,  
 523 8.9% of the bones are complete, and 12.9% nearly complete, in layer 2 these percentages falls to 7.5 and  
 524 12.1%, then to 5.6 and 9.6% in layer 3, and finally to 3.9 and 7.7% in layer 4. The average percentage of  
 525 completeness of the bones is also slightly higher in layers 1 and 2 (36 and 37%) than in the layers 3 and 4  
 526 (34 and 28%). These differences are only significant between the layers 1 and 4 (Chi<sup>2</sup> test; p.  
 527 value<0.01).The anatomical distribution of the remains present strong variations between the layers (Fig.

528 4) but the sizes of the assemblages are also very dissimilar with the bone samples of the layers 1 and 4  
529 containing only 503 and 439 remains and those of the layers 2 and 3 containing 3614 and 1184 bone  
530 fragments. Distributions in small samples could be more strongly impacted by random effects than larger  
531 assemblages and a strict comparison of the four layers might not make sense at all. Some general trends  
532 can, however, still be noted just as the overall PR which is between 27 and 24% in all the layers at the  
533 exception of the first in which it is slightly higher (36%). Nearly all the anatomical parts are present in  
534 every layer but the skulls, vertebrae, and extremities are nearly absent. The stylopods (humerus and  
535 femur) are the best represented bones in the richest layers. They are also well represented in layers 1  
536 and 2 but are outnumbered by some specific carapace plates. Girdles and zeugopods are also present but  
537 in smaller number. Regarding the carapace and the plastron, no clear pattern emerges at the exception  
538 of the nearly systematic lower representation of the peripheral plates of the bridge (PR=25-21%) in  
539 respect to the other peripheral plates (mean PR=52-29%). This could be explained by an identification  
540 bias related to the lower mean completion rate of peripheral plates of the bridge in respect to the others.  
541 Burning (black) and carbonization (grey/white) traces were observed on 325 bones (5.7% of the total  
542 NISP). Such traces were present indiscriminately on every anatomical element and every side of the  
543 carapace and skeleton parts. They were recorded on the internal side of several peripheral plates that  
544 were still in anatomic connection at the moment of the excavation (Fig. 3-F). Such observations were,  
545 however, not repeated on the rest of the material. Cut marks were observed on only three bones: one  
546 peripheral plate, one nuchal plate, and one xiphiplastron. Among the full assemblage 61 fragments of  
547 carapaces were still in anatomical connection at the moment of the excavation. These elements were  
548 distributed mostly in the layers 2 and 3 but also in the lower part of the layer 1 at a depth of 70-80 cm.

549

#### 550 Size of *Indotestudo elongata* individuals

551 The measurements recorded on the *I. elongata* archaeological material of Moh Khiew Cave  
552 enabled for the reconstruction of 201 SCL estimations included between 98 and 310 mm and with a mean  
553 of 193 mm (Figure. 5-B) and corresponding to at least 59 individuals. The mean size of the tortoises is  
554 similar in all layers and no statistically significant differences was noted (student T-test; p.val>0.01). In  
555 layer 1 (N=25) the mean size was 198 mm, 194 mm in layer 2 (N=169), 187 mm in layer 3 (N=52), and  
556 finally 195 mm in layer 4 (N=19). The largest observed specimen was in the layer 1. The global  
557 distribution of these sizes was not unimodal (Hartigans' dip test, p.val>0.01) and mixture models indicate  
558 it is most likely bimodal with a best represent group of individuals around 220 mm and a second group of  
559 smaller specimens around 150 mm.

560

#### 561 *Khao Ta Phlai Cave*

#### 562 Taxonomic composition

563 A total of 3763 bone remains of herpetofauna weighting 7239 gr. and representing at least 43  
564 individuals were analyzed from the two excavated test-pits of the site of Khao Ta Phlai (Table. 6). Most of  
565 these bones correspond to turtle or tortoises in term of WR (87% in TP1 and 90% in TP2), NISP (81% in  
566 TP1 and 86% in TP2), and MNI (65% in TP1 and 70% in TP2). The second most represented herpetofaunal  
567 group is the Monitor lizards (12 and 9% of the WR, 16 and 12% of the NISP, and 23 and 22% of the MNI)  
568 followed by some rare remains of snakes as well as few amphibian bones in the TP2 only.

569

570 **Table 6** - Number of Identified Skeletal Parts (NISP), Weight of the Remains (WR), Number of  
571 Remains (NR), and Minimal Number of Individual (MNI) corresponding to the different  
572 herpetofaunal taxa identified in the herpetofaunal bone assemblages collected in the two test-pits  
573 of the site of Khao Ta Phlai.



	TP1			TP2		
	NISP	WR	NMI	NISP	WR	NMI
Amphibians				3	0.51	1
Snakes	38	33.4	2	37	69.3	2
Turtle/Tortoise	1335	2509.7	19	1825	3901	18
Monitor lizards	274	347.7	6	251	377.9	6
<b>Total</b>	<b>1647</b>	<b>2891</b>	<b>27</b>	<b>2116</b>	<b>4349</b>	<b>27</b>

574  
575

576 Although the distribution of the material between the two TPs is somewhat homogenous, the  
577 repartition of the bones across the two main periods documented (metal ages and Neolithic) is quite  
578 different. Indeed, in the TP2, most of the material (95% of the NISP) is located in the Neolithic layers  
579 while in TP1 the bone are more evenly distributed (55% of the NISP in the Metal Ages layers and 44% of  
580 the NISP in the Neolithic levels).

581 The distribution of the main herpetofaunal taxa across the layers does not present strong  
582 variations (Tab. 7) as only the very poor metal age layer of the TP2 significantly differ from the other  
583 levels (Chi<sup>2</sup>; p.value <0.01). Turtle bones are always the best represented (between 71 and 87% of the  
584 NISP) but Monitor lizards seems a bit better represented in Metal Ages layers from TP1 and 2 (18 and  
585 20% compared to Neolithic layers (14 and 11% of the NISP). These tendencies are, however, not  
586 statistically significant and difficult to interpret in the absence of a study of the complete faunal  
587 assemblages.

588

589 **Table 7** - Number of Identified Skeletal Parts (NISP), Weight of the Remains (WR), and Minimal  
590 Number of Individual (MNI) corresponding to the different herpetofaunal taxa identified in the  
591 herpetofaunal bone assemblages of the two periods identified in the two test-pits of the site of  
592 Khao Ta Phlai.

	TP1-Metal Age			TP1-Neolithic			TP2-Metal Age			TP2-Neolithic		
	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI
Amphibians										3	0.51	1
Snakes	33	22.4	2	5	11	2	9	13.4	2	28	55.9	2
Turtle/Tortoise	720	1011.8	14	615	1497.8	9	70	131.2	2	1755	3769.8	21
Monitor lizards	166	146.2	1	108	201.5	1	20	36.2	1	231	341.7	3
<b>Total</b>	<b>919</b>	<b>1180</b>	<b>17</b>	<b>728</b>	<b>1710</b>	<b>12</b>	<b>99</b>	<b>180.8</b>	<b>5</b>	<b>2017</b>	<b>4168</b>	<b>27</b>

593  
594

595 The identification rate of non-marine turtles bones was lower in the TP1 (41% of the WR and 25%  
596 of the NR) than in TP2 (57% of the WR and 35% of the NR). Regarding the bones attributed to a given  
597 family (Tab. 8), TP1 provided nearly as much Testudinidae as Geoemydidae in terms of WR and NISP  
598 while *Indotestudo elongata* is much more represented than the latter in the TP 2 (61% of the WR and  
599 69% of the NISP). These differences are statistically significant (Chi<sup>2</sup> test; p.value<0.01). Trionychidae are  
600 present in the two TPs. If the chronological phases are considered (Tab. 9), Geoemydidae and  
601 Trionychidae are significantly better represented in the upper layers of the TP2 and TP1 corresponding to  
602 the metal ages. These two layers do not significantly differ in term of family composition (Chi<sup>2</sup> test;  
603 p.value>0.01) but significantly differ from the two Neolithic layers (Chi<sup>2</sup> test; p.value<0.01). This trend to  
604 a more important exploitation of freshwater turtles during the metal ages in regard to the Neolithic  
605 period is for now difficult to interpret considering the possible issues of chronological associations  
606 between the layers of the two TPs and the possibility of a spatial variation in the distribution of the  
607 remains inside the site. Indeed freshwater turtles are also better represented in the Neolithic layer of the  
608 TP1 compare to TP2.

609

610  
611  
612

**Table 8** - Number of Identified Skeletal Parts (NISP), Weight of the Remains (WR), and Minimal Number of Individual (MNI) corresponding to the different turtle/tortoise taxa identified in the bone assemblages collected in the two test-pits of the site of Khao Ta Phlai.

	TP1			TP2		
	NISP	WR	NMI	NISP	WR	NMI
<i>Indotestudo elongata</i>	175	435.5	8	441	1357	14
Geoemydidae	150	581.9	5	191	838.6	3
Trionychidae	10	14.1	1	9	32.2	1
Turtle ind.	1000	1478		1184	1673	
<b>Total</b>	<b>1335</b>	<b>2510</b>	<b>14</b>	<b>1825</b>	<b>3901</b>	<b>18</b>

613  
614  
615  
616  
617  
618


**Table 9** - Number of Identified Skeletal Parts (NISP), Weight of the Remains (WR), and Minimal Number of Individual (MNI) corresponding to the different turtle/tortoise taxa identified in the bone assemblages collected in the different chronological phases of the two test-pits of the site of Khao Ta Phlai.

	TP1-MA			TP1-NE			TP2-MA			TP2-NE		
	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI
<i>Indotestudo elongata</i>	86	149	7	89	286.5	6	14	40.2	1	427	1316.6	12
Geoemydidae	81	243.2	2	69	338.7	3	13	28.7	1	178	809.9	4
Trionychidae	10	14.1	1				3	5.2	1	6	27	1
Turtle ind.	543	605		457	872.63		70	57.1		1144	1616.3	
<b>Total</b>	<b>720</b>	<b>1011</b>	<b>10</b>	<b>615</b>	<b>1498</b>	<b>9</b>	<b>100</b>	<b>131.2</b>	<b>3</b>	<b>1755</b>	<b>3770</b>	<b>17</b>

619  
620  
621  
622  
623  
624  
625  
626  
627  
628

#### Taphonomy of the turtle/tortoise bone assemblage

Among the 3 160 bone fragments attributed to turtle/tortoises in the material of Khao Ta Phlai, 221 were complete elements (6.9%), and 370 were nearly complete (at least 90% of the complete bone) while 760 (24%) were small fragments representing less than 5% of the complete anatomical part. The average percentage of completion of the bones is 29%. This value is similar in the Neolithic layers of TP1 and TP2 (29%). It is slightly lower in the metal ages layer of the TP1 (25%), and higher in the same period layers from TP2 (39%) but the small size of this assemblage does not allow to consider this result as significant.



629  
 630  
 631  
 632  
 633

**Figure 6** - Anatomical distributions of the non-marine turtle remains collected in the different test-pits and layers of the site of Khao Ta Phlai: (A) Test-pit 1 – Metal Ages, (B) Test-pit 1– Neolithic, (C), Test-pit 2– Neolithic. The percentage of representation (PR) is considered here to provide a graphical visualization of the different values observed for the different anatomical elements.

634  
 635  
 636  
 637  
 638  
 639  
 640  
 641  
 642  
 643  
 644  
 645  
 646  
 647  
 648  
 649  
 650  
 651

The anatomical distributions of the turtle bone elements present strong variations among the different layers (Fig. 6). The distribution observed in the Neolithic layers of the TP2 is fairly homogenous (mean PR=33%) with a representation of all the anatomical part at the exception of the smallest elements (phalanges, carpal and tarsal articulations, and vertebrae), and the skull. The most robust anatomical parts are the best represented (peripheral plates, epiplastron, entoplastron, and nuchal plate -PR>38%) and the most fragile the least represented (zeugopods, and most girdles elements - PR<5%). The only exception to that pattern is the peripheral plates of the bridge (PR=15%) which are least represented than the other elements of the carapace and other peripherals (mean PR=73%). The distribution pattern is quite less homogenous in TP1 where the mean PR is lower (24% in the Neolithic layers and 15% in the metal ages layers) but the Neolithic layers follow the same general pattern as the one of the TP2 with a lower global representation due to the strong presence of a single peripheral plate rank. In this last layer, another difference with TP2 is that the stylopods are also better represented than the elements from the carapace. The anatomical distribution of the bones collected in the metal ages layers of TP1 is very different with a very strong representation of the stylopods (mean PR=76%) compare to the most robust elements of the carapace (PR=29-15%). Otherwise, the same general observations applies with a lack of skull and extremities elements, a better representation of the most robust elements of the carapace, and a lower representation of the peripheral plates of the bridge (PR=5%) in regard of other peripheral plates (Mean PR=18%).

652  
 653

Most of the material (72% of the NISP) was covered by a veil of calcite which made very difficult the observation of the surface alterations of the bones. Despite this important limitation, nine bones

654 were recorded as presenting traces of dissolution under the effect of flowing water, and 182 as bearing  
 655 traces of burning and carbonization. During the study we also recorded 22 associations of bones from the  
 656 same context being in anatomical connection in the two TPs below 120 cm of depth in the TP 2 and 135  
 657 cm in the TP 1. This indicates that the material from the deepest layers was not strongly disturbed since  
 658 its deposition. Nine combination of bone in anatomical connection linked together by concretion were  
 659 also found in the same layers.

660

661 Size of *Indotestudo elongata* individuals

662 The measurements recorded on the *I. elongata* archaeological material of the Khao Ta Phlai site  
 663 have enabled for the reconstruction of 219 SCL estimations included between 108 and 252 mm and with  
 664 a mean of 184 mm (Figure. 5-D) and corresponding to at least 26 individuals. Most of the size estimations  
 665 are from the Neolithic layers of the TP2 (n=158) and TP1 (n=31) while only 29 estimations were obtained  
 666 from the metal ages layers (mean size = 178 mm). The strong disparities of the distributions of the size  
 667 estimations between the archaeological contexts do not allow for an individual comparison of the  
 668 different layers. The Neolithic layers provided mean SCL values of 185 and 186 mm. The global  
 669 distribution of these sizes was not unimodal (Hartigans' dip test, p.val>0.01) and mixture models indicate  
 670 it is most likely bimodal with a group of individuals around 210 mm and a second group of smaller  
 671 specimens around 165 mm.

672

673 Laang Spean Cave

674 Composition of the herpetofaunal assemblage

675 The complete herpetofaunal assemblage of Laang Spean consists of 9533 bone fragments  
 676 weighting 18 804 gr. and representing at least 115 individuals (Tab. 10). Most of them come from the  
 677 Hoabinhian layer accounting for 76% of the NISP, 78% of the WR, and 62% of the MNI. The “Neolithic”  
 678 and “sepulture” assemblages are of similar sizes and account respectively for 10.5 and 13.2% of the NR.  
 679 The material corresponds nearly exclusively to non-marine turtle remains which account for 92% of the  
 680 NISP, 95% of the WR, and 70% of the MNI. Monitor lizards represent 5% of the NISP of the assemblage,  
 681 snakes are rare (2% of the NISP), and the occurrence of amphibians and smaller lizards is insignificant  
 682 (below 1%). The distribution of the taxa is not statistically different across the sub-assemblages (Chi<sup>2</sup> test;  
 683 p.value>0.01).

684

685 **Table 10** - Number of Identified Skeletal Parts (NISP), Weight of the remains (WR), and Minimal  
 686 Number of Individuals (MNI) studied in the complete herpetofaunal assemblage from the different  
 687 layers of the Laang Spean Cave.

	Hoabinhian layer			Neolithic layer			Sepulture layers			Total		
	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI
Turtle/Tortoise	6650	13893	56	904	1854	12	1195	2057	12	<b>8749</b>	<b>17804</b>	<b>80</b>
Monitor lizard	406	497	3	30	48	2	39	56	3	<b>475</b>	<b>601</b>	<b>8</b>
Snake	158	311	3	16	37	1	19	30	2	<b>193</b>	<b>378</b>	<b>6</b>
Amphibian	45	9	7	33	6	5	6	1	2	<b>84</b>	<b>16</b>	<b>14</b>
Small lizards	13	2	3	16	2	3	3	1	1	<b>32</b>	<b>5</b>	<b>7</b>
<b>Total</b>	<b>7272</b>	<b>14712</b>	<b>72</b>	<b>999</b>	<b>1947</b>	<b>23</b>	<b>1262</b>	<b>2145</b>	<b>20</b>	<b>9533</b>	<b>18804</b>	<b>115</b>

688

689

690 Regarding the identifications of the turtle and tortoise bone fragments (Tab. 11), only 33% of the  
 691 NISP and 52% of the WR have been attributed at least to a family. *Indotestudo elongata* bone remains  
 692 account for most of the identified turtle/tortoise bones with 86% of the NISP, 86% of the WR, and 87% of

693 the MNI. Geoemydidae are rare with only 14% of the NISP, 13% of the WR, and 9% of the MNI, while the  
694 occurrence of Trionychidae is anecdotic (less than 1% of the NR and WR, and 3.5% of the MNI).

695 **Table 11** - Number of Identified Skeletal Parts (NISP), Weight of the remains (WR), and Minimal  
696 Number of Individuals (MNI) identified in the turtle/tortoise bones assemblage from the different  
697 layers of Laang Spean Cave.


	Hoabinhian layer			Neolithic layer			Sepulture layers			Total		
	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI
<i>Indotestudo elongata</i>	2046	6706	51	204	649	12	260	692	12	2510	8047	75
Geoemydidae	325	955	4	34	115	2	47	162	2	406	1232	8
Trionychidae	7	20.5	1	2	7	1	5	16	1	14	43.5	3
Turtle ind.	4272	6212		664	1084		883	1187		5819	8483	0
<b>Total</b>	<b>6650</b>	<b>13894</b>	<b>56</b>	<b>904</b>	<b>1855</b>	<b>15</b>	<b>1195</b>	<b>2057</b>	<b>15</b>	<b>8749</b>	<b>17806</b>	<b>86</b>

698  
699

700 Taphonomy of the turtle/tortoise bone assemblage

701 Regarding the taphonomy of the turtle bones collected in the different squares, the mean  
702 completion rate is slightly lower in the “sepulture” assemblage (32%) than in the “Neolithic” and  
703 “Hoabinhian” assemblages (39.2 and 36.7%). The general fragmentation pattern is otherwise similar in all  
704 layers. The complete bones constitute between 9.3% of the assemblages for the Hoabinhian assemblage,  
705 and 7.2-6.2% for the “Neolithic” and “Sepulture” assemblages while nearly complete elements account  
706 for 14.2% of the “Hoabinhian”, 13.8% of the “Neolithic”, and 11% of the “Sepulture” assemblages. Most  
707 anatomical parts are represented in the different assemblages but the extremities, vertebrae, and skull  
708 remains are very rare with a PR below 4% in all assemblages (Fig. 7). There is also a global tendency to  
709 the lower representation of the peripheral plates of the bridge (11%-31%) in respect to the other  
710 peripheral plates (58-70%) although this trend is more strongly marked in the Hoabinhian layer (11% vs.  
711 70%). The peripheral plates of the bridge are also systematically more fragmented (59-67% of mean  
712 completion) than the others (83-87% of mean completion). Outside of these common trends, significant  
713 strong differences emerge between the “Hoabinhian” assemblages and the two other layers. Indeed,  
714 although the general PR is similar in the different assemblages (41% for the Hoabinhian assemblage, 37%  
715 for the Neolithic assemblage, and 34% for the Sepulture assemblage) the carapace elements are  
716 dramatically better represented in the Hoabinhian bones (Fig. 7-A) compare to the two other  
717 assemblages (Fig. 7-B, C) (Chi<sup>2</sup> test; p.value<0.01). Indeed, although the PR of the carapace and plastron  
718 bones are more or less similar in the Hoabinhian assemblage (57.8% vs. 67%) the plastron elements are  
719 mostly missing in the other assemblages (54% vs. 20.8% in the Neolithic assemblage and 53% vs. 15.1% in  
720 the Sepulture assemblage). The stylopods are also better represented in the Hoabinhian squares (84%)  
721 compare to the other assemblages (50.4% and 28%).





722  
723  
724  
725

**Figure 7** - Anatomical distributions of the turtle remains collected in the different layers of the site of Lang Spean. The percentage of representation (PR) is considered here to provide a graphical visualization of the different values observed for the different anatomical elements.

726 The observation of traces on the bones is made very challenging by the fact that 61% of them are covered  
727 by a veil of calcite. Interestingly, this calcite deposit was more frequent in the Hoabinhian squares where  
728 it covered 65% of the bones but was scarcer in the Neolithic and sepulture squares where it covered  
729 respectively 56% and 38% of the bones. This is probably related to the position of the remains in the  
730 cave, more or less close to the walls, which influenced their exposition to water flows during the rainy  
731 season. There is also a possibility that the calcite veil might be more frequent on the oldest remains. The  
732 presence of water flow in the site is also indicated by the occurrence of 61 bones having been polished by  
733 water flows. Porcupine gnawing traces were observed on only 13 elements distributed in several areas  
734 and layers of the site and digestion traces on only one. This clearly indicates a minor impact of animal  
735 species on the integrity of the archaeological assemblage. Putative burning traces were observed 10% of  
736 the remains. These traces were better represented in the Hoabinhian squares where they were present  
737 on 12% of the bones while they only occur on 2.8 and 3.8% of the bones recovered in the Neolithic and  
738 Sepulture squares. The characterization of burning traces was made very difficult by the fact that the  
739 material present a strong variability of surface color probably related to post-depositional chemical  
740 alteration. The occurrence of these traces might thus have been underestimated given the fact that we  
741 choose to record them only when their nature was undisputable. No cut mark was observed.  
742 Among the full assemblage 327 fragments of carapaces (3.7% of the NR) were still in anatomical  
743 connection at the moment of the excavation. These elements are mostly from the Hoabinhian squares  
744 (N=292) where they account for 4.3% of the turtle remains. Elements in anatomical connection are  
745 scarcer in the other assemblages with only 35 occurrences (1.7% of the turtle NR). This indicates that the  
746 Hoabinhian squares have been indeed less disturbed than the "Neolithic" and "Sepulture" squares.



747 Size of *Indotestudo elongata* individuals

748 The measurements recorded on the *I. elongata* archaeological material of Laang Spean Cave  
749 enabled for the reconstruction of 688 SCL estimations included between 68 and 345 mm and with a mean  
750 of 201 mm (Figure. 5-C) and corresponding to at least 75 individuals. Most of the data (N=564) are from  
751 Hoabinhian layers while the Neolithic squares only provided 124 SCL data but no significant difference  
752 emerged from the comparison of these two assemblages (Student t-test,  $p.val > 0.05$ ). The global  
753 distribution (all squares) of these sizes is unimodal (Hartigans' dip test,  $p.val > 0.05$ ) with a peak of  
754 specimens around 200 mm SCL. In this site, small specimens below 170 mm represent only 16% of the  
755 population, and specimens below 140 mm only 4.7%.

756 **Discussion**

757 **Taxonomic composition of the herpetofaunal assemblages**

758 In all the assemblages the distribution of the herpetofaunal groups in the four investigated sites  
759 present strong similarities. Non-marine turtles are nearly systematically the best represented  
760 herpetofaunal group (between 59 and 91% of the NISP) before Monitor lizards (between 6 and 25% of  
761 the NISP), snakes (below 3.5% of the NISP), and amphibians. The only exception to this trend is the layer  
762 1 of Moh Khiew cave in which Monitor Lizards (25% of the NISP), and snakes (23% of the NISP) bone  
763 remains are more numerous than turtle skeletal elements (20% of the NISP). This layer is disturbed and  
764 not dated so interpreting this observation is impossible for now. However, the other sites follow a clear  
765 pattern showing that hunter-gatherers groups have preferentially exploited turtles over other reptile and  
766 amphibian taxa. This follow the general regional pattern previously observed in similar zooarchaeological  
767 assemblages (Conrad 2015). Regarding the proportion of the turtle/tortoise families in the assemblages,  
768 Testudinidae (*Indotestudo elongata*) is always the most represented group as it account for between 52  
769 and 89% of the turtle bones NISPs. The proportions of Geoemydidae turtles vary a lot between 48 and  
770 11% of the same NISPs. As most of the species from this group are aquatic freshwater turtles this  
771 variability could be explained by the accessibility of streams, rivers, and lakes by the inhabitants of the  
772 sites. In most sites the proportion of Geoemydidae is around 30% of the NISP but these turtles are much  
773 less prevalent in Laang Spean cave, and the best represented in the TP1 of the site of Khao Ta Phlai. The  
774 published data regarding the faunal assemblage from Laang Spean indicate a weak contribution of  
775 freshwater taxa (mussels and fish) to the overall diet (Forestier *et al.* 2015; Frère *et al.* 2018) which is in  
776 accordance with our observation of the scarcity of freshwater turtles in the site. Regarding the  
777 prevalence of Geoemydidae species in the TP1 of Khao Ta Phlai, this could indicate a stronger reliance on  
778 freshwater resources than in the other sites. However, considering the chronology of the two test-pits of  
779 the site is not yet fully resolved and that general importance of aquatic resources in this assemblage still  
780 need to be estimated, this fact cannot be related to a cultural/chronological trend for now. From a  
781 general point of view, the data regarding the herpetofaunal assemblages, however, point to strong  
782 similarities between assemblages of different ages and from very different environmental settings. This  
783 should be considered at the light of the studies regarding the mammal bone assemblages of the same  
784 sites to test the hypothesis of a putative homogeneity of the Hoabinhian subsistence strategies in  
785 continental Southeast Asia.

786 **Taphonomy of the turtle assemblages**

787 The fragmentation rate of the bones is fairly homogenous among the sites. The average  
788 percentage of completeness of the bones is between 37 and 28%. The material from three first layers of  
789 Moh Khiew Cave and Laang Spean are the less fragmented (average percentage of completeness above  
790 33%). The layer 4 of Moh Khiew cave provided the most fragmented material (average percentage of  
791 completeness of 28%). The presence of large limestone blocks in this layer may be indicative of crumbling

792 that may have altered the faunal material. Regarding the anatomical distribution of the turtle remains  
793 the sites present strong differences with mean PR between 41% (Laang Spean) and 15% (Khao Ta Phlai  
794 metal ages layer from TP1). This means that the anatomical representation of the bone remains is more  
795 or less biased toward some elements. Two main cases occur in the assemblages: sites in which stylopds  
796 are the best represented parts (Khao Ta Phlai metal ages layer from TP1, Doi Pha Kan, layers 2 to 4 of  
797 Moh Khiew cave), and sites in which the most robust parts of the carapace are the best represented  
798 elements. The mean PR is systematically higher in the assemblages where the stylopds are the most  
799 numerous. This indicates that these assemblages are the least altered by post-depositional phenomena.  
800 Indeed, a natural alteration would rather lead to the situation observed in the other assemblages in  
801 which the elements that are the most often founds are the most robust and thus would have the highest  
802 survival rates. However this is not sufficient to explain an overrepresentation of long bones that are  
803 supposed to preserve less well than carapaces elements. Considering that all the sediment of the studied  
804 deposit has been screened, a major recovery bias is unlikely although some of the smallest elements  
805 might have been missed. A post-depositional sorting of the material could also be ruled out as we shown  
806 no evidence of differential fragmentation and no abundant trace of water circulation in the different  
807 deposits studied. The most likely hypothesis is thus that human inhabitants of some of the sites  
808 discarded or transported for further use some of the carapaces of the consumed animals and left on  
809 place the smallest elements among which the largest and toughest (humerus and femurs) have been  
810 recovered and identified. This behavior would in any case not be systematic as the anatomical  
811 distributions indicate that complete individuals have been brought on the sites. The absence of the head  
812 of the specimens could be either related to an identification bias or a removal of these parts outside of  
813 the site. Humerus and femur put apart, the anatomical distributions of turtle bones follow a global  
814 pattern where the most robust anatomical elements are better represented than the more fragile ones.  
815 The only exception to this trend concern the peripheral plates of the bridge which are always less  
816 represented than the other peripheral. This is very likely to be related to an identification bias itself  
817 related to the nearly complete absence of complete pieces of such element in the material. This is  
818 undoubtedly linked to the separation of the carapace from the plastron by the inhabitants of the sites  
819 who have broken the bones in the area that links both part of the shell, a step that is mandatory to access  
820 to most of the meat content of the animal.

821 The observations of the surface traces on the bones indicate a nearly complete lack of predation  
822 and digestion traces which combined with the general weak fragmentation of the material allow to  
823 completely ruling out a putative role of non-human predators in the constitution of the studied  
824 assemblages. This is not surprising as although some predators, including Monitor lizards, are known to  
825 hunt juvenile tortoise individuals, adults' specimens probably have few non-human predators although  
826 some modern specimen bear traces of predation attempts (Ihlow *et al.* 2016) and that predation on other  
827 Southeast Asian tortoises species have been reported (Platt *et al.* 2021). Large felids (Emmons 1989), and  
828 eagles (Gil-Sánchez *et al.* 2022) are known to be able to hunt adults tortoises but such predator would  
829 undoubtedly have left predation traces on the subfossil bone assemblages we studied. Some very rare  
830 bones bearing porcupine traces indicate that these animals had a minor impact on some of the  
831 assemblages but not enough to impact the zooarchaeological interpretations. However, although it  
832 seems fairly evident that the animals present in the sites have been hunted to be consumed as there is  
833 no trace of bone industry in the assemblages, finding direct traces of culinary preparation on the bones is  
834 very challenging. Indeed, in Khao Ta Phlai and Laang Spean, some remains (72 and 37%) were covered by  
835 a veil of calcite making impossible to observe the surface of the bones. In addition, very few cut marks  
836 have been characterized on the bones of the different sites. Many burned bones were observed in all the  
837 sites but linking these to a cooking technique is very questionable. Indeed, these traces do not seem to  
838 be located on specific part of the bones (i. e. external side of the carapace) and appear randomly on

839 every area of every anatomical part. It is likely that these traces are related to post-depositional events  
840 unrelated to the cooking of the animals. The frequency of fire traces combined with the strong  
841 fragmentation of the large vertebrate remains in most sites (C. Griggo; C. Bochaton pers. obs) could  
842 indicate the use of the bones as combustible (Villa *et al.* 2002). Such a use is unlikely for the turtle  
843 skeletons considering their small size and the very good preservation stage of their remains but a  
844 proximity to the fire places (Bennett 1999) could explain the random occurrence of fire traces on their  
845 bone elements.

#### 846 **Size of *Indotestudo elongata* archaeological specimens**

847 The size of *I. elongata* individuals observed in the four archaeological deposits (Fig. 5) shows  
848 common patterns but also some differences. The distributions of the estimated sizes are bimodal in all  
849 sites except Laang Spean. In all the sites, most of the estimations correspond to adult size specimens  
850 above 170 mm SCL reaching maximums of 270-345 mm SCL. These specimens all fall in the size range of  
851 the modern representatives of the species. Yet, all sites present a variable proportion of smaller, likely  
852 immature individuals. The representation of this second group is the lowest in Laang Spean (16% of the  
853 total number of estimations) but is important enough in the other sites to make their distributions  
854 bimodal with 35%-33% in Doi Pha Kan and Khao Ta Phlai, and 24% in Moh Khiew cave. Specimens below  
855 140mm SCL are rare in all sites as they account for less than 10% of the estimations in all sites but  
856 represent more than 15% of the Doi Pha Kan population.

857 Interpreting the size distribution of the archaeological tortoises is a difficult task as it first  
858 requires an idea of what the size structure of a wild population would look like and basic biological data  
859 (season of birth, activity pattern, growth speed...) regarding modern and past *I. elongata* populations.  
860 However, these data are currently mostly missing which make very challenging a detailed interpretation  
861 of the collected archaeological data. The recovery of size distribution data in a natural modern population  
862 is always challenging as it could be influenced by many factors (i. e. climate, environment, seasonality,  
863 behaviors, and sizes of the individuals) that could bias the observations by making one or several size  
864 classes more difficult to observe than the others. In addition, the history and specific conditions of a wild  
865 population itself could have a dramatic impact on its size structure. In these conditions, defining a  
866 modern comparison point to the documented archaeological populations is difficult. To our knowledge,  
867 the only data collected on *I. elongata* concern the population of the Ban Kok Village (Khon Kaen Province,  
868 Thailand). This study shows that the pre-adult individuals have a low survivability rate as their population  
869 was mostly composed of newly born and old adult individuals (Sriprateep *et al.* 2013). The authors  
870 suggest that this strongly biased structure could be related to an absence of predation on the large  
871 individuals, and partly to several phenomena having a stronger impact on small specimens (e. g.  
872 predation, trampling of domestic bovids) but admit the main cause is still unknown while not discussing a  
873 potential poaching of the smaller individuals. Similarly, another publication about *I. travancorica*  
874 indicates a lack of juvenile specimen in the population but highlight that it could be related to a seasonal  
875 activity specific pattern. Indeed, juvenile specimens were much more commonly found at the beginning  
876 of the rainy season than during the dry season when their study has been conducted (Ramesh 2008).  
877 Other published distributions from other tortoises' species also indicate a strong representation of adult  
878 size individuals of different ages having completed their growth but also a much more balanced  
879 distribution of juvenile specimens of all sizes (Hailey & Coulson 1999; Znari *et al.* 2005; Rouag *et al.* 2007).  
880 In all these distributions the juvenile specimens are way scarcer than adult ones which make sense as  
881 adult class specimens correspond to individuals of very different ages having reached their final size. The  
882 only case in which this situation would be reversed is a population in which adult individuals would be  
883 subject to a strong predation pressure superior to the pressures imposed on the smaller individuals.

884 The site of Laang Spean presents a unimodal size distribution in which juvenile specimens are mostly  
885 excluded. In that sense, this distribution is very different from that of a natural population and indicates a  
886 strong selection on adult specimens of moderate to large size. This is clearly indicative of a very selective  
887 hunting strategy that may have been enabled by the abundance of resources in the vicinity of the site.  
888 Such a selection, although visible in other deposits is less marked as juvenile specimens composed more  
889 important parts of the assemblages, especially in Doi Pha Kan. In these sites, it is impossible to estimate  
890 whether or not the proportions of juvenile specimens present in the assemblages are similar to those of  
891 the exploited natural populations and thus to estimate the exact intensity of the selection toward large  
892 size individuals. In any case, it is the sign of an opportunistic foraging as such a combination of juvenile  
893 specimens has been observed on modern hunter-gatherer population actively collecting tortoises this  
894 way (Mena *et al.* 2000). But this might also be influenced by the hunting method in the case a direct  
895 selection by the hunter is not made, for instance with the use of trapping that was also hypothesized in  
896 Doi Pha Kan for the hunting of monitor lizards (Bochaton *et al.* 2019). This technique is also the most  
897 used to hunt tortoise in the Amazon as it is the most efficient method before active searching (Santos *et*  
898 *al.* 2020). This implies no selection on the specimens in the wild although the type of trap used (i. e. size  
899 of the ground hole) might induce some size bias. The use of traps could thus explain the strong  
900 representation of smaller individuals present in the archaeological assemblages and indicate a very  
901 opportunistic strategy indicative of either a poor selection by the hunter and/or a relative scarcity of the  
902 tortoises in the environments making harder the collect of large individuals. The hunting season could  
903 also be an explanation to the stronger or weaker presence of juvenile specimens in the assemblages.  
904 Indeed, during the dry season tortoises are less active and harder to find which could led the hunter to be  
905 less selective especially in the case of a use of non-selective hunting methods allowing to find these  
906 animals. Theobald (1868) mentions the hunts of tortoises by Burmese hunters in the dry season by  
907 clearing grass lands and forests with fire in order to destroy their shelters and locate them. At the  
908 opposite, smaller tortoises are more active in the rainy season during which dogs are more used to track  
909 them (Blythe 1854; Theobald 1868). Ultimately, both seasonal hypotheses could explain the occurrence  
910 of small individual using different explanations (hunting method vs. activity season). Only the use of non-  
911 traditional approaches such as skeletochronology (Ehret 2007) could help to clarify this question by  
912 estimating the season of death of the tortoise individuals as well as the occupation seasonality of the  
913 different sites given the absence of other seasonality makers in the materials.

914 Tortoise populations are vulnerable to intensive exploitation focused on the larger mature  
915 individuals. As such, and their exploitation has been considered as a marker of small scale hunting and  
916 thus of small human groups (Stiner *et al.* 2000). In the studied sites, the strong focus on a single turtle  
917 species (*I. elongata*) and the focus placed on large individuals would undoubtedly damage the natural  
918 populations and lead to a size reduction of the individuals (Close & Seigel 1997). Such exploitation would  
919 be sustainable only if it was not intensive meaning that not many individuals were collected in order to  
920 feed a putatively small size human group. It is difficult to address the question of the overall importance  
921 of the tortoises in the diet of the studied Southeast Asian hunter-gatherer groups as we still lack a  
922 complete and quantified study of the mammal fauna of the sites as well as strong data regarding the use  
923 of the studied sites. It is, however, clear that the investigated prehistoric population have exploited  
924 tortoises which composed a significant part of their meat diet. This is not surprising as turtle species are  
925 supposed to represent an important biomass in the ecosystems (Iverson 1982) and are also fairly easy to  
926 collect. This behavior have subsisted until nowadays in continental Southeast Asia (Hansel 2004) although  
927 all population does not choose to exploit reptile species (Tungittiakornl & Dearden 2002).

928

## Conclusion and perspective

929           The present work has been built as a stepping stone to provide the basic data and study tools  
930 enabling the study of tortoise's assemblages from continental Southeast Asia. As an effect, the true  
931 potential of this work will only be reached through the use of its analytical protocol for future studies and  
932 comparison with additional assemblages. We were however able to reach several conclusions as we  
933 demonstrated a putative strong similarities between the exploitation of the herpetofaunal taxa in the  
934 different sites as well as in the taphonomy of the non-marine turtle assemblages in different  
935 chronological and environmental settings. These data thus open many interesting questions regarding  
936 the trends of hunter-gatherer subsistence strategies in continental Southeast Asia during the Pleistocene  
937 and through the Holocene. However, much work remains to be done to reach a satisfactory  
938 zooarchaeological documentation level regarding these prehistoric human groups. As we demonstrate it  
939 in the introduction of this paper, many of the previously excavated Hoabinhian archaeological deposits of  
940 continental Southeast Asia, including sites which are known to have provided rich assemblages of non-  
941 marine turtle bone remains (i. e. Lang Rongrien), have not benefited from quantified zooarchaeological  
942 analyses. The complete study of these sites will be important to provide additional relevant comparison  
943 points to the present study. The non-herpetofaunal taxa of the sites included in this study should also be  
944 investigated to estimate the relative part of the reptile and amphibian exploitation in the global diet of  
945 these hunter-gatherer populations. Such studies should however be carried out in combination with the  
946 elaboration of appropriate study protocols regarding the estimation of the size/weight of the exploited  
947 individuals of large mammal species. Much needed is also the elaboration of identification methods,  
948 should they be morphological or molecular, designed for Southeast Asian species to complement the  
949 existing works (Pritchard *et al.* 2009; Bochaton *et al.* 2019). Only at the cost of such investment the  
950 zooarchaeology of Southeast Asia will be on par with the rich literature existing on the material  
951 productions of prehistoric groups but this will also require the local development of a strong research  
952 community interested in that discipline that is still lacking at the present.

953  
954

### **Acknowledgments**

955           The authors are grateful to all the excavation teams who collected the studied remains as well to  
956 all the technical staff who helped us in the study of the different assemblages. We are especially grateful  
957 to Silpakorn University who host C. B. during most of the studies as well as to the Fine Art Departement  
958 of Thailand. We also thank S. Coleman and the Florida Museum of Natural History for welcoming us in  
959 their herpetology collections.

960

### **Data, scripts, code, and supplementary information availability**

961           Supplementary information is available online: XXXXDOI of the webpage hosting the data  
962 <https://doi.org/10.5802/fake3.doi>  
963

964

### **Conflict of interest disclosure**

965           The authors declare that they comply with the PCI rule of having no financial conflicts of interest  
966 in relation to the content of the article.

967

### **Funding**

968



969 This work has been done thanks to the support of several funding agencies: the FYSEN foundation, the  
970 DIM-MAP funded by the Region île-de-France, the French Ministry of Europe and Foreign Affairs, and the  
971 IRN PalBioDivASE 0846 funded by the CNRS. It was also supported by the Mission Préhistorique Franco-  
972 Cambodgienne and the Mission Paléolithique Franco-Thaïe of the French Ministry of Europe and Foreign  
973 Affairs (MEAE, Paris).  
974

## 975 **References**

- 976 ADI B.T.H. 2000. — Archaeology of Ulu Kelantan PhD thesis. Canberra, *Australian National University*.  
977 297 p.
- 978 AMPHANSRI A. 2011. — Temporal and Spatial Analysis of Faunal Remains from Tham Lod Rockshelter  
979 Site, Pang Mapha District, Mae Hong Son Province M. A. Thesis. Bangkok, Thailand, *Silpakorn*  
980 *University*.
- 981 ANDERSON D.D. 1990. — *Lang Rongrien Rockshelter: A Pleistocene, Early Holocene Archaeological Site*  
982 *from Krabi, Southwestern Thailand*. Philadelphia, University of Pennsylvania Press. 108 p.
- 983 AUETRAKULVIT P. 2004. — Faunes du pléistocène final à l'holocène de Thaïlande : approche  
984 archéozoologique PhD thesis. Marseille, *Aix Marseille 1*. 433 p.
- 985 AUETRAKULVIT P., FORESTIER H., KHAOKHIEW C. & ZEITOUN V. 2012. — New excavation at MohKhiew site  
986 (Southern Thailand), in BONATZ D., REINECKE A. & TJOA-BONATZ M.L. (eds.), *Crossing Borders in*  
987 *Southeast Asian Archaeology*. p. 62–74.
- 988 AUFFENBERG W. 1974. — Checklist of Fossil Land Tortoises (Testudinidae). *Bulletin of the Florida State*  
989 *Museum Biological Sciences* 18 (3): 122–247
- 990 AVERY G., KANDEL A.W., KLEIN R.G., CONARD N.J. & CRUZ-URIBE K. 2004. — Tortoises as food and  
991 taphonomic elements in palaeo « landscapes » in: XXIVe Rencontres internationales d'archéologie et  
992 d'histoire d'Antibes, Antibes, France. p. 147–161.
- 993 BENNETT J.L. 1999. — Thermal Alteration of Buried Bone. *Journal of Archaeological Science* 26 (1): 1–  
994 8. <https://doi.org/10.1006/jasc.1998.0283>
- 995 BITON R., SHARON G., ORON M., STEINER T. & RABINOVICH R. 2017. — Freshwater turtle or tortoise? The  
996 exploitation of testudines at the Mousterian site of Nahal Mahanayeem Outlet, Hula Valley, Israel.  
997 *Journal of Archaeological Science: Reports* 14: 409–419.  
998 <https://doi.org/10.1016/j.jasrep.2017.05.058>
- 999 BLASCO R. 2008. — Human consumption of tortoises at Level IV of Bolomor Cave (Valencia, Spain).  
1000 *Journal of Archaeological Science* 35 (10): 2839–2848. <https://doi.org/10.1016/j.jas.2008.05.013>
- 1001 BLASCO R., ROSELL J., SMITH K.T., MAUL L.C., SAÑUDO P., BARKAI R. & GOPHER A. 2016. — Tortoises as a  
1002 dietary supplement: A view from the Middle Pleistocene site of Qesem Cave, Israel. *Quaternary*  
1003 *Science Reviews* 133: 165–182. <https://doi.org/10.1016/j.quascirev.2015.12.006>
- 1004 BLYTHE E. 1854. — Notices and descriptions of various reptiles, new or little-known. Part I. *The journal*  
1005 *of the Asiatic Society of Bengal* 22 (7): 639–655
- 1006 BOCHATON C., HANOT P., FRÈRE S., CLAUDE J., NAKSRI W., AUETRAKULVIT P. & ZEITOUN V. 2019. — Size and  
1007 weight estimations of subfossil monitor lizards (*Varanus* sp. Merrem 1820) with an application to the  
1008 Hoabinhian assemblage of Doi Pha Kan (Late Pleistocene, Lampang province, Thailand). *Annales de*  
1009 *Paléontologie* 4 (105): 295–304. <https://doi.org/10.1016/j.annpal.2019.05.003>
- 1010 BOCHATON C., IVANOV M. & CLAUDE J. 2019. — Osteological criteria for the specific identification of  
1011 Monitor lizards (*Varanus* Merrem, 1820) remains in subfossil deposits of Sundaland and continental  
1012 Southeast Asia. *Amphibia-Reptilia* 40 (2): 219–232. <https://doi.org/10.1163/15685381-20181101>
- 1013 BULBECK F. 2003. — Hunter-Gatherer Occupation of the Malay Peninsula from the Ice Age to the Iron  
1014 Age, *Under the Canopy: The Archaeology of Tropical Rain Forests*. Brunswick, Rutgers University  
1015 Press. p. 119–160.

- 1016 CELIBERTI V., FORESTIER H., AUETRAKULVIT P. & ZEITOUN V. 2018. — Doi Pha Kan, Lampang Province, North  
1017 of Thailand: A New Lithic Assemblage in the Hoabinhian Chrono-Cultural Context, *in* TAN N. (ed.),  
1018 *Advancing Southeast Asian Archaeology*. p. 91–99, 316–319.
- 1019 CHUNG T.N. 2008. — Research on Hoa Binh culture in Viet Nam, Laos and Cambodia. *Vietnam*  
1020 *Archaeology* 3: 19–32
- 1021 CLAUDE J., AUETRAKULVIT P., NAKSRI W., BOCHATON C., ZEITOUN V. & TONG H. 2019. — The recent fossil  
1022 turtle record of the central plain of Thailand reveals local extinctions. *Annales de Paléontologie* 4  
1023 (105): 305–315. <https://doi.org/10.1016/j.annpal.2019.04.005>
- 1024 CLOSE L.M. & SEIGEL R. 1997. — Differences in body size among populations of red-eared sliders  
1025 (*Trachemys scripta elegans*) subjected to different levels of harvesting. *Chelonian Conservation and*  
1026 *Biology* 2 (4): 563–566
- 1027 CODRON D., HOLT S., WILSON B. & HORWITZ L.K. 2022. — Skeletal allometries in the leopard tortoise  
1028 (*Stigmochelys pardalis*): Predicting chelonian body size and mass distributions in archaeozoological  
1029 assemblages, Quaternary Environments and Archaeology of the Northern Cape (South Africa).  
1030 *Quaternary International* 614: 59–72. <https://doi.org/10.1016/j.quaint.2021.07.021>
- 1031 COLLECTIF 1932. — *Premier congrès des préhistoriens d'Extrême-Orient*. Hanoi, Imprimerie d'Extrême-  
1032 Orient. 154 p.
- 1033 CONRAD C. 2015. — Archaeozoology in Mainland Southeast Asia: Changing Methodology and  
1034 Pleistocene to Holocene Forager Subsistence Patterns in Thailand and Peninsular Malaysia. *Open*  
1035 *Quaternary* 1 (1). <https://doi.org/10.5334/oq.af>
- 1036 CONRAD C., HIGHAM C., EDA M. & MARWICK B. 2016. — Palaeoecology and Forager Subsistence  
1037 Strategies during the Pleistocene – Holocene Transition: A Reinvestigation of the Zooarchaeological  
1038 Assemblage from Spirit Cave, Mae Hong Son Province, Thailand. *Asian Perspectives* 55 (1): 2–27.  
1039 <https://doi.org/10.1353/asi.2016.0013>
- 1040 DAS I. 2010. — *A field guide to the reptiles of South-East Asia*. London, New Delhi, New York, Sydney,  
1041 Bloomsbury Publishing. 376 p.
- 1042 DODSON P. & WEXLAR D. 1979. — Taphonomic Investigations of Owl Pellets. *Paleobiology* 5 (3): 275–  
1043 284
- 1044 DUNN F.L. 1964. — Excavations at Gua Kechil, Pahang. *Journal of the Malaysian Branch of the Royal*  
1045 *Asiatic Society* 37 (2 (206)): 87–124
- 1046 EBERLING G. 2011. — Haltung und Nachzucht von *Indotestudo elongata* (Blythe, 1853). *Draco* 2: 61–72
- 1047 EHRET D. 2007. — Skeletochronology: A Method for Determining the Individual Age and Growth of  
1048 Modern and Fossil Tortoises (Reptilia: Testudines). *Bulletin of the Florida Museum of Natural History*  
1049 47 (2): 49–72
- 1050 EMMONS L.H. 1989. — Jaguar Predation on Chelonians. *Journal of Herpetology* 23 (3): 311–314.  
1051 <https://doi.org/10.2307/1564460>
- 1052 ESKER D.A., FORMAN S.L. & BUTLER D.K. 2019. — Reconstructing the mass and thermal ecology of North  
1053 American Pleistocene tortoises. *Paleobiology* 45 (2): 363–377. <https://doi.org/10.1017/pab.2019.6>
- 1054 FORESTIER H. 2003. — Des outils nés de la forêt : l'importance du végétal en Asie du Sud-Est dans  
1055 l'imagination et l'invention technique aux périodes préhistoriques, *in* FROMENT A., GUFFROY J., &  
1056 PEUPELEMENTS ANCIENS ET ACTUELS DES FORÊTS TROPICALES : SÉMINAIRE-ATELIER, ORLÉANS (FRA), 1998/10/15-  
1057 16 (eds.), *Peuplements anciens et actuels des forêts tropicales : actes du séminaire-atelier*. Paris, IRD.  
1058 p. 315–337. (Colloques et Séminaires).
- 1059 FORESTIER H., GRIGGO C. & SOPHADY H. 2023. — L'Hoabinhien ou le paradigme égaré de la modernité  
1060 européenne en Extrême-Orient : l'exemple de la Préhistoire du Cambodge *In*: 29e Congrès  
1061 préhistorique de France, Toulouse. Société préhistorique française. p. 59–76.
- 1062 FORESTIER H., SOPHADY H. & CELIBERTI V. 2017. — Le techno-complexe hoabinhien en Asie du Sud-est  
1063 continentale : L'histoire d'un galet qui cache la forêt. *Journal of Lithic Studies* 4 (2): 305–349.  
1064 <https://doi.org/10.2218/jls.v4i2.2545>
- 1065 FORESTIER H., SOPHADY H., PUAUD S., CELIBERTI V., FRÈRE S., ZEITOUN V., MOURER-CHAUVIRÉ C., MOURER R.,  
1066 THAN H. & BILLAULT L. 2015. — The Hoabinhian from Laang Spean Cave in its stratigraphic,

- 1067 chronological, typo-technological and environmental context (Cambodia, Battambang province).  
1068 *Journal of Archaeological Science: Reports* 3: 194–206. <https://doi.org/10.1016/j.jasrep.2015.06.008>  
1069 FORESTIER H., ZEITOUN V., WINAYALAI C. & MÉTAIS C. 2013. — The open-air site of Huai Hin  
1070 (Northwestern Thailand): Chronological perspectives for the Hoabinhian. *Comptes Rendus Palevol* 12  
1071 (1): 45–55. <https://doi.org/10.1016/j.crvp.2012.09.003>  
1072 FORESTIER H., ZHOU Y., AUETRAKULVIT P., KHAOKHIEW C., LI Y., JI X. & ZEITOUN V. 2021. — Hoabinhian  
1073 variability in Mainland Southeast Asia revisited: The lithic assemblage of Moh Khiew Cave,  
1074 Southwestern Thailand. *Archaeological Research in Asia* 25: 100236.  
1075 <https://doi.org/10.1016/j.ara.2020.100236>  
1076 FORESTIER H., ZHOU Y., VIALLET C., AUETRAKULVIT P., LI Y. & SOPHADY H. 2022. — Reduction Sequences  
1077 During the Hoabinhian Technocomplex in Cambodia and Thailand: A New Knapping Strategy in  
1078 Southeast Asia from the Terminal Upper Pleistocene to mid Holocene. *Lithic Technology* 47 (2): 147–  
1079 170. <https://doi.org/10.1080/01977261.2021.1981654>  
1080 FRÈRE S., AUETRAKULVIT P., ZEITOUN V., SOPHADY H. & FORESTIER H. 2018. — Doi Pha Kan (Thailand), Ban  
1081 Tha Si (Thailand) and Laang Spean (Cambodia) Late Paleolithic animal bone assemblages. A new  
1082 perception of meat supply strategies for Early Holocene Mainland South-east Asia?, in TAN N.H. (ed.),  
1083 *Advancing Southeast Asian Archaeology*. p. 100–108, 320–323.  
1084 GARBIN R.C., ASCARRUNZ E. & JOYCE W.G. 2018. — Polymorphic characters in the reconstruction of the  
1085 phylogeny of geoemydid turtles. *Zoological Journal of the Linnean Society* 184 (3): 896–918.  
1086 <https://doi.org/10.1093/zoolinnea/zlx106>  
1087 GIL-SÁNCHEZ J.M., RODRÍGUEZ-CARO R.C., MOLEÓN M., MARTÍNEZ-PASTOR M.C., LEÓN-ORTEGA M., EGUÍA S.,  
1088 GRACIÁ E., BOTELLA F., SÁNCHEZ-ZAPATA J.A., MARTÍNEZ-FERNÁNDEZ J., ESTEVE-SELMA M.A. & GIMÉNEZ A.  
1089 2022. — Predation impact on threatened spur-thighed tortoises by golden eagles when main prey is  
1090 scarce. *Scientific Reports* 12 (1): 17843. <https://doi.org/10.1038/s41598-022-22288-9>  
1091 GLOVER I.C. 1977. — The Hoabinhian hunter-gatherers or early agriculture list in Southeast Asia, in  
1092 MEGAW J.V.S. (ed.), *Hunter Gatherers and First Farmer beyond Europe*. Leicester, UK, Leicester  
1093 University Press. p. 145–166.  
1094 GORMAN C. 1971. — The Hoabinhian and After: Subsistence Patterns in Southeast Asia during the  
1095 Late Pleistocene and Early Recent Periods. *World Archaeology* 2 (3): 300–320  
1096 GORMAN C.F. 1970. — Excavations at Spirit Cave, North Thailand: SOME INTERIM INTERPRETATIONS.  
1097 *Asian Perspectives* 13: 79–107  
1098 GORMAN C.F. 1969. — Hoabinhian: A Pebble-Tool Complex with Early Plant Associations in Southeast  
1099 Asia. *Science* 163 (3868): 671–673. <https://doi.org/10.1126/science.163.3868.671>  
1100 GOULD S.J. 1966. — Allometry and size in ontogeny and phylogeny. *Biological Reviews* 41 (4): 587–  
1101 640  
1102 GUERIN C. & MOURER-CHAUVIRÉ C. 1969. — Le Rhinoceros sondaicus Desmarest du gisement  
1103 néolithique de Loang Spean, province de Battambang, Cambodge. *Ann. Fac. Sc. Phnom Penh, 1970* 2:  
1104 261–274  
1105 HAILEY A. & COULSON I. 1999. — The growth pattern of the African tortoise *Geochelone pardalis* and  
1106 other chelonians. *Canadian Journal of Zoology* 77: 181–193. <https://doi.org/10.1139/cjz-77-2-181>  
1107 HANSEL T. 2004. — Observations on subsistence hunting along the Phu Yai Mountain Range,  
1108 Xanakham District, Vientiane Province, Lao PDR. *Natural History Bulletin of the Siam Society* 52: 195–  
1109 200  
1110 HIGHAM C. 2002. — *Early Cultures of Mainland Southeast Asia*. Art Media Resources. 380 p.  
1111 HIGHAM C.F.W. 1977. — Economic Change in Prehistoric Thailand, in REED C. (ed.), *Origins of*  
1112 *Agriculture*. The Hague, De Gruyter Mouton. p. 385–412.  
1113 <https://doi.org/10.1515/9783110813487.385>  
1114 HUXLEY J.S. 1932. — *Problems of relative growth*. London, Methuen & co. 319 p.  
1115 IHLOW F., DAWSON J.E., HARTMANN T. & SOM S. 2016. — *Indotestudo elongata* (Blyth 1854) – Elongated  
1116 Tortoise, Yellow-headed Tortoise, Yellow Tortoise. *Chelonian Conservation and Biology* 5: 1–14.  
1117 <https://doi.org/10.3854/crm.5.096.elongata.v1.2016>

- 1118 IHLOW F. & HANDSCHUH M. 2011. — Auswilderung von *Indotestudo elongata* im Kulen Promtep  
1119 Wildlife Sanctuary im Norden Kambodschas. *Marginata* 8 (4): 16–23
- 1120 IMDIRAKPHOL S., ZAZZO A., AUETRAKULVIT P., TIAMTINKRIT C., PIERRET A., FORESTIER H. & ZEITOUN V. 2017. —  
1121 The perforated stones of the Doi Pha Kan burials (Northern Thailand): A Mesolithic singularity?  
1122 *Comptes Rendus Palevol* 16 (3): 351–361. <https://doi.org/10.1016/j.crpv.2016.12.003>
- 1123 IVERSON J.B., SPINKS P.Q., SHAFFER B.H., MCCORD W.P. & DAS I. 2001. — Phylogenetic relationships  
1124 among the asian tortoises of the genus *Indotestudo* (Reptilia: Testudines: testudinidae). *Hamadryad*  
1125 26 (2): 283–286
- 1126 IVERSON J.B. 1982. — Biomass in turtle populations: A neglected subject. *Oecologia* 55 (1): 69–76.  
1127 <https://doi.org/10.1007/BF00386720>
- 1128 JI X., KUMAN K., CLARKE R.J., FORESTIER H., LI Y., MA J., QIU K., LI H. & WU Y. 2016. — The oldest  
1129 Hoabinhian technocomplex in Asia (43.5 ka) at Xiaodong rockshelter, Yunnan Province, southwest  
1130 China, Peking Man and related studies. *Quaternary International* 400: 166–174.  
1131 <https://doi.org/10.1016/j.quaint.2015.09.080>
- 1132 KLEIN R.G. & CRUZ-URIBE K. 1983. — Stone Age Population Numbers and Average Tortoise Size at  
1133 Byneskranskop Cave 1 and DieKelders Cave 1, Southern Cape Province, South Africa. *The South*  
1134 *African Archaeological Bulletin* 38 (137): 26–30
- 1135 MASOJC M., LE H.D., GRALAK T., MICHAŁEC G., APOLINARSKA K., BADURA M., CENDROWSKA M., GALAS A.,  
1136 KRUPA-KURZYŃOWSKA J., MIAZGA B., OSPINSKA M., ROZOK Z. & VIET N. 2023. — The Early Holocene  
1137 Hoabinhian (8300-8000 cal BC) occupation from Hiem Cave, Vietnam. *Comptes Rendus Palevol* 22  
1138 (5): 59–76
- 1139 MEDWAY, LORD 1969. — Excavations At Gua Kechil, Pahang III. Animal Remains. *Journal of the*  
1140 *Malaysian Branch of the Royal Asiatic Society* 42 (2 (216)): 197–205
- 1141 MENA P., STALLINGS J.R., REGALADO J. & CUEVA R. 2000. — The Sustainability of Current Hunting Practices  
1142 by the Huaorani. In *Hunting for Sustainability in Tropical Forests*: 57–78
- 1143 MOSER J. 2001. — *Hoabinhian: Geographie und Chronologie eines steinzeitlichen Technokomplexes in*  
1144 *Südostasien*. Germany, Linden Soft. 194 p. (AVA - Forschungen; 6).
- 1145 MOURER-CHAUVIRÉ C. & MOURER R. 1970. — The Prehistoric Industry of Laang Spean, Province of  
1146 Battambang, Cambodia. *Archaeology and Physical Anthropology in Oceania* 5: 128–146.  
1147 <https://doi.org/10.1002/j.1834-4453.1970.tb00110.x>
- 1148 MOURER-CHAUVIRÉ C., MOURER R. & THOMMERET Y. 1970. — Premières datations absolues de l’habitat  
1149 préhistorique de la grotte de Laang Spean, province of Battambang (Cambodge). *C. R. Acad. Sci.*  
1150 *Paris, Ser. D* 270: 471–473
- 1151 MUDAR K. & ANDERSON D.D. 2007. — New Evidence for Southeast Asian Pleistocene Foraging  
1152 Economies: Faunal Remains from the Early Levels of Lang Rongrien Rockshelter, Krabi, Thailand.  
1153 *Asian Perspectives* 46 (2): 298–334
- 1154 NABAIS M., BONETA I. & SOARES R. 2019. — Chelonian use in Portugal: Evidence from Castelo Velho de  
1155 Safara. *Journal of Archaeological Science: Reports* 28: 102054.  
1156 <https://doi.org/10.1016/j.jasrep.2019.102054>
- 1157 NABAIS M. & ZILHÃO J. 2019. — The consumption of tortoise among Last Interglacial Iberian  
1158 Neanderthals, Neanderthals: Ecology and Evolution. *Quaternary Science Reviews* 217: 225–246.  
1159 <https://doi.org/10.1016/j.quascirev.2019.03.024>
- 1160 NAKSRI W. 2013. — Origin and changes in continental turtle diversity from the Miocene to Holocene  
1161 in Thailand PhD thesis. Maha Sarakham, Thailand, *Mahasarakham University*.
- 1162 NAKSRI W. 2007. — Appraisal of the evolution of testudinoid turtle diversity from the Oligocene and  
1163 Neogene of Thailand Master thesis. Maha Sarakham, Thailand, *Mahasarakham University*. 230 p.
- 1164 NAKSRI W., TONG H., LAUPRASERT K., SUTEETHORN V. & CLAUDE J. 2013. — A new species of *Cuora*  
1165 (Testudines: Geoemydidae) from the Miocene of Thailand and its evolutionary significance.  
1166 *Geological Magazine* 150 (5): 908–922. <https://doi.org/10.1017/S0016756812001082>
- 1167 PLATT S.G., AUNG S.H.N., SOE M.M., LWIN T., PLATT K., WALDE A.D. & RAINWATER T.R. 2021. — Predation  
1168 on Translocated Burmese Star Tortoise (*Geochelone platynota*) by Asiatic Jackals (*Canis aureus*) and



- 1169 Wild Pigs (*Sus scrofa*) at a Wildlife Sanctuary in Myanmar. *Chelonian Conservation and Biology* 20 (1):  
1170 133–138. <https://doi.org/10.2744/CCB-1461.1>
- 1171 POOKAJORN S. 2001. — New perspectives for Palaeolithic research in Thailand, *Origine des*  
1172 *peuplements et chronologie des cultures paléolithiques dans le sud-est asiatique*. Paris, France,  
1173 Semenanjung. p. 167–188.
- 1174 PRITCHARD P.C.H., RABETT R.J. & PIPER P.J. 2009. — Distinguishing species of Geoemydid and Trionychid  
1175 turtles from shell fragments: evidence from the Pleistocene at Niah Caves, Sarawak. *International*  
1176 *Journal of Osteoarchaeology* 19 (4): 531–550. <https://doi.org/10.1002/oa.1038>
- 1177 R CORE TEAM 2020. — R: A language and environment for statistical computing
- 1178 RABETT R.J. 2012. — Tropical Subsistence Strategies at the End of the Last Glacial, *Human Adaptation*  
1179 *in the Asian Palaeolithic: Hominin Dispersal and Behaviour during the Late Quaternary*. Cambridge,  
1180 UK, Cambridge University Press. p. 208–265.
- 1181 RAMESH M. 2008. — Relative Abundance and Morphometrics of the Travancore Tortoise, *Indotestudo*  
1182 *travancorica*, in the Indira Gandhi Wildlife Sanctuary, Southern Western Ghats, India. *Chelonian*  
1183 *Conservation and Biology* 7 (1): 108–113
- 1184 RHODIN A.G.J. 1992. — Chelonian Zooarchaeology of Eastern New England: Turtle bone remains from  
1185 Cedar Swamp and other prehistoric sites. *Bulletin of the Massachusetts Archaeological Society* 53  
1186 (1): 21–30
- 1187 RHODIN A.G.J., IVERSON J., BOUR R., FRITZ U., GEORGES A., SHAFFER B.H. & VAN DIJK P.P. 2021. — *Turtles of*  
1188 *the World: Annotated Checklist and Atlas of Taxonomy, Synonymy, Distribution, and Conservation*  
1189 *Status (9th Ed.)*. United States of America, Chelonian Research Foundation and Turtle Conservancy.  
1190 482 p. (Chelonian Research Monograph; 8).
- 1191 ROUAG R., BENYACOUB S., LUISELLI L., EL MOUDEN E.H., TIAR G. & FERRAH C. 2007. — Population structure  
1192 and demography of an Algerian population of the Moorish tortoise, *Testudo graeca*. *Animal Biology*  
1193 57: 267–279. <https://doi.org/10.1163/157075607781753065>
- 1194 SANTOS A., MAYOR P., LOUREIRO L., GILMORE M., PEREZ PEÑA P., BOWLER M., LEMOS L., SVENSSON M., NEKARIS  
1195 K.A., NIJMAN V., VALSECCHI J. & MORCATTY T. 2020. — Widespread Use of Traditional Techniques by  
1196 Local People for Hunting the Yellow-Footed Tortoise (*Chelonoidis denticulatus*) Across the Amazon.  
1197 *Journal of Ethnobiology* 40: 268. <https://doi.org/10.2993/0278-0771-40.2.268>
- 1198 SHOOCONGDEJ R. 1996. — Forager mobility organization in seasonal tropical environments: A view  
1199 from Lang Kamnan Cave, western Thailand. Thesis.
- 1200 SOM S. & COTTET M. 2016. — Rescue and relocation programme of turtles and tortoises and  
1201 elongated tortoise monitoring programme in the Nam Theun 2 Reservoir (Laos). *Hydroécologie*  
1202 *Appliquée* 19: 383–406. <https://doi.org/10.1051/hydro/2015007>
- 1203 SOPHADY H. 2016. — Archeo-stratigraphy of Laang Spean prehistoric site (Battambang province): a  
1204 contribution to Cambodian prehistory PhD thesis. Paris, France, *Muséum national d'Histoire*  
1205 *naturelle*. 474 p.
- 1206 SOPHADY H., FORESTIER H., ZEITOUN V., PUAUD S., FRÈRE S., CELIBERTI V., WESTAWAY K., MOURER R., MOURER-  
1207 CHAUVIRÉ C., THAN H., BILLAULT L. & TECH S. 2016. — Laang Spean cave (Battambang province): A tale of  
1208 occupation in Cambodia from the Late Upper Pleistocene to Holocene, Southeast Asia: human  
1209 evolution, dispersals and adaptation. *Quaternary International* 416: 162–176.  
1210 <https://doi.org/10.1016/j.quaint.2015.07.049>
- 1211 SPETH J.D. & TCHERNOV E. 2002. — Middle Paleolithic Tortoise Use at Kebara Cave (Israel). *Journal of*  
1212 *Archaeological Science* 29 (5): 471–483. <https://doi.org/10.1006/jasc.2001.0740>
- 1213 SRIPRATEEP K., ARANYAVALAI V., AOWPHOL A. & THIRAKHUPT K. 2013. — Population Structure and  
1214 Reproduction of the Elongated Tortoise *Indotestudo elongata* (Blyth, 1853) at Ban Kok Village,  
1215 Northeastern Thailand. *Tropical Natural History* 13 (1): 21–37
- 1216 STINER M.C., MUNRO N.D. & SUROVELL T.A. 2000. — The Tortoise and the Hare: Small-Game Use, the  
1217 Broad-Spectrum Revolution, and Paleolithic Demography. *Current Anthropology* 41 (1): 39–79.  
1218 <https://doi.org/10.1086/300102>




- 1219 TAYLOR E.H. 1970. — The turtles and crocodiles of Thailand and adjacent waters; with a synoptic  
1220 herpetological bibliography. *The University of Kansas Science Bulletin* 49 (3): 87–179  
1221 THEOBALD W. 1868. — Catalogue of the Reptiles of British Birma, embracing the Provinces of Pegu,  
1222 Martaban, and Tenasserim; with descriptions of new or little-known species. *Journal of the Linnean*  
1223 *Society of London, Zoology* 10 (41): 4–67. <https://doi.org/10.1111/j.1096-3642.1868.tb02007.x>  
1224 THOMPSON J.C. & HENSHILWOOD C.S. 2014. — Tortoise taphonomy and tortoise butchery patterns at  
1225 Blombos Cave, South Africa. *Journal of Archaeological Science* 41: 214–229.  
1226 <https://doi.org/10.1016/j.jas.2013.08.017>  
1227 TREERAYAPIWAT C. 2005. — Patterns of Habitation and Burial Activity in the Ban Rai Rock Shelter,  
1228 Northwestern Thailand. *Asian Perspectives* 44 (1): 231–245. <https://doi.org/10.1353/asi.2005.0001>  
1229 TUNGITPLAKORN W. & DEARDEN P. 2002. — Hunting and wildlife use in some Hmong communities in  
1230 northern Thailand. *Natural History Bulletin of the Siam Society* 50: 57–73  
1231 VAN DIJK P.P. 1998. — The Natural History of the Elongated Tortoise *Indotestudo Elongata* (Blyth,  
1232 1853) (Reptilia: Testudines) in a Hill Forest Mosaic in Western Thailand with Notes on Sympatric  
1233 Turtle Species Ph.D. dissertation. Galway, Ireland, *National University of Ireland*. 255 p.  
1234 VAN VLACK H. 2014. — Forager Subsistence Regimes in the Thai-Malay Peninsula: An Environmental  
1235 Archaeological Case Study of Khao Toh Chong Rockshelter, Krabi Master thesis. 4484. San Jose, *San*  
1236 *José State University*. 188 p.  
1237 VILLA P., BON F. & CASTEL J.-C. 2002. — Fuel, fire and fireplaces in the Palaeolithic of western Europe.  
1238 *Review of Archaeology* 23: 33–42  
1239 VU T.L. 1994. — The Hoabinhian animal complex. *Vietnam Soc. Sci.* 5 (43): 70–74  
1240 WHITE J.C. 2011. — Emergence of cultural diversity in mainland Southeast Asia: A view from  
1241 prehistory, in ENFIELD N.J. (ed.), *Dynamics of Human Diversity: The Case of Mainland Southeast Asia*.  
1242 Canberra, Australian National University. p. 9–46.  
1243 YEN E. 1977. — Hoabinhian horticulture? The evidence and the questions from Northwest Thailand,  
1244 in ALLEN J.G. & JONES R. (eds.), *Sunda and Sahul. Prehistoric Studies in Southeast Asia, Melanesia and*  
1245 *Australia*. London, Academic Press. p. 567–599.  
1246 ZEITOUN V., AUETRAKULVIT P., ZAZZO A., PIERRET A., FRÈRE S. & FORESTIER H. 2019. — Discovery of an  
1247 outstanding Hoabinhian site from the Late Pleistocene at Doi Pha Kan (Lampang province, northern  
1248 Thailand). *Archaeological Research in Asia* 18: 1–16. <https://doi.org/10.1016/j.ara.2019.01.002>  
1249 ZEITOUN V., FORESTIER H., SOPHADY H., PUAUD S. & BILLAULT L. 2012. — Direct dating of a Neolithic burial  
1250 in the Laang Spean cave (Battambang Province, Cambodia): First regional chrono-cultural  
1251 implications. *Comptes Rendus Palevol* 11 (7): 529–537. <https://doi.org/10.1016/j.crpv.2012.06.006>  
1252 ZEITOUN V., FORESTIER H. & SUPAPORN N. 2008. — *Préhistoires au sud du Triangle d'Or* 252 p.  
1253 ZEITOUN V., HENG S. & FORESTIER H. 2021. — The funeral cave of Laang Spean In: SEAMEO SPAFA  
1254 International Conference on Southeast Asian Archaeology and Fine Arts, 13-17 December 2021,  
1255 Bangkok, Thailand. SEAMEO SPAFA. p. 1–14. <https://doi.org/10.26721/spafa.pqcnu8815a-14>  
1256 ZNARI M., GERMANO D.J. & MACÉ J.-C. 2005. — Growth and population structure of the Moorish  
1257 Tortoise (*Testudo graeca graeca*) in Westcentral Morocco: possible effects of over-collecting for the  
1258 tourist trade. *Journal of Arid Environments* 62 (1): 55–74.  
1259 <https://doi.org/10.1016/j.jaridenv.2004.11.013>  
1260 ZURAINA M. 1994. — *The Excavation of Gua Gunung Runtuh and the Discovery of the Perak Man in*  
1261 *Malaysia*. Kuala Lumpur, Department of Museums and Antiquity Malaysia. 174 p.  
1262

1263







## Appendix

1264 Appendix 1: All measurements recorded on the modern turtle skeletons.




1265











Nuchal plate  
(dorsal view)

GL



Peripheral plate  
(lateral view)


GH



Pygal plate  
(posterior view)

GW


GH



Neural plate  
(dorsal view)

GW


GL



Entoplastron  
(ventral view)


GW

GL




Supra-pygale plate 2  
(posterior view)

GH



Epiplastron  
(ventral view)

GL




Hyoplastron  
(ventral view)

GmL


GW

GmL



Hypoplastron  
(ventral view)

GW




Xiphiplastron  
(ventral view)

GW


GL

GpdvW



Humerus  
(Posterior view)


GpapW



(Dorsal view)

GL

GplmW



Radius

GL

GplmW



Ulna


GL

GpapW

GdapW


GdapW

GpdvW



Femur  
(Posterior view)

GpapW




GdapW

(Dorsal view)

GL

GplmW




GdlmW

Tibia

(Posterior view)


GpapW



GdapW

(Lateral view)

GL



GdlmW

Fibula

(Lateral view)




GdapW






(Posterior view)

GL




























### Kao Tah Plai Test Pit 1 - Metal Ages



### Kao Tah Plai Test Pit1 - Neolithic



### Kao Tah Plai Test Pit 2 - Neolithic

