



HAL
open science

Chronic exposure to tebuconazole alters thyroid hormones and plumage quality in house sparrows (*Passer domesticus*)

Pauline Bellot, François Brischoux, Hélène Budzinski, Sophie Dupont, Clémentine Fritsch, Sydney Hope, Bruno Michaud, Marie Pallud, Charline Parenteau, Louise Prouteau, et al.

► To cite this version:

Pauline Bellot, François Brischoux, Hélène Budzinski, Sophie Dupont, Clémentine Fritsch, et al.. Chronic exposure to tebuconazole alters thyroid hormones and plumage quality in house sparrows (*Passer domesticus*). *Environmental Science and Pollution Research*, 2023, 10.1007/s11356-023-28259-5 . hal-04145278

HAL Id: hal-04145278

<https://hal.science/hal-04145278>

Submitted on 23 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Chronic exposure to tebuconazole alters thyroid hormones and plumage quality in house sparrows (*Passer domesticus*)

Pauline Bellot ^{1,*}, François Brischoux ¹, H  l  ne Budzinski ², Sophie M Dupont.^{3,4}, Cl  mentine Fritsch ⁵, Sydney F Hope.¹, Bruno Michaud ¹, Marie Pallud ¹, Charline Parenteau ¹, Louise Prouteau ², Steffi Rocchi ⁵, Fr  d  ric Angelier ¹

1 Centre d'Etudes Biologiques de Chiz  , CNRS-La Rochelle Universit  , UMR 7372, 79360 Villiers en Bois, France

2 University of Bordeaux, CNRS-EPOC, UMR 5805, LPTC Research Group, 33400 Talence, France

3 BOREA, MNHN, CNRS 8067, SU, IRD 207, UCN, UA, 97233 Schoelcher Martinique, France

4 LIENSs, UMR 7266 CNRS-La Rochelle Universit  , 2 Rue Olympe de Gouges, 17000, La Rochelle, France

5 Laboratoire Chrono-Environnement, UMR 6249 CNRS / Universit   de Franche-Comt  , F-25000, Besan  on, France

Abstract

Triazoles belong to a family of fungicides that are ubiquitous in agroecosystems due to their widespread use in crops. Despite their efficiency in controlling fungal diseases, triazoles are also suspected to affect non-target vertebrate species through the disruption of key physiological mechanisms. Most studies so far have focused on aquatic animal models, and the potential impact of triazoles on terrestrial vertebrates has been overlooked despite their relevance as sentinel species of contaminated agroecosystems. Here, we examined the impact of tebuconazole on the thyroid endocrine axis, associated phenotypic traits (plumage quality and body condition) and sperm quality in wild-caught house sparrows (*Passer domesticus*). We experimentally exposed house sparrows to realistic concentrations of tebuconazole under controlled conditions and tested the impact of this exposure on the levels of thyroid hormones (T3 and T4), feather quality (size and density), body condition, and sperm morphology. We found that exposure to tebuconazole caused a significant decrease in T4 levels, suggesting that this azole affects the thyroid endocrine axis, although T3 levels did not differ between control and exposed sparrows. Importantly, we also found that exposed females had an altered plumage structure (larger but less dense feathers) relative to control females. The impact of tebuconazole on body condition was dependent on the duration of exposure and the sex of individuals. Finally, we did not show any effect of exposure to tebuconazole on sperm morphology. Our study demonstrates for the first time that exposure to tebuconazole can alter the thyroid axis of wild birds, impact their plumage quality, and potentially affect their body condition. Further endocrine and transcriptomic studies are now needed to understand not only the underlying mechanistic effects of tebuconazole on these variables, but also to further investigate their ultimate consequences on performance (*i.e.*, reproduction and survival).

Keywords: passerine birds, agroecosystem, fungicides, sublethal effects, thyroid hormones, body condition, sperm quality

Introduction

It is estimated that, worldwide, only fourteen crops make up the majority of the human diet. However, all of these crops are susceptible to diseases while growing in the field or after the harvest (Strange & Scott, 2005), and fungal diseases represent a major risk not only to crop yield (Fisher et al. 2012), but also to food safety (Strange & Scott, 2005). Many pesticides, and more specifically fungicides, are massively used in agriculture to circumvent these problems and to reduce the economic costs and health issues of fungal pests. For example, fungicides accounted for more than 40% of total pesticide sales from 2011 to 2018 in the European Union (Zubrod et al. 2019). Fungicides can even represent up to 90% of the pesticides that are used for certain crops (*e.g.*, orchards, vineyards, Zubrod et al. 2019).

Triazoles are a large family of systemic, broad-spectrum fungicides used in agriculture since the 1970s-80s (Ribas e Ribas et al. 2016). They are designed to interfere with lanosterol 14 α -demethylase (CYP51), an enzyme of the cytochrome P450 monooxygenase (CYP) superfamily (Becher & Wirsal, 2012). In fungi, CYP51 plays a key role in the biosynthesis of ergosterol, a sterol located in fungal membranes and involved in membrane permeability and fluidity (Daum et al. 1998). One of the main problems associated with agricultural triazoles is their potential effect on non-target species, through non-target physiological mechanisms (*e.g.*, Fernández-Vizcaíno et al. 2020; Lopez-Antia et al. 2021; Bellot et al. 2022). In general, the effects of fungicides, and more particularly triazoles, are understudied in wildlife. Indeed, most studies focus on acute ecotoxicity tests and with model species commonly used in toxicology (*e.g.*, zebrafish, earthworm, Binev et al. 2005; Sancho et al. 2010; Andreu-Sánchez et al. 2012; Raby et al. 2019; Rico et al. 2016). Although these studies elucidate the biological targets of triazoles and their modes of action, they are not necessarily representative of the effects of triazoles at concentrations found in the environment. In addition, model species are often laboratory-raised species, and thus, are not always representative of the ecosystems where triazoles are used. For wildlife, acute exposures to high concentrations of triazoles are scarce and correspond to sporadic events, such as accidental contamination (*e.g.*, Capel et al. 1988). Therefore, to better understand the sublethal effects that triazoles may have on non-target species, it is necessary to study the impact of long-term exposure of triazoles on wildlife, and especially vertebrate sentinel species of agroecosystems.

Among vertebrates, birds are particularly relevant to study this question because many bird species exploit or live near agricultural areas and are potentially exposed to triazoles through

multiple routes of exposure (Robinson, 2004; Lopez-Antia et al. 2016; Fernández-Vizcaíno et al. 2020, 2022; Ortiz-Santaliestra et al. 2020; Angelier et al. 2023). This contamination is likely associated with chronic exposure to low concentrations of triazoles, for example in water, soil, air and seeds (*e.g.*, Kahle et al. 2008; Rokbani et al. 2019; Pelosi et al. 2021; Lopez-Antia et al. 2016, 2021). In this context, specific bird species (*e.g.*, non-migratory species) can be relevant sentinel species of agroecosystems because they rely on this specific environment to complete their annual life cycle. As such, they should be chronically exposed to specific triazoles that are used for agricultural purposes and, thus, represent a powerful model to better understand the impact of environmental concentrations of triazoles on key organismal traits. Experimental studies have shown that chronic or subchronic exposure to triazoles can induce reprotoxic effects in birds (Grote et al. 2008; Lopez-Antia et al. 2018). These effects may arise from an interaction of triazoles with non-fungal sterol synthesis (Zarn et al 2003). Thus, triazoles can act as major endocrine disruptors by modulating the expression and activity of enzymes such as cytochrome P450 aromatase (Goetz et al. 2007; Taxvig et al. 2007; Jacobsen et al. 2012; Warrilow et al. 2013; Yu et al. 2013) with potential implications for multiple metabolic pathways, and also for reproduction (Poulsen et al. 2015; Huang et al. 2022). Indeed, pesticides, and possibly triazoles can have a direct impact on reproduction by causing testicular abnormalities that affect spermatogenesis and the mobility and quality of germ cells (carbamates and neonicotinoids: Tokumoto et al. 2013; Mohanty et al. 2017; Humann-Guillemint et al. 2019; triazoles: Machado-Neves et al. 2018). For example, triazoles could alter the fertility of individuals, notably through a decrease in sperm quantity and quality (*e.g.*, Taxvig et al. 2007; Grote et al. 2008; Pereira et al 2019).

More recently, triazoles have also been suspected to affect key endocrine mechanisms, such as the Hypothalamic-Pituitary-Thyroid (HPT) axis and the resulting regulation of thyroid hormones (THs). These hormones also play a key role in gonadal development and the regulation of seasonal reproduction in mammals and birds (McNabb, 2007; Mohanty et al. 2017). The first evidence highlighting the effect of triazoles on thyroid disruption comes from the study of aquatic models (*e.g.*, zebrafish, Yu et al. 2013; Li et al. 2019). However, the potential effects ofazole fungicides on thyroid functioning have been overlooked in non-aquatic vertebrates. In adults birds, the HPT axis is also a key physiological system, which is, for example, involved in (i) seasonal processes, notably the moulting process that allows birds to renew their plumage, and (ii) metabolism and thermoregulation (reviewed by McNabb, 2007). An increase in THs, such as triiodothyronine (T3) and thyroxine (T4), triggers the initiation of

moulting (Groscolas & Leloup, 1986; Groscolas & Cherel, 1992; Jenni-Eiermann et al. 2002), as experimentally shown in the white-crowned sparrow (*Zonotrichia leucophrys gambelii*) (Pérez et al. 2018). Similarly, TH levels have been linked to metabolism in multiple bird species and increased TH levels are associated with increased maintenance costs (basal metabolic rate, Chastel et al. 2003; Elliott et al. 2013).

In this study, we experimentally tested the impact of environmental concentration of tebuconazole on key traits of a sentinel bird species of agroecosystems, the house sparrow (*Passer domesticus*). Tebuconazole is the most widely used triazole fungicides in the world and it is found in multiple agroecosystems, especially in air (Désert et al. 2018), soil (Kalogridi et al. 2014), and water (Kahle et al. 2008). To date, only a few studies have evaluated the impact of tebuconazole on terrestrial non-target vertebrate species: partridges (Fernández-Vizcaíno et al. 2020; Lopez-Antia et al. 2021), bats (Machado-Neves et al. 2018), and amphibians (Poulsen et al. 2015; Bernabò et al. 2020). However, none of these experimental studies have tested the impact of tebuconazole on the levels of TH and related physiological traits. Thus, the aim of this study was to experimentally test the impact of exposure to tebuconazole on the HPT axis in wild-caught house sparrows that were kept in captivity under controlled conditions. We also examined the impact of tebuconazole on plumage quality and body condition, two phenotypic traits that are closely related to the HPT axis, as well as on sperm quality.

To do this, we randomly chose half of our wild-caught captive house sparrows to be exposed to tebuconazole through their drinking water ('exposed sparrows'), while the other sparrows were not exposed ('control sparrows'). We then examined the impact of this experimental treatment on the (1) levels of THs (T3 and T4), (2) plumage quality, using the morphometric characteristics of feathers, (3) body condition (body mass corrected for body size: body condition index (BCI)), and (4) the morphology of the sperm. Plumage quality, body condition and sperm quality are closely related to individual performance and fitness. Indeed, feathers enable birds to perform vital functions such as flight, signalling and thermoregulation (Swaddle et al. 1996; Dawson et al. 2000; Jenni & Winkler, 2020). Similarly, the BCI is a commonly used proxy of the physiological state of an individual, and disruption of metabolism usually translates into an abnormal BCI. Finally, sperm morphology may be an indicator of sperm quality and reproductive success (Helfenstein, Podelvin, & Richner, 2010). We predicted that exposure to tebuconazole would (1) disrupt the HPT axis and result in reduced TH levels (T3 and T4), as previously shown in laboratory aquatic model species (Li et al. 2019); (2) affect moult, with a detrimental impact on plumage quality (smaller and less dense feathers in exposed

sparrows); (3) result in an increased BCI, because tebuconazole should lower metabolism (Bellot et al. 2022), and thus maintenance costs, through its impact on TH levels (reduced TH levels, prediction 1), and (4), reduce sperm quality by altering sperm morphology.

Material and Method

Study species

This study was conducted on captive adult's house sparrows (*Passer domesticus*, Linnaeus, 1758). House sparrows are an appropriate sentinel species for assessing the contamination of agroecosystems because they are sedentary, strongly linked to agricultural environments (feeding, nesting, and breeding), and can be impacted by agricultural practices and landscapes (Swaileh and Sansur, 2006; Humann-Guillemint et al. 2019). A total of 39 sparrows were used in this experiment ($n_{females} = 20$ and $n_{males} = 19$) that were captured in Western France in 2019 (Secondigné-sur-Belle). Birds were acclimated to captivity for two weeks before the start of the experiment. All sparrows were maintained in captivity in 4 similar outdoor aviaries (length: 3 m, width: 2 m, height: 2 m). The birds were maintained in outdoor aviaries to keep them under natural seasonal temperature cycles. Birds were split in the aviaries by treatment type and sex: control birds were split into two aviaries for a total of 9 females and 9 males. Exposed birds were split into the two other aviaries for a total of 11 females and 10 males. As females and males were kept in separate aviaries for the duration of the experiment, no eggs were laid. These sample sizes differ slightly between the two treatments, as some of the birds did not adapt well to captivity and were released.

Each aviary was equipped with artificial shelter boxes (*i.e.*, one box per bird). The sparrows were fed *ad libitum* with a mixture of commercial seeds, millet sprays, grit, and salt/mineral blocks. In all aviaries, water was dispensed *ad libitum* through a drinking trough and was changed on a weekly basis.

Experimental design

Individuals were exposed to tebuconazole through their drinking water a source of exposure that is thus far less studied in comparison to the ingestion of treated seeds (e.g., Lopez-Antia et al. 2013; Lopez-Antia et al. 2018; Lopez-Antia et al. 2021; Fernández-Vizcaíno et al. 2022). In

the two control aviaries, control individuals had access to regular tap drinking water without tebuconazole. In the two exposed aviaries, tebuconazole (Sigma-Aldrich, CAS. No: 107534-96-3, purity: $\geq 98.0\%$) was diluted with tap water using a precision balance ($\pm 0.0001\text{ g}$) to reach a concentration of $550\ \mu\text{g}\cdot\text{L}^{-1}$. As in the control group, the water was changed once per week. Based on the physico-chemical characteristics of tebuconazole (<http://sitem.herts.ac.uk/aeru/ppdb/en/Reports/610.htm>), it is considered that the weekly concentration in the drinking water should vary negligibly (i.e., non-photodegradable, non-volatile, thermostable and stable compound in water). Considering that a sparrow drinks, on average, the equivalent of 30% of its body weight per day (Bartholomew & Cade, 1963), a 25 g individual would consume approximately $4.1\ \mu\text{g}$ of tebuconazole *per day* via drinking water, corresponding to an exposure concentration of $0.164\ \text{mg}\ \text{kg}^{-1}\ \text{bw}\ \text{d}^{-1}$. Comparatively, the median lethal dose (LD50) and the chronic (21 days) No-Observed-Effect Level (NOEL) of tebuconazole reported in Northern Bobwhite (*Colinus virginianus*) were $1988\ \text{mg}\cdot\text{kg}^{-1}$ and $5.8\ \text{mg}\ \text{kg}^{-1}\ \text{bw}\ \text{d}^{-1}$, respectively (<http://sitem.herts.ac.uk/aeru/ppdb/en/Reports>). Therefore, our exposure concentration is estimated to be 36 times lower than the NOEL. The plasma concentration of tebuconazole in the sparrows was checked analytically (see Bellot et al. 2022), and showed that the individuals from the control group were free of contamination and that the individuals from the exposed group had an average plasma concentration of $59.7\ \text{pg}\cdot\text{g}^{-1}$ tebuconazole. Moreover, we chose this water concentration of tebuconazole because we have previously shown that it results in an environmentally realistic plasma contamination (Angelier et al. 2023). Exposure to tebuconazole started on 6 November 2019 and lasted until 29 June 2020 (i.e., until the end of sampling, see below, **Fig 1**). We specifically decided to expose individuals to tebuconazole during seven months to test the impact of a long-term exposure to environmental doses of tebuconazole on TH and key physiological parameters. Although the duration of exposure to tebuconazole is rather long in our experiment, such a long-term chronic exposure can be found in some agroecosystems that rely on multiple treatments of fungicides (e.g., vineyards where tebuconazole is used from April to August), and also in mixed landscapes with multiple crops that used triazoles during different periods (e.g., cereal crops in autumn and winter, orchards, vegetables, vineyards in spring and summer). In order to verify that there was no major difference in drinking behaviour due to treatment, we analysed the hydration state (plasma osmolarity, Vapro2 osmometer, Elitech group, France) of several control and exposed sparrows (for individuals with sufficient plasma left, $n = 9$ for control sparrows and $n = 9$ for exposed sparrows). This parameter has been shown to vary according to sparrow drinking behaviour (Brischoux et al. 2020). We found no difference in plasma osmolarity ($F_{(1,16)} = 0.64$,

$p = 0.436$), suggesting that there is no major difference in drinking behaviour between control and exposed sparrows. Tebuconazole concentrations of exposed and control drinking water were assayed by an independent accredited analytical laboratory (QUALYSE laboratory, La Rochelle, France). Tebuconazole was undetectable in control water while tebuconazole concentrations were $550 \mu\text{g}\cdot\text{L}^{-1}$ in the water of exposed aviaries (by Liquid Chromatography Tandem Mass Spectrometry (LC-MS/MS)).

Feather morphometry

We used a standardized protocol to study plumage quality of exposed and control sparrows. In April 2020, we plucked the two inner rectrices (central tail feathers) of all sparrows, **Fig 1**. These plucked feathers had grown before birds were captured and before the experimental exposure to tebuconazole (*i.e.*, in August/September 2019 during the annual post-nuptial moult) and they, therefore, did not allow us to test the impact of tebuconazole exposure on plumage quality. However, feather plucking stimulates the growth of new feathers, and all inner rectrices had regrown within a month in most of individuals ($n = 30$). Then, in May 2020, after 6 months of exposure to tebuconazole, we plucked these two inner rectrices that had regrown while birds were either exposed to tebuconazole or not in order to test the impact of tebuconazole exposure on plumage quality, **Fig 1**. Five morphometric parameters were then assessed on the plucked tail feathers of control and exposed house sparrows (initial and regrown). The morphometric parameters of both rectrices of each individual were averaged for each measurement timepoint. Only individuals with data for both measurement times (initial: measure 1, regrown: measure 2, **Fig 1**) were considered for statistical analysis ($n = 30$ sparrows, with exposed and control individuals equally distributed). The first three morphometric parameters were measured using a high-resolution balance (feather mass, $\pm 0.01 \text{ mg}$) and digital caliper (total feather length and diameter of the calamus $\pm 0.01 \text{ mm}$). Standardized pictures of the feather were also taken, and feather area was calculated using a dedicated image software (ImageJ, v1.53). Lastly, the density of feathers was calculated by dividing the mass of the feathers by their area.

Thyroid hormones

After 6 months of exposure, on 26 and 27 May, blood samples ($100\text{-}150\mu\text{L}$) were successfully collected from all but one individual by puncture of the brachial vein to assess the effect of the treatment on total triiodothyronine (T3) and total thyroxine (T4) of the house sparrows ($n = 37$, **Fig 1**). All blood samples were centrifuged, and red blood cells and plasma were separated and

stored at $-20\text{ }^{\circ}\text{C}$ until the assay. Plasma concentrations of total T3 and T4 were determined using radioimmunoassay at the Centre d'Études Biologiques de Chizé (CEBC) as previously described by Chastel et al. (2003). The lower limits of detection for T3 and T4 were 0.08 ng.mL^{-1} and 0.602 ng.mL^{-1} respectively. For each hormone, all samples were run in duplicates in a single assay. The intra-assay CVs were 9.49% and 11.36% for T3 and T4, respectively.


Individual condition

To assess the effect of the treatment on the body condition of the sparrows, body mass was taken three times during the experiment ($n = 38$, **Fig 1**): once at the start of the experiment (M1), a second time after 5 months (M2), and a third time after 6 months of experimentation (M3, **Fig 1**). BCI was calculated by extracting the residuals from the linear regression of individual body mass as a function of tarsus length ($F_{(1,112)} = 44.28$, $p < 0.001$). Control and exposed sparrows did not differ in body size (tarsus, $F_{(1,36)} < 0.01$, $p = 0.932$).

Sperm morphology

In order to test the effect of the experimental treatment on the sperm morphology of the sparrows, we took a sperm sample during the breeding season (10 June 2020, **Fig 1**). These sperm samples were obtained by faecal and abdominal massage described in detail by Girndt et al. (2017). This method allowed us to have a successful collection for 17 males including 8 males from the control group and 9 males from the exposure group. Each sperm sample was collected directly into a 0.5 ml microtube containing 200 μl of 5% formalin. At the laboratory, 10 μl of each sample was put on a slide to perform microscopic observation of sperm cells at 400x magnification on a bright field setting. For each individual, 100 photos of distinct intact sperm were taken (based on the method of Girndt et al. 2017; 2019). The different components of the sperm (*i.e.*, head, including acrosome, flagellum, including midpiece) were measured by a single observer from digital images using ImageJ software (v1.53). Total sperm length was calculated as the sum of head and flagellum measurements (Girndt et al. 2017; 2019).

Fig 1. Timeline describing the different steps of the experiments. The five dots highlight a key event during the experiment. The two red borders highlight the period of exposure of the experimental individuals to tebuconazole.



Statistical analysis

All statistical analyses were carried out in Rstudio (R version 4.2.1). To test the effect of the experimental treatment (*i.e.*, exposed to tebuconazole or control) on the levels of TH, feather morphometry, body condition (BCI), and sperm length, linear mixed models (LMM) and linear models (LM) were used. All models were selected using the backward stepwise selection method by eliminating non-significant terms ($p > 0.100$). After conducting LMMs, specific comparisons were conducted using contrast procedures with the Satterthwaite approximation to describe the differences that were found between different groups of birds. The contrasts were made on the selected minimal model. We checked that all models met the assumptions of normality and homoscedasticity of residuals.

First, the plasma levels of T3 and T4 were analysed using linear models (LM) with two explanatory variables (treatment: control or exposed; sex: female or male) and their interaction. A log-transformed function was applied to the T4 thyroid hormone data to improve the distribution and homogeneity of the model residuals.

Second, because the morphometric parameters of the feathers were correlated, a principal component analysis (PCA) was performed. The first principal component (PC1) of the PCA explained 57.4% of the variation. PC1 was significantly and positively correlated with total

feather length, ($r = 0.92$, $p < 0.001$), feather mass ($r = 0.91$, $p < 0.001$), feather area ($r = 0.80$, $p < 0.001$), diameter of the calamus ($r = 0.60$, $p < 0.001$), and feather density ($r = 0.42$, $p < 0.001$). The second principal component (PC2) explained 28.9% of the variation. PC2 was significantly and positively correlated with feather density ($r = 0.87$, $p < 0.001$) and feather mass ($r = 0.38$, $p < 0.005$). Overall, PC1 represented the size of the feather while PC2 represented the density of the feather. For each individual, PC1 and PC2 values were extracted and then analysed using an LMM. The initial model included 3 fixed factors: (treatment: control or exposed; sex: female or male; and time: measurement 1 of initial feathers or measurement 2 of newly formed feathers, **Fig 1**), and their interactions to test the effect of tebuconazole on feather morphometry (PC1 or PC2). The identity of individuals was added as a random effect in the models because individuals were sampled twice (measures 1 and 2).

In addition, an LMM with three fixed effects (treatment: control or exposed; sex: female or male; time of mass measurements: M1, M2 or M3, **Fig 1**.) and their interactions was carried out to test the effect of tebuconazole on BCI. Individual identity was added as a random effect in the model because the mass of individuals was measured three times.

Finally, the sperm length was analysed by LMMs with treatment as an explanatory variable (control or exposed). Specifically, three response variables were assessed: head length, flagellum length and total sperm length. The identity of the individual was added as a random effect in each model as 100 sperm were measured for each individual.

Results

Thyroid Hormones (T3 and T4)

The “Treatment” factor did not significantly influence the T3 levels of sparrows (**Table 1**), whereas there was a significant effect of the “Sex” factor (**Table 1**), with higher T3 concentrations in females than in males (**Fig 2.A**). The treatment had a significant effect on the levels of T4 (**Table 1**). Individuals exposed to tebuconazole had significantly lower T4 levels than controls (**Fig 2.B**). T4 levels were not influenced by sex (**Table 1**). The “Treatment x Sex” interaction had no significant effect on T3 or T4 levels (**Table 1**), demonstrating that the influence of the experimental treatment on TH did not differ significantly between sexes (**Fig 2.A.B**).

Fig 2. Plasma concentrations of T3 (2.A) and T4 (2.B) in ng.mL⁻¹ according to treatment (control or exposed to tebuconazole) and sex (female or male).

Control and exposed individuals are represented in black and gray, respectively. Females and males are represented by circles and triangles, respectively. The concentration of T3 did not differ significantly between exposed and control sparrows ('ns', $p > 0.100$), whereas the concentration of T4 was significantly lower for exposed sparrows compared to control sparrows ('*', $p < 0.050$). Means \pm SE are presented.




Table 1: Effect of treatment (control or exposed to tebuconazole) and sex (female or male) on the levels of T3 and T4 of house sparrows.

A log-transformed function was applied to the T4 data in statistical models. Significant terms are indicated by the following codes: '****' $p < 0.001$ '***' $p < 0.010$ '**' $p < 0.050$ '.' $p < 0.100$. All models were selected using the backward stepwise selection method by eliminating non-significant terms ($p > 0.100$). The abbreviations used are as follows: Degrees of freedom in numerator (NumDF), degrees of freedom in denominator (DenDF), Fisher's statistics (F-value), the p-value associated (Pr (> F)).

Response variables	Explanatory factors	NumDF	DenDF	F-value	Pr (>F)
T3	Treatment	1	34	1.00	0.324
	Sex	1	35	5.16	0.029 *
	Treatment : Sex	1	33	0.09	0.770
T4	Treatment	1	35	5.51	0.025 *
	Sex	1	34	0.09	0.767
	Treatment : Sex	1	33	0.04	0.838

Feather morphometrics

PC1 and PC2 scores significantly varied according to the “Time” factor (**Table 2**). Specifically, the regrown feathers were overall smaller and less dense relative to the initial feathers (**Fig 3**).

Although there was no significant effect of either the “Sex” or “Treatment” factors on PC1 and PC2 scores (**Table 2**), there was a significant effect of multiple interactions on both PC1 and PC2 scores (**Table 2**), demonstrating that the influence of the treatment on PC1 and PC2 was complex and differed between sexes and groups.

Regarding PC1 scores, the “Time x Sex” and “Time x Treatment x Sex” interactions were significant (**Table 2**), demonstrating that the influence of the experimental treatment on this parameter differed significantly depending on the time of measurement (*i.e.*, before or after feather regrowth) and the sex of the individuals (**Fig 3.A.B**). Specifically, the regrown feathers of exposed females were significantly larger than those of control females (measure 2, contrast: $t_{(41.2)} = -2.72$, $p = 0.009$, **Fig 3.A**), whereas this difference was not significant for the initial feathers (measure 1, contrast: $t_{(41.2)} = -1.00$, $p = 0.322$, **Fig 3.A**). For males, there was no difference in feather size between exposed and control individuals at any time (measure 2, contrast: $t_{(41.2)} = 0.64$, $p = 0.524$ and measure 1, contrast: $t_{(41.2)} = -0.68$, $p = 0.500$, **Fig 3.B**).

Regarding PC2 scores, the “Treatment x Sex”, and “Time x Treatment” interactions were significant (**Table 2**), demonstrating that the influence of the experimental treatment on this parameter differed significantly depending on the time of measurement (*i.e.*, before or after feather regrowth) and the sex of the individuals (**Fig 3.C.D**). Interestingly the regrown feathers of exposed females (measure 2) were significantly less dense than those of control females (contrast: $t_{(48.5)} = 2.34$, $p = 0.024$), whereas this difference was not significant for the initial feathers (measure 1, contrast: $t_{(48.5)} = -0.60$, $p = 0.549$, **Fig 3.C**). The feathers of exposed males were overall denser than those of the control males (contrast: $t_{(26.0)} = -2.08$, $p = 0.048$, **Fig 3.D**). Specifically, feathers of exposed males were denser than those of control males before exposure to tebuconazole (contrast: $t_{(48.5)} = -1.94$, $p = 0.058$) but this difference was not significant for the regrown feathers, *i.e.*, after exposure (contrast: $t_{(48.5)} = -1.37$, $p = 0.178$, **Fig 3.D**).

Fig 3. PC1 scores (feather size, 3.A and 3.B) and PC2 scores (feather density, 3.C and 3.D) as a function of the time of measurement (initial feathers corresponding to measurement 1 and regrown feathers corresponding to measurement 2).

Figures 3.A and 3.C show the results obtained for females and figures 3.B and 3.D show the results obtained for males. Control and exposed individuals are represented in black and gray, respectively. Significant terms are indicated by the following codes: '*' $p < 0.050$, '.' $p < 0.100$ and 'ns' $p > 0.150$. The means \pm SE are presented.




Table 2: Effects of time of measurement on feather morphometric parameters (initial or after regrowth), treatment (control or exposed to tebuconazole), sex (female or male), as well as their interactions on PC1 scores (feather size) and PC2 scores (feather density), respectively.

Individual identity was added as a random factor. Significant terms are indicated by the following codes: '****' $p < 0.001$ '***' < 0.010 '**' $p < 0.050$ '.' $p < 0.100$. All models were selected using the backward stepwise selection method by eliminating non-significant terms ($p > 0.100$). The abbreviations used are as follows: Degrees of freedom in numerator (NumDF), degrees of freedom in denominator (DenDF), the Chi-square statistic (khi²-value), the p-value associated (Pr (> khi²)).

Response variables	Explanatory factors	NumDF	Den DF	khi ² -value	Pr (> khi ²)
PC1 (~ Length)	Time	1	26	35.49	< 0.001 ***
	Treatment	1	26	2.50	0.114
	Sex	1	26	< 0.001	0.976
	Time : Treatment	1	26	0.13	0.718
	Time : Sex	1	26	6.82	0.009 **
	Treatment : Sex	1	26	2.09	0.148
	Time : Treatment : Sex	1	26	4.70	0.030 *
PC2 (~ Density)	Time	1	28	43.43	< 0.001 ***
	Treatment	1	26	0.39	0.534
	Sex	1	26	2.53	0.112
	Time : Treatment	1	28	5.01	0.025 *
	Time : Sex	1	27	1.74	0.187
	Treatment : Sex	1	26	5.11	0.024 *
	Time : Treatment : Sex	1	26	1.72	0.190

Body Condition Index

Although the factors “Sex”, “Time”, and “Treatment” did not significantly influence the BCI of individuals, the “Time x Treatment”, and “Time x Treatment x Sex” interactions had significant effects on the BCI of the individuals (**Table 3**). This showed that the influence of the experimental treatment on the BCI differs significantly according to the sex of the individual and the time of the measurement (M1, M2 or M3, **Fig 4**). Specifically, the BCI of control females varied over time, where they had a higher BCI at time M3 compared to time M1 (contrast: $t_{(68.0)} = -4.45$, $p < 0.001$, **Fig 4.A**). However, the BCI of exposed females did not vary over time (contrasts: all $p > 0.345$ **Fig 4.A**). Moreover, exposed females had a greater BCI than control females at time M1, *i.e.*, before exposure to tebuconazole (contrast: $t_{(59.1)} = -2.31$, $p = 0.024$) whereas this difference became marginal at time M2, *i.e.*, after 5 months of exposure (contrast: $t_{(59.1)} = -1.97$, $p = 0.054$) and non-significant at time M3, *i.e.*, after 6 months of exposure (contrast: $t_{(59.1)} = 1.20$, $p = 0.235$, **Fig 4.A**). In control and exposed males, there was no significant variation in BCI among times M1, M2 and M3 (contrasts: all $p > 0.089$, **Fig 4.B**).

Fig 4. Body condition index of individuals (BCI) as a function of the timing of mass measurement (M1, M2 or M3), treatment (control or exposed to tebuconazole) and the sex of individuals (female 4A or male 4B).

Control and exposed individuals are represented in black and gray, respectively. Significant terms are indicated by the following codes: '*' $p < 0.050$, '.' $p < 0.100$ and 'ns' $p > 0.150$. Means \pm SE are presented.




Table 3: Effects of the timing of mass measurement (M1, M2 or M3), treatment (control or exposed to tebuconazole), sex (female or male), and their interactions on the BCI of sparrows.

Individual identity was added as a random factor. Significant terms are indicated by the following codes: '****' < 0.001 '***' < 0.010 '**' < 0.050 '.' < 0.100. All models were selected using the backward stepwise selection method by eliminating non-significant terms ($p > 0.100$). The abbreviations used are as follows: Degrees of freedom in numerator (NumDF), degrees of freedom in denominator (DenDF), the Chi-square statistic (khi²-value), the p-value associated (Pr (> khi²)).

Response variable	Explanatory factors	NumDF	DenDF	khi ² -value	Pr (> khi ²)
Body Condition Index	Treatment	1	34	3.81	0.051 .
	Sex	1	34	2.41	0.121
	Time	2	68	2.04	0.360
	Treatment : Sex	1	34	0.10	0.748
	Treatment : Time	2	68	9.44	0.009 **
	Sex : Time	2	68	4.70	0.095 .
	Treatment : Sex : Time	2	68	10.26	0.006 **

Sperm morphology

Sperm head length, flagellum length, and total length were not significantly affected by treatment (control or exposed, Table 4). Sperm head length averaged (mean \pm SE): 14.85 $\mu\text{m} \pm 0.033$ for controls and 14.68 $\mu\text{m} \pm 0.027$ for exposed sparrows. The length of the sperm flagellum was on average 86.95 $\mu\text{m} \pm 0.15$ for the controls and 87.09 ± 0.12 for the exposed sparrows. Finally, the total length of the spermatozoa was 101.80 $\mu\text{m} \pm 0.16$ for the controls and 101.77 $\mu\text{m} \pm 0.12$ for the exposed sparrows.

Table 4: Effects of treatment (control or exposed to tebuconazole) on the length of the different sperm components (head, flagellum and total length).

Individual identity was added as a random factor. Significant terms are indicated by the following codes: '****' < 0.001 '***' < 0.010 '**' < 0.050 '.' < 0.100. The abbreviations used are as follows: Degrees of freedom in numerator (NumDF), degrees of freedom in denominator (DenDF), Fisher's statistics (F-value), the p-value associated (Pr (> F)).

Response variables	Explanatory factors	NumDF	DenDF	F-value	Pr (> F)
Head length	Treatment	1	14.4	1.68	0.215
Flagellum length	Treatment	1	15.0	0.15	0.700
Total length	Treatment	1	15.0	0.04	0.844

Discussion

In this study, we showed for the first time in wild-caught birds that chronic exposure to a widely used triazole fungicide (tebuconazole) could significantly reduce the levels of T4. However, we did not find any effect of such exposure to tebuconazole on the levels of T3. In female sparrows, exposure to tebuconazole also affected the structure of the feathers that grew during the experiment, with larger but less dense feathers for exposed individuals compared to controls. In addition, body condition did not vary considerably throughout the experiment, except for the control females that experienced a significant increase in their body condition towards the end of the experiment. Finally, sperm morphology was not affected by the treatment. Overall, we demonstrated sublethal effects of tebuconazole on key physiological parameters, such as the HPT axis and one other related physiological trait (plumage quality) in vertebrates that are commonly found in agroecosystems.

Thyroid hormones

Although T3 levels were not significantly affected by the chronic exposure to tebuconazole, our experimental treatment was associated with a significant decrease in circulating T4 levels. In agreement with our results, previous studies have shown that an exposure to tebuconazole could induce an alteration of the HPT axis, and notably a decrease in T4 levels in laboratory

model species (Yu et al. 2013; Li et al. 2019). Indeed, zebrafish showed a significant decrease in T4 levels when exposed to tebuconazole concentrations of 200 and 500 $\mu\text{g.L}^{-1}$ (Li et al. 2019). Similar effects were also found in zebrafish larvae, with a significant decrease in T4 levels when larvae were exposed to a tebuconazole concentration of 4 mg.L^{-1} (Yu et al. 2013), although this effect was not found at lower concentrations (1 and 2 mg.L^{-1}). Furthermore, and interestingly, Bernabò et al. (2016) found an effect of tebuconazole on the success and speed of metamorphosis in tadpoles, suggesting a disrupting effect of tebuconazole on the HPT axis.

Here, we showed for the first time that such dysregulation of the HPT axis may also occur in wild-caught terrestrial vertebrates. Importantly, the activity of the thyroid gland is under the control of the pituitary, which governs the biosynthesis and secretion of the thyroid stimulating hormone (TSH, encoded by *tsh α* and *tsh β* , **Fig 5**, Hall et al. 1970; Harris et al. 1978; Capen, 1998; Stoker et al. 2006). When TH levels become insufficient, the levels of TSH increase in order to stimulate the secretion of THs by the thyroid and restore homeostasis. Some studies have shown that exposure to azoles (difenoconazole, hexaconazole and tebuconazole, Liang et al. 2015; Yu et al. 2013) cause a significant increase in the transcription of genes related to TSH production and secretion (*i.e.*, *tsh β*), which could result from a stimulation of the HPT axis in response to decreased T4 levels (Liang et al. 2015; Pirahanchi et al. 2018). To test this hypothesis, it would be interesting now to measure the transcriptomic activity of *tsh β* to know if it is inversely proportional to the decrease in T4 observed in the sparrows exposed to tebuconazole or not.

Contrary to our prediction, we did not find any effect of tebuconazole on the levels of T3. Li et al. (2019) showed that exposing zebrafish to tebuconazole caused a significant decrease in T3 levels (dose: 200 and 500 $\mu\text{g.L}^{-1}$) but Yu et al. (2013) reported in another study that exposure to tebuconazole resulted in a significant increase in T3 levels in zebrafish larvae (4 mg.L^{-1}). Interestingly, Liang et al. (2015), found similar results to our study. Indeed, after exposing zebrafish larvae to difenoconazole (another triazole), a significant decrease in T4 levels was observed while no significant effect was observed on T3 levels. Altogether, these results suggest that the effect of tebuconazole on T3 levels may depend on the dose of tebuconazole, life-history stages, and/or species.


At the mechanistic level, the decrease in T4 levels may be enhanced by an increased activation of type II deiodase (encoded by *dio2*, **Fig 5.A**). Type II deiodase is involved in the catalytic

conversion of T4 to its more active form T3 (Bianco & Kim, 2006), so its activation could lead to a decrease in the stock of T4 (as suggested by Pandey & Mohanty, 2015, **Fig 5.A**). Supporting this idea, Liang et al. (2015) found that difenoconazole can cause an increase in *dio2* transcription. In agreement with the results of Liang et al. (2015), we did not observe any change in the levels of T3, although an increase in the conversion of T4 to T3 by type II deiodase should logically lead to an increase in T3 levels. However, the levels of T3 also depend on the activity of TH-degrading enzymes, such as type III deiodase (*dio3*, Bianco & Kim, 2006), which activates the conversion of T3 into inactive forms (diiodothyronine, T2, Dentice & Salvatore, 2011). Therefore, tebuconazole could affect the activation of these enzymes (*dio2* and *dio3*) and result in decreased T4 levels, but stable T3 levels (Dentice & Salvatore, 2011; Pandey & Mohanty, 2015). Alternatively, tebuconazole could also affect T4 transcription. If the transcriptomic activity of T4 is decreased but the levels of T3 are maintained, it is possible that exposed individuals show an increased biosynthesis of T3 independently of T4 in order to compensate for the decrease in T4 levels (**Fig 5.B**). Under normal circumstances, about 80% of T3 is formed from T4, while only 20% is produced directly by the thyroid (Pirahanchi et al. 2018). Thus, as suggested by Pandey & Mohanty (2015), the induction of a T4-independent T3 biosynthetic pathway could explain why T4 levels decrease but T3 levels are maintained (**Fig 5.B**).

Therefore, there is no clear consensus on how tebuconazole affects the thyroid system. Our study shows that tebuconazole seems to disrupt the HPT axis in wild terrestrial vertebrates, as previously found in aquatic models (Yu et al. 2013; Liang et al. 2015; Li et al. 2019). However, its mode of action could be complex and may involve several metabolic pathways (**Fig. 5**, see also Kjærstad et al. 2007; Yang et al. 2013; Yu et al. 2013; Liang et al. 2015; Li et al. 2019; Zhang et al. 2019). To further understand the impact of tebuconazole on the HPT axis, it would be relevant to use a transcriptomic approach to identify the biological targets responsible for this disruption of the thyroid axis.

Fig 5. Schematic representation of the different scenarios proposed for the mode of action of tebuconazole on the house sparrow endocrine system.

(A) By induction of type II deiodase activity resulting in increased conversion of T4 to T3 to ensure production of active TH at the expense of a decrease in T4 stock. (B) By inhibition of T4 transcription or release leading to compensation of T3 production by a T4 independent biosynthetic pathway. These examples are not exhaustive and hypotheses (A) and (B) are non-mutually exclusive. The flash symbol represents potential biological targets of tebuconazole in the negative feedback loop of thyroid hormones.



Feather quality

After exposure to tebuconazole, exposed individuals had larger but less dense feathers than the controls, but this effect was only found in females. To our knowledge, our study is the first to provide evidence for an effect of triazoles on plumage quality. Although we could not demonstrate that these effects were linked to a disruption of the HPT axis, this is plausible because the relationship between TH and the moulting process is well established (Davis & Davis, 1954; Wilson & Farner, 1960; Groscolas & Cherel, 1992; Cherel et al. 2004; McNabb, 2007). Previous studies have shown that moulting can be induced by the administration of exogenous TH, while the removal of the thyroid gland leads to the inhibition of moult or regrowth of artificially plucked feathers (Voitkevich, 1966; Hahn et al. 1992; Kuenzel, 2003; Pérez et al. 2018).

We found that tebuconazole caused a decrease in circulating T4 levels in exposed sparrows and several studies have demonstrated that T4 plays a key role in the moulting process of birds (Groscolas & Cherel, 1992; Cherel et al. 2004; McNabb, 2007). For example, Groscolas & Leloup (1986), showed a positive correlation between peak plasma T4 and the onset of moult in birds, suggesting a more important role of T4 than T3 in this process (Groscolas & Leloup, 1986). These findings are supported by several studies in multiple bird species (Decuyper &

Verheyen, 1986; Cherel et al. 1988, 2004; Jenni-Eiermann et al. 2002; Vézina et al. 2009). According to these studies and the results that we obtained on T4, we can hypothesise that the interference of tebuconazole with the HPT axis may have caused a change in the quality of the feathers that regrew during the phase of exposure to tebuconazole. However, it is important to note that we are not aware of any study that has linked circulating TH levels with feather quality because most studies focused on the onset or the rate of molting rather than on the quality of the feathers produced (*e.g.*, Decuyper, 1986; Cherel et al. 2004; Pérez et al. 2018). In addition, we found an effect of tebuconazole on feather quality mainly in females, while the T4 levels of both males and females were affected by tebuconazole. This suggests that the observed effects on feather quality could also be related to the effect of tebuconazole on other mechanisms. For example, feather quality and moult processes have been also linked to levels of glucocorticoids (corticosterone in birds) and testosterone (Nolan et al. 1992; Stoehr & Hill, 2001; DesRochers et al. 2009; Siefferman et al. 2013; Jenni-Eiermann et al. 2015). Previous studies have found that triazoles can also affect the synthesis of these two steroids in vertebrates through their action on cytochrome P450 enzymes (Goetz et al. 2007; Taxvig et al. 2007; Yang et al. 2018; Draskau et al. 2019). Therefore, disruption of sex hormones, such as testosterone, could potentially explain the sex-dependent effect of tebuconazole on feather quality. Further studies of these endocrine axes are therefore required to better understand the impact of triazoles on molt processes and feather quality.

The regrown feathers of the exposed females were overall larger and less dense relative to control females. This structural modification of feathers can potentially have implications for individual performance (*i.e.*, survival and reproduction) through an impact on flight quality, the ability of individuals to thermoregulate, or the ability to express secondary sexual signals through plumage (Swaddle et al. 1996; Dawson et al. 2000; Jenni & Winkler, 2020). Future studies should now examine the impact of tebuconazole on these feather-related traits.

Body condition index

Contrary to our prediction, we did not find any major effects of tebuconazole on the body condition of individuals. Most individuals maintained their body condition throughout the experiment, independently of the treatment. The only significant effect was an increase in BCI for control females towards the end of the experiment (M3), whereas this increase was not observed in exposed females or control and exposed males. This puzzling result may be an artifact, although our sample size is quite robust for such an experimental study. Indeed, most

previous studies have not reported any significant effect of exposure to triazoles on the body condition of birds (Lopez-Antia et al. 2013, 2018; Pandey & Mohanty, 2015; Fernández-Vizcaíno et al. 2020). However, it could also be linked to a biological process: control females may have accumulated body reserves at the end of the experiment in preparation for breeding. Indeed, females usually increase their body mass a few days/weeks before laying their eggs (e.g., Kvalnes et al. 2013; Hennin et al. 2015) while this phenomenon is not detectable in males. Tebuconazole exposure may have delayed or even suppressed the initiation of reproduction in females by interfering with seasonal endocrine processes, explaining then this absence of body weight gain in exposed females. Accordingly, studies have reported that triazoles may impair reproduction in birds (e.g., Fernandez-Vizcaino et al 2020; Lopez-Antia et al. 2021). Moreover, triazoles are also known to interfere with cytochrome P450 enzymes, which are associated with the regulation of sex hormones (e.g., CYP19, which is involved in the conversion of androgens to estrogens, Poulsen et al. 2015; Saxena et al. 2015). Since estrogens (such as estradiol) are known to control feeding behaviour, fat deposition and body condition in vertebrates, it is possible that deregulation of the steroid axis during the reproductive period could result in a sex-dependent effect of treatment on body condition. Interestingly, Pandey & Mohanty (2015) found a significant decrease in the mass of exposed birds in response to fungicides, but only during the breeding phase, suggesting that the effects of fungicides on the body condition of individuals may be more apparent during the energetically demanding reproductive stage (Pandey & Mohanty, 2015). We do not know the exact age of our adult sparrows as they were caught in the wild, but it would also be interesting to see if age could influence the response to tebuconazole exposure. We must, however, remain cautious because exposed birds and controls unexpectedly differed in body condition at the beginning of the experiment, making it difficult to draw strong conclusions. Moreover, the sparrows had access to food *ad libitum* in our experiment and captivity limits mobility and therefore energy expenditure. These ideal nutritional conditions may have masked a potential impact of tebuconazole on metabolism and/or food consumption, and individuals may have been able to maintain their BCI despite the tebuconazole treatment. Finally, it would be interesting to further test how tebuconazole may affect body condition. For example, it is possible that tebuconazole has an impact on food intake, water consumption, metabolism, or/and activity, all of which influence body condition. However, Lopez-Antia et al. (2021) showed no effect of tebuconazole on food intake in red-legged partridges suggesting that there is no effect on palatability or hunger. Future studies should therefore examine the impact of tebuconazole on body mass under realistic food

situations and test whether exposure to tebuconazole affects food consumption, activity and reproduction.

Sperm morphology

Chronic exposure to tebuconazole prior to the breeding season did not result in any significant effect on sperm length in exposed sparrows. This result is contrary to our prediction as tebuconazole is known to interfere with steroid hormones and catalase and to have a negative impact on the reproduction of individuals (Poulsen et al. 2015; Lopez-Antia et al. 2021; Huang et al. 2022). In addition, some studies have shown that pesticides can affect male germ cells, such as causing a decrease in the quantity and density of these cells, a reduction in testicular mass and an interruption of spermatogenesis (Tokumoto et al. 2013; Mohanty et al. 2017; Humann-Guilleminot et al. 2019). Machado-Neves et al. (2018) also showed that bats treated with 1 ml.L⁻¹ of tebuconazole for 30 days had morphometric changes in their testes and epididymis suggesting a potential effect of tebuconazole on the gonadal axis. Importantly, it would be interesting to complement our measurements with an assessment of sperm motility, sperm count, abnormalities, and oxidative damage, because fungicides may affect these variables despite no effect on sperm morphology. Therefore, further studies would be needed to understand how tebuconazole impacts fertility and reproductive success in wildlife.

Conclusion

Overall, our results showed physiological effects of tebuconazole on a sentinel species of agroecosystems, the house sparrow. These effects include a significant decrease in T4 levels, and an alteration of feather structure, which altogether suggests a dysregulation of thyroid homeostasis by tebuconazole. Therefore, our study highlights the importance of assessing the potential risks of chronic exposure to tebuconazole for wild vertebrates living in agroecosystems.

REFERENCES

- Andreu-Sánchez, O., Parafba, L. C., Jonsson, C. M., & Carrasco, J. M. (2012). Acute toxicity and bioconcentration of fungicide tebuconazole in zebrafish (*Danio rerio*). *Environmental Toxicology*, 27(2), 109-116. <https://doi.org/10.1002/tox.20618>
- Angelier, F., Prouteau, L., Brischoux, F., Chastel, O., Devier, M.-H., Le Menach, K., Martin, S., Mohring, B., Pardon, P., & Budzinski, H. (2023). High contamination of a sentinel vertebrate species by azoles in vineyards : A study of common blackbirds (*Turdus merula*) in multiple habitats in western France. *Environmental Pollution*, 316, 120655. <https://doi.org/10.1016/j.envpol.2022.120655>
- Bartholomew, G. A., & Cade, T. J. (1963). The Water Economy of Land Birds. *The Auk*, 80(4), 504-539. <https://doi.org/10.2307/4082856>
- Becher, R., & Wirsal, S. G. R. (2012). Fungal cytochrome P450 sterol 14 α -demethylase (CYP51) and azole resistance in plant and human pathogens. *Applied Microbiology and Biotechnology*, 95(4), 825-840. <https://doi.org/10.1007/s00253-012-4195-9>
- Bellot, P., Dupont, S. M., Brischoux, F., Budzinski, H., Chastel, O., Fritsch, C., Lourdaï, O., Prouteau, L., Rocchi, S., & Angelier, F. (2022). Experimental Exposure to Tebuconazole Affects Metabolism and Body Condition in a Passerine Bird, the House Sparrow (*Passer domesticus*). *Environmental Toxicology and Chemistry*, 41(10), 2500-2511. <https://doi.org/10.1002/etc.5446>
- Bernabò, I., Guardia, A., Macirella, R., Sesti, S., Crescente, A., & Brunelli, E. (2016). Effects of long-term exposure to two fungicides, pyrimethanil and tebuconazole, on survival and life history traits of Italian tree frog (*Hyla intermedia*). *Aquatic Toxicology*, 172, 56-66. <https://doi.org/10.1016/j.aquatox.2015.12.017>
- Bernabò, I., Guardia, A., Macirella, R., Sesti, S., Tripepi, S., & Brunelli, E. (2020). Tissues injury and pathological changes in *Hyla intermedia* juveniles after chronic larval exposure to tebuconazole. *Ecotoxicology and Environmental Safety*, 205, 111367. <https://doi.org/10.1016/j.ecoenv.2020.111367>
- Bianco, A. C., & Kim, B. W. (2006). Deiodinases : Implications of the local control of thyroid hormone action. *Journal of Clinical Investigation*, 116(10), 2571-2579. <https://doi.org/10.1172/JCI29812>
- Binev, R., Simeonov, R., Todorov, R., & Nikolov, Y. (s. d.). *Pathomorphological studies following experimental acute intoxication with the triazole fungicide triticonazole in pigs*. 3.
- Brischoux, F., Beaugeard, E., Mohring, B., Parenteau, C., & Angelier, F. (2020). Short-term dehydration influences baseline but not stress-induced corticosterone levels in the house sparrow (*Passer domesticus*). *Journal of Experimental Biology*, 223(3), jeb216424. <https://doi.org/10.1242/jeb.216424>
- Capel, P. D., Giger, Walter., Reichert, Peter., & Wanner, Oskar. (1988). Accidental input of pesticides into the Rhine River. *Environmental Science & Technology*, 22(9), 992-997. <https://doi.org/10.1021/es00174a001>
- Capen, C. C. (1998). Correlation of mechanistic data and histopathology in the evaluation of selected toxic endpoints of the endocrine system. *Toxicology Letters*, 102-103, 405-409. [https://doi.org/10.1016/S0378-4274\(98\)00244-6](https://doi.org/10.1016/S0378-4274(98)00244-6)
- Cherel, Yves., Leloup, Jacques., & Le Maho, Yvon. (1988). Fasting in king penguin. II. Hormonal and metabolic changes during molt. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 254(2), R178-R184.
- Chastel, O., Lacroix, A., & Kersten, M. (2003). Pre-breeding energy requirements : Thyroid hormone, metabolism and the timing of reproduction in house sparrows *Passer domesticus*. *Journal of Avian Biology*, 34(3), 298-306. <https://doi.org/10.1034/j.1600-048X.2003.02528.x>
- Cherel, Y., Durant, J. M., & Lacroix, A. (2004). Plasma thyroid hormone pattern in king penguin chicks : A semi-altricial bird with an extended posthatching developmental period. *General and Comparative Endocrinology*, 136(3), 398-405. <https://doi.org/10.1016/j.ygcen.2004.02.003>
- Daum, G., Lees, N. D., Bard, M., & Dickson, R. (1998). Biochemistry, cell biology and molecular biology of lipids of *Saccharomyces cerevisiae*. *Yeast*, 14(16), 1471-1510. [https://doi.org/10.1002/\(SICI\)1097-0061\(199812\)14:16<1471::AID-YEA353>3.0.CO;2-Y](https://doi.org/10.1002/(SICI)1097-0061(199812)14:16<1471::AID-YEA353>3.0.CO;2-Y)
- Davis, J., & Davis, B. S. (1954). The Annual Gonad and Thyroid Cycles of the English Sparrow in Southern California. *The Condor*, 56(6), 328-345. <https://doi.org/10.2307/1365019>

- Dawson, A., Hinsley, S. A., Ferns, P. N., Bonser, R. H. C., & Eccleston, L. (2000). Rate of moult affects feather quality : A mechanism linking current reproductive effort to future survival. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 267(1457), 2093-2098. <https://doi.org/10.1098/rspb.2000.1254>
- De Cock, M., de Boer, M. R., Lamoree, M., Legler, J., & van de Bor, M. (2014). Prenatal exposure to endocrine disrupting chemicals in relation to thyroid hormone levels in infants – a Dutch prospective cohort study. *Environmental Health*, 13(1), 106. <https://doi.org/10.1186/1476-069X-13-106>
- Decuyper, E., & Verheyen, G. (1986). Physiological basis of induced moulting and tissue regeneration in fowls. *World's Poultry Science Journal*, 42(1), 56-68. <https://doi.org/10.1079/WPS19860006>
- Dentice, M., & Salvatore, D. (2011). Local impact of thyroid hormone inactivation. *Journal of Endocrinology*, 209(3), 273-282. <https://doi.org/10.1530/JOE-11-0002>
- Désert, M., Ravier, S., Gille, G., Quinapallo, A., Armengaud, A., Pochet, G., ... & Quivet, E. (2018). Spatial and temporal distribution of current-use pesticides in ambient air of Provence-Alpes-Côte-d'Azur Region and Corsica, France. *Atmospheric environment*, 192, 241-256. <https://doi.org/10.1016/j.atmosenv.2018.08.054>
- DesRochers, D. W., Reed, J. M., Awerman, J., Kluge, J. A., Wilkinson, J., van Griethuijsen, L. I., Aman, J., & Romero, L. M. (2009). Exogenous and endogenous corticosterone alter feather quality. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 152(1), 46-52. <https://doi.org/10.1016/j.cbpa.2008.08.034>
- Draskau, M. K., Boberg, J., Taxvig, C., Pedersen, M., Frandsen, H. L., Christiansen, S., & Svingen, T. (2019). In vitro and in vivo endocrine disrupting effects of the azole fungicides triticonazole and flusilazole. *Environmental Pollution*, 255, 113309. <https://doi.org/10.1016/j.envpol.2019.113309>
- Elliott, K. H., Welcker, J., Gaston, A. J., Hatch, S. A., Palace, V., Hare, J. F., Speakman, J. R., & Anderson, W. G. (2013). Thyroid hormones correlate with resting metabolic rate, not daily energy expenditure, in two charadriiform seabirds. *Biology Open*, 2(6), 580-586. <https://doi.org/10.1242/bio.20134358>
- Fernández-Vizcaíno, E., Fernández de Mera, I. G., Mougeot, F., Mateo, R., & Ortiz-Santaliestra, M. E. (2020). Multi-level analysis of exposure to triazole fungicides through treated seed ingestion in the red-legged partridge. *Environmental Research*, 189, 109928. <https://doi.org/10.1016/j.envres.2020.109928>
- Fernández-Vizcaíno, E., Ortiz-Santaliestra, M. E., Fernández-Tizón, M., Mateo, R., Camarero, P. R., & Mougeot, F. (2022). Bird exposure to fungicides through the consumption of treated seeds : A study of wild red-legged partridges in central Spain. *Environmental Pollution*, 292, 118335. <https://doi.org/10.1016/j.envpol.2021.118335>
- Fisher, M. C., Henk, Daniel. A., Briggs, C. J., Brownstein, J. S., Madoff, L. C., McCraw, S. L., & Gurr, S. J. (2012). Emerging fungal threats to animal, plant and ecosystem health. *Nature*, 484(7393), 186-194. <https://doi.org/10.1038/nature10947>
- Girndt, A., Cockburn, G., Sánchez-Tójar, A., Løvlie, H., & Schroeder, J. (2017). Method matters : Experimental evidence for shorter avian sperm in faecal compared to abdominal massage samples. *PLOS ONE*, 12(8), e0182853. <https://doi.org/10.1371/journal.pone.0182853>
- Girndt, A., Cockburn, G., Sánchez-Tójar, A., Hertel, M., Burke, T., & Schroeder, J. (2019). Male age and its association with reproductive traits in captive and wild house sparrows. *Journal of Evolutionary Biology*, 32(12), 1432-1443. <https://doi.org/10.1111/jeb.13542>
- Goetz, A. K., Ren, H., Schmid, J. E., Blystone, C. R., Thillainadarajah, I., Best, D. S., Nichols, H. P., Strader, L. F., Wolf, D. C., Narotsky, M. G., Rockett, J. C., & Dix, D. J. (2007). Disruption of Testosterone Homeostasis as a Mode of Action for the Reproductive Toxicity of Triazole Fungicides in the Male Rat. *Toxicological Sciences*, 95(1), 227-239. <https://doi.org/10.1093/toxsci/kfl124>
- Groscolas, R., & Cherel, Y. (1992). How to Molt While Fasting in the Cold : The Metabolic and Hormonal Adaptations of Emperor and King Penguins. *Ornis Scandinavica*, 23(3), 328. <https://doi.org/10.2307/3676657>
- Groscolas, R., & Leloup, J. (1986). The endocrine control of reproduction and molt in male and female emperor (*Aptenodytes forsteri*) and adélie (*Pygoscelis adeliae*) penguins. *General and Comparative Endocrinology*, 63(2), 264-274. [https://doi.org/10.1016/0016-6480\(86\)90164-4](https://doi.org/10.1016/0016-6480(86)90164-4)
- Grote, K., Niemann, L., Selzsam, B., Haider, W., Gericke, C., Herzler, M., & Chahoud, I. (2008). epoxiconazole causes changes in testicular histology and sperm production in the japanese quail (*Coturnix coturnix japonica*). *Environmental Toxicology and Chemistry*, 27(11), 2368. <https://doi.org/10.1897/08-048.1>

- Hahn, T. P., Swingle, J., Wingfield, J. C., & Ramenofsky, M. (1992). Adjustments of the prebasic molt schedule in birds. *Ornis Scandinavica*, 314-321.
- Hall, R., Amos, J., Garry, R., & Buxton, R. L. (1970). Thyroid-stimulating Hormone Response to Synthetic Thyrotrophin-releasing Hormone in Man. *BMJ*, 2(5704), 274-277. <https://doi.org/10.1136/bmj.2.5704.274>
- Harris, A. R. C., Christianson, D., Smith, M. S., Fang, S.-L., Braverman, L. E., & Vagenakis, A. G. (1978). The Physiological Role of Thyrotrophin-Releasing Hormone in the Regulation of Thyroid-Stimulating Hormone and Prolactin Secretion in the Rat. *Journal of Clinical Investigation*, 61(2), 441-448. <https://doi.org/10.1172/JCI108955>
- Helfenstein, F., Podelin, M., & Richner, H. (2010). Sperm morphology, swimming velocity, and longevity in the house sparrow *Passer domesticus*. *Behavioral Ecology and Sociobiology*, 64(4), 557-565. <https://doi.org/10.1007/s00265-009-0871-x>
- Hennin, H. L., Legagneux, P., Bêty, J., Williams, T. D., Grant Gilchrist, H., Baker, T. M., & Love, O. P. (2015). Pre-breeding energetic management in a mixed-strategy breeder. *Oecologia*, 177(1), 235-243. <https://doi.org/10.1007/s00442-014-3145-x>
- Huang, T., Zhao, Y., He, J., Cheng, H., & Martyniuk, C. J. (2022). Endocrine disruption by azole fungicides in fish: A review of the evidence. *Science of The Total Environment*, 822, 153412. <https://doi.org/10.1016/j.scitotenv.2022.153412>
- Humann-Guilleminot, S., Clément, S., Desprat, J., Binkowski, Ł. J., Glauser, G., & Helfenstein, F. (2019). A large-scale survey of house sparrows feathers reveals ubiquitous presence of neonicotinoids in farmlands. *Science of the Total Environment*, 660, 1091-1097. <https://doi.org/10.1016/j.scitotenv.2019.01.068>
- Humann-Guilleminot, S., Tassin de Montaigu, C., Sire, J., Grünig, S., Gning, O., Glauser, G., Vallat, A., & Helfenstein, F. (2019). A sublethal dose of the neonicotinoid insecticide acetamiprid reduces sperm density in a songbird. *Environmental Research*, 177, 108589. <https://doi.org/10.1016/j.envres.2019.108589>
- Jacobsen, P. R., Axelstad, M., Boberg, J., Isling, L. K., Christiansen, S., Mandrup, K. R., Berthelsen, L. O., Vinggaard, A. M., & Hass, U. (2012). Persistent developmental toxicity in rat offspring after low dose exposure to a mixture of endocrine disrupting pesticides. *Reproductive Toxicology*, 34(2), 237-250. <https://doi.org/10.1016/j.reprotox.2012.05.099>
- Jenni, L., & Winkler, R. (2020). *The Biology of Molt in Birds*. Bloomsbury Publishing.
- Jenni-Eiermann, S., Helfenstein, F., Vallat, A., Glauser, G., & Jenni, L. (2015). Corticosterone : Effects on feather quality and deposition into feathers. *Methods in Ecology and Evolution*, 6(2), 237-246. <https://doi.org/10.1111/2041-210X.12314>
- Jenni-Eiermann, S., Jenni, L., & Piersma, T. (2002). Temporal uncoupling of thyroid hormones in Red Knots : T3 peaks in cold weather, T4 during moult. *Journal for Ornithologie*, 143(3), 331-340. <https://doi.org/10.1046/j.1439-0361.2002.02011.x>
- Kahle, M., Buerge, I. J., Hauser, A., Müller, M. D., & Poiger, T. (2008). Azole Fungicides : Occurrence and Fate in Wastewater and Surface Waters. *Environmental Science & Technology*, 42(19), 7193-7200. <https://doi.org/10.1021/es8009309>
- Kalogridi, E. C., Christophoridis, C., Bizani, E., Drimaropoulou, G., & Fytianos, K. (2014). Part II: temporal and spatial distribution of multiclass pesticide residues in lake sediments of northern Greece: application of an optimized MAE-LC-MS/MS pretreatment and analytical method. *Environmental Science and Pollution Research*, 21, 7252-7262. <https://doi.org/10.1007/s11356-014-2794-y>
- Kjærstad, M. B., Taxvig, C., Hass, U., Axelstad, M., Metzdorff, S., & Vinggaard, A. M. (2007). *Effects of azole fungicides on the function of sex and thyroid hormones*.
- Kuenzel, W. J. (2003). Neurobiology of molt in avian species. *Poultry Science*, 82(6), 981-991.
- Kvalnes, T., Ringsby, T. H., Jensen, H., & Sæther, B.-E. (2013). Correlates of egg size variation in a population of house sparrow *Passer domesticus*. *Oecologia*, 171(2), 391-402. <https://doi.org/10.1007/s00442-012-2437-2>
- Li, S., Wu, Q., Sun, Q., Coffin, S., Gui, W., & Zhu, G. (2019). Parental exposure to tebuconazole causes thyroid endocrine disruption in zebrafish and developmental toxicity in offspring. *Aquatic Toxicology*, 211, 116-123. <https://doi.org/10.1016/j.aquatox.2019.04.002>

- Liang, X., Yu, L., Gui, W., & Zhu, G. (2015). Exposure to difenoconazole causes changes of thyroid hormone and gene expression levels in zebrafish larvae. *Environmental Toxicology and Pharmacology*, 40(3), 983-987. <https://doi.org/10.1016/j.etap.2015.10.005>
- Lopez-Antia, A., Ortiz-Santaliestra, M. E., Mougeot, F., & Mateo, R. (2013). Experimental exposure of red-legged partridges (*Alectoris rufa*) to seeds coated with imidacloprid, thiram and difenoconazole. *Ecotoxicology*, 22(1), 125-138. <https://doi.org/10.1007/s10646-012-1009-x>
- Lopez-Antia, A., Feliu, J., Camarero, P. R., Ortiz-Santaliestra, M. E., & Mateo, R. (2016). Risk assessment of pesticide seed treatment for farmland birds using refined field data. *Journal of Applied Ecology*, 53(5), 1373-1381. <https://doi.org/10.1111/1365-2664.12668>
- Lopez-Antia, A., Ortiz-Santaliestra, M. E., Mougeot, F., Camarero, P. R., & Mateo, R. (2018). Brood size is reduced by half in birds feeding on flutriafol-treated seeds below the recommended application rate. *Environmental Pollution*, 243, 418-426. <https://doi.org/10.1016/j.envpol.2018.08.078>
- Lopez-Antia, A., Ortiz-Santaliestra, M. E., Mougeot, F., Camarero, P. R., & Mateo, R. (2021). Birds feeding on tebuconazole treated seeds have reduced breeding output. *Environmental Pollution*, 271, 116292. <https://doi.org/10.1016/j.envpol.2020.116292>
- Machado-Neves, M., Neto, M. J. O., Miranda, D. C., Souza, A. C. F., Castro, M. M., Sertorio, M. N., Carvalho, T. F., Matta, S. L. P., & Freitas, M. B. (2018). Dietary Exposure to Tebuconazole Affects Testicular and Epididymal Histomorphometry in Frugivorous Bats. *Bulletin of Environmental Contamination and Toxicology*, 101(2), 197-204. <https://doi.org/10.1007/s00128-018-2377-6>
- McNabb, F. M. A. (2007). The Hypothalamic-Pituitary-Thyroid (HPT) Axis in Birds and Its Role in Bird Development and Reproduction. *Critical Reviews in Toxicology*, 37(1-2), 163-193. <https://doi.org/10.1080/10408440601123552>
- Mohanty, B., Pandey, S. P., & Tsutsui, K. (2017). Thyroid disrupting pesticides impair the hypothalamic-pituitary-testicular axis of a wildlife bird, *Amandava amandava*. *Reproductive Toxicology*, 71, 32-41. <https://doi.org/10.1016/j.reprotox.2017.04.006>
- Nolan, V., Ketterson, E. D., Ziegenfus, C., Cullen, D. P., & Chandler, C. R. (1992). Testosterone and Avian Life Histories : Effects of Experimentally Elevated Testosterone on Prebasic Molt and Survival in Male Dark-Eyed Juncos. *The Condor*, 94(2), 364-370. <https://doi.org/10.2307/1369209>
- Ortiz-Santaliestra, M. E., Alcaide, V., Camarero, P. R., Mateo, R., & Mougeot, F. (2020). Egg overspray with herbicides and fungicides reduces survival of red-legged partridge chicks. *Environmental Science & Technology*, 54(19), 12402-12411. <https://doi.org/10.1021/acs.est.0c04203>
- Pandey, S. P., & Mohanty, B. (2015). The neonicotinoid pesticide imidacloprid and the dithiocarbamate fungicide mancozeb disrupt the pituitary–thyroid axis of a wildlife bird. *Chemosphere*, 122, 227-234. <https://doi.org/10.1016/j.chemosphere.2014.11.061>
- Pelosi, C., Thiel, P., Bart, S., Amossé, J., Jean-Jacques, J., Thoisy, J.-C., & Crouzet, O. (2021). The contributions of enchytraeids and earthworms to the soil mineralization process in soils with fungicide. *Ecotoxicology*, 30(9), 1910-1921. <https://doi.org/10.1007/s10646-021-02452-z>
- Pereira, V. R., Pereira, D. R., de Melo Tavares Vieira, K. C., Ribas, V. P., Constantino, C. J. L., Antunes, P. A., & Favareto, A. P. A. (2019). Sperm quality of rats exposed to difenoconazole using classical parameters and surface-enhanced Raman scattering: classification performance by machine learning methods. *Environmental Science and Pollution Research*, 26, 35253-35265. <https://doi.org/10.1007/s11356-019-06407-0>
- Pérez, J. H., Meddle, S. L., Wingfield, J. C., & Ramenofsky, M. (2018). Effects of thyroid hormone manipulation on pre-nuptial molt, luteinizing hormone and testicular growth in male white-crowned sparrows (*Zonotrichia leucophrys gambelii*). *General and Comparative Endocrinology*, 255, 12-18. <https://doi.org/10.1016/j.ygcen.2017.09.025>
- Pirahanchi, Y., Toro, F., & Jialal, I. (2018). Physiology, Thyroid Stimulating Hormone. In *StatPearls*. StatPearls Publishing. <http://www.ncbi.nlm.nih.gov/books/NBK499850/>
- Poulsen, R., Luong, X., Hansen, M., Styrihave, B., & Hayes, T. (2015). Tebuconazole disrupts steroidogenesis in *Xenopus laevis*. *Aquatic Toxicology*, 168, 28-37. <https://doi.org/10.1016/j.aquatox.2015.09.008>

- Raby, M., Maloney, E., Poirier, D. G., & Sibley, P. K. (2019). Acute Effects of Binary Mixtures of Imidacloprid and Tebuconazole on 4 Freshwater Invertebrates. *Environmental Toxicology and Chemistry*, 38(5), 1093-1103. <https://doi.org/10.1002/etc.4386>
- Ribas e Ribas, A. D., Spolti, P., Del Ponte, E. M., Donato, K. Z., Schrekker, H., & Fuentefria, A. M. (2016). Is the emergence of fungal resistance to medical triazoles related to their use in the agroecosystems? A mini review. *Brazilian Journal of Microbiology*, 47(4), 793-799. <https://doi.org/10.1016/j.bjm.2016.06.006>
- Rico, A., Sabater, C., & Castillo, M.-Á. (2016). Lethal and sub-lethal effects of five pesticides used in rice farming on the earthworm *Eisenia fetida*. *Ecotoxicology and Environmental Safety*, 127, 222-229. <https://doi.org/10.1016/j.ecoenv.2016.02.004>
- Robinson, R. A. (2004). The diet of seed-eating birds on lowland farmland. *British Birds*.
- Rokbani, O., Fattouch, S., Chakir, A., & Roth, E. (2019). Heterogeneous oxidation of two triazole pesticides (diniconazole and tebuconazole) by OH-radicals and ozone. *Science of The Total Environment*, 694, 133745. <https://doi.org/10.1016/j.scitotenv.2019.133745>
- Sancho, E., Villarroel, M. J., Fernández, C., Andreu, E., & Ferrando, M. D. (2010). Short-term exposure to sublethal tebuconazole induces physiological impairment in male zebrafish (*Danio rerio*). *Ecotoxicology and Environmental Safety*, 73(3), 370-376. <https://doi.org/10.1016/j.ecoenv.2009.09.020>
- Saxena, A. K., Devillers, J., Bhunia, S. S., & Bro, E. (2015). Modelling inhibition of avian aromatase by azole pesticides. *SAR and QSAR in Environmental Research*, 26(7-9), 757-782. <https://doi.org/10.1080/1062936X.2015.1090749>
- Siefferman, L., Liu, M., Navara, K. J., Mendonça, M. T., & Hill, G. E. (2013). Effect of Prenatal and Natal Administration of Testosterone on Production of Structurally Based Plumage Coloration. *Physiological and Biochemical Zoology*, 86(3), 323-332. <https://doi.org/10.1086/670383>
- Stoehr, A. M., & Hill, G. E. (2001). The effects of elevated testosterone on plumage hue in male House Finches. *Journal of Avian Biology*, 32(2), 153-158. <https://doi.org/10.1034/j.1600-048X.2001.320208.x>
- Stoker, T. E., Ferrell, J. M., Laws, S. C., Cooper, R. L., & Buckalew, A. (2006). Evaluation of ammonium perchlorate in the endocrine disruptor screening and testing program's male pubertal protocol : Ability to detect effects on thyroid endpoints. *Toxicology*, 228(1), 58-65. <https://doi.org/10.1016/j.tox.2006.08.026>
- Strange, R. N., & Scott, P. R. (2005). Plant Disease : A Threat to Global Food Security. *Annual Review of Phytopathology*, 43(1), 83-116. <https://doi.org/10.1146/annurev.phyto.43.113004.133839>
- Swaddle, J. P., Witter, M. S., Cuthill, I. C., Budden, A., & McCowen, P. (1996). Plumage Condition Affects Flight Performance in Common Starlings: Implications for Developmental Homeostasis, Abrasion and Moulting. *Journal of Avian Biology*, 27(2), 103. <https://doi.org/10.2307/3677139>
- Swaileh, K. M., & Sansur, R. (2006). Monitoring urban heavy metal pollution using the House Sparrow (*Passer domesticus*). *Journal of Environmental Monitoring*, 8(1), 209-213. DOI: [10.1039/B510635D](https://doi.org/10.1039/B510635D)
- Taxvig, C., Hass, U., Axelstad, M., Dalgaard, M., Boberg, J., Andeasen, H. R., & Vinggaard, A. M. (2007). Endocrine-Disrupting Activities In Vivo of the Fungicides Tebuconazole and Epoxiconazole. *Toxicological Sciences*, 100(2), 464-473. <https://doi.org/10.1093/toxsci/kfm227>
- Tokumoto, J., Danjo, M., Kobayashi, Y., Kinoshita, K., Omotehara, T., Tatsumi, A., Hashiguchi, M., Sekijima, T., Kamisoyama, H., Yokoyama, T., Kitagawa, H., & Hoshi, N. (2013). Effects of Exposure to Clothianidin on the Reproductive System of Male Quails. *Journal of Veterinary Medical Science*, 75(6), 755-760. <https://doi.org/10.1292/jvms.12-0544>
- Vézina, F., Gustowska, A., Jalvingh, K. M., Chastel, O., & Piersma, T. (2009). Hormonal correlates and thermoregulatory consequences of molting on metabolic rate in a northerly wintering shorebird. *Physiological and biochemical zoology*, 82(2), 129-142.
- Voitkevich, A. A. E. (1966). The feathers and plumage of birds. *The feathers and plumage of birds*.
- Warrilow, A. G., Parker, J. E., Kelly, D. E., & Kelly, S. L. (2013). Azole Affinity of Sterol 14 α -Demethylase (CYP51) Enzymes from *Candida albicans* and *Homo sapiens*. *Antimicrobial Agents and Chemotherapy*, 57(3), 1352-1360. <https://doi.org/10.1128/AAC.02067-12>
- Wilson, A. C., & Farner, D. S. (1960). The Annual Cycle of Thyroid Activity in White-Crowned Sparrows of Eastern Washington. *The Condor*, 62(6), 414-425. <https://doi.org/10.2307/1365589>

- Yang, H., Zhang, W., Kong, Q., Liu, H., Sun, R., Lin, B., Zhang, H., & Xi, Z. (2013). Effects of pubertal exposure to thiazole-Zn on thyroid function and development in female rats. *Food and Chemical Toxicology*, *53*, 100-104. <https://doi.org/10.1016/j.fct.2012.11.003>
- Yang, J.-D., Liu, S.-H., Liao, M.-H., Chen, R.-M., Liu, P.-Y., & Ueng, T.-H. (2018). Effects of tebuconazole on cytochrome P450 enzymes, oxidative stress, and endocrine disruption in male rats. *Environmental Toxicology*, *33*(8), 899-907. <https://doi.org/10.1002/tox.22575>
- Yu, L., Chen, M., Liu, Y., Gui, W., & Zhu, G. (2013). Thyroid endocrine disruption in zebrafish larvae following exposure to hexaconazole and tebuconazole. *Aquatic Toxicology*, *138-139*, 35-42. <https://doi.org/10.1016/j.aquatox.2013.04.001>
- Zarn, J. A., Brüschweiler, B. J., & Schlatter, J. R. (2003). Azole fungicides affect mammalian steroidogenesis by inhibiting sterol 14 alpha-demethylase and aromatase. *Environmental Health Perspectives*, *111*(3), 255-261. <https://doi.org/10.1289/ehp.5785>
- Zhang, W., Chen, L., Xu, Y., Deng, Y., Zhang, L., Qin, Y., Wang, Z., Liu, R., Zhou, Z., & Diao, J. (2019). Amphibian (*Rana nigromaculata*) exposed to cyproconazole : Changes in growth index, behavioral endpoints, antioxidant biomarkers, thyroid and gonad development. *Aquatic Toxicology*, *208*, 62-70. <https://doi.org/10.1016/j.aquatox.2018.12.015>
- Zubrod, J. P., Bundschuh, M., Arts, G., Brühl, C. A., Imfeld, G., Knäbel, A., Payraudeau, S., Rasmussen, J. J., Rohr, J., Scharmüller, A., Smalling, K., Stehle, S., Schulz, R., & Schäfer, R. B. (2019). Fungicides : An Overlooked Pesticide Class? *Environmental Science & Technology*, *53*(7), 3347-3365. <https://doi.org/10.1021/acs.est.8b04392>

Statements & Declarations

Acknowledgments and Funding

We thank E. Beaugeard, E. Daniaud, B. Mohring, P. Pardon, L. Lièvre and N. Imbert for their help during the course of this study. This research has been funded by the Agence Nationale de la Recherche (ANR project VITIBIRD ANR-20-CE34-0008 to F.A.), by the Région Nouvelle-Aquitaine (MULTISTRESS project), by the Agence nationale de sécurité sanitaire de l'alimentation, de l'environnement et du travail (ANSES project BiodiTox #2019/1/031 to F.B.), by the Centre National de la Recherche Scientifique and by MITI-PEPSAN (mission for transversal and interdisciplinary initiatives to C.F.). LIENS laboratory of the University of La Rochelle for their technical support. The platform PLATINE (<https://www.u-bordeaux.fr/Recherche/Plateformes/Sciences-de-l-environnement/Plateforme-de-chimie-analytique-organique-environnementale-PLATINE>) of the University of Bordeaux and UMR EPOC is acknowledged for technical and analytical support.

Competing Interests

The authors have no relevant financial or non-financial interests to disclose.

Credit author statement

Pauline Bellot: Data collection, Data curation, Formal analysis, Writing – Original Draft, Visualization. **François Brischoux:** Conceptualization, Data collection, Validation, Writing – Review & Editing, Funding acquisition. **Hélène Budzinski:** Data collection. **Sophie Dupont:** Writing – Review & Editing. **Clémentine Fritsch:** Conceptualization, Writing – Review & Editing, Funding acquisition. **Sydney Hope:** Data collection, Writing – Review & Editing. **Bruno Michaud:** Data collection, Data curation. **Marie Pallud:** Data collection, Data curation. **Charline Parenteau:** Data collection, Data curation. **Louise Prouteau:** Data collection, Data curation. **Steffi Rocchi:** Writing – Review & Editing. **Frédéric Angelier:** Conceptualization, Data collection, Data curation, Validation, Ressources, Writing – Review & Editing, Supervision, Funding acquisition.