



**HAL**  
open science

# On a Galois property of fields generated by the torsion of an abelian variety

Sara Checcoli, Gabriel A. Dill

► **To cite this version:**

Sara Checcoli, Gabriel A. Dill. On a Galois property of fields generated by the torsion of an abelian variety. 2023. hal-04148573

**HAL Id: hal-04148573**

**<https://hal.science/hal-04148573>**

Preprint submitted on 3 Jul 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# On a Galois property of fields generated by the torsion of an abelian variety

S. Checcoli and G. A. Dill

June 22, 2023

## Abstract

In this article, we study a certain Galois property of subextensions of  $k(A_{\text{tors}})$ , the minimal field of definition of all torsion points of an abelian variety  $A$  defined over a number field  $k$ . Concretely, we show that each subfield of  $k(A_{\text{tors}})$  which is Galois over  $k$  (of possibly infinite degree) and whose Galois group has finite exponent is contained in an abelian extension of some finite extension of  $k$ . As an immediate corollary of this result and a theorem of Bombieri and Zannier, we deduce that each such field has the Northcott property, *i.e.* does not contain any infinite set of algebraic numbers of bounded height.

## 1 Introduction

In [BZ01], Bombieri and Zannier introduce two properties of fields of algebraic numbers with respect to the absolute logarithmic Weil height (see [BG06, bottom of p. 16] for a definition).

Throughout this article, we let  $\bar{\mathbb{Q}}$  be a fixed, once and for all, algebraic closure of  $\mathbb{Q}$  and all algebraic extensions of  $\mathbb{Q}$  that appear will be assumed to be subfields of  $\bar{\mathbb{Q}}$ . We say that an algebraic extension  $F/\mathbb{Q}$  has the *Bogomolov property (B)* if there exists a constant  $c = c(F) > 0$  such that all elements of  $F \setminus \{0\}$  which are not roots of unity have height at least  $c$ . We say that  $F$  has the *Northcott property (N)* if it does not contain any infinite set of algebraic numbers of bounded height.

By Kronecker's theorem (see [BG06, Theorem 1.5.9]), property (N) implies property (B) and by Northcott's theorem (see [Nor49, Theorem 1] and [BG06, Theorem 1.6.8]), every number field  $L$  has both property (N) as well as property (B). In the last decades, there has been a lot of work

around deciding the validity of these properties for infinite extensions of  $\mathbb{Q}$ .

In particular, property (B) holds for any subfield of  $k^{\text{ab}}$ , the maximal abelian extension of a number field  $k$ : after a question posed by Zannier, this was shown in [AD00] for  $k = \mathbb{Q}$ , and then worked out for general number fields in [AZ00], and further in [AZ10]. The result was later generalised by the main theorem in [ADZ14], which together with [Che13, Theorem 1.2] implies, in particular, that property (B) is satisfied by any Galois extension  $F/k$  of a number field  $k$  such that the quotient of  $\text{Gal}(F/k)$  by its center has finite exponent, *i.e.* such that the orders of the elements of this group are finite and uniformly bounded.

In [Hab13], Habegger proved an elliptic analogue of the above-mentioned result in [AD00] by showing that property (B) holds for  $\mathbb{Q}(E_{\text{tors}})$ , the smallest subfield of  $\bar{\mathbb{Q}}$  over which all torsion points of an elliptic curve  $E$ , defined over  $\mathbb{Q}$ , are defined. We remark that, if  $E$  has no complex multiplication, the extension  $\mathbb{Q}(E_{\text{tors}})/\mathbb{Q}$  is very far from being abelian.

The lower bound for the height of non-roots of unity in  $\mathbb{Q}(E_{\text{tors}}) \setminus \{0\}$  was later made explicit in [Fre21]. Moreover, in [Fre22], property (B) was proved to hold, under the same assumptions on  $E$ , for extensions of the form  $F(E_{\text{tors}})$  for some class of possibly infinite Galois extensions  $F/\mathbb{Q}$ .

It is an open problem to decide whether, given a general abelian variety  $A$  of dimension at least 2 defined over a number field  $k$ , property (B) holds for  $k(A_{\text{tors}})$ , the smallest subfield of  $\bar{\mathbb{Q}}$  containing  $k$  over which all torsion points of  $A$  are defined.

As for property (N), it was for instance proved in [BZ01] to hold for the field  $k_{\text{ab}}^{(d)}$ , which is the compositum of all abelian extensions of  $k$  of degree at most  $d$ . Clearly,  $k_{\text{ab}}^{(d)}/k$  is an abelian extension of  $k$  such that  $\text{Gal}(k_{\text{ab}}^{(d)}/k)$  has finite exponent. Using the proof of [CZ11, Proposition 2.1], one can show that any Galois extension  $F/k$  with  $F \subseteq k^{\text{ab}}$  and such that  $\text{Gal}(F/k)$  has finite exponent is contained in  $k_{\text{ab}}^{(d)}$  for some positive integer  $d$ . So, in particular, property (N) holds for such extensions.

It is an open problem whether  $\text{Gal}(F/k)$  having finite exponent is enough to ensure that the field  $F$  has property (N); for instance, it is unknown whether property (N) holds for the compositum  $k^{(d)}$  of all extensions of  $k$  of degree at most  $d$  when  $d \geq 3$ . We note that  $k_{\text{ab}}^{(d)}$  is the maximal subextension of  $k^{(d)}$  which is abelian over  $k$  (as originally defined in [BZ01]).

In this setting, it is quite natural to ask about the analogue of the above-mentioned result of [BZ01] for abelian varieties:

**Question 1.** *Let  $A$  be an abelian variety defined over a number field  $k$  and let  $k(A_{\text{tors}})$  be the smallest subfield of  $\bar{\mathbb{Q}}$  containing  $k$  over which all torsion*

points of  $A$  are defined. Is it true that if  $F \subseteq k(A_{\text{tors}})$ ,  $F/k$  is Galois, and  $\text{Gal}(F/k)$  has finite exponent, then  $F$  has property (N)?

In this article, we answer Question 1 in the affirmative by showing that, although for an abelian variety  $A$  over a number field  $k$ , the fields  $k^{\text{ab}}$  and  $k(A_{\text{tors}})$  in general have very different Galois groups, their Galois subextensions whose Galois groups have finite exponent look the same up to replacing  $k$  by another number field  $M$  containing  $k$ . Our main theorem is the following:

**Theorem 1.** *Let  $A$  be an abelian variety defined over a number field  $k$ . Let  $e \geq 1$  be an integer. Then there exists a number field  $M$  containing  $k$  and depending only on  $A$ ,  $k$ , and  $e$  such that for any subfield  $F \subseteq k(A_{\text{tors}})$  with  $F/k$  Galois and  $\text{Gal}(F/k)$  of exponent at most  $e$ , one has  $F \subseteq M^{\text{ab}}$ .*

We now briefly sketch the proof of Theorem 1, which is carried out in Section 3. Notice that it suffices to prove Theorem 1 if the extension  $F/k$  is finite and therefore contained in some  $k(A[n])$  where  $A[n]$  denotes the kernel of multiplication by  $n$  on  $A(\bar{\mathbb{Q}})$ . In Subsection 3.1, we show furthermore how to reduce the proof of Theorem 1 to the construction of two subgroups  $W'_n$  and  $W''_n$  of  $\text{Gal}(k(A[n])/k)$  satisfying certain conditions. As we recall in Proposition 1 in Section 2, up to replacing  $k$  by a fixed finite extension, we can write  $\text{Gal}(k(A[n])/k)$  canonically as the product of the Galois groups  $\text{Gal}(k(A[p^t])/k)$  for  $p^t$  exactly dividing  $n$ . Moreover, for each prime  $p$ , the Galois representation on the  $p$ -adic Tate module of  $A$  yields an algebraic group  $\mathcal{G}_p$  over  $\mathbb{Z}_p$ , which has nice properties for all primes larger than some constant  $c$ . We then construct  $W'_n$  and  $W''_n$  in such a way that we can check the required conditions working “prime by prime”. For small primes  $p$  (i.e. those less than or equal to  $\max\{c, e\}$ ), we use the  $p$ -adic exponential map associated to  $\mathcal{G}_p$  while for large primes  $p$ , we use a group-theoretic property of  $\mathcal{G}_p(\mathbb{F}_p)$  (see Proposition 2 in Section 2).

The following result, which finally answers Question 1, is an immediate corollary of the result in [BZ01] together with Theorem 1 and is proved in Section 4:

**Corollary 1.** *Let  $G = \mathbb{G}_m$  or  $G = A$  with  $A$  an abelian variety defined over a number field  $k$ . Let  $k(G_{\text{tors}})$  denote the smallest subfield of  $\bar{\mathbb{Q}}$  containing  $k$  over which every torsion point of  $G$  is defined. Then every subfield of  $k(G_{\text{tors}})$  which is Galois over  $k$  and whose Galois group has finite exponent has property (N).*

## 2 Preliminaries on abelian varieties

Let  $A$  be an abelian variety defined over a number field  $K \subseteq \bar{\mathbb{Q}}$ .

We fix some notation. Let  $K_n = K(A[n])$  be the smallest subfield of  $\bar{\mathbb{Q}}$  containing  $K$  over which every point of  $A[n]$ , the group of torsion points of  $A$  of order dividing  $n$ , is defined. Then the extension  $K_n/K$  is finite and Galois. Set  $g = \dim A$  and  $G_n = \text{Gal}(K_n/K)$ , which we identify with a subgroup of  $\text{GL}_{2g}(\mathbb{Z}/n\mathbb{Z})$  after having fixed once and for all a compatible system of isomorphisms  $A[n] \simeq (\mathbb{Z}/n\mathbb{Z})^{2g}$ .

For a prime number  $p$ , we consider the representation  $\rho_p : \text{Gal}(\bar{\mathbb{Q}}/K) \rightarrow \text{GL}_{2g}(\mathbb{Z}_p)$ , coming from the  $p$ -adic Tate module

$$\varprojlim A[p^n] \simeq \varprojlim (\mathbb{Z}/p^n\mathbb{Z})^{2g} \simeq \mathbb{Z}_p^{2g}$$

of  $A$ . We let  $\mathcal{G}_p$  denote the Zariski closure of  $\rho_p(\text{Gal}(\bar{\mathbb{Q}}/K))$  in  $\text{GL}_{2g, \mathbb{Z}_p}$ , regarded as an algebraic group over  $\mathbb{Z}_p$ .

Set now  $K_{p^\infty} = \bigcup_{n \geq 0} K_{p^n}$ , it is a Galois extension of  $K$ . Let  $G_{p^\infty}$  denote its Galois group over  $K$ . Note that  $\rho_p$  factors through the restriction homomorphism  $\text{Gal}(\bar{\mathbb{Q}}/K) \rightarrow G_{p^\infty}$ . Furthermore, the induced homomorphism  $G_{p^\infty} \rightarrow \text{GL}_{2g}(\mathbb{Z}_p)$  is injective and we identify  $G_{p^\infty}$  with its image in  $\text{GL}_{2g}(\mathbb{Z}_p)$ .

We collect here some well-known important classical results from the theory of Galois representations attached to abelian varieties over number fields. Recall that a smooth algebraic group  $G$  over  $\mathbb{Z}_p$  is called *reductive* if both  $G \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$  and  $G \otimes_{\mathbb{Z}_p} \mathbb{F}_p$  are reductive algebraic groups in the usual sense.

**Proposition 1.** *Let  $A$  be an abelian variety defined over a number field  $k$ . There exist a finite extension  $K/k$  and a constant  $c = c(A, K) > 0$  such that, with notation as above, the following properties hold true:*

- (i) *for any coprime natural numbers  $n'$  and  $n''$ , the canonical homomorphism  $G_{n'n''} \rightarrow G_{n'} \times G_{n''}$  is an isomorphism,*
- (ii) *the algebraic group  $\mathcal{G}_p$  over  $\mathbb{Z}_p$  is connected for every prime  $p$ ,*
- (iii)  *$\mathcal{G}_p$  is smooth and reductive for every prime  $p > c$ , and*
- (iv) *for every prime  $p$ , the index  $[\mathcal{G}_p(\mathbb{Z}_p) : \rho_p(\text{Gal}(\bar{\mathbb{Q}}/K))]$  is bounded from above by  $c$ .*

*Proof.* Property (i) follows from [Ser13, Théorème 1 and §3.1] (see also [Ser86]), while (ii) is a result from [Ser81]. Property (iii) follows from [Win02, Théorème 1 and §2.1] while the proof of (iv) is detailed in [Zyw19, Theorem 1.2 (a), Remark 1.3, and §6] and is based on, amongst others, results in [Win02] and [Ser86].  $\square$

*Remark 2.* Our proof does not use any properties of abelian varieties over number fields other than that Proposition 1 holds for the associated  $p$ -adic Galois representations: we can start with any system of continuous

Galois representations  $\rho_p : \text{Gal}(\bar{\mathbb{Q}}/K) \rightarrow \text{GL}_{g(p)}(\mathbb{Z}_p)$  ( $p$  prime,  $g(p) \in \mathbb{N} = \{1, 2, \dots\}$ ) and define  $K_n$  as the fixed field of  $\bigcap_{i=1}^r \ker(\pi_{p_i, k_i} \circ \rho_{p_i})$  where  $n = \prod_{i=1}^r p_i^{k_i}$  is the prime factorization of  $n \in \mathbb{N}$  and  $\pi_{p_i, k_i} : \text{GL}_{g(p_i)}(\mathbb{Z}_{p_i}) \rightarrow \text{GL}_{g(p_i)}(\mathbb{Z}/p_i^{k_i}\mathbb{Z})$  is the canonical projection ( $i = 1, \dots, r$ ). Then, if the analogue of Proposition 1 holds in this situation, our proof will go through and show that any Galois subextension of  $\bigcup_{n=1}^{\infty} K_n$  whose Galois group has finite exponent is contained in the maximal abelian extension of some number field.

We will need the following fact, which is certainly well-known to experts. We give a proof which is heavily modeled after two answers to a question on MathOverflow (see [Hum13, Mar13]) and uses, for instance, classical results in [BT71, Lan56, Ste68].

**Proposition 2.** *Let  $q$  be a power of a prime  $p$  and let  $G$  be a smooth, connected, reductive algebraic group over the finite field  $\mathbb{F}_q$ . Let  $G(\mathbb{F}_q)^+$  be the normal subgroup of  $G(\mathbb{F}_q)$  generated by its  $p$ -Sylow subgroups. Then the quotient  $G(\mathbb{F}_q)/G(\mathbb{F}_q)^+$  is abelian.*

*Proof.* Let  $H$  be an arbitrary smooth, connected, reductive algebraic group over  $\mathbb{F}_q$ . We define  $H(\mathbb{F}_q)^+$  analogously to  $G(\mathbb{F}_q)^+$ . By [Mil17, Theorem 14.5 and Proposition 14.14], a subgroup  $\Gamma$  of  $H(\mathbb{F}_q)$  is equal to the  $\mathbb{F}_q$ -points of a unipotent algebraic subgroup  $U$  of  $H$  if and only if the order of  $\Gamma$  is a power of  $p$ . By [BT71, Corollaire 3.7],  $H(\mathbb{F}_q)^+$  is then the subgroup of  $H(\mathbb{F}_q)$  generated by the  $U(\mathbb{F}_q)$ 's, where  $U$  varies through the unipotent radicals of the minimal parabolic subgroups of  $H$  that are defined over  $\mathbb{F}_q$ . Note that the minimal parabolic subgroups of  $H$  are precisely the Borel subgroups of  $H$  by [Mil17, Aside 17.73 and Proposition 17.99], where the latter proposition is a consequence of the main result in [Lan56]. In particular,  $H$  has a Borel subgroup. Note that  $k$  should be assumed to be finite in [Mil17, Proposition 17.99] and the morphism  $\sigma$  that occurs in the proof of [Mil17, Proposition 17.99] is the Frobenius endomorphism relative to  $k$ .

Let  $G'$  denote the derived subgroup of  $G$ , which is a semisimple algebraic group over  $\mathbb{F}_q$ , and let  $\omega : \tilde{G}' \rightarrow G'$  denote the universal cover of  $G'$  (as an algebraic group; for the existence, see [Mil17, Remark 18.27]). Then  $\tilde{G}'$  is also semisimple by [Mil17, Lemma 19.14]. Since  $G/G'$  is a torus by [Mil17, Proposition 12.46 and Corollary 21.50], the preceding paragraph together with [Mil17, Corollary 14.18] implies that  $G(\mathbb{F}_q)^+ = G'(\mathbb{F}_q)^+$ . By [Ive76, Lemma 2.5], the preimage of a Borel subgroup  $B$  of  $G'$  under  $\omega$  is a Borel subgroup  $\tilde{B}$  of  $\tilde{G}'$  (Iversen works over an algebraically closed field, but we can use [Mil17, 17.66] to descend to  $\mathbb{F}_q$ ). Let  $\tilde{U}$  denote the unipotent radical of  $\tilde{B}$  and let  $U$  denote the unipotent radical of  $B$ . We have that  $\tilde{U} \subseteq \omega^{-1}(U)$  since  $B/U$  is of multiplicative type by [Mil17,

Theorem 16.33]. Thus,  $\omega^{-1}(U)/\tilde{U} \subseteq \tilde{B}/\tilde{U}$  is of multiplicative type while  $\omega^{-1}(U)/\ker \omega \simeq U$  is unipotent. By [Mil17, Lemma 16.44], the multiplication morphism  $(\ker \omega) \times \tilde{U} \rightarrow \omega^{-1}(U)$  is an isomorphism. It follows that  $\omega|_{\tilde{U}} : \tilde{U} \rightarrow U$  is an isomorphism and so  $U(\mathbb{F}_q) = \omega(\tilde{U}(\mathbb{F}_q))$ . This implies that the map  $\tilde{G}'(\mathbb{F}_q)^+ \rightarrow G'(\mathbb{F}_q)^+$  is surjective. Thus, it suffices to show that  $G(\mathbb{F}_q)/\omega(\tilde{G}'(\mathbb{F}_q)^+)$  is abelian.

For an algebraic group  $H$  over  $\mathbb{F}_q$ , we denote its center by  $Z(H)$ . The commutator morphism  $G \times G \rightarrow G$  factors through the projection  $G \times G \rightarrow (G/Z(G)) \times (G/Z(G))$ . At the same time, we have a chain of canonical isomorphisms  $\tilde{G}'/Z(\tilde{G}') \simeq G'/Z(G') \simeq G/Z(G)$ , the first of which is induced by  $\omega$ , by [Mil17, Example 19.25 and Remark 19.30] (the universal covering of  $G'^{\text{ad}} := G'/Z(G')$  is the composition of the universal covering of  $G'$  with the projection  $G' \rightarrow G'^{\text{ad}}$  because of [Mil17, Proposition 18.2]). This implies that the commutator morphism  $G \times G \rightarrow G$  decomposes as the induced morphism  $G \times G \rightarrow (\tilde{G}'/Z(\tilde{G}')) \times (\tilde{G}'/Z(\tilde{G}'))$  followed by first the morphism  $(\tilde{G}'/Z(\tilde{G}')) \times (\tilde{G}'/Z(\tilde{G}')) \rightarrow \tilde{G}'$  that is induced by the commutator morphism of  $\tilde{G}'$  and then  $\omega$ . It follows that  $\omega(\tilde{G}'(\mathbb{F}_q))$  contains the commutator subgroup of  $G(\mathbb{F}_q)$ .

Combining the two preceding paragraphs shows that it suffices to prove that  $\tilde{G}'(\mathbb{F}_q)^+ = \tilde{G}'(\mathbb{F}_q)$ .

By [Mil17, Remark 18.27],  $\tilde{G}'$  is still simply connected when base changed to an algebraic closure  $\bar{\mathbb{F}}_q$  of  $\mathbb{F}_q$  and, by [Mil17, Proposition 23.59], over  $\bar{\mathbb{F}}_q$ , the definitions of simply connectedness in [Mil17, Definition 18.5] and [Ste68, §6.4] are equivalent. We can then apply [Ste68, Theorem 12.4] to  $\tilde{G}'$  over  $\bar{\mathbb{F}}_q$  with  $\sigma$  chosen to be the Frobenius endomorphism relative to  $\mathbb{F}_q$  to conclude. □

## 3 Proof of Theorem 1

### 3.1 Some reduction steps

Let  $F/k$  be a (possibly infinite) Galois extension with  $F \subseteq k(A_{\text{tors}})$  and  $\text{Gal}(F/k)$  having finite exponent  $\exp(\text{Gal}(F/k)) \leq e$  for some positive integer  $e \geq 1$ . As  $F$  equals the compositum of all its subextensions which are finite and Galois over  $k$ , it is enough to prove that Theorem 1 holds true when  $[F : k]$  is finite.

So, let  $F/k$  be finite and Galois with  $\text{Gal}(F/k)$  of exponent at most  $e$  and suppose  $F \subseteq k(A_{\text{tors}})$ .

Let  $K/k$  be the finite extension from Proposition 1 and let  $L = KF \subseteq$

$K(A_{\text{tors}})$  be the compositum of  $F$  and  $K$ . Note that

$$\text{Gal}(L/K) \simeq \text{Gal}(F/F \cap K) \leq \text{Gal}(F/k),$$

hence  $\exp(\text{Gal}(L/K)) \leq e$ .

We are going to prove that  $L$  is contained in the maximal abelian extension of a fixed number field containing  $K$  and depending only on  $A$ ,  $K$ , and  $e$ . This of course will prove Theorem 1 for the extension  $F/k$ .

As before, we denote by  $K_n = K(A[n])$  the smallest subfield of  $\bar{\mathbb{Q}}$  containing  $K$  over which every point of  $A[n]$ , the group of all torsion points of  $A$  of order dividing  $n$ , is defined. We set  $g = \dim A$  and  $G_n = \text{Gal}(K_n/K)$ , which we identify with a subgroup of  $\text{GL}_{2g}(\mathbb{Z}/n\mathbb{Z})$ , having fixed once and for all a compatible system of isomorphisms  $A[n] \simeq (\mathbb{Z}/n\mathbb{Z})^{2g}$ .

We fix an integer  $n$  such that  $L \subseteq K_n$  and we set  $H_n = \text{Gal}(K_n/L) \leq G_n$ . Our goal is to prove that there exists a positive integer  $m = m(A, K, e)$  and subgroups  $W'_n$  and  $W''_n$  of  $G_n$  such that:

- (a)  $W''_n \trianglelefteq G_n$  and  $G_n/W''_n$  is abelian,
- (b)  $W'_n$  contains the kernel of the projection from  $G_n$  onto  $G_{\text{gcd}(n,m)}$ , and
- (c)  $W'_n \cap W''_n \subseteq H_n$ .

From these properties it follows that

$$L = K_n^{H_n} \subseteq K_n^{W'_n \cap W''_n} = K_n^{W'_n} K_n^{W''_n}$$

where we use the usual Galois-theoretic notation for the fixed field of a subgroup of the Galois group. We conclude that

$$L \subseteq K_m K^{\text{ab}} \subseteq K_m^{\text{ab}}$$

and the statement of Theorem 1 holds with  $M = K_m$ .

## 3.2 Construction of the auxiliary subgroups

We let  $c = c(A, K)$  be the constant from Proposition 1 and set

$$\delta = e!c!$$

In the following,  $\mathbb{I}_{2g}$  will denote the  $2g \times 2g$  identity matrix. By abuse of notation, we use the same symbol regardless of the ring of coefficients of the matrix.

We write  $n = n'n''$  where  $n'$  and  $n''$  are the unique coprime positive integers satisfying that  $p \mid n'$  if and only if  $p \mid \text{gcd}(n, \delta)$  for every prime  $p$ .



Writing  $n' = \prod_{i=1}^r p_i^{a_i}$  and  $n'' = \prod_{j=1}^s q_j^{b_j}$  with  $p_i$  and  $q_j$  distinct primes and  $a_i$  and  $b_j$  positive integers ( $i = 1, \dots, r, j = 1, \dots, s$ ), we can apply Proposition 1.(i) to identify

$$G_n = G_{n'} \times G_{n''} = \left( \prod_{i=1}^r G_{p_i^{a_i}} \right) \times \left( \prod_{j=1}^s G_{q_j^{b_j}} \right). \quad (3.1)$$

Consider, under this identification, the subgroups of  $G_n$  given by

$$U'_n = H_n \cap (G_{n'} \times \{(\mathbb{I}_{2g}, \dots, \mathbb{I}_{2g})\})$$

and

$$U''_n = H_n \cap (\{(\mathbb{I}_{2g}, \dots, \mathbb{I}_{2g})\} \times G_{n''})$$

and let  $H'_n$  and  $H''_n$  be their isomorphic images in  $G_{n'}$  and  $G_{n''}$  respectively. Let finally

$$W'_n = H'_n \times G_{n''}$$

and

$$W''_n = G_{n'} \times H''_n.$$

We are going to prove that these groups satisfy properties (a), (b), and (c) from above. This will conclude the proof of Theorem 1.

Note that, under the identification (3.1), one has that the element

$$\gamma = (\gamma'_1, \dots, \gamma'_r, \gamma''_1, \dots, \gamma''_s)$$

of  $G_n$  belongs to  $W'_n \cap W''_n$  if and only if the two elements

$$\gamma' = (\gamma'_1, \dots, \gamma'_r, \mathbb{I}_{2g}, \dots, \mathbb{I}_{2g})$$

and

$$\gamma'' = (\mathbb{I}_{2g}, \dots, \mathbb{I}_{2g}, \gamma''_1, \dots, \gamma''_s)$$

are both in  $H_n$ . As clearly  $\gamma = \gamma' \gamma''$ , we get that  $W'_n \cap W''_n \subseteq H_n$  and so (c) holds.

**Claim 1.** *The group  $W''_n$  satisfies condition (a).*

*Proof of Claim 1.* Since  $L/K$  is Galois, the subgroup  $H_n$  is normal in  $G_n$ , so  $H''_n$  is normal in  $G_{n''}$  and hence  $W''_n$  is normal in  $G_n$ . Note that

$$G_n/W''_n \simeq G_{n''}/H''_n \simeq (\{(\mathbb{I}_{2g}, \dots, \mathbb{I}_{2g})\} \times G_{n''})/U''_n$$

and we are going to prove that the last quotient is abelian.

For  $j = 1, \dots, s$ , set

$$U_{q_j^{b_j}} = U''_n \cap (\{(\mathbb{I}_{2g}, \dots, \mathbb{I}_{2g})\} \times G_{q_j^{b_j}} \times \{(\mathbb{I}_{2g}, \dots, \mathbb{I}_{2g})\}),$$

where of course the only non-trivial factor occurs at the  $(r+j)$ -th position, and let  $H_{q_j}^{b_j}$  denote its isomorphic image in  $G_{q_j}^{b_j}$ . Using the identification (3.1), we have that

$$\{(\mathbb{I}_{2g}, \dots, \mathbb{I}_{2g})\} \times \prod_{j=1}^s H_{q_j}^{b_j} \subseteq U_n''.$$

It follows that  $(\{(\mathbb{I}_{2g}, \dots, \mathbb{I}_{2g})\} \times G_{n''})/U_n''$  is isomorphic to a quotient of

$$(\{(\mathbb{I}_{2g}, \dots, \mathbb{I}_{2g})\} \times \prod_{j=1}^s G_{q_j}^{b_j}) / (\{(\mathbb{I}_{2g}, \dots, \mathbb{I}_{2g})\} \times \prod_{j=1}^s H_{q_j}^{b_j}),$$

which, in turn, is isomorphic to the product

$$\prod_{j=1}^s (G_{q_j}^{b_j} / H_{q_j}^{b_j}).$$

Hence it suffices to show that each factor  $G_{q_j}^{b_j} / H_{q_j}^{b_j}$  is abelian for  $j = 1, \dots, s$ . Note that this group has finite exponent at most  $e$  since it admits an injective homomorphism into the group  $G_n / H_n$  of finite exponent at most  $e$ .

From now on, we fix  $j$  and we write  $q, b$  instead of  $q_j, b_j$  to ease the reading.

Let  $\mathcal{G}_q$  be the algebraic group over  $\mathbb{Z}_q$  defined in Section 2 and consider the homomorphism

$$\varphi : \mathcal{G}_q(\mathbb{Z}/q^b\mathbb{Z}) \rightarrow \mathcal{G}_q(\mathbb{Z}/q\mathbb{Z})$$

given by reduction modulo  $q$ . We have that  $\varphi$  is surjective by Proposition 1.(iii) and Hensel's lemma. Furthermore, the order of the kernel of  $\varphi$  is a power of  $q$ .

Let  $\mathcal{G}_q(\mathbb{Z}/q\mathbb{Z})^+$  denote the normal subgroup of  $\mathcal{G}_q(\mathbb{Z}/q\mathbb{Z})$  that is generated by its  $q$ -Sylow subgroups.

Note that  $\mathcal{G}_q(\mathbb{Z}/q\mathbb{Z})^+$  is generated by elements  $g$  whose order is a power of  $q$  and any lift  $\gamma \in \mathcal{G}_q(\mathbb{Z}/q^b\mathbb{Z})$  of such an element  $g$  will have order equal to a power of  $q$  as well. Proposition 1.(iv) together with Hensel's lemma implies that  $\gamma^{cl} \in G_{q^b}$  for any such  $\gamma$ .

Also, as  $\exp(G_{q^b}/H_{q^b}) \leq e$ ,  $\gamma^\delta \in H_{q^b}$  and finally, as  $\gamma$  has order equal to a power of  $q$  and  $q \nmid \delta$ , we have that  $\gamma \in H_{q^b}$ . It follows that  $\ker \varphi \subseteq H_{q^b}$  and  $\mathcal{G}_q(\mathbb{Z}/q\mathbb{Z})^+ \subseteq \varphi(H_{q^b})$ . Hence, we have that  $\varphi^{-1}(\mathcal{G}_q(\mathbb{Z}/q\mathbb{Z})^+) \subseteq H_{q^b}$ .

Thus  $G_{q^b}/H_{q^b}$  is a quotient of the group  $G_{q^b}/\varphi^{-1}(\mathcal{G}_q(\mathbb{Z}/q\mathbb{Z})^+)$ , which admits an injective homomorphism into the group  $\mathcal{G}_q(\mathbb{Z}/q\mathbb{Z})/\mathcal{G}_q(\mathbb{Z}/q\mathbb{Z})^+$ . This last group is abelian by Proposition 2 and so is therefore  $G_{q^b}/H_{q^b}$ .  $\square$

**Claim 2.** *The group  $W'_n$  satisfies condition (b).*

*Proof of Claim 2.* For  $i = 1, \dots, r$ , set

$$U_{p_i}^{a_i} = U'_n \cap (\{\mathbb{I}_{2g}, \dots, \mathbb{I}_{2g}\} \times G_{p_i}^{a_i} \times \{\mathbb{I}_{2g}, \dots, \mathbb{I}_{2g}\}),$$

where now the only non-trivial factor occurs at the  $i$ -th position, and let  $H_{p_i}^{a_i}$  denote its isomorphic image in  $G_{p_i}^{a_i}$ . Note that, as before, the quotient group  $G_{p_i}^{a_i}/H_{p_i}^{a_i}$  has exponent at most  $e$  since it admits an injective homomorphism into the group  $G_n/H_n$  of exponent at most  $e$ . Using the identification (3.1), we have

$$\prod_{i=1}^r H_{p_i}^{a_i} \times G_{n''} \subseteq W'_n.$$

Thus, thanks to Proposition 1.(i), in order to prove Claim 2 it suffices to show that, for each  $i = 1, \dots, r$ , there exists some  $t_i = t_i(A, K, e) \in \mathbb{N}$  such that  $H_{p_i}^{a_i}$  contains the kernel of the projection map

$$\pi_{p_i}^{t_i} : G_{p_i}^{a_i} \rightarrow G_{p_i}^{\min\{t_i, a_i\}}.$$

From now on, we fix  $i$  and write  $p, a, t$  instead of  $p_i, a_i, t_i$ .

Let  $G_{p^\infty} = \text{Gal}(K_{p^\infty}/K) \subseteq \mathcal{G}_p(\mathbb{Z}_p)$  be the group defined in Section 2. Let

$$\lambda : G_{p^\infty} \rightarrow G_{p^a}$$

be the canonical surjective restriction homomorphism and set  $H_{p^\infty} = \lambda^{-1}(H_{p^a})$ .

Note that it is enough to show that there exists some  $t = t(A, K, e) \in \mathbb{N}$  such that  $H_{p^\infty}$  contains the kernel of the projection map

$$\tilde{\pi}_{p^t} : \mathcal{G}_p(\mathbb{Z}_p) \rightarrow \mathcal{G}_p(\mathbb{Z}/p^t\mathbb{Z}).$$

Indeed this would imply that  $\ker(\pi_{p^t}) = \lambda(\ker(\tilde{\pi}_{p^t})) \subseteq H_{p^a}$ .

As  $H_{p^a}$  is normal in  $G_{p^a}$ , the subgroup  $H_{p^\infty}$  is normal in  $G_{p^\infty}$ . Since  $G_{p^\infty}/H_{p^\infty} \simeq G_{p^a}/H_{p^a}$  has finite exponent at most  $e$  and

$$[\mathcal{G}_p(\mathbb{Z}_p) : G_{p^\infty}] = [\mathcal{G}_p(\mathbb{Z}_p) : \rho_p(\text{Gal}(\bar{\mathbb{Q}}/K))] \leq c$$

by Proposition 1.(iv), we have that, if  $\gamma \in \mathcal{G}_p(\mathbb{Z}_p)$ , then  $\gamma^\delta$  lies in  $H_{p^\infty}$ .

We now want to show that  $t$  can be chosen such that every element in  $\ker(\tilde{\pi}_{p^t})$  is of the form  $\gamma^\delta$  for some  $\gamma \in \mathcal{G}_p(\mathbb{Z}_p)$ . This will complete the proof of Claim 2 and of Theorem 1.

Let  $\mathbf{L}_p$  denote the Lie algebra of  $\mathcal{G}_p \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$ . By [Bou72, Ch. III, §7, No. 2, Proposition 3], there is an open subgroup  $\mathcal{U} \subseteq \mathbf{L}_p$  and a map

$$\varphi : \mathcal{U} \rightarrow \mathcal{G}_p(\mathbb{Q}_p),$$

called exponential map and satisfying  $\varphi(\ell u) = \varphi(u)^\ell$  for all  $u \in \mathcal{U}$  and  $\ell \in \mathbb{Z}$  and such that  $\mathcal{V} = \varphi(\mathcal{U})$  is an open subgroup of  $\mathcal{G}_p(\mathbb{Q}_p)$  homeomorphic to  $\mathcal{U}$  via  $\varphi$ . After shrinking  $\mathcal{U}$ , we can and will assume without loss of generality that  $\varphi(\mathcal{U}) \subseteq \mathcal{G}_p(\mathbb{Z}_p)$  since  $\mathcal{G}_p(\mathbb{Z}_p)$  is an open subgroup of  $\mathcal{G}_p(\mathbb{Q}_p)$ .

Note that

$$\delta\mathcal{U} = \{\delta u \mid u \in \mathcal{U}\}$$

is open in  $\mathcal{U}$  as multiplication by  $\delta$  is an automorphism of  $\mathbf{L}_p$  as a topological group. Furthermore, since  $\varphi$  is a homeomorphism and  $\varphi(0) = \mathbb{I}_{2g}$ , we have that  $\varphi(\delta\mathcal{U})$  is open in  $\mathcal{V}$  and contains  $\mathbb{I}_{2g}$ .

Therefore there exists  $t = t(A, K, e) \in \mathbb{N}$  such that  $\varphi(\delta\mathcal{U})$  contains the open ball (with respect to the maximum norm on  $\mathcal{G}_p(\mathbb{Z}_p) \subseteq \mathrm{GL}_{2g}(\mathbb{Z}_p) \subseteq \mathbb{Q}_p^{4g^2}$ ) centered at  $\mathbb{I}_{2g}$  of radius  $p^{-t+1}$ . Thus,  $\ker(\tilde{\pi}_{p^t}) \subseteq \varphi(\delta\mathcal{U}) \subseteq \mathcal{V}$ .

We set  $\psi = \varphi^{-1} : \mathcal{V} \rightarrow \mathcal{U}$ . Then, if  $v \in \ker(\tilde{\pi}_{p^t})$ , we have that  $\psi(v) \in \delta\mathcal{U}$  and so  $\psi(v) = \delta u$  for some  $u \in \mathcal{U}$ . Hence

$$v = \varphi(\psi(v)) = \varphi(\delta u) = \varphi(u)^\delta = \gamma^\delta$$

with  $\gamma = \varphi(u) \in \mathcal{G}_p(\mathbb{Z}_p)$  and we are done.  $\square$

## 4 Proof of Corollary 1

We recall that, from the proof of [CZ11, Proposition 2.1], it follows that any abelian extension of  $\mathbb{Q}$  having Galois group of exponent at most  $e$  is contained in  $\mathbb{Q}_{\mathrm{ab}}^{(e)}$ . We repeat that proof here, the only difference being that the base field  $\mathbb{Q}$  is replaced by any number field  $L$ .

Let  $F/L$  be a possibly infinite abelian extension of a number field  $L$  and suppose that  $G = \mathrm{Gal}(F/L)$  has exponent at most  $e$ . Clearly,  $F$  equals the compositum of all its Galois subextensions  $L \subseteq F' \subseteq F$  with  $G' = \mathrm{Gal}(F'/L)$  finite and abelian of exponent at most  $e$ . We can thus write  $G' = \prod_{i=1}^n U_i$  as a direct product of finite cyclic groups  $U_i$  of order at most  $e$ . Let  $H_i$  be the subgroup of  $G'$  defined by  $H_i := \prod_{j \neq i} U_j$ . We have that  $[G' : H_i] = |U_i| \leq e$  for all  $i$ 's and  $\bigcap_{i=1}^n H_i = \{\mathrm{id}\}$ . Thus,  $F'$  equals the compositum of the fields  $F'^{H_1}, \dots, F'^{H_n}$ , which are abelian extensions of  $L$  of degrees  $[F'^{H_i} : L] = [G' : H_i] \leq e$ . Hence  $F' \subseteq L_{\mathrm{ab}}^{(e)}$  for every  $F'$  as above, which implies that  $F \subseteq L_{\mathrm{ab}}^{(e)}$ .

Now let  $F$  be a subfield of  $k(G_{\mathrm{tors}})$  such that  $F/k$  is Galois and  $\mathrm{Gal}(F/k)$  has exponent at most  $e$ .

If  $G = \mathbb{G}_m$ , then  $\mathrm{Gal}(F/k)$  is abelian and, by the above-mentioned results,  $F \subseteq k_{\mathrm{ab}}^{(e)}$ .

If  $G = A$ , then, by Theorem 1, there exists a finite extension  $M/k$ , depending only on  $A$ ,  $k$ , and  $e$ , such that  $F \subseteq M^{\mathrm{ab}}$ . Setting  $L = F \cap M$ ,

we have that  $\text{Gal}(F/L)$  is abelian and has exponent at most  $e$  as a subgroup of  $\text{Gal}(F/k)$ . So again  $F \subseteq L_{\text{ab}}^{(e)}$ .

We now conclude the proof using the fact that fields of the form  $L_{\text{ab}}^{(e)}$  have property (N) by [BZ01, Theorem 1].

## Acknowledgements

The authors thank Michel Brion, Philipp Habegger, Samuel Le Fourn, David Loeffler, Davide Lombardo, Thomas Scanlon, and Umberto Zannier for useful discussions and correspondence and for their kind interest in this work.

This material is mostly based upon work supported by the National Science Foundation under Grant Nos. 1440140 and DMS-1928930 while the authors were in residence at the Mathematical Sciences Research Institute in Berkeley, California, during the Spring 2023 semester. We thank the organisers of the “Diophantine Geometry” program for giving us the opportunity to participate in this program and the MSRI staff for providing an excellent environment for mathematical collaborations.

Sara Checcoli’s work has been also supported by the French National Research Agency in the framework of the *Investissements d’avenir* program (ANR-15-IDEX-02, IRGA project *ProBoNo*). Gabriel Dill has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement n° 945714).

## References

- [AD00] F. Amoroso and R. Dvornicich, *A lower bound for the height in abelian extensions*. J. Number Theory 80, 260–272 (2000).
- [ADZ14] F. Amoroso, S. David, and U. Zannier, *On fields with property (B)*. Proc. Amer. Math. Soc. 142, 1893–1910 (2014).
- [AZ00] F. Amoroso and U. Zannier, *A relative Dobrowolski lower bound over abelian extensions*. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 29, 711–727 (2000).
- [AZ10] F. Amoroso and U. Zannier, *A uniform relative Dobrowolski’s lower bound over abelian extensions*. Bull. London Math. Soc. 42, 489–498 (2010).
- [BG06] E. Bombieri and W. Gubler, *Heights in Diophantine Geometry*. Cambridge University Press, Cambridge (2006).

- [BZ01] E. Bombieri and U. Zannier, *A note on heights in certain infinite extensions of  $\mathbb{Q}$* . Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl. 12, 5–14 (2001).
- [BT71] A. Borel and J. Tits, *Éléments unipotents et sous-groupes paraboliques de groupes réductifs. I*. Invent. Math. 12, 95–104 (1971).
- [Bou72] N. Bourbaki, *Éléments de mathématique. Groupes et algèbres de Lie, Chapitres 2 et 3*. Hermann, Paris (1972).
- [CZ11] S. Checcoli and U. Zannier, *On fields of algebraic numbers with bounded local degrees*. C. R. Acad. Sci. Paris 349, no. 1–2, 11–14 (2011).
- [Che13] S. Checcoli, *Fields of algebraic numbers with bounded local degrees and their properties*. Trans. Amer. Math. Soc. 365, 2223–2240 (2013).
- [Fre21] L. Frey, *Explicit small heights in infinite non-abelian extensions*. Acta Arith. 199, 111–133 (2021).
- [Fre22] L. Frey, *Small heights in large non-abelian extensions*. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) 23, 1357–1393 (2022).
- [Hab13] P. Habegger, *Small height and infinite nonabelian extensions*. Duke Math. J. 162, 2027–2076 (2013).
- [Hum13] J. Humphreys (<https://mathoverflow.net/users/4231/jim-humphreys>), *Reductive groups over finite fields*. URL (version: 2013-10-15): <https://mathoverflow.net/q/144932>.
- [Ive76] B. Iversen, *The geometry of algebraic groups*. Advances in Math. 20, 57–85 (1976).
- [Lan56] S. Lang, *Algebraic groups over finite fields*. Amer. J. Math. 78, 555–563 (1956).
- [Mar13] Marguax (<https://mathoverflow.net/users/39487/marguax>), *Reductive groups over finite fields*. URL (version: 2013-10-16): <https://mathoverflow.net/q/144946>.
- [Mil17] J. S. Milne, *Algebraic groups*. Cambridge University Press, Cambridge (2017).
- [Nor49] D. G. Northcott, *An inequality in the theory of arithmetic on algebraic varieties*. Proc. Cambridge Philos. Soc. 45, 502–509 (1949).
- [Ser81] J.-P. Serre, *Letters to K. Ribet, January 1st and 29th, 1981*. In: Œuvres. Collected papers. IV. Springer-Verlag, Berlin (2000).
- [Ser86] J.-P. Serre, *Letter to K. Ribet, March 7th, 1986*. In: Œuvres. Collected papers. IV. Springer-Verlag, Berlin (2000).

- [Ser13] J.-P. Serre, *Un critère d'indépendance pour une famille de représentations  $\ell$ -adiques*. Comment. Math. Helv. 88, 541–554 (2013).
- [Ste68] R. Steinberg, *Endomorphisms of linear algebraic groups*. American Mathematical Society, Providence, R.I. (1968).
- [Win02] J.-P. Wintenberger, *Démonstration d'une conjecture de Lang dans des cas particuliers*. J. Reine Angew. Math. 553, 1–16 (2002).
- [Zyw19] D. Zywina, *An effective open image theorem for abelian varieties*. Preprint, ArXiv:1910.14171 (2019).

ADDRESSES:

S. Checcoli: Institut Fourier, UMR 5582, Laboratoire de Mathématiques, Université Grenoble Alpes, CS 40700, 38058 Grenoble cedex 9, France.  
sara.checcoli@univ-grenoble-alpes.fr

G. A. Dill: Institut für Algebra, Zahlentheorie und Diskrete Mathematik, Leibniz Universität Hannover, Welfengarten 1, 30167 Hannover, Germany.  
dill@math.uni-hannover.de