



HAL
open science

Contrasting aerosol composition in and out of Paris plume during the ACROSS campaign at the Rambouillet supersite

Joel Brito, Paola Formenti, Sébastien Dusanter, Jamar Marina, Alexandre Tomas, Laurent Alleman, Esperanza Perdrix, Pablo Espina Martin, Véronique Riffault, Chenjie Yu, et al.

► To cite this version:

Joel Brito, Paola Formenti, Sébastien Dusanter, Jamar Marina, Alexandre Tomas, et al.. Contrasting aerosol composition in and out of Paris plume during the ACROSS campaign at the Rambouillet supersite. European Geosciences Union General Assembly 2023, Apr 2023, Vienna, Austria. hal-04148986

HAL Id: hal-04148986

<https://hal.science/hal-04148986v1>

Submitted on 7 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

EGU23-17306, updated on 07 Sep 2023
<https://doi.org/10.5194/egusphere-egu23-17306>
EGU General Assembly 2023
© Author(s) 2023. This work is distributed under
the Creative Commons Attribution 4.0 License.



Contrasting aerosol composition in and out of Paris plume during the ACROSS campaign at the Rambouillet supersite

Joel Ferreira de Brito¹, Paola Formenti², Sebastien Dusanter¹, Marina Jamar¹, Alexandre Tomas¹, Laurent Alleman¹, Esperanza Perdrix¹, Pablo Espina¹, Veronique Riffault¹, Chenjie Yu², Claudia Di Biagio², Aline Gratien², Ludovico Di Antonio³, Lelia Hawkins⁴, Barbara D'Anna⁵, Julien Kammer⁵, Anne Monod⁵, Jean-Eudes Petit⁶, Shravan Deshmukh⁷, Laurent Poulain⁷, and the ACROSS Leadership*

¹Centre for Energy and Environment, IMT Nord Europe, Institut Mine-Télécom, Université de Lille, Lille, 59000, France

²Université Paris Cité and Univ Paris Est Creteil, CNRS, LISA, F-75013 Paris, France

³Univ Paris Est Creteil and Université Paris Cité, CNRS, LISA, F-94010 Créteil, France

⁴Department of Chemistry, Harvey Mudd College, 301 Platt Blvd, Claremont, California 91711, United States

⁵Laboratoire Chimie-Environnement, Université Aix-Marseille / CNRS, Marseille

⁶Laboratoire des Sciences du Climat et de l'Environnement, CEA-CNRS-UVSQ, IPSL, Université Paris-Saclay, 91191 Gif-sur-Yvette, France

⁷Leibniz-Institut für Troposphärenforschung e.V. (TROPOS), Permoserstraße 15, 04318 Leipzig, Germany

*A full list of authors appears at the end of the abstract

Based on a combination of ground-sites and aircraft observations, the ACROSS (Atmospheric ChemistRy Of the Suburban foreSt) project seeks to better understand how the pollution from Paris affects air quality over large regional scales, particularly through interaction with biogenic species. A field campaign took place in June-July 2022 at three sites: a forested remote site (Rambouillet, 50 km southwest from downtown Paris) a peri-urban (SIRTA) and an urban ones (PRG). The extensive instrumentation deployed at those sites allows to gain insights into the transformation of gaseous and particulate species from local to regional scales. Besides, the Rambouillet site has not only served as a receptor site from Paris plume but also led to detailed observations of freshly emitted biogenic compounds and their role in local chemistry, reactivity, and aerosol properties.

This work focuses on the real-time aerosol composition at the Rambouillet site. High resolution time-of-flight aerosol mass spectrometry sampled air from below (3 m above ground level) and above the canopy (40 m), and has been complemented by a suite of aerosol instruments including PTRMS-CHARON (targeting semi-volatile species above the canopy), and the PEGASUS mobile facility sampling below the canopy. Furthermore, data will be combined with SIRTA and PRG observations for case studies where Paris plume was transported into the supersite, or conversely, clean air masses were first observed at the remote site, and then advected towards the urban area. During the six-week observation period, the average PM₁ (non-refractory PM₁ + equivalent BC) was 8.4 µg m⁻³, with significant variability (10th-90th percentiles of 2.3 and 17.9 µg m⁻³). The

dominant component was organics with an average contribution to PM₁ of 62%, and peak contribution during periods of warm temperature of up to 90%. The organic aerosol was fairly oxidized at the site (average O:C ratio of 0.6), as expected for such an environment, with a negative trend of oxidation with organic aerosol loading, indicating primary or relatively fresh particles contributing the most to polluted episodes. Secondary inorganic aerosols (sulfate, nitrate, and ammonium) represented about 34% of PM₁, with the highest contribution during cold, rainy periods, and equivalent BC only had a minor contribution of about 4%. This presentation aims at providing a large picture of aerosol composition in the Ile-de-France region during the ACROSS experiment.

Acknowledgments. This work is supported by the French national research agency (ANR-20-CE01-0010), the Labex CaPPA (ANR-11-LABX-005-01) and the French national program LEFE/CHAT INSU.

ACROSS Leadership: Christopher Cantrell (3), Vincent Michoud (2)