



3D model related to the publication: An eosimiid primate of South Asian affinities in the Paleogene of Western Amazonia and the origin of New World monkeys

Laurent Marivaux, Francisco R Negri, Ana Maria Ribeiro

► To cite this version:

Laurent Marivaux, Francisco R Negri, Ana Maria Ribeiro. 3D model related to the publication: An eosimiid primate of South Asian affinities in the Paleogene of Western Amazonia and the origin of New World monkeys. *MorphoMuseum*, 2023, 9 (3), pp.e188. <10.18563/journal.m3.188>. <hal-04154006>

HAL Id: hal-04154006

<https://hal.science/hal-04154006v1>

Submitted on 6 Jul 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

3D model related to the publication: An eosimiid primate of South Asian affinities in the Paleogene of Western Amazonia and the origin of New World monkeys

Laurent Marivaux^{1*}, Francisco Ricardo Negri², Ana Maria Ribeiro^{3,4}

¹Laboratoire de Paléontologie, Institut des Sciences de l'Évolution de Montpellier (ISE-M, UMR 5554, CNRS/UM/IRD/EPHE), c.c. 064, Université de Montpellier, place Eugène Bataillon, 34095 Montpellier Cedex 05, France

²Laboratório de Paleontologia, Campus Floresta, Universidade Federal do Acre, Estrada do Canela Fina, Km 12, 69980-000, Cruzeiro do Sul, Acre, Brazil

³Seção de Paleontologia, Museu de Ciências Naturais, Secretaria do Meio Ambiente e Infraestrutura, Av. Dr. Salvador França 1427, 90690-000, Porto Alegre, Rio Grande do Sul, Brazil

⁴Universidade Federal do Rio Grande do Sul, Av. Bento Gonçalves 9500, 91501-970, Porto Alegre, Rio Grande do Sul, Brazil

*Corresponding author: Laurent.Marivaux@UMontpellier.fr

Abstract

This contribution contains the three-dimensional digital model of one isolated fossil tooth of an anthropoid primate (*Ashaninkacebus simpsoni*), discovered in sedimentary deposits located on the upper *Rio Juruá* in State of Acre, Brazil (Western Amazonia). This fossil was described, figured and discussed in the following publication: Marivaux et al. (2023), An eosimiid primate of South Asian affinities in the Paleogene of Western Amazonia and the origin of New World monkeys. Proceedings of the National Academy of Sciences USA. <https://doi.org/10.1073/pnas.2301338120>

Keywords: Brazilian Amazonia, early Anthropoidea, Eosimiidae, Paleobiogeography, Platyrrhini

Submitted:2023-01-31, published online:2023-07-04. <https://doi.org/10.18563/journal.m3.188>

INTRODUCTION

We present here the three-dimensional (3D) digital model of a single isolated tooth of a fossil primate (Fig. 1; Table 1) that was unearthed in allochthonous detrital Holocene sediments (PRJ-33', transported blocks of microconglomerate) deposited on the left bank of the *Rio Juruá* near Foz do Breu, Brazilian Amazonia (State of Acre, Brazil). The upriver provenance and stratigraphic context of the UFAC-CS 066 primate tooth from PRJ-33' remain so far unknown. However, biochronological inferences deriving from associated rodent teeth indicate a Paleogene age, likely around the Eocene/Oligocene transition (EOT; i.e., ca. 34 Ma; see, Kerber et al., 2017; Marivaux et al., 2023). The dental specimen (a pristine right upper M1; Fig. 1) has allowed the description of a new anthropoid primate, *Ashaninkacebus simpsoni* Marivaux et al. (2023), which was a diminutive species (230 g), equivalent in size to some small living marmoset callitrichine platyrrhines (e.g., *Callithrix jacchus*, 236–256 g after Ford, 1994). *Ashaninkacebus* is among the earliest primates to be known in South America. If recent fossil discoveries in Western Amazonia (Peru) revealed that two distinct anthropoid primate clades of African origin (Oligopithecidae-like primate and Parapithecidae; Bond et al., 2015; Seiffert et al., 2020) colonized South America near the Eocene/Oligocene transition, *Ashaninkacebus* demonstrates that a third clade of basal anthropoids was involved in the Paleogene colonization of South America by primates. This discovery provides increasingly puzzling insights into the origin and historical biogeography of New World monkeys, as this new taxon has strong affinities with stem anthropoid primates not of African but of South Asian origin:

Inv nr.	Taxon	Description
UFAC-CS066	<i>Ashaninkacebus simpsoni</i>	Right first upper molar (rM1).

Table 1. List of models. Collection: Paleontology Laboratory, Universidade Federal do Acre, Cruzeiro do Sul, Acre, Brazil

the Eosimiidae. Whatever the most ancient primates of South America known to date (*Ashaninkacebus*, *Perupithecus*, and *Ucayalipithecus*), they bear little adaptive resemblance to later Oligocene-early Miocene “true” platyrrhines, and the scarcity of paleontological data so far assembled precludes elucidating firmly their affinities with or within Platyrrhini. Measurement of the M1 buccal cutting-edge development of *Ashaninkacebus* (lengths of the buccal shearing crests were measured on the 3D digital model of the UFAC-CS 066 M1, using Avizo 2020.2 measurement tools) revealed a positive and high Shearing Quotient (SQ) value. The high SQ and the very small body size of this primate, indicate a diet with primarily insect and probably fruit consumption, but lacking exudates and leaves (after Kay et al., 2019, fig. 6B). Interestingly, such life-history traits (very small body size and insectivory) might have increased the chances of survival on a natural floating island during the extraordinary transatlantic journey to South America from Africa.

METHODS

The dental specimen documenting *Ashaninkacebus simpsoni* from the PRJ-33' fossil-bearing sediments (Fig. 1) was scanned with a resolution of 5.64 µm, using a high energy µCT-scanning station SkyScan 1173. AVIZO 2020.2 (Visualization Sciences




Figure 1. Dental remain of *Ashaninkacebus simpsoni* (UFAC-CS 066, Holotype). Images are renderings of a 3D digital model of the fossil specimen (right upper M1), obtained by a X-ray micro-computed (μ CT) surface reconstruction (rendering of the segmented surface). A) occlusal view; B) lingual view; C) buccal view.

Group) software was used for visualization, segmentation and three-dimensional (3D) rendering. This isolated tooth was prepared within a “labelfield” module of AVIZO, using the segmentation threshold selection tool. The crown and roots of the tooth were virtually delimited by manual segmentation. The 3D model is provided in “.ply” format, and thus can be opened with a wide range of software programs (e.g., MorphoDig 1.6.5., an open-source 3D freeware (Lebrun, 2018; <https://morphomuseum.com/Pages/morphodig>).

ACKNOWLEDGEMENTS

The 3D data presented in this work was produced through the technical facilities of the *Instituto de Petróleo e dos Recursos Naturais, Pontfícia Universidade Católica do Rio Grande do Sul, Brazil*. We particularly thank Adolpho Herbert Augustin (Institute of Petroleum and Natural Resources, Porto Alegre, Brazil) for μ CT-scan acquisition, treatment and reconstruction. This research was supported by the Brazilian *Conselho Nacional de Desenvolvimento Científico e Tecnológico* (CNPq 310023/2021-1 and 310948/2021-5, CNPq/MCTI/CONFAP-FAPS PROTAX 22/2020 441626/2020-3, and Fapergs 21/2551-0000781-8), the São Paulo Research Foundation (FAPESP, process n. 2011/14080-0) and the French *Agence Nationale de la Recherche* (ANR) in the framework of the LabEx CEBA (ANR-10-LABX-25-01; strategic project EMERGENCE). ISEM publication n° 2023-113 SUD.

BIBLIOGRAPHY

Bond, M., Tejedor, M. F., Campbell Jr, K. E., Chornogubsky, L., Novo, N., Goin, F. J., 2015. Eocene primates of South America and the African origins of New World monkeys. *Nature* 520, 538–541. <https://doi.org/10.1038/nature14120>

Ford, S. M., 1994. Evolution of sexual dimorphism in body weight in platyrrhines. *American Journal of Primatology* 34, 221–244.

Kay, R. F., Gonzales, L. A., Salenbien, W., Martinez, J.-N., Cooke, S. B., Valdivia, L. A., Rigsby, C., Baker, P. A., 2019. *Parvimico materdei* gen. et sp. nov.: a new platyrrhine from the Early Miocene of the Amazon Basin, Peru. *Journal of Human Evolution* 134, 102628. <https://doi.org/10.1016/j.jhevol.2019.05.016>

Kerber, L., Negri, F. R., Ribeiro, A. M., Nasif, N., Pereira de Souza-Filho, J., Ferigolo, J., 2017. Tropical fossil caviomorph rodents from the southwestern Brazilian Amazonia in the context of the South American faunas: systematics, biochronology, and paleobiogeography. *Journal of Mammalian Evolution* 24, 57–70. <https://doi.org/10.1007/s10914-016-9340-2>

Lebrun, R., 2018. MorphoDig, an open-source 3D freeware dedicated to biology. 5th International Paleontological Congress (IPC5) – *The Fossil Week*, July 9–13th, 2018 (Paris, France). Abstract volume, 399.

Marivaux, L., Negri, F. R., Antoine, P.-O., Stutz, N., Condamine, F. L., Kerber, L., Pujos, F., Ventura Santos, R., Alvim, A. M. V., Hsiou, A. S., Bissaro Júnior, M. C., Adami-Rodrigues, K., Ribeiro, A. M., 2023. An eosimiid primate of South Asian affinities in the Paleogene of Western Amazonia and the origin of New World monkeys. *Proceedings of the National Academy of Sciences USA*. <https://doi.org/10.1073/pnas.2301338120>

Seiffert, E. R., Tejedor, M. F., Fleagle, J. G., Novo, N. M., Cornejo, F. M., Bond, M., de Vries, D., Campbell Jr, K. E., 2020. A parapithecoid stem anthropoid of African origin in the Paleogene of South America. *Science* 368, 194–197. <https://doi.org/10.1126/science.aba1135>