



HAL
open science

Bile acids regulate MAdCAM-1 expression to link the gut microbiota to cancer immunosurveillance

Marine Fidelle, Ai-Ling Tian, Laurence Zitvogel, Guido Kroemer

► To cite this version:

Marine Fidelle, Ai-Ling Tian, Laurence Zitvogel, Guido Kroemer. Bile acids regulate MAdCAM-1 expression to link the gut microbiota to cancer immunosurveillance. *OncoImmunology*, 2023, 12 (1), 10.1080/2162402X.2023.2224672 . hal-04154967

HAL Id: hal-04154967

<https://hal.science/hal-04154967>

Submitted on 7 Jul 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Bile acids regulate MAdCAM-1 expression to link the gut microbiota to cancer immunosurveillance

Marine Fidelle^{a,b,c}, Ai-Ling Tian^{b,d}, Laurence Zitvogel^{a,c,e,f}, and Guido Kroemer^{b,d,g}

^aMetabolomics and Cell Biology Platforms, Gustave Roussy, Villejuif, France; ^bMetabolomics and Cell Biology Platforms, Gustave Roussy Cancer Center, Université Paris Saclay, Villejuif, France; ^cEquipe Labellisée Par la Ligue Contre le Cancer, Inserm U1015, Villejuif, France; ^dCentre de Recherche des Cordeliers, Equipe Labellisée Par la Ligue Contre le Cancer, Université de Paris, Sorbonne Université, Inserm U1138, Institut Universitaire de France, Paris, France; ^eFaculty of Medicine, Université Paris Saclay, France; ^fDepartment of Biology, Center of Clinical Investigations in Biotherapies of Cancer (CICBT) BOTHERIS, Villejuif, France; ^gDepartment of Biology, Institut du Cancer Paris Carpem, Hôpital Européen Georges Pompidou, AP-HP, Paris, France

ABSTRACT

In a recent paper in *Science*, Fidelle et al. unravel a gut immune checkpoint that is subverted by antibiotic treatment. Post-antibiotic dysbiosis of the ileum causes an increase in bile acids that downregulate MAdCAM-1, thereby triggering the exodus of immunosuppressive T cells from gut-associated lymphoid tissues toward tumors.

ARTICLE HISTORY

Received 29 May 2023

Revised 9 June 2023

Accepted 9 June 2023

KEYWORDS

Cancer immunotherapy; microbiota; immunosuppression; MAdCAM-1; bile acids; *Enterocloster*; gut

Over the past few years, it has become increasingly recognized that cancer is more than a disease caused by malignant cells. Rather, cancer results from a disequilibrium of the bodywide ecosystem with its metabolic, neuroendocrine and immune circuitries that also implies the microbiota.¹ Intestinal dysbiosis is a general feature of poor health and aging.² Dysbiosis also participates to carcinogenesis, tumor progression and oncoterapeutic failure, not only in gastrointestinal but also in distal cancers. This latter point has been particularly well documented for immunotherapy targeting the PD-1/PD-L1 interaction, where the prolonged (≥ 7 days) use of broad-spectrum antibiotics during the 2 months before and 1 month after initiation of the treatment is associated with poor outcome.³

In mechanistic terms, gut dysbiosis may be expected to result in an increase in the abundance of noxious (immunosuppressive and proinflammatory) microbes or – alternatively or in addition – the depletion of beneficial (immunostimulatory and anti-inflammatory) microbes.³ In a recent *Science* paper,⁴ we reported the finding that treatment with broad-spectrum antibiotics, followed by their discontinuation, leads to the recolonization of the gut by harmful bacteria from the *Enterocloster* genus including *E. clostridioformis*, both in mice and in cancer patients. Indeed, oral gavage of tumor-bearing mice with *E. clostridioformis* is sufficient to block the therapeutic response to PD-1 blockade.⁴ Intrigued by this finding, we engaged in a combination of hypothesis-driven and systematic studies to understand how transient treatment with antibiotics subverts therapeutically induced immunosurveillance.

We emitted the hypothesis that dysbiosis caused by antibiotics might affect the long-range trafficking of immune cells

from the gut through the lymphatic and cardiovascular systems. Indeed, we observed in mice that administration of antibiotics causes an increase in the trafficking of a particular immunosuppressive T cell subpopulation from the lamina propria of the ileum through the mesenteric lymph node to the tumor microenvironment and the tumor-draining lymph node. Extensive phenotyping of this T cell subset revealed that they bear both characteristics of regulatory T cells and proinflammatory T helper 17 cells (with the simultaneous expression of two master transcription factors, Foxp3 and Ror γ t) leading to their designation as T_{reg}17 cells. In addition, the mechanisms through which such T_{reg}17 cells are released from the gut was unraveled. Indeed, high endothelial cells present in the ileum normally express mucosal addressin cell adhesion molecule-1 (MAdCAM-1), which interacts with a specific integrin heterodimer ($\alpha 4\beta 7$) that is expressed on some immune cell types present in the gut including T_{reg}17 cells, hence retaining them locally. However, dysbiosis is linked to the downregulation of MAdCAM-1, thus unleashing $\alpha 4^{\beta 7+}$ T_{reg}17 cells from their local confinement and allowing them to travel to tumors. Indeed, knockout of the genes coding for MAdCAM-1 or the integrin $\beta 7$, as well antibodies blocking MAdCAM-1 or the $\alpha 4\beta 7$ heterodimer, are sufficient to cause the translocation of $\alpha 4^{\beta 7+}$ T_{reg}17 cells from the gut to tumors, and to compromise the efficacy of PD-1 blockade in vivo. These observations led us to the conclusion that the downregulation of ileal MAdCAM-1 expression explains why gut dysbiosis compromises immunosurveillance. In favor of this hypothesis, oral gavage of *E. clostridioformis* caused the downregulation of ileal MAdCAM-1.⁴

The gavage with *E. clostridioformis* was accompanied by major shifts in the ileal abundance of bile acids,⁴ which collectively may affect MAdCAM-1 expression⁵ and mediate systemic immunosuppression.⁶ To identify which particular bile acid species regulate MAdCAM-1, we engineered two distinct mouse endothelial cell lines to express green fluorescent protein (GFP) under the control of the *Madcam1* promoter. These cell lines downregulated GFP (indicative of the inhibition of the *Madcam1* promoter) in response to *E. clostridioformis* in vitro. Moreover, they upregulated GFP in response to two inflammatory cytokines (interleukin-1 β and tumor necrosis factor- α), and this upregulation was inhibited by individual bile acids, in particular lithocholic acid (LCA), and ursodeoxycholic acid (UDCA). Subsequent in vivo experiments confirmed that gavage with either LCA and UDCA is sufficient to trigger the reduction of *Madcam1* mRNA expression in vivo, in the ileum, Peyer's patches and mesenteric lymph nodes.⁷ However, at this point, it remains to be clarified how these effects of LCA and UDCA are achieved at the mechanistic level, likely through an action on one or several bile acid receptors.⁸ Moreover, it will be interesting to explore the possibility that dietary fibers mediate their reported immunotherapy-stimulatory effects⁹ through effects on toxic bile acids.¹⁰

Of note among these two bile acids, UDCA is clinically used for the treatment of primary biliary cholangitis, an autoimmune disease affecting cholangiocytes,¹¹ as well as for the avoidance of biliary complications after liver transplantation.¹² It will be interesting to explore the possibility that this liver-specific

immunosuppressive effect is achieved through the UDCA-induced downregulation of ileal MAdCAM-1, followed by the homing of T_{reg}17 cells from the gut-associated lymphoid tissue into the inflamed liver, where MAdCAM-1 is expressed on endothelial cells.¹³ Irrespective of this conjecture, we found that transgene-enforced expression of MAdCAM-1 in the liver (mostly on hepatocytes) increased the local infiltration by T_{reg}17 cells in mice subjected to transient antibiotic treatment and simultaneously reduced the frequency of T_{reg}17 cells in the tumor bed.⁷ Thus, MAdCAM-1 expression by the liver can lead to the interception of T_{reg}17 cells during their voyage from the gut to the cancer.

Clinically, the dosage in the serum of the soluble form of MAdCAM-1 (sMAdCAM-1) helped to evaluate its expression in the gut. Indeed, we found that patients who took antibiotics showed a significant decrease in sMAdCAM-1, in accordance with the decrease of *Madcam1* expression assessed on ileal biopsies from antibiotics-treated patients.⁷ Furthermore, sMAdCAM-1 was a strong prognostic factor in cancer patients treated with immunotherapies. Low levels of sMAdCAM-1 before starting the immunotherapy were associated with poor clinical outcome, in lung, renal and bladder cancer patients.⁷ Also, patients with low sMAdCAM-1 showed an enrichment of their gut microbiota with *E. clostridioformis*.⁷

Altogether, the aforementioned results plead in favor of a scenario in which gut dysbiosis causes the expansion of *E. clostridioformis*, which downregulates ileal expression of MAdCAM-1, thus allowing $\alpha 4^{\beta 7} T_{reg}17$ cells to travel from




Figure 1. Summary of the pathogenic cascade linking antibiotic therapy to deficient anticancer immunotherapy responses. The central pathogenic pathway can be mimicked by experimental interventions (red) or interrupted by countermeasures (green).

the gut to the mesenteric lymph node and then to tumors to cause local immunosuppression (Figure 1).

It should be noted that dysbiosis is difficult to be defined in global terms due to major geographic differences in the composition of the normal microbiota across continents, countries and regions.¹⁴ For this reason, it would not be surprising that other bacterial species than *E. clostridioformis* would mediate similar effects as those observed in our (French and Canadian) cohorts. Indeed, it is our intuition that, cancer- and immune-relevant products of the microbiota, including metabolites and ligands of pattern recognition receptors, should be more easily discernible biomarkers of dysbiosis than shifts in the metagenomic characteristics of the microbiota. As a caveat, however, such functional products should not be investigated in feces but rather at their major site of action, most likely the ileum.

Acknowledgments

MF is supported by the Seerave Foundation and the European Union's Horizon Europe research and innovation programme under grant agreement number 101095604 - GAP-101095604 -[project acronym: PREVALUNG-EU, project title: Personalized lung cancer risk assessment leading to stratified Interception]. This work was supported by SEERAVE Foundation, European Union Horizon 2020:Project Number: 825410 and Project Acronym: ONCOBIOME, Institut National du Cancer (INCa), ANR Ileobiome - 19-CE15-0029-01, ANR RHU5 "ANR-21-5 RHUS-0017" IMMUNOLIFE", MAdCAM INCA_ 16698, Ligue contre le cancer, LABEX OncoImmunology, la direction generale de l'offre de soins (DGOS), Universite Paris-Sud, SIRIC SOCRATE (INCa/DGOS/INSERM 6043), and PACRI network. Ai-L is supported by the ANR RHU5 "ANR-21-5 RHUS-0017" IMMUNOLIFE". LZ and GK are supported by the Ligue contre le Cancer (équipes labellisées, Program "Equipe labellisée LIGUE"; no. EL2016.LNCC (VT/PLP)); Agence Nationale de la Recherche (ANR) – Projets blancs; ANR under the frame of E-Rare-2, the ERA-Net for Research on Rare Diseases; AMMiCa US23/CNRS UMS3655; Association pour la recherche sur le cancer (ARC); Association "Le Cancer du Sein, Parlons-en!"; Cancéropôle Ile-de-France; Chancellerie des universités de Paris (Legs Poix), Fondation pour la Recherche Médicale (FRM); a donation by Elior; European Research Area Network on Cardiovascular Diseases (ERA-CVD, MINOTAUR); Gustave Roussy Odyssey, the European Union Horizon 2020 Project Oncobiome; Fondation Carrefour; INCa; Inserm (HTE); Institut Universitaire de France; LeDucq Foundation; the LabEx Immuno-Oncology (ANR-18-IDEX-0001); the RHU Torino Lumière; the Seerave Foundation; the SIRIC Stratified Oncology Cell DNA Repair and Tumor Immune Elimination (SOCRATE); and the SIRIC Cancer Research and Personalized Medicine (CARPEM).

Disclosure statement

LZ is a cofounder of everImmune, and holds patents covering the treatment of cancer and the therapeutic manipulation of the microbiota. LZ has held research contracts with Glaxo Smyth Kline, Incyte, Lytix, Kaleido, Pileje, Transgene, 9 m, Tusk Pharma, Merus, Roche and Innovate Pharma, and now has current research support from Biomérieux, Daiichi Sankyo, everImmune, Pileje, and 9 meters. LZ is in the SAB of Hookipa. LZ was on the Board of Directors of Transgene. GK has been holding research contracts with Daiichi Sankyo, Eleor, Kaleido, Lytix Pharma, PharmaMar, Osasuna Therapeutics, Samsara Therapeutics, Sanofi, Sotio, Tollys, Vascage and Vasculox/Tioma. GK has been consulting for Reithera. GK is on the Board of Directors of the Bristol Myers Squibb Foundation France. GK is a scientific co-founder of everImmune, Osasuna Therapeutics, Samsara Therapeutics and Therafast Bio. GK is the inventor of patents covering therapeutic targeting of aging, cancer, cystic fibrosis and metabolic disorders. GK's brother, Romano Kroemer, was an employee of Sanofi and now consults for Boehringer-Ingelheim.

The funders had no role in the design of the study; in the writing of the manuscript, or in the decision to publish the results.

Funding

The work was supported by the Agence Nationale de la Recherche [Projets blancs]; Association pour la Recherche sur le Cancer Cancéropôle Ile de France Fondation Leducq Fondation pour la Recherche Médicale Institut National Du Cancer Institut Universitaire de France Ligue Contre le Cancer [EL2016.LNCC (VT/PLP)]; European Research Area Network on Cardiovascular Diseases (ERA-CVD, MINOTAUR) Gustave Roussy Odyssey the European Union Horizon 2020 Project Oncobiome Fondation Carrefour Inserm (HTE) the LabEx Immuno-Oncology [ANR-18-IDEX-0001]; the RHU Torino Lumière a donation by Elior Chancellerie des universités de Paris (Legs Poix) Association "Le Cancer du Sein, Parlons-en!" AMMiCa US23/CNRS UMS3655 ANR under the frame of E-Rare-2, the ERA-Net for Research on Rare Diseases Agence Nationale de la Recherche (ANR) – Projets blancs the SIRIC Cancer Research and Personalized Medicine (CARPEM) Equipe labellisée LIGUE [EL2016.LNCC (VT/PLP)]; Immunolife [ANR-21-5 RHUS-0017]; the SIRIC Stratified Oncology Cell DNA Repair and Tumor Immune Elimination (SOCRATE) Seerave Foundation.

ORCID

Guido Kroemer  <http://orcid.org/0000-0002-9334-4405>

Data availability statement

All data that led to the conclusions in this manuscript have been referenced and all sources have been described.

References

- Kroemer G, McQuade JL, Merad M, André F, Zitvogel L. Bodywide ecological interventions on cancer. *Nat Med.* 2023;29(1):59–74. doi:10.1038/s41591-022-02193-4.
- López-Otin C, Pietrocola F, Roiz-Valle D, Galluzzi L, Kroemer G. Meta-hallmarks of aging and cancer. *Cell Metab.* 2023;35(1):12–35. doi:10.1016/j.cmet.2022.11.001.
- Derosa L, Routy B, Desilets A, Daillère R, Terrisse S, Kroemer G, Zitvogel L. Microbiota-centered interventions: the next breakthrough in immuno-oncology? *Cancer Discov.* 2021;11(10):2396–2412. doi:10.1158/2159-8290.CD-21-0236.
- Fidelle M, Rauber C, Alves Costa Silva C, Tian A-L, Lahmar I, de La Varenne A-L, Zhao L, Thelemaque C, Lebhar I, Messaoudene M, et al. A microbiota-modulated checkpoint directs immunosuppressive intestinal T cells into cancers. *Science.* 2023;380(6649):eabo2296. doi:10.1126/science.abo2296.
- Gao RY, Shearn CT, Orlicky DJ, Battista KD, Alexeev EE, Cartwright IM, Lanis JM, Kostelccky RE, Ju C, Colgan SP, et al. Bile acids modulate colonic MAdCAM-1 expression in a murine model of combined cholestasis and colitis. *Mucosal Immunol.* 2021;14(2):479–490. doi:10.1038/s41385-020-00347-6.
- Liu J, Fei Y, Zhou T, Ji H, Wu J, Gu X, Luo Y, Zhu J, Feng M, Wan P, et al. Bile acids impair vaccine response in children with biliary atresia. *Front Immunol.* 2021;12:12. doi:10.3389/fimmu.2021.642546.
- Fidelle M, Rauber C, Alves Costa Silva C, Tian A-L, Lahmar I, Mallard de La Varenne A-L, Zhao L, Thelemaque C, Lebhar I, Messaoudene M, et al. A microbiota-modulated checkpoint directs immunosuppressive intestinal T cells into cancers. *Science.* 2023;380(6649):380. doi:10.1126/science.abo2296.
- Lin S, Wang S, Wang P, Tang C, Wang Z, Chen L, Luo G, Chen H, Liu Y, Feng B, et al. Bile acids and their receptors in regulation of gut health and diseases. *Prog Lipid Res.* 2023;89:101210. doi:10.1016/j.plipres.2022.101210.

9. Spencer CN, McQuade JL, Gopalakrishnan V, McCulloch JA, Vetizou M, Cogdill AP, Khan MAW, Zhang X, White MG, Peterson CB, et al. Dietary fiber and probiotics influence the gut microbiome and melanoma immunotherapy response. *Science*. 2021;374(6575):1632–1640. doi:10.1126/science.aaz7015.
10. Saito Y, Sagae T. Defecation status, intestinal microbiota, and habitual diet are associated with the fecal bile acid composition: a cross-sectional study in community-dwelling young participants. *Eur J Nutr* 2023 Mar 7. doi:10.1007/s00394-023-03126-8.
11. Paillet J, Plantureux C, Lévesque S, Le Naour J, Stoll G, Sauvat A, Caudana P, Tosello Boari J, Bloy N, Lachkar S, et al. Autoimmunity affecting the biliary tract fuels the immunosurveillance of cholangiocarcinoma. *J Exp Med*. 2021;218(10):e20200853. doi:10.1084/jem.20200853.
12. Pedersen MR, Greenan G, Arora S, Murali AR, Mayo MJ. Ursodeoxycholic acid decreases incidence of primary biliary cholangitis and biliary complications after liver transplantation: a meta-analysis. *Liver Transpl*. 2021;27(6):866–875. doi:10.1002/lt.25935.
13. Graham JJ, Mukherjee S, Yuksel M, Sanabria Mateos R, Si T, Huang Z, Huang X, Abu Arqoub H, Patel V, McPhail M, et al. Aberrant hepatic trafficking of gut-derived T cells is not specific to primary sclerosing cholangitis. *Hepatology*. 2022;75(3):518–530. doi:10.1002/hep.32193.
14. McCulloch JA, Davar D, Rodrigues RR, Badger JH, Fang JR, Cole AM, Balaji AK, Vetizou M, Prescott SM, Fernandes MR, et al. Intestinal microbiota signatures of clinical response and immune-related adverse events in melanoma patients treated with anti-PD-1. *Nat Med*. 2022;28(3):545–556. doi:10.1038/s41591-022-01698-2.