



**HAL**  
open science

# The Jacobs–Keane theorem from the $\mathcal{S}$ -adic viewpoint

Felipe Arbulú, Fabien Durand, Bastián Espinoza

► **To cite this version:**

Felipe Arbulú, Fabien Durand, Bastián Espinoza. The Jacobs–Keane theorem from the  $\mathcal{S}$ -adic viewpoint. 2023. hal-04163989

**HAL Id: hal-04163989**

**<https://hal.science/hal-04163989>**

Preprint submitted on 17 Jul 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# THE JACOBS–KEANE THEOREM FROM THE $\mathcal{S}$ -ADIC VIEWPOINT

FELIPE ARBULÚ, FABIEN DURAND, AND BASTIÁN ESPINOZA

ABSTRACT. In the light of recent developments of the  $\mathcal{S}$ -adic study of subshifts, we revisit, within this framework, a well-known result on Toeplitz subshifts due to Jacobs–Keane giving a sufficient combinatorial condition to ensure discrete spectrum. We show that the notion of coincidences, originally introduced in the '70s for the study of the discrete spectrum of substitution subshifts, together with the  $\mathcal{S}$ -adic structure of the subshift allow to go deeper in the study of Toeplitz subshifts. We characterize spectral properties of the factor maps onto the maximal equicontinuous topological factors by means of coincidences density. We also provide an easy to check necessary and sufficient condition to ensure unique ergodicity for constant length  $\mathcal{S}$ -adic subshifts.

## 1. INTRODUCTION

A series of recent works shows that the underlying  $\mathcal{S}$ -adic structures of zero-entropy subshifts shed new light on the study of their properties as shown by [BSTY19, BCB<sup>+</sup>21, DDMP21, EM22, Esp22, GLL22, AD23, Esp23]. This approach, which goes back to [LV92, Fer96], consists in finding a representation of the subshift in terms of an infinite sequence of morphisms defined on finitely generated monoids. For more references on this subject, see the general references [BD14, DP22].

In the present work we investigate the spectral properties of Toeplitz subshifts through the  $\mathcal{S}$ -adic perspective. For a sequence  $x = (x_n)_{n \in \mathbb{Z}} \in \mathcal{A}^{\mathbb{Z}}$  on some finite alphabet  $\mathcal{A}$  and  $p \geq 0$ , we denote by  $\text{Per}_p(x)$  the set of integers  $n$  such that  $(x_{n+kp})_{k \in \mathbb{Z}}$  is constant. The aperiodic part of  $x$  is the set

$$\text{Aper}(x) = \mathbb{Z} \setminus \bigcup_{p \geq 0} \text{Per}_p(x).$$

A Toeplitz sequence is a non periodic sequence  $x$  such that  $\text{Aper}(x) = \emptyset$  and a Toeplitz subshift is the subshift generated by a Toeplitz sequence  $x$ , *i.e.*, the shift orbit closure of  $x$ .

Toeplitz sequences  $x$  always have a *periodic structure*, that is, a sequence  $(p_n)_{n \geq 0}$  such that for all  $n \geq 0$  we have

- $\text{Per}_{p_n}(x) \neq \emptyset$  and  $\text{Per}_{p_n}(x) \neq \text{Per}_q(x)$  for all  $0 \leq q < p_n$ ;
- $p_n$  divides  $p_{n+1}$ ; and
- $\bigcup_{n \geq 0} \text{Per}_{p_n}(x) = \mathbb{Z}$ .

---

*Date:* July 17, 2023.

*2020 Mathematics Subject Classification.* Primary: 37B10; Secondary: 37B20, 37A05.

*Key words and phrases.* Toeplitz subshifts,  $\mathcal{S}$ -adic subshifts, coincidences, odometer, discrete spectrum.

This results, topologically, in a maximal equicontinuous topological factor which is the odometer [Wil84]. A famous result of Jacobs and Keane [JK69, Mar75, Dow05] gives a sufficient condition by means of this periodic structure to ensure that the Toeplitz subshift has *discrete spectrum*, that is, there exists a measure theoretical isomorphism between the subshift and a translation on a compact Abelian group.

In this paper we revisit this result through the  $\mathcal{S}$ -adic framework. Indeed from [GJ00, Theorem 8] (see also [DP22, Theorem 6.1.7]) we know that Toeplitz subshifts are generated by proper, constant length, primitive and recognizable  $\mathcal{S}$ -adic sequences. We will precise this can be relaxed changing “proper” by “having coincidences” in Proposition 7.

Observe that when a minimal dynamical system  $(X, T)$  continuously factorizes onto a group rotation then the factor map is unique up to a translation. Thus below we will speak about “the” factor map instead of “a” factor map. We denote by  $\pi_{\text{meq}}$  this continuous factor map.

In [JK69, Theorem 6] the authors show that for a Toeplitz subshift the equivalent conditions in Item 2 and Item 3 below imply the discrete spectrum of the system. However, these conditions are not necessary for the discrete spectrum. Indeed, they are equivalent to the strictly stronger condition that the subshift is a *regular extension* of its maximal equicontinuous topological factor (the union of the singleton  $\pi_{\text{meq}}$ -fibers has full measure). For more details see Section 2.1. There exist discrete spectrum Toeplitz subshifts that do not fulfill this condition (see Section 6).

The difference between regularity and having discrete spectrum has been studied in detail in [DG16, GR17, HLSY21, GRJY21] for topological dynamical systems. There are given various strictly different notions, from topological to more measurable ones, all implying the discrete spectrum, such as mean equicontinuity or  $\mu$ -mean equicontinuity (which is equivalent to the discrete spectrum relative to  $\mu$ ).

In our work we introduced the point of view of coincidences to propose a more computable way to see the discrete spectrum. This allows us to build many examples having various spectral and ergodic properties.

The notion of coincidence is quantified below by the term “coinc” that is defined in Section 2.4. Coincidences have shown to be very relevant to detect the pure discrete spectrum. For example, in the context of constant length substitution subshifts [Dek78], two-symbol Pisot substitutions [BD02, HS03] or more general substitutions [Liv87, Que87]. Moreover, this notion has been intensively used to tackle the Pisot substitution conjecture [CS01, Fog02, BK06, BBK06].

For a subset  $A \subseteq \mathbb{Z}$ , the *density* of  $A$  is  $\text{dens}(A) = \limsup_{N \rightarrow +\infty} \frac{|A \cap [0, N]|}{N}$ .

**Theorem 1.** *Let  $(X, S)$  be the subshift generated by the Toeplitz sequence  $x$  with periodic structure  $(p_n)_{n \geq 0}$ . Then,  $X$  is defined by some constant length, recognizable directive sequence with coincidences  $\boldsymbol{\tau} = (\tau_n)_{n \geq 0}$  and the following are equivalent:*

- (1)  $(X, S)$  is a regular extension of its maximal equicontinuous topological factor: the union of the singleton  $\pi_{\text{meq}}$ -fibers has full measure.
- (2)  $\text{dens}(\text{Per}_{p_n}(x)) \rightarrow 1$  as  $n \rightarrow \infty$ .
- (3)  $\text{dens}(\text{Aper}(y)) = 0$  for every  $y \in X$ .
- (4) For every  $n \geq 0$ , we have

$$\lim_{N \rightarrow +\infty} \frac{|\text{coinc}(\tau_{[n, N]})|}{|\tau_{[n, N]}|} = 1.$$

(5) We have

$$\lim_{N \rightarrow +\infty} \frac{|\text{coinc}(\tau_{[0,N]})|}{|\tau_{[0,N]}|} = 1.$$

(6) There is a contraction  $\tau' = (\tau'_k)_{k \geq 0}$  of  $\tau$  such that

$$\sum_{k \geq 0} \frac{|\text{coinc}(\tau'_k)|}{|\tau'_k|} = +\infty.$$

Moreover, any of these conditions implies that  $(X, S)$  is uniquely ergodic and that  $\pi_{\text{meq}}$  is a measure theoretical isomorphism between  $(X, S)$  and its maximal equicontinuous topological factor.

It is observed in [GRJY21, Theorem 4.12][Theorem 54]Gar17 that for a minimal topological dynamical system the regularity condition is equivalent to the *diam-mean equicontinuity property*.

The  $\mathcal{S}$ -adic approach also allows us to study measure-theoretic aspects of a subshift. This is done by considering the notion of *adapted* directive sequences. Roughly speaking, when considering adapted sequences we get rid of parts of the system that are negligible.

We present a variation of Theorem 1 in terms of coincidence densities relative to the non negligible part of the system. This result is stated for subshifts generated by constant length directive sequences. It includes the case of Toeplitz subshifts as shown by Proposition 7, but also cases of non Toeplitz, non uniquely ergodic subshifts.

**Theorem 2.** *Let  $X$  be generated by the constant length, recognizable and primitive directive sequence  $\tau = (\tau_n : \mathcal{A}_{n+1}^* \rightarrow \mathcal{A}_n^*)_{n \geq 0}$ . Suppose that  $\tau$  is  $(\mathcal{A}'_n)_{n \geq 0}$ -adapted to an ergodic measure  $\mu$  of  $X$ . The following conditions are equivalent.*

- (1) *The map  $\pi_{\text{meq}} : (X, S, \mu) \rightarrow (\mathbb{Z}_{(|\tau_{[0,n]}|)_{n \geq 0}}, +1, \nu)$  defines a measure-theoretic isomorphism.*
- (2) *For every  $n \geq 0$ , we have*

$$\limsup_{N \rightarrow +\infty} \frac{|\text{coinc}_{\mathcal{A}'_N}(\tau_{[n,N]})|}{|\tau_{[n,N]}|} = 1.$$

We illustrate below, as applications of our results, various behaviours of Toeplitz subshifts with respect to discrete spectrum. They are detailed in Section 6.

Let  $\mathcal{A} = \{a, b\}$  and consider the directive sequence  $\tau = (\tau_n : \mathcal{A}^* \rightarrow \mathcal{A}^*)_{n \geq 0}$  such that for all  $n \geq 0$  the morphism  $\tau_n : \mathcal{A}^* \rightarrow \mathcal{A}^*$  is defined by

$$a \mapsto a^{n^2} b a a^{n^2}, \quad b \mapsto b^{n^2} a a b^{n^2}.$$

The Toeplitz subshift  $(X_\tau, S)$  is not a regular extension of its maximal equicontinuous topological factor, it has two ergodic measures  $\mu_a$  and  $\mu_b$ . Moreover, the map  $\pi_{\text{meq}}$  defines a measure theoretical isomorphism to the maximal equicontinuous topological factor for both  $(X_\tau, S, \mu_a)$  and  $(X_\tau, S, \mu_b)$  (see Section 6.3).

Now let  $\mathcal{A} = \{a, b, c\}$  and consider the directive sequence  $\tau = (\tau_n : \mathcal{A}^* \rightarrow \mathcal{A}^*)_{n \geq 0}$  such that for all  $n \geq 0$  the morphism  $\tau_n : \mathcal{A}^* \rightarrow \mathcal{A}^*$  is defined by

$$a \mapsto (ab)^{n^2+1} a c (ab)^{n^2+1}, \quad b \mapsto (ab)^{n^2+1} b c (ab)^{n^2+1}, \quad c \mapsto c (ab)^{n^2} c c c c (ab)^{n^2} c.$$

The Toeplitz subshift  $(X_\tau, S)$  is not a regular extension of its maximal equicontinuous topological factor, it possesses a unique invariant probability measure  $\mu$

such that  $\pi_{\text{meq}}$  is a measure theoretical isomorphism to its maximal equicontinuous topological factor (see Section 6.4). As far as we know, the first such example was given in [DK15, Section 5].

Other interesting examples are the following. Let  $\mathcal{A} = \{a, b, c\}$ ,  $m : \mathbb{N} \rightarrow \mathbb{N}$  defined by  $2m(n) + 3 = 3^n$  for each  $n \geq 0$  and consider the directive sequence  $\tau = (\tau_n : \mathcal{A}^* \rightarrow \mathcal{A}^*)_{n \geq 0}$  such that for all  $n \geq 0$  the morphism  $\tau_n : \mathcal{A}^* \rightarrow \mathcal{A}^*$  is

$$a \mapsto (ab)^{m(n)}abc, \quad b \mapsto a(ab)^{m(n)}ac, \quad c \mapsto (ab)^{m(n)}cbc.$$

The Toeplitz subshift  $(X_\tau, S)$  is such that its maximal equicontinuous topological factor is  $(\mathbb{Z}_3, +1)$ . Moreover it has a unique ergodic measure  $\mu$ . Nevertheless  $\pi_{\text{meq}}$  is not a measure theoretical isomorphism, but  $(X_\tau, S, \mu)$  has discrete spectrum: it is measure theoretically isomorphic to  $(\mathbb{Z}_3 \times \mathbb{Z}/2\mathbb{Z}, +(1, 1))$  (see Section 6.5).

Let  $\mathcal{A} = \{1, 2, a, b, c\}$ ,  $s : \mathbb{N} \rightarrow \mathbb{N}$  a nonnegative function such that  $2s(n) + 5 = 3^n$  for  $n \geq 0$  and consider the directive sequence  $\tau = (\tau_n : \mathcal{A}^* \rightarrow \mathcal{A}^*)_{n \geq 0}$  such that for all  $n \geq 0$  the morphism  $\tau_n : \mathcal{A}^* \rightarrow \mathcal{A}^*$  is defined by

$$\begin{aligned} 1 &\mapsto a(12)^{s(n)}12bc, & a &\mapsto (ab)^{s(n)}ab12c \\ 2 &\mapsto a(12)^{s(n)}21bc, & b &\mapsto a(ab)^{s(n)}b12c \\ & & c &\mapsto (ab)^{s(n)}ab12c. \end{aligned}$$

The Toeplitz subshift  $(X_\tau, S)$  has two ergodic measures  $\mu$  and  $\nu$  and its maximal equicontinuous topological factor is  $(\mathbb{Z}_3, +1)$ . The map  $\pi_{\text{meq}}$  is a measure theoretical isomorphism for the measure  $\mu$  whereas it is not for the measure  $\nu$ . Nevertheless, the system  $(X_\tau, S, \nu)$  has discrete spectrum: it is measure theoretically isomorphic to  $(\mathbb{Z}_3 \times \mathbb{Z}/2\mathbb{Z}, +(1, 1))$  (see Section 6.6).

Let  $\mathcal{A} = \{1, 2\}$ ,  $s : \mathbb{N} \rightarrow \mathbb{N}$  a nonnegative function such that  $3s(n) + 1 = 5^{2n}$  for  $n \geq 0$  and consider the directive sequence  $\tau = (\tau_n : \mathcal{A}^* \rightarrow \mathcal{A}^*)_{n \geq 0}$  such that for all  $n \geq 0$  the morphism  $\tau_n : \mathcal{A}^* \rightarrow \mathcal{A}^*$  is defined by

$$1 \mapsto (121)^{s(n)}2, \quad 2 \mapsto 1(121)^{s(n)}.$$

The Toeplitz subshift  $(X_\tau, S)$  has a unique ergodic measure  $\mu$ . However,  $(X_\tau, S, \mu)$  does not have discrete spectrum (see Section 6.7).

Finally we consider an example of a minimal non Toeplitz subshift generated by a constant length directive sequence that has positive entropy and where the conclusion of Theorem 2 holds (see Section 6.8).

#### ACKNOWLEDGEMENT

The first author is supported by ANID (ex CONICYT) Doctorado Becas Chile 72210185 grant. The third author thanks Doctoral Fellowship CONICYT-PFCHA Doctorado Nacional 2020 – 21202229. All authors are supported by ANR-22-CE40-0011 project Inside Zero Entropy Systems.

## 2. BACKGROUND

**2.1. Basics in topological dynamics.** A *topological dynamical system* (or just a system) is a pair  $(X, T)$  where  $X$  is a compact metric space and  $T : X \rightarrow X$  is a homeomorphism. The system  $(X, T)$  is *minimal* if for every point  $x \in X$  the orbit  $\{T^n x : n \in \mathbb{Z}\}$  is dense in  $X$ .

Let  $(X, T)$  and  $(Y, S)$  be two topological dynamical systems. We say that  $(Y, S)$  is a *topological factor* of  $(X, T)$  if there exists a continuous and onto map  $\pi : X \rightarrow Y$  such that

$$(1) \quad \pi \circ T = S \circ \pi.$$

In this case, we say that  $\pi$  is a *factor map* and that  $(X, T)$  is an *extension* of  $(Y, S)$ . If in addition the map  $\pi$  in (1) is a homeomorphism, we say that it is a *topological conjugacy* and that  $(X, T)$  and  $(Y, S)$  are *topologically conjugate*.

Let  $(X, T)$  be a uniquely ergodic system and denote by  $\mu$  its unique invariant probability measure. We say that  $(X, T)$  is a *regular extension* of  $(Y, S)$  if there exists a factor map  $\pi : X \rightarrow Y$  such that

$$(2) \quad \mu(\{x \in X : |\pi^{-1}(\{\pi(x)\})| = 1\}) = 1.$$

Any topological dynamical system  $(X, T)$  possesses a *maximal equicontinuous topological factor* [EG60], that is, there exists an equicontinuous system  $(Y, S)$  and a factor map  $\pi : X \rightarrow Y$  such that every other equicontinuous factor of  $(X, T)$  is a factor of  $(Y, S)$ . The system  $(Y, S)$  is unique up to topological conjugacy. Moreover, if  $(X, T)$  is minimal, then  $(Y, S)$  corresponds to a rotation on a compact abelian group and we usually denote by  $\nu$  the Haar measure on  $Y$ .

In the minimal case, if  $(X, T)$  is a regular extension of its maximal equicontinuous topological factor  $(Y, S)$ , then condition (2) is equivalent to

$$(3) \quad \nu(\{y \in Y : |\pi^{-1}(\{y\})| = 1\}) = 1.$$

Moreover, since any two such factor maps differ only by a translation, condition (3) is independent of the choice of the factor map  $\pi$  and of the representation of  $(Y, S)$  in its conjugacy class.

**2.2. Subshifts.** Let  $\mathcal{A}$  be a finite and nonempty set that we call *alphabet*. Elements in  $\mathcal{A}$  are called *letters* or *symbols*. The number of letters of  $\mathcal{A}$  is denoted by  $|\mathcal{A}|$ . The set of finite sequences or *words* of length  $\ell \in \mathbb{N}$  with letters in  $\mathcal{A}$  is denoted by  $\mathcal{A}^\ell$ . The *full shift*  $\mathcal{A}^{\mathbb{Z}}$  is the set of all bi-infinite sequences  $(x_n)_{n \in \mathbb{Z}}$  with  $x_n \in \mathcal{A}$  for all  $n \in \mathbb{Z}$ .

A word  $w = w_0 w_1 \dots w_{\ell-1} \in \mathcal{A}^\ell$  can be seen as an element of the free monoid  $\mathcal{A}^*$  endowed with the operation of concatenation (whose neutral element is  $\varepsilon$ , the empty word). The integer  $\ell$  is the *length* of the word  $w$  and is denoted by  $|w| = \ell$ ; the length of the empty word is 0. We say that a word  $w$  *occurs* in a sequence  $x = (x_n)_{n \in \mathbb{Z}} \in X$  if there exists  $m \in \mathbb{Z}$  such that  $w = x_m \dots x_{m+|w|-1}$ . We use the same notion for words. For such a word  $w$  in  $\mathcal{A}^*$  and a letter  $a$  in  $\mathcal{A}$ , the number of occurrences of  $a$  in  $w$  is denoted by  $|w|_a$ . A nonempty word  $w = w_0 w_1 \dots w_{\ell-1} \in \mathcal{A}^*$  *starts* (resp. *ends*) with a nonempty word  $u \in \mathcal{A}^*$  if  $u = w_0 \dots w_{i-1}$  for some  $i \leq \ell$  (resp.  $u = w_j \dots w_{\ell-1}$  for some  $j \geq 0$ ).

The *shift map*  $S : \mathcal{A}^{\mathbb{Z}} \rightarrow \mathcal{A}^{\mathbb{Z}}$  is defined by  $S((x_n)_{n \in \mathbb{Z}}) = (x_{n+1})_{n \in \mathbb{Z}}$ . A *subshift* is a subset  $X$  of a fullshift  $\mathcal{A}^{\mathbb{Z}}$  which is closed for the product topology and invariant under the shift map. Thus  $(X, S)$  is a topological dynamical system that we also call subshift.

Let  $(X, S)$  be a subshift. For  $x \in X$  and  $i, j \in \mathbb{Z}$  with  $i < j$  we define  $x_{[i,j]} = x_i x_{i+1} \dots x_{j-1}$ . The *language* of  $(X, S)$  is the set  $\mathcal{L}(X)$  containing all words  $w \in \mathcal{A}^*$  such that  $w = x_{[m, m+|w|]}$  for some  $x = (x_n)_{n \in \mathbb{Z}} \in X$  and  $m \in \mathbb{Z}$ . In this case, we also say that  $w$  is a *factor* (also called *subword*) of  $x$ . Given  $x \in X$ , the language  $\mathcal{L}(x)$  is the set of all words that occur in  $x$ . For two words  $u, v \in \mathcal{L}(X)$ , the *cylinder*

set  $[u.v]$  is the set  $\{x \in X : x_{[-|u|,|v|]} = uv\}$ . When  $u$  is the empty word we only write  $[v]$ , erasing the dot. We remark that cylinder sets are clopen sets and they form a base for the topology of the subshift.

**2.3. Morphisms and substitutions.** By a *morphism* we simply mean a morphism  $\tau : \mathcal{A}^* \rightarrow \mathcal{B}^*$  between the free monoids  $\mathcal{A}^*$  and  $\mathcal{B}^*$  for some finite alphabets  $\mathcal{A}$  and  $\mathcal{B}$ . We say that  $\tau$  is *erasing* whenever there exists  $a$  in  $\mathcal{A}$  such that  $\tau(a)$  is the empty word. Otherwise we say it is *nonerasing*.

The morphism  $\tau$  is *positive* if for all  $a$  in  $\mathcal{A}$  every letter of  $\mathcal{B}$  occurs in  $\tau(a)$ . The morphism  $\tau$  is *proper* if there exist  $p$  and  $s$  in  $\mathcal{B}$  such that  $\tau(a)$  starts with  $p$  and ends with  $s$  for every letter  $a$  in  $\mathcal{A}$ . The morphism  $\tau$  is of *constant length* if the length of the word  $\tau(a)$  does not depend on  $a$ . In that case, we denote by  $|\tau|$  the length of the word  $\tau(a)$  for any  $a$  in  $\mathcal{A}$ . Observe that if  $\tau : \mathcal{A}^* \rightarrow \mathcal{B}^*$  and  $\sigma : \mathcal{B}^* \rightarrow \mathcal{C}^*$  are morphisms of constant length, then the morphism  $\sigma \circ \tau$  is of constant length and

$$|\sigma \circ \tau| = |\sigma||\tau|.$$

When it is nonerasing,  $\tau$  extends naturally to maps from  $\mathcal{A}^{\mathbb{N}}$  to itself and from  $\mathcal{A}^{\mathbb{Z}}$  to itself in the obvious way by concatenation (in the case of a two-sided sequence we apply  $\tau$  to positive and negative coordinates separately and we concatenate the results at the coordinate zero).

To the morphism  $\tau : \mathcal{A}^* \rightarrow \mathcal{B}^*$  we associate a *composition matrix*  $M_\tau$  indexed by  $\mathcal{B} \times \mathcal{A}$  such that its entry at position  $(b, a)$  is the number of occurrences of  $b$  in  $\tau(a)$  for every  $a \in \mathcal{A}$  and  $b \in \mathcal{B}$ .

**2.4. Coincidences of constant length morphisms.** Throughout this article we will use the notion of *coincidences* of constant length morphisms. Let  $\tau : \mathcal{A}^* \rightarrow \mathcal{B}^*$  be a constant length morphism. A *coincidence* of  $\tau$  relative to  $\mathcal{A}' \subseteq \mathcal{A}$  is an integer  $i \in [0, |\tau|)$  such that the map that sends a letter  $a$  in  $\mathcal{A}'$  to the  $i$ -th letter of  $\tau(a)$  is constant. We denote by  $\text{coinc}_{\mathcal{A}'}(\tau)$  the set of such integers. When  $\mathcal{A}' = \mathcal{A}$  we just say these integers are coincidences and denote the set of such integers by  $\text{coinc}(\tau)$ .

The following lemma states how coincidences behave when morphisms are composed. The elementary proof is left to the reader.

**Lemma 3.** *Let  $\tau : \mathcal{A}^* \rightarrow \mathcal{B}^*$  and  $\sigma : \mathcal{B}^* \rightarrow \mathcal{C}^*$  be two morphisms of constant length. Then the set  $\text{coinc}(\sigma \circ \tau)$  contains the disjoint union*

$$\begin{aligned} & \{i|\sigma| + j : i \in \text{coinc}(\tau), 0 \leq j < |\sigma|\} \\ & \cup \{i|\sigma| + j : i \in [0, |\tau|) \setminus \text{coinc}(\tau), j \in \text{coinc}(\sigma)\} \end{aligned}$$

and we have

$$(4) \quad 1 - \frac{|\text{coinc}(\sigma \circ \tau)|}{|\sigma \circ \tau|} \leq \left(1 - \frac{|\text{coinc}(\sigma)|}{|\sigma|}\right) \left(1 - \frac{|\text{coinc}(\tau)|}{|\tau|}\right).$$

Moreover, if  $|\mathcal{A}| = |\mathcal{B}| = 2$  then inequality (4) is an equality between both terms.

**2.5. S-adic subshifts.** A *directive sequence*  $\boldsymbol{\tau} = (\tau_n : \mathcal{A}_{n+1}^* \rightarrow \mathcal{A}_n^*)_{n \geq 0}$  is a sequence of morphisms. From now on, we only consider morphisms that are nonerasing. When all morphisms  $\tau_n$  for  $n \geq 1$  are proper we say that  $\boldsymbol{\tau}$  is *proper*; and when all morphisms  $\tau_n$  for  $n \geq 1$  are of constant length we say that  $\boldsymbol{\tau}$  is of *constant length*.

For  $0 \leq n < N$ , we denote by  $\tau_{[n,N]}$  or  $\tau_{[n,N-1]}$  the morphism  $\tau_n \circ \tau_{n+1} \circ \dots \circ \tau_{N-1}$ . We say  $\tau$  is *primitive* if for any  $n \in \mathbb{N}$  there exists  $N > n$  such that  $M_{\tau_{[n,N]}} > 0$ , i.e., all letters in  $\mathcal{A}_n$  occur in  $\tau_{[n,N]}(a)$  for all  $a \in \mathcal{A}_N$ .

For  $n \in \mathbb{N}$ , the *language*  $\mathcal{L}^{(n)}(\tau)$  of level  $n$  associated with  $\tau$  is defined by

$$\mathcal{L}^{(n)}(\tau) = \{w \in \mathcal{A}_n^* : w \text{ occurs in } \tau_{[n,N]}(a) \text{ for some } a \in \mathcal{A}_N \text{ and } N > n\}$$

and  $X_\tau^{(n)}$  is the set of points  $x \in \mathcal{A}_n^{\mathbb{Z}}$  such that  $\mathcal{L}(x) \subseteq \mathcal{L}^{(n)}(\tau)$ . When nonempty, this set clearly defines a subshift that we call the *subshift generated by*  $\mathcal{L}^{(n)}(\tau)$ .

We set  $X_\tau = X_\tau^{(0)}$  and call  $(X_\tau, S)$  the  $\mathcal{S}$ -adic subshift generated by the directive sequence  $\tau$ . See [DP22, Chapter 6] for more details.

A *contraction* of  $\tau = (\tau_n : \mathcal{A}_{n+1}^* \rightarrow \mathcal{A}_n^*)_{n \geq 0}$  is a directive sequence of the form

$$\tau' = (\tau'_k = \tau_{[n_k, n_{k+1}]} : \mathcal{A}_{n_{k+1}}^* \rightarrow \mathcal{A}_{n_k}^*)_{k \geq 0},$$

where the sequence  $(n_k)_{k \geq 0}$  is such that  $n_0 = 0$  and  $n_k < n_{k+1}$  for all  $k \geq 0$ . Observe that any contraction of  $\tau$  generates the same  $\mathcal{S}$ -adic subshift  $X_\tau$ .

Inspired by the definition of substitutions with coincidences in [Dek78], we say a constant length directive sequence  $\tau$  has *coincidences* if there is a contraction  $\tau' = (\tau'_k)_{k \geq 0}$  of  $\tau$  such that  $\text{coinc}(\tau'_k) \neq \emptyset$  for all  $k \geq 0$ .

One says that  $\tau$  has *finite alphabet rank* when  $\liminf_{n \rightarrow +\infty} |\mathcal{A}_n| < +\infty$ . This notion is related to the *topological rank* defined more generally for minimal Cantor systems. We refer to [DDMP21] for more details about their interaction.

**2.6. Recognizability and dynamical partitions.** We now present the recognizability property for morphisms and directive sequences in terms of topological partitions. The usual definition and a more general discussion can be found in [BSTY19, BPR23, BPRS23].

Let  $\tau : \mathcal{A}^* \rightarrow \mathcal{B}^*$  be a nonerasing morphism and let  $X \subseteq \mathcal{A}^{\mathbb{Z}}$  be a subshift. We say that  $\tau$  is *recognizable* on  $X$  if

$$\mathcal{P} = \{S^k \tau([a]) : a \in \mathcal{A}, 0 \leq k < |\tau(a)|\}$$

defines a partition of the subshift  $\bigcup_{k \in \mathbb{Z}} S^k \tau(X)$ .

We say that a directive sequence  $\tau = (\tau_n : \mathcal{A}_{n+1}^* \rightarrow \mathcal{A}_n^*)_{n \geq 0}$  is recognizable if

$$(5) \quad \mathcal{P}_n = \{S^k \tau_{[0,n]}([a]) : a \in \mathcal{A}_n, 0 \leq k < |\tau_{[0,n]}(a)|\}, \quad n \geq 0,$$

defines a sequence of partitions of  $X_\tau$ . We have that  $\tau$  is recognizable if and only if for each  $n \geq 0$  the morphism  $\tau_n$  is recognizable on  $X_\tau^{(n+1)}$  [BSTY19, Section 4].

It will be convenient, in order to manipulate these partitions, to consider the following definitions. For  $n \geq 0$  and  $a \in \mathcal{A}_n$ , we denote by

$$(6) \quad \mathcal{T}_n(a) = \bigcup_{0 \leq k < |\tau_{[0,n]}(a)|} S^k \tau_{[0,n]}([a])$$

the *tower* indexed by  $a$  and by  $B_n(a)$  the *base*  $\tau_{[0,n]}([a])$  of this tower. The *base of*  $\mathcal{P}_n$  is

$$(7) \quad B_n = \bigcup_{a \in \mathcal{A}_n} B_n(a).$$

We make here two important observations about  $(\mathcal{P}_n)_{n \geq 0}$ . For any  $0 \leq n < m$ ,

- (1) the partition  $\mathcal{P}_m$  is finer than the partition  $\mathcal{P}_n$ ; and



(2) we have that

$$(8) \quad |\{k \in [0, |\tau_{[0,m]}(b)|) : S^k B_m(b) \subseteq B_n(a)\}| = |\tau_{[n,m]}(b)|_a, \quad a \in \mathcal{A}_n, \quad b \in \mathcal{A}_m.$$

We now state a fundamental theorem for the ergodic and topological study of  $\mathcal{S}$ -adic subshifts.

**Theorem 4** ([BSTY19, Lemma 6.3]). *Let  $\tau = (\tau_n : \mathcal{A}_{n+1}^* \rightarrow \mathcal{A}_n^*)_{n \geq 0}$  be a recognizable directive sequence and  $\mu$  an invariant measure of  $(X_\tau, \mathcal{S})$ . Then, for  $\mu$ -almost all  $x \in X_\tau$ , the set  $\bigcap_{n \geq 0} S^{k_n} \tau_{[0,n]}([a_n])$  is a singleton, where  $(k_n, a_n)_{n \geq 0}$  for  $n \geq 0$  is the  $(\mathcal{P}_n)_{n \geq 0}$ -address of  $x$ , that is, the unique sequence satisfying  $x \in S^{k_n} \tau_{[0,n]}([a_n])$ ,  $n \geq 0$ .*

We end this section with the following remark: if in addition to recognizability the sequence  $\tau$  is assumed to be primitive and proper, then the atoms of  $\bigcup_{n \geq 0} \mathcal{P}_n$  generate the topology of  $X_\tau$  [DL12, Proposition 2.2].

**2.7. Invariant measures.** Let  $\tau = (\tau_n : \mathcal{A}_{n+1}^* \rightarrow \mathcal{A}_n^*)_{n \geq 0}$  be a recognizable directive sequence and  $\mu$  be an invariant measure of  $(X_\tau, \mathcal{S})$ . In this context, Theorem 4 implies the following.

**Corollary 5.** *The sequence  $(\mathcal{P}_n)_{n \geq 0}$   $\mu$ -generates the Borel  $\sigma$ -algebra of  $X_\tau$ , that is,  $\mu$  is uniquely determined by the values it assigns to  $B_n(a)$ ,  $a \in \mathcal{A}_n$ ,  $n \geq 0$ .*

This motivates the following. For  $n \geq 0$  we define the column vector  $\mu_n$  by

$$\mu_n = (\mu_n(a) : a \in \mathcal{A}_n), \quad \text{where } \mu_n(a) = \mu(B_n(a)).$$

Since  $\tau$  is recognizable, we have

$$(9) \quad \mu_m = M_{\tau_{[m,n]}} \mu_n, \quad 0 \leq m < n.$$

The measure of the tower indexed by  $a$  is

$$(10) \quad \mu(\mathcal{T}_n(a)) = \mu(B_n(a)) |\tau_{[0,n]}| = \mu_n(a) |\tau_{[0,n]}|, \quad n \geq 0.$$

When dealing with measure theoretical arguments, it is natural to get rid of small portions of the space. We will need the definition of *adapted* directive sequences to state our main results, where we distinguish those towers with small measure and those with a measure uniformly bounded away from zero.

For each  $n \geq 0$  let  $\mathcal{A}'_n \subseteq \mathcal{A}_n$  be a nonempty alphabet and set

$$\mathcal{T}'_n = \bigcup_{a \in \mathcal{A}'_n} \mathcal{T}_n(a), \quad \mathcal{T}'_{\geq n} = \bigcap_{m \geq n} \mathcal{T}'_m, \quad \mathcal{T}' = \bigcup_{n \geq 0} \mathcal{T}'_{\geq n}.$$

Observe that  $\mathcal{T}' = \liminf_{n \rightarrow +\infty} \mathcal{T}'_n$ .

We say that  $\tau$  is  $(\mathcal{A}'_n)_{n \geq 0}$ -*weakly-adapted* to  $\mu$  if

$$(11) \quad \sum_{n \geq 0} \mu(X_\tau \setminus \mathcal{T}'_n) < +\infty.$$

If in addition there exists  $\delta > 0$  such that

$$(12) \quad \mu(\mathcal{T}_n(a)) \geq \delta, \quad a \in \mathcal{A}'_n, \quad n \geq 0,$$

we say that  $\tau$  is  $(\mathcal{A}'_n)_{n \geq 0}$ -*adapted* to  $\mu$  (for the constant  $\delta$ ). Observe that in such case  $\sup_{n \geq 0} |\mathcal{A}'_n|$  is bounded by  $1/\delta$ .

When  $\tau$  has finite alphabet rank, up to a contraction and a renaming of the alphabets, there exists  $\mathcal{A}_\mu \subseteq \mathcal{A}_n$  for all  $n \geq 0$  such that  $\tau$  is  $(\mathcal{A}_\mu)$ -adapted to  $\mu$ .

The previous definitions are inspired by the definition of *clean Bratteli diagrams* in [BDM10] (see also [BKMS13, Section 3]).

In the weakly-adapted case, the Borel–Cantelli lemma implies that

$$(13) \quad \mu(\mathcal{T}') = 1.$$

**2.8. Convergence of frequencies.** It is a well known fact that for substitutions systems the measures  $\mu(B_n(a))$  correspond to the letter frequencies of the iterates of the substitution [Que87, Chapter 5]. A similar result was proved in [BKMS13, Proposition 5.1] in the more general context of uniquely ergodic Bratteli–Vershik systems.

In this paper, we will need an extension of these results that is valid for the class of non uniquely ergodic systems we later consider.

**Proposition 6.** *Let  $\tau = (\tau_n : \mathcal{A}_{n+1}^* \rightarrow \mathcal{A}_n^*)_{n \geq 0}$  be a primitive and recognizable directive sequence, let  $\mu$  an ergodic measure for  $(X_\tau, S)$  and let  $(b_m)_{m \geq 0}$  a sequence of letters with  $b_m$  in  $\mathcal{A}_m$ . Suppose that any of the following conditions hold:*

- (1) *there exists a  $\mu$ -generic point  $x$  in  $X_\tau$  such that  $x \in \mathcal{T}_m(b_m)$  for all  $m \geq 0$ ;*
- (2)  *$\liminf_{m \rightarrow +\infty} \mu(\mathcal{T}_m(b_m)) > 0$ ;*
- (3)  *$\tau$  is  $(\mathcal{A}'_n)_{n \geq 0}$ -adapted to  $\mu$  and  $b_m$  belongs to  $\mathcal{A}'_m$  for all  $m \geq 0$ ;*
- (4) *the system  $(X_\tau, S)$  is uniquely ergodic.*

Then, for any  $n \geq 0$  and  $a \in \mathcal{A}_n$ , we have

$$(14) \quad \lim_{m \rightarrow +\infty} \frac{|\tau_{[n,m]}(b_m)|_a}{|\tau_{[0,m]}(b_m)|} = \mu(B_n(a)).$$

*Proof.* Let  $x$  be a  $\mu$ -generic point such that  $x$  belongs to  $\mathcal{T}_m(b_m)$  for all  $m \geq 0$ . We define  $i_m = \max\{k \leq 0 : S^k x \in B_m\}$  and  $j_m = \min\{k > 0 : S^k x \in B_m\}$ . Observe that  $j_m - i_m = |\tau_{[0,m]}(b_m)|$ . Since  $\tau$  is recognizable, from (8) we get, for  $n < m$ ,

$$|\{k \in [0, |\tau_{[0,m]}(b_m)|] : S^k B_m(b_m) \subseteq B_n(a)\}| = |\tau_{[n,m]}(b_m)|_a, \quad b_m \in \mathcal{A}_m.$$

This implies, as  $S^{i_m} x$  and  $S^{j_m} x$  are in  $B_m(b_m)$  and  $\mathcal{P}_m$  is finer than  $\mathcal{P}_n$ , that

$$|\{k \in [i_m, j_m) : S^k x \in B_n(a)\}| = |\tau_{[n,m]}(b_m)|_a.$$

The left-hand side corresponds to the ergodic sum  $\sum_{k \in [i_m, j_m)} 1_{B_n(a)}(S^k x)$  of the indicator function  $1_{B_n(a)}$ . Hence, since  $x$  is  $\mu$ -generic, we conclude that

$$\lim_{m \rightarrow +\infty} \frac{|\tau_{[n,m]}(b_m)|_a}{|\tau_{[0,m]}(b_m)|} = \lim_{m \rightarrow +\infty} \frac{1}{j_m - i_m} \sum_{k \in [i_m, j_m)} 1_{B_n(a)}(S^k x) = \mu(B_n(a))$$

and (14) holds.

Now suppose that  $\liminf_{m \rightarrow +\infty} \mu(\mathcal{T}_m(b_m)) > 0$ . We assume, with the aim to obtain a contradiction, that there exist  $n \geq 0$  and  $a \in \mathcal{A}_n$  such that  $\frac{|\tau_{[n,m]}(b_m)|_a}{|\tau_{[0,m]}(b_m)|}$  is bounded away from  $\mu(B_n(a))$  for all  $m$  in an infinite set  $E \subseteq \mathbb{N}$ .

Let

$$K = \limsup_{\substack{m \rightarrow +\infty \\ m \in E}} \mathcal{T}_m(b_m).$$

From our hypothesis, we have  $\mu(K) > 0$  and hence that there is a  $\mu$ -generic point  $x$  in  $K$ . We have that  $x$  is in  $\mathcal{T}_m(b)$  for all  $m$  belonging to an infinite set  $E' \subseteq E$ .

But then from Item 1 in Proposition 6 the sequence  $\left( \frac{|\tau_{[n,m]}(b_m)|_a}{|\tau_{[0,m]}(b)|} : m \in E' \right)$  is

arbitrarily close to  $\mu(B_n(a))$ , which contradicts our previous assumption. This shows that (14) holds.

Next, we suppose that  $\tau$  is  $(\mathcal{A}'_n)_{n \geq 0}$ -adapted to  $\mu$  and  $b_m \in \mathcal{A}'_m$  for all  $m \geq 0$ . Then, from (12) we get  $\liminf_{m \rightarrow +\infty} \mu(\mathcal{T}_m(b_m)) > 0$  and thus (14) holds by Item 2 of this proposition.

Finally, if the system  $(X_\tau, S)$  is uniquely ergodic, then any point is  $\mu$ -generic and (14) holds for any sequence of letters  $b_m$  in  $\mathcal{A}_m$  from Item 1 in Proposition 6.  $\square$

**2.9. Toeplitz sequences as  $\mathcal{S}$ -adic subshifts.** In [GJ00, Theorem 8] it is shown that the class of Toeplitz subshifts coincides, up to topological conjugacy, with the class of expansive Bratteli–Vershik systems defined on diagrams with the *equal path number property*. This is the starting point of the link between Toeplitz subshifts and  $\mathcal{S}$ -adic subshifts that is summarized in the proposition below. This will be used throughout the article. We follow the proof in [DP22, Theorem 6.1.7].

**Proposition 7.** *Let  $(X, S)$  be a subshift. The following are equivalent.*

- (1)  $(X, S)$  is a Toeplitz subshift;
- (2)  $(X, S)$  is generated by a constant length, primitive, proper and recognizable directive sequence; and
- (3)  $(X, S)$  is generated by a constant length, primitive and recognizable directive sequence with coincidences.

*Proof.* It is clear that Item 2 implies Item 3. We now prove that Item 3 implies Item 1. Let  $\tau = (\tau_n : \mathcal{A}_{n+1}^* \rightarrow \mathcal{A}_n^*)_{n \geq 0}$  be a constant length, primitive, proper, and recognizable directive sequence. For each  $n \geq 0$ , let  $k_n$  be a coincidence of  $\tau_n$ ,  $\ell_n = \lfloor |\tau_{[0,n]}|/2 \rfloor$ ,  $r_n = |\tau_{[0,n]}| - \ell_n$  and  $x$  a point belonging to  $\bigcap_{n \geq 0} \tau_{[0,n]}([\mathcal{A}_n \cdot \mathcal{A}_n])$ . If we define the sequence of points  $x_n = S^{k_n |\tau_{[0,n]}| + \ell_n} x$ ,  $n \geq 0$ , we observe that

$$(x_n)_{[-\ell_n, r_n]} = (x_n)_{[-\ell_n + k \cdot |\tau_{[0, n+1]}|, r_n + k \cdot |\tau_{[0, n+1]}|]}, \quad k \in \mathbb{Z}.$$

This implies that any accumulation point of the sequence  $(x_n)_{n \geq 0}$  is a Toeplitz sequence, and hence  $(X_\tau, S)$  is a Toeplitz subshift.

It remains to prove that Item 1 implies Item 2.

**Claim 8.** *Let  $x \in \mathcal{A}^{\mathbb{Z}}$  be a Toeplitz sequence. Then there exists a nonempty alphabet  $\mathcal{B}$ , a constant length and positive morphism with coincidences  $\tau : \mathcal{B}^* \rightarrow \mathcal{A}^*$  and a Toeplitz sequence  $y \in \mathcal{B}^{\mathbb{Z}}$  such that  $x = \tau(y)$ . Moreover, the morphism  $\tau$  is recognizable on the subshift  $Y$  generated by  $y$ .*

*Proof.* Let  $m \in \mathbb{N}$  be large enough so that all letters that occur in  $x$  also occur in  $x_{[-m, m]}$ . Define  $u = x_{[-m, 0]}$  and  $v = x_{[0, m]}$ . As  $x$  is a Toeplitz sequence, there exists a minimal  $p > 2m$  such that

$$x_{[-m+kp, m+kp]} = x_{[-m, m]} = uv, \quad k \in \mathbb{Z}.$$

Define  $\mathcal{B}'$  be the set of words  $\{x_{[kp, (k+1)p]} : k \in \mathbb{Z}\} \subseteq \mathcal{A}^*$ . Let  $\tau : \mathcal{B} \rightarrow \mathcal{B}'$  be a bijection, where  $\mathcal{B}$  is an alphabet. The map  $\tau$  of course defines a morphism from  $\mathcal{B}^*$  to  $\mathcal{A}^*$ .

We have that  $\tau$  is positive, with coincidences and  $|\tau| = p$ . It is clear that there exists a unique point  $y \in \mathcal{B}^{\mathbb{Z}}$  such that  $x = \tau(y)$ . Since  $x$  is Toeplitz, we have that  $y$  is Toeplitz. We are left to prove that  $\tau$  is recognizable on the subshift  $Y$  generated by  $y$ .

Consider the cylinder  $U = [u.v]$  in  $X$  and define

$$\text{Per}_q(x', U) = \{n \in \mathbb{Z} : S^{n+kq}(x') \in U, k \in \mathbb{Z}\}, \quad x' \in X, \quad q \in \mathbb{N}.$$

Since  $p$  is minimal we have  $\text{Per}_q(x, U) = \emptyset$  if  $q < p$  and  $\text{Per}_p(x, U) \neq \emptyset$ . The set  $\{x' \in X : \text{Per}_p(x', U) \neq \emptyset\}$  is nonempty, closed and  $S$ -invariant, hence it is equal to  $X$  as  $(X, S)$  is minimal. From this we see that if

$$C = \{x' \in X : \text{Per}_p(x', U) = \text{Per}_p(x, U)\}$$

then  $\{S^i C : 0 \leq i < p\}$  is a clopen partition of  $X$ . Finally, since  $(X, S)$  is minimal one has  $C = \bigcup_{b \in \mathcal{B}} \tau([b])$ , where the cylinders  $[b]$ ,  $b \in \mathcal{B}$ , are considered in  $Y$ . This implies that

$$\{S^k \tau([b]) : b \in \mathcal{B}, 0 \leq k < p\}$$

is a clopen partition of  $X$ , and thus  $\tau$  is recognizable on  $(Y, S)$ , finishing the proof of the claim.  $\square$

By the claim, we inductively construct a sequence of alphabets  $(\mathcal{B}_n)_{n \geq 0}$  with  $\mathcal{B}_0 = \mathcal{A}$ ; a sequence of Toeplitz sequences  $(y_n)_{n \geq 0}$  with  $y_n \in \mathcal{B}_n^{\mathbb{Z}}$  for  $n \geq 0$  such that  $y_0 = x$ ; and a directive sequence of morphisms  $\tau = (\tau_n : \mathcal{B}_{n+1}^* \rightarrow \mathcal{B}_n^*)_{n \geq 0}$  which is of constant length, primitive, with coincidences and such that for all  $n \geq 0$  the morphism  $\tau_n$  is recognizable on the subshift generated by  $y_n$ . Since we have  $y_n = \tau_{[n, N]}(y_N)$  for all  $0 \leq n < N$ , we see that  $y_n$  generates the subshift  $X_\tau^{(n)}$ . In particular,  $\tau$  is recognizable and  $X = X_\tau$ , which finishes the proof.  $\square$

## 2.10. The maximal equicontinuous topological factor.

2.10.1. *Odometers.* For a sequence of positive integers  $(p_n)_{n \geq 0}$  such that  $p_n$  divides  $p_{n+1}$  for  $n \geq 0$ , the *odometer* given by  $(p_n)_{n \geq 0}$  is the system  $(\mathbb{Z}_{(p_n)_{n \geq 0}}, +1)$ , where

$$\mathbb{Z}_{(p_n)_{n \geq 0}} = \varprojlim \mathbb{Z}/p_n \mathbb{Z} = \left\{ (x_n)_{n \geq 0} \in \prod_{n \geq 0} \mathbb{Z}/p_n \mathbb{Z} : x_{n+1} \equiv x_n \pmod{p_n}, n \geq 0 \right\}$$

and the map “+1” is given by

$$(x_n)_{n \geq 0} \mapsto (x_n + 1 \pmod{p_n})_{n \geq 0}.$$

If  $p_n = p^n$  for some  $p \geq 1$  and all  $n \geq 0$ , we simply denote  $\mathbb{Z}_{(p_n)_{n \geq 0}}$  by  $\mathbb{Z}_p$ .

Odometers are, in particular, minimal rotations on compact abelian groups. They are equicontinuous and uniquely ergodic, where the unique ergodic measure is the Haar measure. In particular, the measurable system given by the odometer has discrete spectrum.

For  $\ell \geq 0$  and  $z = (z_n)_{n \geq 0} \in \mathbb{Z}_{(p_n)_{n \geq 0}}$  we define the cylinder

$$[z_0, z_1, \dots, z_{\ell-1}] = \{(x_n)_{n \geq 0} \in \mathbb{Z}_{(p_n)_{n \geq 0}} : x_k = z_k, 0 \leq k < \ell\}.$$

The Haar measure  $\nu$  on  $(\mathbb{Z}_{(p_n)_{n \geq 0}}, +1)$  satisfies

$$(15) \quad \nu([z_0, z_1, \dots, z_{\ell-1}]) = \frac{1}{p_0 p_1 \dots p_{\ell-1}}.$$

We finally remark that the odometer is *coalescent* as a measure theoretical system, that is, every measurable automorphism of the odometer is an isomorphism. This follows since it has discrete spectrum [HP68].

2.10.2. *The maximal equicontinuous topological factor.* Let  $\tau = (\tau_n)_{n \geq 0}$  be a constant length, primitive, and recognizable directive sequence with coincidences. It is a well-known fact that the maximal equicontinuous topological factor of  $(X_\tau, S)$  corresponds to the odometer  $(\mathbb{Z}_{(|\tau_{[0,n]}|)_{n \geq 0}}, +1)$ . Indeed, this follows from Proposition 7 and [Wil84, Theorem 2.2].

The factor map  $\pi_{\text{meq}} : X_\tau \rightarrow \mathbb{Z}_{(|\tau_{[0,n]}|)_{n \geq 0}}$  can be described as follows. Let  $x \in X_\tau$ . By recognizability of  $\tau$ , for every  $n \geq 0$  there exists  $k_n(x)$  with  $0 \leq k_n(x) < |\tau_{[0,n]}|$  such that  $x \in S^{k_n(x)} B_n$ , where  $B_n$  is the base given by (7). Then we define

$$(16) \quad \pi_{\text{meq}}(x) = (k_n(x))_{n \geq 0}.$$

It can be observed that  $k_{n+1}(x) \equiv k_n(x) \pmod{|\tau_{[0,n]}|}$  for  $x \in X_\tau$ . In order to help the reader, it is important to notice the following relation

$$(17) \quad \pi_{\text{meq}}^{-1}([z_0, z_1, \dots, z_\ell]) = S^{z_\ell} B_\ell, \quad (z_n)_{n \geq 0} \in \mathbb{Z}_{(|\tau_{[0,n]}|)_{n \geq 0}}, \quad \ell \geq 0.$$

The following lemma will be useful.

**Lemma 9.** *Let  $0 \leq n < m$  and  $A_{n,m}$  be a subset of  $[0, |\tau_{[n,m]}|)$ . Then*

$$\nu(\{z \in \mathbb{Z}_{(|\tau_{[0,n]}|)_{n \geq 0}} : z_m \in z_n + |\tau_{[0,n]}| A_{n,m}\}) = \frac{|A_{n,m}|}{|\tau_{[n,m]}|}.$$

*Proof.* Observe that, from (15), the set  $\{z \in \mathbb{Z}_{(|\tau_{[0,n]}|)_{n \geq 0}} : z_m \in z_n + |\tau_{[0,n]}| A_{n,m}\}$  is a disjoint union of  $|\tau_{[0,n]}| \cdot |A_{n,m}|$  cylinders of measure  $1/|\tau_{[0,m]}|$ . We obtain

$$\nu(\{z \in \mathbb{Z}_{(|\tau_{[0,n]}|)_{n \geq 0}} : z_m \in z_n + |\tau_{[0,n]}| A_{n,m}\}) = \frac{|A_{n,m}|}{|\tau_{[n,m]}|}.$$

□

### 3. ABOUT UNIQUE ERGODICITY OF $\mathcal{S}$ -ADIC SUBSHIFTS OF CONSTANT LENGTH

In this section we give a necessary and sufficient condition for unique ergodicity in the specific case of  $\mathcal{S}$ -adic subshifts generated by constant length and recognizable directive sequences. This condition is in terms of the matrices of the directive sequence and can be checked relatively easily.

**Lemma 10.** *Let  $(a_{n,N} : 0 \leq n < N)$  be a doubly indexed sequence in  $[0, 1]$  that satisfies*

$$(18) \quad a_{l,n} \leq a_{l,m} a_{m,n}, \quad l < m < n.$$

*The following are equivalent.*

(1)

$$\lim_{N \rightarrow +\infty} a_{n,N} = 0, \quad \text{for all } n \geq 0;$$

(2) *there exists an increasing sequence  $(n_k)_{k \geq 0}$  such that*

$$\lim_{K \rightarrow +\infty} a_{n_k, n_K} = 0, \quad \text{for all } k \geq 0; \text{ and}$$

(3) *there exists an increasing sequence  $(n_k)_{k \geq 0}$  such that*

$$\sum_{k \geq 0} a_{n_k, n_{k+1}} < +\infty.$$

*Proof.* We first prove that Item 1 is equivalent to Item 2. It is clear that Item 1 implies Item 2. Let now  $n \geq 0$  and  $(n_k)_{k \geq 0}$  be such that Item 2 holds. From (18), we deduce that the sequence  $(a_{n,N} : N > n)$  is nonincreasing in  $N$ . In particular, since it is bounded, we have  $\lim_{N \rightarrow +\infty} a_{n,N} = \inf_{N > n} a_{n,N}$ . Then, for any integers  $0 \leq k < K$  such that  $n < n_k < n_K$ , Equation (18) yields

$$0 \leq \inf_{N > n} a_{n,N} \leq a_{n,n_K} \leq a_{n_k,n_K}.$$

By letting  $K \rightarrow +\infty$ , we obtain Item 1.

Item 3 can be obtained from Item 2 by considering a sequence  $(n_k)_{k \geq 0}$  such that  $a_{n_k,n_{k+1}} \leq 2^{-k}$  for  $k \geq 0$ . It is left to prove that Item 3 implies Item 2. Let  $k \geq 0$ . By using (18) and the inequality  $\log(x) \leq x - 1$  for  $x \in [0, 1]$ , we obtain

$$a_{n_k,n_K} \leq \prod_{k \leq j < K} a_{n_j,n_{j+1}} \leq \exp\left(\sum_{k \leq j < K} a_{n_j,n_{j+1}} - K + k\right), \quad 0 \leq k < K,$$

from which we deduce Item 2 if we let  $K \rightarrow +\infty$ .  $\square$

We now prove the main result of this section. Denote by  $|v|$  the sum of the entries of a vector  $v$ .

**Theorem 11.** *Let  $\tau = (\tau_n : \mathcal{A}_{n+1}^* \rightarrow \mathcal{A}_n^*)_{n \geq 0}$  be a constant length and recognizable directive sequence. The following are equivalent.*

- (1) *There exists a contraction  $\tau' = (\tau'_k = \tau'_{[n_k, n_{k+1}]})_{k \geq 0}$  of  $\tau$  such that the vectors  $(v_k)_{k \geq 0}$  given by*

$$v_k(a) = \min_{b \in \mathcal{A}_{n_{k+1}}} |\tau'_k(b)|_a, \quad a \in \mathcal{A}_{n_k}, \quad k \geq 0,$$

*satisfy*

$$(19) \quad \sum_{k \geq 0} \frac{|v_k|}{|\tau'_k|} = +\infty.$$

- (2) *The system  $(X_\tau, S)$  is uniquely ergodic.*

*Proof.* Let  $\tau' = (\tau'_k = \tau'_{[n_k, n_{k+1}]} : \mathcal{A}_{n_{k+1}}^* \rightarrow \mathcal{A}_{n_k}^*)_{k \geq 0}$  be a contraction of  $\tau$  and  $(v_k)_{k \geq 0}$  be the vectors such that Item 1 in Theorem 11 holds. Observe that, with this definition of the vectors  $(v_k)_{k \geq 0}$ , we have

$$(20) \quad M_{\tau'_k} = v_k \mathbf{1}_{k+1} + M'_k, \quad k \geq 0,$$

where the matrix  $M'_k$  is nonnegative and  $\mathbf{1}_{k+1}$  is the row vector of ones in  $\mathbb{Z}^{\mathcal{A}_{n_{k+1}}}$ .

**Claim 12.** *Define the doubly indexed sequences of vectors  $(v_{k,K} : 0 \leq k < K)$  and of matrices  $(M'_{k,K} : 0 \leq k < K)$  inductively by  $v_{k,k+1} = v_k$ ;  $M'_{k,k+1} = M'_k$  for  $k \geq 0$ ; and*

$$\begin{aligned} v_{k,K+1} &= |v_K| v_{k,K} + M'_{k,K} v_K + (|\tau'_K| - |v_K|) v_{k,K} \\ M'_{k,K+1} &= M'_{k,K} M'_K, \quad K > k. \end{aligned}$$

*Then we have*

$$(21) \quad M_{\tau'_{[k,K]}} = v_{k,K} \mathbf{1}_K + M'_{k,K}, \quad 0 \leq k < K.$$

*Proof.* From (20) we have  $1_k M'_k = (|\tau'_k| - |v_k|)1_{k+1}$  for all  $k \geq 0$ . We prove the claim by induction on  $K$ . Equation (21) holds if  $K = k + 1$  by (20). If (21) holds, then

$$\begin{aligned} M'_{\tau'_{[k, K+1]}} &= (v_{k, K} 1_K + M'_{k, K})(v_K 1_{K+1} + M'_K) \\ &= (|v_K| v_{k, K} + M'_{k, K} v_K + (|\tau'_K| - |v_K|) v_{k, K}) 1_{K+1} + M'_{k, K} M'_K \\ &= v_{k, K+1} 1_{K+1} + M'_{k, K+1}, \end{aligned}$$

thus proving the claim by induction.  $\square$

Observe that from the previous claim we have

$$1_k M'_{k, K} = (|\tau'_{[k, K]}| - |v_{k, K}|) 1_K$$

and hence

$$\begin{aligned} |v_{k, K+1}| &= |v_K| |v_{k, K}| + (1_k M'_{k, K}) v_K + (|\tau'_K| - |v_K|) |v_{k, K}| \\ &= |v_K| |v_{k, K}| + (|\tau'_{[k, K]}| - |v_{k, K}|) |v_K| + (|\tau'_K| - |v_K|) |v_{k, K}| \\ &= (|\tau'_K| - |v_K|) |v_{k, K}| + |\tau'_{[k, K]}| |v_K|, \quad 0 \leq k < K. \end{aligned}$$

After rearranging terms, we obtain

$$1 - \frac{|v_{k, K+1}|}{|\tau'_{[k, K+1]}|} = \left(1 - \frac{|v_{k, K}|}{|\tau'_{[k, K]}|}\right) \left(1 - \frac{|v_K|}{|\tau'_K|}\right), \quad 0 \leq k < K,$$

therefore

$$1 - \frac{|v_{k, K}|}{|\tau'_{k, K}|} = \prod_{k \leq \ell < K} \left(1 - \frac{|v_\ell|}{|\tau'_\ell|}\right), \quad 0 \leq k < K.$$

Define

$$a_{k, K} = 1 - \frac{|v_{k, K}|}{|\tau'_{k, K}|}, \quad 0 \leq k < K.$$

Since Equation (19) holds, Lemma 10 applied to  $(a_{k, K} : 0 \leq k < K)$  gives

$$(22) \quad \lim_{K \rightarrow +\infty} \frac{|v_{k, K}|}{|\tau'_{[k, K]}|} = 1, \quad k \geq 0.$$

We now prove that Item 2 in Theorem 11 holds. It is enough to prove that  $(X_{\tau'}, S)$  is uniquely ergodic. Let  $\mu$  be an invariant measure of  $(X_{\tau'}, S)$  and define the sequence of column vectors  $(\mu_k)_{k \geq 0}$  by

$$\mu_k = \mu(\tau'_{[0, k]}([a])), \quad a \in \mathcal{A}_{n_k}, \quad k \geq 0.$$

From Corollary 5, it is enough to prove that

$$(23) \quad \mu_k = \lim_{K \rightarrow +\infty} \frac{v_{k, K}}{|\tau'_{[0, K]}|}, \quad k \geq 0.$$

Let  $0 \leq k < K$ . By recognizability of  $\tau'$ , we have that  $|\mu_k| = |\tau'_{[0, k]}|^{-1}$  and  $\mu_k = M'_{\tau'_{[k, K]}} \mu_K$ . Hence, from (21) we deduce

$$\mu_k = \frac{v_{k, K}}{|\tau'_{[0, K]}|} + M'_{k, K} \mu_K.$$

If we multiply both sides by  $|\tau'_{[0, k]}|$  and sum over all letters  $a \in \mathcal{A}_{n_k}$ , we obtain

$$\frac{|v_{k, K}|}{|\tau'_{[k, K]}|} + |\tau'_{[0, k]}| |M'_{k, K} \mu_K| = 1.$$

From (22), we have  $|M'_{k,K}\mu_K| \rightarrow 0$  as  $K \rightarrow +\infty$  and, since the matrix  $M'_{k,K}$  is nonnegative, we finally deduce (23).

Now assume that Item 2 in Theorem 11 holds. Define  $n_0 = 0$  and denote by  $\mu$  the unique invariant probability measure of  $(X_\tau, S)$ . From Proposition 6 we inductively construct an increasing sequence  $(n_k)_{k \geq 0}$  such that

$$(24) \quad \frac{\min_{b \in \mathcal{A}_{n_{k+1}}} |\tau_{[n_k, n_{k+1})}(b)|_a}{|\tau_{[n_k, n_{k+1})}|} \geq \frac{1}{2} |\tau_{[0, n_k)}| \mu_{n_k}(a), \quad a \in \mathcal{A}_{n_k}, \quad k \geq 0$$

Define the contraction  $\tau' = (\tau'_k = \tau'_{[n_k, n_{k+1}))}_{k \geq 0}$  of  $\tau$ . By summing over all letters  $a \in \mathcal{A}_{n_k}$  in (24), we deduce that (19) holds for the contraction  $\tau'$ . This shows that Item 2 implies Item 1 in Theorem 11 and finishes the proof.  $\square$

The following corollary shows that the conditions stated in Theorem 1 implies the unique ergodicity.

**Corollary 13.** *Let  $\tau = (\tau_n : \mathcal{A}_{n+1}^* \rightarrow \mathcal{A}_n^*)_{n \geq 0}$  be a constant length and recognizable directive sequence. If there exists a contraction  $\tau' = (\tau'_k)_{k \geq 0}$  of  $\tau$  such that*

$$(25) \quad \sum_{k \geq 0} \frac{|\text{coinc}(\tau'_k)|}{|\tau'_k|} = +\infty,$$

*then the system  $(X_\tau, S)$  is uniquely ergodic.*

Let us make some observations about Theorem 11. There are in the literature other such conditions implying unique ergodicity. Let  $m_n$  and  $M_n$  be the smallest and greatest entry of  $M_{\tau_n}$ , respectively. It comes from [Sen81] that if

$$\sum_{n \geq 0} \frac{m_n}{M_n} = +\infty$$

then  $(X_\tau, S)$  is uniquely ergodic. This implies the well-known fact that, when dealing with square matrices, not necessarily in the constant length case, if a matrix with positive coefficients occurs infinitely many times in the sequence  $(M_{\tau_n} : n \geq 0)$ , then  $(X_\tau, S)$  is uniquely ergodic. This criteria for unique ergodicity can be tracked down to [Fur60].

For the constant length case, we do not need all coefficients to be positive.

**Corollary 14.** *Let  $\tau = (\tau_n : \mathcal{A}_{n+1}^* \rightarrow \mathcal{A}_n^*)_{n \geq 0}$  be a constant length and recognizable directive sequence. Suppose that there exists a morphism  $\tau : \mathcal{A}^* \rightarrow \mathcal{B}^*$  which satisfies:*

- (1) *There exists a letter  $b$  in  $\mathcal{B}$  which occurs in  $\tau(a)$  for every  $a \in \mathcal{A}$ ; and*
- (2) *We have  $\tau_n = \tau$  for infinitely many values of  $n$ .*

*Then  $(X_\tau, S)$  is uniquely ergodic.*

Our result also implies unique ergodicity when matrices are upper or lower triangular with positive coefficients (in the corresponding triangular and diagonal parts of the matrices) and one of each type occurring infinitely many times.

Theorem 11 covers more situations in the constant length case than classical criteria. For example, one cannot deduce from the Seneta criteria the unique ergodicity when

$$M_{\tau_n} = \begin{pmatrix} 1 & n^2 \\ 2n^2 - 1 & n^2 \end{pmatrix}, \quad n \geq 0$$



whereas it is uniquely ergodic as  $|v_n| = 1 + n^2$ . Moreover, in our framework we do not require  $(|\mathcal{A}_n| : n \geq 0)$  to be bounded, which is a constraint in [Sen81].

Let us mention that when

$$M_{\tau_n} = \begin{pmatrix} a_n & 1 \\ 1 & a_n \end{pmatrix}, \text{ where } a_n > 0$$

it is shown in [FFT09] that  $(X_\tau, S)$  is uniquely ergodic if and only if  $\sum_n 1/a_n$  diverges.

We finally remark in this section that there exist other result about ergodic measures in the context of constant length directive sequences. However, these results go in the direction of characterizing when the system possesses exactly  $k$  ergodic measures, where  $k$  is the rank of every composition matrix in the directive sequence [ABKK17, Theorem 3.7].

#### 4. PRELIMINARY DEFINITIONS AND RESULTS

In this section we include all main preliminary lemmas that allow us to deduce our main results in the next section.

Let  $\tau = (\tau_n : \mathcal{A}_{n+1}^* \rightarrow \mathcal{A}_n^*)_{n \geq 0}$  be a constant length and recognizable directive sequence and let  $\mu$  be an ergodic measure of  $(X_\tau, S)$ . For each  $n \geq 0$ , let  $\mathcal{A}'_n \subseteq \mathcal{A}_n$  be a nonempty alphabet. For convenience, we set

$$\text{coinc}'(\tau_{[n,N]}) = \text{coinc}_{\mathcal{A}'_N}(\tau_{[n,N]}), \quad N > n$$

where  $\text{coinc}_{\mathcal{A}'_N}(\tau_{[n,N]})$  is the set of coincidences of the morphism  $\tau_{[n,N]}$  relatively to  $\mathcal{A}'_N$  introduced in Section 2.4.

From now on, we suppose that  $\mu$  is  $(\mathcal{A}'_n)_{n \geq 0}$ -adapted to  $\mu$ . We recall the definition of the sequence of partitions  $(\mathcal{P}_n)_{n \geq 0}$  given in Section 2.6 as well as the definition of the sets  $B'_n, \mathcal{T}'_n, \mathcal{T}'_{\geq n}$  for  $n \geq 0$  and  $\mathcal{T}'$  given in Section 2.7.

In the following we aim to control the set (and their measure) of  $z \in \mathbb{Z}_{(|\tau_{[0,n]}|)_{n \geq 0}}$  having a unique  $\pi_{\text{meq}}$ -preimage. We set

$$\begin{aligned} \mathcal{I}' &= \{z \in \mathbb{Z}_{(|\tau_{[0,n]}|)_{n \geq 0}} : |\pi_{\text{meq}}^{-1}(\{z\}) \cap \mathcal{T}'| \leq 1\} \text{ and} \\ \mathcal{I}'_m &= \{z \in \mathbb{Z}_{(|\tau_{[0,n]}|)_{n \geq 0}} : \pi_{\text{meq}}^{-1}(\{z\}) \cap \mathcal{T}'_{\geq m} \text{ is contained in some atom of } \mathcal{P}_m\} \end{aligned}$$

for each  $m \geq 0$ .

We recall that  $\nu$  is the Haar measure on  $\mathbb{Z}_{(|\tau_{[0,n]}|)_{n \geq 0}}$ .

**Lemma 15.** *The set  $\mathcal{I}'$  is equal to  $\bigcap_{n \geq 0} \mathcal{I}'_n$  up to a set of  $\nu$ -measure zero.*

*Proof.* Observe that the inclusion  $\mathcal{I}' \subseteq \bigcap_{n \geq 0} \mathcal{I}'_n$  follows directly from the definitions. Thus, it is enough to prove that

$$(26) \quad \nu(\bigcap_{n \geq 0} \mathcal{I}'_n) \leq \nu(\mathcal{I}').$$

We observe that if  $z$  belongs to  $\mathcal{I}'_n$  for every  $n \geq 0$ , then all points in  $\pi_{\text{meq}}^{-1}(\{z\}) \cap \mathcal{T}'$  share the same  $(\mathcal{P}_n)_{n \geq 0}$ -address. Therefore, by Theorem 4 we deduce

$$\mu(\pi_{\text{meq}}^{-1}(\bigcap_{n \geq 0} \mathcal{I}'_n)) \leq \mu(\pi_{\text{meq}}^{-1}(\mathcal{I}')),$$

which implies (26).  $\square$

The next lemma allows us to interpret coincidences in topological terms. The proof is left to the reader since it is straightforward from the definitions.

**Lemma 16.** *Let  $0 \leq n < m$ . Then  $k$  belongs to  $\text{coinc}'(\tau_{[n,m]})$  if and only if there exists a letter  $a \in \mathcal{A}_n$  such that  $S^{k|\tau_{[0,n]}|}B'_m \subseteq B_n(a)$ .*

The next lemma is a first step to control the measure of the set of points having a unique  $\pi_{\text{meq}}$ -preimage.

**Lemma 17.** *We have that*

$$\nu(\mathcal{I}'_n) \geq \sup_{m>n} \frac{|\text{coinc}'(\tau_{[n,m]})|}{|\tau_{[n,m]}|}, \quad n \geq 0.$$

*Proof.* Let  $n \geq 0$  and define the sets

$$D_{n,m} = \{z \in \mathbb{Z}_{(|\tau_{[0,n]}|)_{n \geq 0}} : z_m \in z_n + |\tau_{[0,n]}| \text{coinc}'(\tau_{[n,m]})\}, \quad m > n.$$

We first prove that  $D_{n,m}$  is contained in  $\mathcal{I}'_n$  for all  $m > n$ . Indeed, if  $z \in D_{n,m}$  from Lemma 16 there exists a letter  $a \in \mathcal{A}_n$  such that

$$S^{z_m} B'_m \subseteq S^{z_n} B_n(a)$$

Thus if  $x$  is in  $\pi_{\text{meq}}^{-1}(\{z\}) \cap \mathcal{T}'_{\geq n}$  then  $x$  is in the atom  $S^{z_n} B_n(a)$  and thus  $z$  is in  $\mathcal{I}'_n$ .

From Lemma 9, we obtain

$$\nu(\mathcal{I}'_n) \geq \sup_{m>n} \nu(D_{n,m}) = \sup_{m>n} \frac{|\text{coinc}'(\tau_{[n,m]})|}{|\tau_{[n,m]}|}.$$

□

As it is more easy to control the positions that are not coincidences, we give the following definition. For  $0 \leq n < m$  and letters  $a$  and  $b$  in  $\mathcal{A}_n$  we define

$$C_{n,m}(a, b) = \{k \in [0, |\tau_{[n,m]}|) : S^{k|\tau_{[0,n]}|}B'_m \cap B_n(c) \neq \emptyset, \quad \forall c \in \{a, b\}\}.$$

Observe that, since the partition  $\mathcal{P}_m$  is finer than  $\mathcal{P}_n$ , for all  $k$  in  $C_{n,m}(a, b)$  and  $c$  in  $\{a, b\}$  there exists a letter  $c'$  in  $\mathcal{A}'_m$  such that

$$(27) \quad S^{k|\tau_{[0,n]}|}B_m(c') \subseteq B_n(c).$$

The following lemma will allow us define two disjoint sets in  $X_\tau$  with the same projection under  $\pi_{\text{meq}}$  and whose  $\nu$ -measure is controlled in terms of coincidences.

**Lemma 18.** *Let  $n \geq 0$ . Then there exist distinct letters  $a$  and  $b$  in  $\mathcal{A}_n$  and an increasing sequence  $(N_\ell)_{\ell \geq 0}$  such that*

$$(28) \quad \frac{|C_{n,N_\ell}(a, b)|}{|\tau_{[n,N_\ell]}|} \geq \frac{1}{2|\mathcal{A}_n|^2} \left( 1 - \limsup_{\ell \rightarrow +\infty} \frac{|\text{coinc}'(\tau_{[n,N_\ell]})|}{|\tau_{[n,N_\ell]}|} \right), \quad \ell \geq 0.$$

*If in addition the directive sequence  $\tau$  is  $(\mathcal{A}'_n)_{n \geq 0}$ -adapted to  $\mu$ , then  $(N_\ell)_{\ell \geq 0}$  can be defined so that it satisfies*

$$(29) \quad \frac{|\tau_{[N_\ell, N_{\ell+1}]}(c)|_d}{|\tau_{[N_\ell, N_{\ell+1}]}|} \leq \frac{1}{2^\ell |\mathcal{A}_{N_\ell}|} + \mu(\mathcal{T}_{N_\ell}(d)), \quad c \in \mathcal{A}'_{N_{\ell+1}}, \quad d \in \mathcal{A}_{N_\ell}, \quad \ell \geq 0.$$

*Proof.* Let  $n \geq 0$  and  $n_0 \geq n$  such that

$$(30) \quad \frac{|[0, |\tau_{[n,N]}|) \setminus \text{coinc}'(\tau_{[n,N]})|}{|\tau_{[n,N]}|} \geq \frac{1}{2} \left( 1 - \limsup_{N \rightarrow +\infty} \frac{|\text{coinc}'(\tau_{[n,N]})|}{|\tau_{[n,N]}|} \right), \quad N > n_0.$$

By definition, for every  $N > n_0$  and  $k \in [0, |\tau_{[n,N]}|) \setminus \text{coinc}'(\tau_{[n,N]})$  there exist distinct letters  $a_{N,k}$  and  $b_{N,k}$  in  $\mathcal{A}_n$  and letters  $a'_{N,k}$  and  $b'_{N,k}$  in  $\mathcal{A}'_N$  such that

the  $k$ -th letter of  $\tau_{[n,N]}(a'_{N,k})$  (resp. of  $\tau_{[n,N]}(b'_{N,k})$ ) is  $a_{N,k}$  (resp.  $b_{N,k}$ ). This translates into

$$S^{k|\tau_{[0,n]}|}B'_N \cap B_n(c) \neq \emptyset, \quad \forall c \in \{a_{N,k}, b_{N,k}\},$$

or  $k \in C_{n,N}(a_{N,k}, b_{N,k})$ . Now, for each  $N > n_0$  we use the Pigeonhole principle to obtain distinct letters  $a_N$  and  $b_N$  in  $\mathcal{A}_n$  and a subset  $C'_{n,N}$  of  $C_{n,N}(a_N, b_N)$  with

$$a_{N,k} = a_N, \quad b_{N,k} = b_N, \quad k \in C'_{n,N},$$

$$(31) \quad |C'_{n,N}| \geq |[0, |\tau_{[n,N]}|) \setminus \text{coinc}'(\tau_{[n,N]})|/|\mathcal{A}_n|^2.$$

Moreover, we can find an increasing sequence  $(N_\ell)_{\ell \geq 0}$  with  $N_0 > n_0$  and distinct letters  $a, b \in \mathcal{A}_n$  such that  $a = a_{N_\ell}$  and  $b = b_{N_\ell}$  for all  $\ell \geq 0$ . Therefore the sets  $C_{n,N_\ell}(a, b)$  satisfy (28) by (30) and (31).

If  $\tau$  is  $(\mathcal{A}'_n)_{n \geq 0}$ -adapted to  $\mu$ , from Proposition 6 we can construct  $(N_\ell)_{\ell \geq 0}$  inductively so that it additionally satisfies (29).  $\square$

The next lemma provides two disjoint sets that, roughly speaking, separate the fibers and whose measures can be controlled by the density of coincidences. It is the main lemma of this section. Its proof is subdivided in some technical claims.

**Lemma 19.** *Let  $n \geq 0$ . There exist sets  $E_n$  and  $E'_n$  in  $X_\tau$  such that*

- (1)  $E_n$  and  $E'_n$  are disjoint;
- (2)  $\pi_{\text{meq}}(E_n) = \pi_{\text{meq}}(E'_n)$ ; and

$$\nu(\pi_{\text{meq}}(E_n)) = \nu(\pi_{\text{meq}}(E'_n)) \geq \frac{1}{2|\mathcal{A}_n|^2} \left( 1 - \limsup_{N \rightarrow +\infty} \frac{|\text{coinc}'(\tau_{[n,N]})|}{|\tau_{[n,N]}|} \right).$$

If in addition the directive sequence  $\tau$  is  $(\mathcal{A}'_n)_{n \geq 0}$ -adapted to  $\mu$  for the constant  $\delta > 0$ , then

- (1)  $\mu(E_n)$  and  $\mu(E'_n)$  are greater than

$$\frac{\delta}{2|\mathcal{A}_n|^2} \left( 1 - \limsup_{N \rightarrow +\infty} \frac{|\text{coinc}'(\tau_{[n,N]})|}{|\tau_{[n,N]}|} \right); \quad \text{and}$$

- (2)  $\pi_{\text{meq}}(E_n \cap U) = \pi_{\text{meq}}(E'_n \cap U) \pmod{\nu}$  for any Borel set  $U$  in  $X_\tau$  such that  $\mu(U) = 1$ .

*Proof.* From Lemma 18, there exist distinct letters  $a$  and  $b$  in  $\mathcal{A}_n$  and an increasing sequence  $(N_\ell)_{\ell \geq 0}$  such that (28) holds. For  $\ell \geq 0$  define  $C_{n,N_\ell} = C_{n,N_\ell}(a, b)$  and observe that if  $k$  is in  $C_{n,N_\ell}$  and  $c$  is in  $\{a, b\}$ , then (27) implies that there exists a nonempty set  $\mathcal{A}_{n,N_\ell,k}(c) \subseteq \mathcal{A}'_{N_\ell}$  such that

$$S^{k|\tau_{[0,n]}|}B_{N_\ell}(c') \subseteq B_n(c), \quad c' \in \mathcal{A}_{n,N_\ell,k}(c).$$

Let  $K \subseteq X_\tau$  be a compact set. For all  $\ell \geq 0$  we set

$$\begin{aligned} F_{n,N_\ell} &= \{z \in \mathbb{Z}_{(|\tau_{[0,n]}|)_{n \geq 0}} : z_{N_\ell} \in z_n + |\tau_{[0,n]}|C_{n,N_\ell}\}; & F_{n,\infty} &= \limsup_{\ell \rightarrow +\infty} F_{n,N_\ell} \\ F_{n,N_\ell,K}(c) &= \{z \in F_{n,N_\ell} : S^{z_{N_\ell}} B'_{N_\ell} \cap S^{z_n} B_n(c) \cap K \neq \emptyset\}; & c &\in \{a, b\} \\ F_{n,\infty,K} &= \limsup_{\ell \rightarrow +\infty} F_{n,N_\ell,K}(a); \\ F'_{n,\infty,K} &= \limsup_{\ell \rightarrow +\infty} F_{n,N_\ell,K}(b) \\ E_{n,N_\ell}(c) &= \bigcup_{k \in C_{n,N_\ell}} \bigcup_{\substack{c' \in \mathcal{A}_{n,N_\ell,k}(c) \\ 0 \leq j < |\tau_{[0,n]}|}} S^{k|\tau_{[0,n]}|+j} B_{N_\ell}(c'); & c &\in \{a, b\} \\ E_n &= \limsup_{\ell \rightarrow +\infty} E_{n,N_\ell}(a); \\ E'_n &= \limsup_{\ell \rightarrow +\infty} E_{n,N_\ell}(b). \end{aligned}$$

Observe that, from the definition of the sets  $C_{n,N_\ell}$ , we have

$$F_{n,\infty,X_\tau} = F'_{n,\infty,X_\tau} = F_{n,\infty}$$

and that

$$E_n \subseteq \mathcal{T}_n(a) \text{ and } E'_n \subseteq \mathcal{T}_n(b).$$

In particular,  $E_n$  and  $E'_n$  are disjoint.

**Claim 20.** *For all compact set  $K \subseteq X_\tau$  we have*

$$\pi_{\text{meq}}(E_n \cap K) = F_{n,\infty,K} \text{ and } \pi_{\text{meq}}(E'_n \cap K) = F'_{n,\infty,K}.$$

*Proof.* By symmetry, it is enough to show that  $\pi_{\text{meq}}(E_n \cap K) = F_{n,\infty,K}$ . Observe that if  $x$  belongs to  $E_{n,N_\ell}(a) \cap K$  then  $\pi_{\text{meq}}(x)$  is in  $F_{n,N_\ell,K}(a)$ ; hence we have the inclusion

$$\pi_{\text{meq}}(E_n \cap K) \subseteq F_{n,\infty,K}.$$

Suppose now that  $z$  belongs to  $F_{n,\infty,K}$ . Then, there is an infinite set  $\Lambda \subset \mathbb{N}$  such that  $z$  belongs to  $F_{n,N_\ell}$  and a point  $x_\ell$  in  $S^{z_{N_\ell}} B'_{N_\ell} \cap S^{z_n} B_n(a) \cap K$  for all  $\ell \in \Lambda$ . Since  $X_\tau$  is compact, we can find an infinite set  $\Lambda' \subseteq \Lambda$  and  $x \in X_\tau$  such that  $x_\ell \rightarrow x$  as  $\ell \rightarrow +\infty$  and  $\ell \in \Lambda'$ . Then  $z = \pi_{\text{meq}}(x)$  and  $x \in E_n \cap K$  since  $K$  is compact, which shows that  $F_{n,\infty,K} \subseteq \pi_{\text{meq}}(E_n \cap K)$ .  $\square$

**Claim 21.** *We have  $\pi_{\text{meq}}(E_n) = \pi_{\text{meq}}(E'_n) = F_{n,\infty}$  and*

$$\nu(F_{n,\infty}) \geq \frac{1}{2|\mathcal{A}_n|^2} \left( 1 - \limsup_{N \rightarrow +\infty} \frac{|\text{coinc}'(\tau_{[n,N]})|}{|\tau_{[n,N]}|} \right).$$

*Proof.* If we put  $K = X_\tau$  in Claim 20 we obtain  $\pi_{\text{meq}}(E_n) = \pi_{\text{meq}}(E'_n) = F_{n,\infty}$ . On the other hand, from Lemma 9 and (28) we obtain

$$\nu(F_{n,\infty}) \geq \limsup_{\ell \rightarrow +\infty} \frac{|C_{n,N_\ell}|}{|\tau_{[n,N_\ell]}|} \geq \frac{1}{2|\mathcal{A}_n|^2} \left( 1 - \limsup_{N \rightarrow +\infty} \frac{|\text{coinc}'(\tau_{[n,N]})|}{|\tau_{[n,N]}|} \right).$$

$\square$

This proves the first part of Lemma 19.

We henceforth assume that  $\tau$  is  $(\mathcal{A}'_n)_{n \geq 0}$ -adapted to  $\mu$  for the constant  $\delta > 0$ .

**Claim 22.**  $\mu(E_n)$  and  $\mu(E'_n)$  are at least

$$\frac{\delta}{2|\mathcal{A}_n|^2} \left( 1 - \limsup_{N \rightarrow +\infty} \frac{|\text{coinc}'(\tau_{[n,N]})|}{|\tau_{[n,N]}|} \right).$$

*Proof.* Observe that, for each  $\ell \geq 0$  and  $c \in \{a, b\}$ , the sets

$$\{S^{k|\tau_{[0,n]}|+j} B_{N_\ell}(c') : k \in C_{n,N_\ell}, \ 0 \leq j < |\tau_{[0,n]}|, \ c' \in \mathcal{A}_{n,N_\ell,k}(c)\}$$

are disjoint. Since  $\mathcal{A}_{n,N_\ell,k}(c) \subseteq \mathcal{A}'_{N_\ell}$ , from (12) we have

$$\min_{c' \in \mathcal{A}_{n,N_\ell,k}(c)} \mu(B_{N_\ell}(c')) \geq \delta/|\tau_{[0,N_\ell]}|.$$

Thus we obtain

$$\mu(E_{n,N_\ell}(c)) \geq |C_{n,N_\ell}| \cdot |\tau_{[0,n]}| \min_{c' \in \mathcal{A}_{n,N_\ell,k}(c)} \mu(B_{N_\ell}(c')) \geq \delta \frac{|C_{n,N_\ell}|}{|\tau_{[n,N_\ell]}|}$$

and (28) implies

$$\mu \left( \limsup_{\ell \rightarrow +\infty} E_{n,N_\ell}(c) \right) \geq \frac{\delta}{2|\mathcal{A}_n|^2} \left( 1 - \limsup_{N \rightarrow +\infty} \frac{|\text{coinc}'(\tau_{[n,N]})|}{|\tau_{[n,N]}|} \right).$$

□

**Claim 23.** For all compact set  $K \subseteq X_\tau$  we have

$$\nu \left( \limsup_{\ell \rightarrow +\infty} F_{n,N_\ell,K}(c) \right) \geq \limsup_{\ell \rightarrow +\infty} \nu(F_{n,N_\ell}) - \frac{1}{\delta} \mu(X_\tau \setminus K), \quad c \in \{a, b\}.$$

*Proof.* Let  $c \in \{a, b\}$  and  $\ell \geq 0$ . We define  $\mathcal{R}_{n,N_\ell,K}(c)$  as the set of coordinates  $z_{N_\ell}$  in  $[0, |\tau_{[0,N_\ell]}|)$  where  $z$  ranges over  $F_{n,N_\ell} \setminus F_{n,N_\ell,K}(c)$ .

From (15) we have

$$(32) \quad \nu(F_{n,N_\ell} \setminus F_{n,N_\ell,K}(c)) \leq \nu \left( \bigcup_{r \in \mathcal{R}_{n,N_\ell,K}(c)} \{z \in \mathbb{Z}_{(|\tau_{[0,n]}|)_{n \geq 0}} : z_{N_\ell} = r\} \right) \leq \frac{|\mathcal{R}_{n,N_\ell,K}(c)|}{|\tau_{[0,N_\ell]}|}.$$

Observe that if  $z$  and  $z'$ , with  $z_{N_\ell} \neq z'_{N_\ell}$ , belong to  $F_{n,N_\ell} \setminus F_{n,N_\ell,K}(c)$ , then  $S^{z_{N_\ell}} B'_{N_\ell} \cap S^{z_n} B_n(c)$  and  $S^{z'_{N_\ell}} B'_{N_\ell} \cap S^{z'_n} B_n(c)$  are disjoint and contained in  $X_\tau \setminus K$ . Hence,

$$|\mathcal{R}_{n,N_\ell,K}(c)| \min_{z \in F_{n,N_\ell} \setminus F_{n,N_\ell,K}(c)} \mu(S^{z_{N_\ell}} B'_{N_\ell} \cap S^{z_n} B_n(c)) \leq \mu(X_\tau \setminus K).$$

Now, if  $z$  belongs to  $F_{n,N_\ell} \setminus F_{n,N_\ell,K}(c)$ , then from (27) we have that  $S^{z_{N_\ell}} B'_{N_\ell} \cap S^{z_n} B_n(c)$  contains  $S^{z_{N_\ell}} B'_{N_\ell}(c')$  for some  $c' \in \mathcal{A}'_{N_\ell}$ . Hence, by (12),

$$\min_{z \in F_{n,N_\ell} \setminus F_{n,N_\ell,K}(c)} \mu(S^{z_{N_\ell}} B'_{N_\ell} \cap S^{z_n} B_n(c)) \geq \min_{c' \in \mathcal{A}'_{N_\ell}} \mu(B_{N_\ell}(c')) \geq \frac{\delta}{|\tau_{[0,N_\ell]}|}.$$

Combining this with (32) yields

$$\nu(F_{n,N_\ell} \setminus F_{n,N_\ell,K}(c)) \leq \frac{1}{\delta} \mu(X_\tau \setminus K),$$

and then we finally obtain

$$\nu \left( \limsup_{\ell \rightarrow +\infty} F_{n,N_\ell,K}(c) \right) \geq \limsup_{\ell \rightarrow +\infty} \nu(F_{n,N_\ell,K}(c)) \geq \limsup_{\ell \rightarrow +\infty} \nu(F_{n,N_\ell}) - \frac{1}{\delta} \mu(X_\tau \setminus K).$$

□

**Claim 24.** *We have*

$$\nu(F_{n,\infty}) = \limsup_{\ell \rightarrow +\infty} \nu(F_{n,N_\ell}).$$

*Proof.* Define

$$\begin{aligned} G_\ell &= \{k \in [0, |\tau_{[N_\ell, N_{\ell+1}]})|) : S^{k|\tau_{[0, N_\ell]}|} B'_{N_{\ell+1}} \subseteq B'_{N_\ell}\} \\ H_\ell &= \{z \in \mathbb{Z}_{(|\tau_{[0, n]}|)_{n \geq 0}} : S^{zN_{\ell+1}} B'_{N_{\ell+1}} \subseteq S^{zN_\ell} B'_{N_\ell}\}, \quad \ell \geq 0 \\ H_{\ell^*, \infty} &= \bigcap_{\ell \geq \ell^*} H_\ell, \quad \ell^* \geq 0. \end{aligned}$$

Observe that  $k$  belongs to  $G_\ell$  if and only if for all  $c$  in  $\mathcal{A}'_{N_{\ell+1}}$  the  $k$ -th letter of  $\tau_{[N_\ell, N_{\ell+1})}(c)$  is in  $\mathcal{A}'_{N_\ell}$ . Hence

$$|[0, |\tau_{[N_\ell, N_{\ell+1})}|) \setminus G_\ell| \leq \sum_{c \in \mathcal{A}'_{N_{\ell+1}}} \sum_{d \in \mathcal{A}_{N_\ell} \setminus \mathcal{A}'_{N_\ell}} |\tau_{[N_\ell, N_{\ell+1})}(c)|_d.$$

From (29) we obtain

$$\begin{aligned} \frac{|[0, |\tau_{[N_\ell, N_{\ell+1})}|) \setminus G_\ell|}{|\tau_{[N_\ell, N_{\ell+1})}|} &\leq \sum_{c \in \mathcal{A}'_{N_{\ell+1}}} \sum_{d \in \mathcal{A}_{N_\ell} \setminus \mathcal{A}'_{N_\ell}} \frac{|\tau_{[N_\ell, N_{\ell+1})}(c)|_d}{|\tau_{[N_\ell, N_{\ell+1})}|} \\ &\leq \sup_{n \geq 0} |\mathcal{A}'_n| \left( \frac{1}{2^\ell} + \mu(X_\tau \setminus \mathcal{T}'_{N_\ell}) \right). \end{aligned}$$

On the other hand, one has  $z \in H_\ell$  if and only if  $zN_{\ell+1} \in zN_\ell + |\tau_{[0, N_\ell]}|G_\ell$ . From Lemma 9 we have

$$\nu(H_\ell) = |G_\ell|/|\tau_{[N_\ell, N_{\ell+1})}|, \quad \ell \geq 0.$$

Therefore, for  $\ell^* \geq 0$  one has

$$\nu\left(X_\tau \setminus \bigcap_{\ell \geq \ell^*} H_\ell\right) \leq \sum_{\ell \geq \ell^*} \sup_{n \geq 0} |\mathcal{A}'_n| \left( \frac{1}{2^\ell} + \mu(X_\tau \setminus \mathcal{T}'_{N_\ell}) \right).$$

Recall that by (12) we have  $\sup_{n \geq 0} |\mathcal{A}'_n| \leq 1/\delta$ , hence by (11)

$$(33) \quad \lim_{\ell^* \rightarrow +\infty} \nu\left(X_\tau \setminus \bigcap_{\ell \geq \ell^*} H_\ell\right) = 0.$$

Now let  $\ell^* \leq \ell < \ell'$ . Observe that if  $z$  belongs to  $F_{n, N_{\ell'}} \cap H_{\ell^*, \infty}$ , then for  $c$  in  $\{a, b\}$  we have  $S^{zN_\ell} B'_{N_\ell} \cap S^{zN_n} B_n(c) \subseteq S^{zN_{\ell'}} B'_{N_{\ell'}} \cap S^{zN_n} B_n(c)$ , which is nonempty. We deduce that  $z$  belongs to  $F_{n, N_\ell}$  and so

$$F_{n, N_{\ell'}} \cap H_{\ell^*, \infty} \subseteq F_{n, N_\ell} \cap H_{\ell^*, \infty},$$

*i.e.*, the sequence of sets  $(F_{n, N_\ell} \cap H_{\ell^*, \infty})_{\ell \geq \ell^*}$  is decreasing. We obtain

$$\begin{aligned} &\nu\left(\limsup_{\ell \rightarrow +\infty} F_{n, N_\ell}\right) \\ &= \nu\left(\limsup_{\ell \rightarrow +\infty} F_{n, N_\ell} \cap H_{\ell^*, \infty}\right) + \nu\left(\limsup_{\ell \rightarrow +\infty} F_{n, N_\ell} \cap (X_\tau \setminus H_{\ell^*, \infty})\right) \\ &\leq \limsup_{\ell \rightarrow +\infty} \nu(F_{n, N_\ell}) + \nu(X_\tau \setminus H_{\ell^*, \infty}). \end{aligned}$$

If we let  $\ell^* \rightarrow +\infty$ , from (33) we deduce

$$\nu(F_{n,\infty}) = \limsup_{\ell \rightarrow +\infty} \nu(F_{n,N_\ell}),$$

which proves the claim.  $\square$

**Claim 25.** *We have*

$$\pi_{\text{meq}}(E_n \cap U) = \pi_{\text{meq}}(E'_n \cap U) \pmod{\nu}$$

for any Borel set  $U \subset X_\tau$  such that  $\mu(U) = 1$ .

*Proof.* By symmetry, it is enough to prove that  $\pi_{\text{meq}}(E_n \cap U) = F_{n,\infty} \pmod{\nu}$ . Since  $\pi_{\text{meq}}(E_n) = F_{n,\infty}$ , we have the inclusion  $\pi_{\text{meq}}(E_n \cap U) \subseteq F_{n,\infty}$ . It remains to prove that  $\nu(\pi_{\text{meq}}(E_n \cap U)) \geq \nu(F_{n,\infty})$ . Let  $\varepsilon > 0$ . Since  $X_\tau$  is compact,  $\mu$  is a regular measure and there exists a compact set  $K_\varepsilon \subseteq U$  with  $\mu(X_\tau \setminus K_\varepsilon) < \varepsilon$ .

From Claim 20 we have

$$\pi_{\text{meq}}(E_n \cap K_\varepsilon) = F_{n,\infty,K_\varepsilon}$$

and from Claim 23 and Claim 24 we deduce

$$\nu(F_{n,\infty,K_\varepsilon}) \geq \limsup_{\ell \rightarrow +\infty} \nu(F_{n,N_\ell}) - \frac{\varepsilon}{\delta} \geq \nu(F_{n,\infty}) - \frac{\varepsilon}{\delta}.$$

Thus, since  $K_\varepsilon$  is a subset of  $U$ , one has

$$\nu(\pi_{\text{meq}}(E_n \cap U)) \geq \nu(F_{n,\infty,K_\varepsilon}) \geq \nu(F_{n,\infty}) - \frac{\varepsilon}{\delta}.$$

If we let  $\varepsilon$  go to 0, we deduce  $\nu(\pi_{\text{meq}}(E_n \cap U)) \geq \nu(F_{n,\infty})$ , which finishes the proof.  $\square$

The achieves the proof of the Lemma 19.  $\square$

## 5. PROOF OF THE MAIN RESULTS

We will freely use the terminology introduced in Section 4.

### 5.1. Proof of Theorem 1.

*Proof.* From Proposition 7 and recognizability of  $\tau$  it is easy to see that Item 2, Item 3, Item 4 and Item 5 in Theorem 1 are equivalent.

We define

$$a_{n,N} = 1 - \frac{|\text{coinc}(\tau_{[n,N]})|}{|\tau_{[n,N]}|}, \quad 0 \leq n < N.$$

From Lemma 3 we deduce that inequality (18) holds. Hence, we obtain that Item 4 and Item 6 are equivalent.

It remains to show that Item 1 and Item 4. We set  $\mathcal{A}'_n = \mathcal{A}_n$  for each  $n \geq 0$ . Observe that, with this choice, the directive sequence  $\tau$  is  $(\mathcal{A}'_n)_{n \geq 0}$ -weakly-adapted to any invariant probability measure  $\mu$  of  $(X_\tau, S)$ . In this case, we have

$$\mathcal{I}' = \{z \in \mathbb{Z}_{(|\tau_{[0,n]}|)_{n \geq 0}} : |\pi_{\text{meq}}^{-1}(\{z\})| = 1\}.$$

Suppose that Item 1 holds. By contradiction, if Item 4 does not hold, there exists  $n \geq 0$  such that

$$\limsup_{N \rightarrow +\infty} \frac{|\text{coinc}(\tau_{[n,N]})|}{|\tau_{[n,N]}|} < 1.$$

Lemma 19 ensures the existence of disjoint sets  $E_n$  and  $E'_n$  in  $X_\tau$  such that  $\nu(\pi_{\text{meq}}(E_n)) > 0$  and if  $z \in \pi_{\text{meq}}(E_n)$ , then  $\pi_{\text{meq}}^{-1}(\{z\}) \cap E_n \neq \emptyset$  and  $\pi_{\text{meq}}^{-1}(\{z\}) \cap$

$E'_n \neq \emptyset$ . Thus  $\mathbb{Z}_{(|\tau_{[0,n]}|)_{n \geq 0}} \setminus \mathcal{I}'$  contains  $\pi_{\text{meq}}(E_n)$ , which contradicts the fact that  $\nu(\mathcal{I}') = 1$ .

Now assume that Item 4 holds. Then, Lemma 15 and Lemma 17 imply that  $\nu(\mathcal{I}') = 1$ . This shows that Item 1 holds and finishes the proof.  $\square$

## 5.2. Proof of Theorem 2.

*Proof.* Suppose that Item 1 in Theorem 2 holds. This implies that there exists a Borel subset  $U$  of  $X_\tau$  and a Borel subset  $V$  of  $\mathbb{Z}_{(|\tau_{[0,n]}|)_{n \geq 0}}$  such that  $\pi_{\text{meq}} : U \rightarrow V$  is a bijection and  $\mu(U) = \nu(V) = 1$ .

By contradiction, if Item 2 does not hold, then there exists  $n \geq 0$  such that

$$\limsup_{N \rightarrow +\infty} \frac{|\text{coinc}_{\mathcal{A}'_N}(\tau_{[n,N]})|}{|\tau_{[n,N]}|} < 1.$$

Lemma 19 then ensures the existence of disjoint sets  $E_n$  and  $E'_n$  in  $X_\tau$  such that  $\mu(E_n) > 0$ ,  $\mu(E'_n) > 0$  and

$$\pi_{\text{meq}}(E_n \cap U) = \pi_{\text{meq}}(E'_n \cap U) \pmod{\nu}.$$

This implies, as  $\pi_{\text{meq}} : U \rightarrow V$  is a bijection, that  $E_n \cap U = E'_n \cap U \pmod{\mu}$ , which contradicts the fact that  $E_n$  and  $E'_n$  are disjoint and of positive measure.

Now assume that Item 2 holds. Define

$$X_\mu = \pi_{\text{meq}}^{-1}(\mathcal{I}') \cap \mathcal{T}'.$$

From Lemma 15, Lemma 17 and (13), we have  $\mu(X_\mu) = 1$  and  $\nu(\pi_{\text{meq}}(X_\mu)) = 1$ . On the other hand, the definition of  $X_\mu$  implies that  $\pi_{\text{meq}} : X_\mu \rightarrow \pi_{\text{meq}}(X_\mu)$  is a bijection. This shows that Item 1 holds and finishes the proof.  $\square$

*Remark 26.* Let us observe that when  $\tau$  is of alphabet rank two, then it follows from Lemma 3 and Item 6 in Theorem 1 that  $(X_\tau, S)$  is a regular extension of its maximal equicontinuous topological factor if and only if

$$\sum_{n \geq 0} \frac{|\text{coinc}(\tau_n)|}{|\tau_n|} = +\infty.$$

Indeed, it follows from Lemma 3 that if  $\tau' = (\tau'_k)_{k \geq 0}$  is a contraction of  $\tau$  then

$$\sum_{k \geq 0} \frac{|\text{coinc}(\tau'_k)|}{|\tau'_k|} \leq \sum_{n \geq 0} \frac{|\text{coinc}(\tau_n)|}{|\tau_n|}.$$

## 6. EXAMPLES AND APPLICATIONS

**6.1. Dekking's theorem revisited.** A *substitution* is a morphism  $\sigma : \mathcal{A}^* \rightarrow \mathcal{A}^*$  that is nonerasing. It naturally defines a directive sequence  $\tau = (\sigma, \sigma, \dots)$  which is of finite alphabet rank and which generates a (possibly empty) substitution subshift  $(X_\sigma, S)$ .

When  $\sigma$  is *primitive* (that is, the sequence  $\sigma$  is primitive) the subshift  $(X_\sigma, S)$  is minimal, has a unique ergodic measure  $\mu$  [Que87, Chapter V],  $\mathcal{A}_\mu = \mathcal{A}$  and  $\sigma$  is recognizable [Mos92, Mos96].

Let  $\sigma : \mathcal{A}^* \rightarrow \mathcal{A}^*$  be a constant length and primitive substitution. We say  $\sigma$  is *pure* if its maximal equicontinuous topological factor is the odometer  $(\mathbb{Z}_{|\sigma|}, +1)$ . We say  $\sigma$  admits a *coincidence* if there exists  $m \geq 1$  such that  $\text{coinc}(\sigma^m) \neq \emptyset$ , *i.e.*, if the directive sequence  $\sigma = (\sigma, \sigma, \dots)$  has coincidences.



As a consequence of Theorem 1 and Theorem 2, we recover the following result.

**Corollary 27** ([Dek78]). *The system  $(X_\sigma, S, \mu)$  where  $\sigma$  is a pure substitution has discrete spectrum if and only if  $\sigma$  admits a coincidence.*

*Proof.* Suppose that  $\sigma$  is pure and admits a coincidence. Let  $m \geq 1$  be such that  $\text{coinc}(\sigma^m) \neq \emptyset$ . From Equation (4) we deduce

$$1 - \frac{|\text{coinc}(\sigma^{mn})|}{|\sigma|^{mn}} \leq \left(1 - \frac{|\text{coinc}(\sigma^m)|}{|\sigma|^m}\right)^n \rightarrow 0 \text{ as } n \rightarrow +\infty$$

which implies by Theorem 1 that  $(X_\sigma, S, \mu)$  has discrete spectrum.

Now suppose that  $(X_\sigma, S, \mu)$  has discrete spectrum. Every measurable eigenvalue for the system  $(X_\sigma, S, \mu)$  is necessarily continuous [Hos86]. This implies, since  $\sigma$  is pure, that there exists a measure theoretical isomorphism  $\pi : (X_\sigma, S, \mu) \rightarrow (\mathbb{Z}_{|\sigma|}, +1, \nu)$ . As  $(\mathbb{Z}_{|\sigma|}, +1, \nu)$  is coalescent (see Section 2.10), the factor map  $\pi_{\text{meq}}$  is a measure theoretical isomorphism. Thanks to Theorem 2 and the fact that  $\mathcal{A}_\mu = \mathcal{A}$ , we deduce

$$\frac{|\text{coinc}(\sigma^n)|}{|\sigma^n|} \rightarrow 1 \text{ as } n \rightarrow +\infty$$

and hence  $\sigma$  has coincidences, finishing the proof.  $\square$

We remark that the previous proof shows that for primitive, pure substitutions  $\sigma$  the three hypothesis  $(X_\sigma, S)$  is a regular extension of  $(\mathbb{Z}_\sigma, +1)$ ;  $\pi_{\text{meq}} : (X_\sigma, S, \mu) \rightarrow (\mathbb{Z}_\sigma, +1, \nu)$  is a measure theoretical isomorphism; and  $(X_\sigma, S, \mu)$  has discrete spectrum are all equivalent.

**6.2. Example 1.** Rank-one transformations have been studied extensively in ergodic theory since the apparition of the first examples in the '60s. In order to study these systems from the topological and symbolic dynamics perspective, rank-one subshifts have been introduced. We refer to [Fer96, Fer97] for a more extensive discussion on these subshifts. Minimal rank-one subshifts are studied from the  $\mathcal{S}$ -adic perspective in [AD23], where the authors prefer to call them *Ferenczi subshifts*. In [AD23, Section 4.3] it is shown that a minimal Ferenczi subshift possesses an induced system that is (conjugate to) a Toeplitz subshift generated by the constant length directive sequence  $\tau = (\tau_n : \mathcal{A}_{n+1}^* \rightarrow \mathcal{A}_n^*)_{n \geq 0}$  which have the form

$$(34) \quad \tau_n(a) = L_n a R_n, \quad a \in \mathcal{A}_{n+1}$$

for some nonempty words  $L_n, R_n$  in  $\mathcal{A}_n^*$ . In particular, from (34) we immediately deduce

$$\frac{|\text{coinc}(\tau_n)|}{|\tau_n|} \geq \frac{2}{3}, \quad n \geq 0$$

and hence from Item 6 in Theorem 1 we obtain that this induced system is a regular extension of its maximal equicontinuous topological factor. This improves the remark made in [AD23] saying that these systems are *mean equicontinuous* (we refer to [GRJY21] for the definition and details about this notion).

**6.3. Example 2.** Let  $\mathcal{A} = \{a, b\}$ ,  $f : \mathbb{N} \rightarrow \mathbb{N}$  a nonnegative function such that for some pair of constants  $s > 1$  and  $c > 0$  we have  $f(n) \geq cn^s$  for  $n \geq 0$  and consider the directive sequence  $\tau = (\tau_n : \mathcal{A}^* \rightarrow \mathcal{A}^*)_{n \geq 0}$  such that for all  $n \geq 0$  the morphism  $\tau_n : \mathcal{A}^* \rightarrow \mathcal{A}^*$  is defined by

$$a \mapsto a^{f(n)}ba, \quad b \mapsto b^{f(n)}aa.$$

Observe that  $\tau$  is a constant length primitive directive sequence with coincidences. Moreover, the sequence  $\tau$  is recognizable since the composition matrix of the morphism  $\tau_n$  for each  $n \geq 0$  is

$$M_{\tau_n} = \begin{pmatrix} f(n) + 1 & 2 \\ 1 & f(n) \end{pmatrix}$$

which is invertible [BSTY19, Theorem 4.6]. Thus  $(X_\tau, S)$  is a Toeplitz subshift.

From Theorem 1 we have that  $(X_\tau, S)$  is not a regular extension of its maximal equicontinuous topological factor. Indeed, since  $\tau$  is of alphabet rank two, from Remark 26 we have

$$\sum_{n \geq 0} \frac{|\text{coinc}(\tau_n)|}{|\tau_n|} = \sum_{n \geq 0} \frac{1}{f(n) + 2} < +\infty$$

However, the hypothesis on  $f$  implies that  $(X_\tau, S)$  has two ergodic measures  $\mu_a$  and  $\mu_b$  [ABKK17, Proposition 3.1]. We set

$$\mathcal{A}_{\mu_a} = \{a\} \text{ and } \mathcal{A}_{\mu_b} = \{b\}.$$

We see that  $\tau$  is  $(\mathcal{A}_{\mu_a})$ -adapted to  $\mu_a$  and  $(\mathcal{A}_{\mu_b})$ -adapted to  $\mu_b$  [BKMS13, Theorem 3.3]. From Theorem 2 we conclude that  $\pi_{\text{meq}}$  defines a measure theoretical isomorphism to the maximal equicontinuous topological factor for both  $(X_\tau, S, \mu_a)$  and  $(X_\tau, S, \mu_b)$ .

**6.4. Example 3.** Let  $\mathcal{A} = \{a, b, c\}$ ,  $f : \mathbb{N} \rightarrow \mathbb{N}$  a nonnegative function such that for some pair of constants  $s > 1$  and  $c > 0$  we have  $f(n) \geq cn^s$  for  $n \geq 0$  and consider the directive sequence  $\tau = (\tau_n : \mathcal{A}^* \rightarrow \mathcal{A}^*)_{n \geq 0}$  such that for all  $n \geq 0$  the morphism  $\tau_n : \mathcal{A}^* \rightarrow \mathcal{A}^*$  is defined by

$$\begin{aligned} a &\mapsto (ab)^{f(n)+1}ac \\ b &\mapsto (ab)^{f(n)+1}bc \\ c &\mapsto c(ab)^{f(n)}ccc. \end{aligned}$$

Observe that  $\tau$  is a constant length primitive directive sequence with coincidences. Moreover, the sequence  $\tau$  is recognizable since the composition matrix of the morphism  $\tau_n$  for each  $n \geq 0$  is

$$(35) \quad M_{\tau_n} = \begin{pmatrix} f(n) + 2 & f(n) + 1 & f(n) \\ f(n) + 1 & f(n) + 2 & f(n) \\ 1 & 1 & 4 \end{pmatrix}$$

which is invertible [BSTY19, Theorem 4.6]. Thus  $(X_\tau, S)$  is a Toeplitz subshift.

From Theorem 1 we have that  $(X_\tau, S)$  is not a regular extension of its maximal equicontinuous topological factor. Indeed, by contradiction assume that

$$1 - \frac{|\text{coinc}(\tau_{[0,N]})|}{|\tau_{[0,N]}|} \rightarrow 0 \text{ as } N \rightarrow +\infty.$$

By induction, it is easy to see that

$$1 - \frac{|\text{coinc}(\tau_{[0,N+1]})|}{|\tau_{[0,N+1]}|} \geq \left(1 - \frac{1}{2f(N) + 4}\right) \left(1 - \frac{|\text{coinc}(\tau_{[0,N]})|}{|\tau_{[0,N]}|}\right), \quad N \geq 1$$

and hence

$$1 - \frac{|\text{coinc}(\tau_{[0,N]})|}{|\tau_{[0,N]}|} \geq \prod_{k=1}^{N-1} \left(1 - \frac{1}{2f(k)+4}\right) \left(1 - \frac{|\text{coinc}(\tau_0)|}{|\tau_0|}\right)$$

but it is standard to check that, under the hypothesis on  $f$ , the infinite product  $\prod_{k=1}^{\infty} \left(1 - \frac{1}{2f(k)+4}\right)$  does not converge to 0. This contradiction shows that  $(X_{\tau}, S)$  is not a regular extension of its maximal equicontinuous topological factor.

On the other hand, from (35) the vectors  $(v_n)_{n \geq 0}$  in the statement of Theorem 11 (for the trivial contraction of  $\tau$ ) satisfy

$$\sum_{n \geq 0} \frac{|v_n|}{|\tau_n|} = \sum_{n \geq 0} \frac{2f(n)+1}{2f(n)+4} = +\infty$$

which implies that the system  $(X_{\tau}, S)$  is uniquely ergodic. Denote by  $\mu$  the unique invariant probability measure and let  $\mathcal{A}_{\mu} = \{a, b\}$ . We claim that  $\tau$  is  $(\mathcal{A}_{\mu})$ -adapted to  $\mu$ . Indeed, from (9) and (10) we have

$$\begin{aligned} \mu(\mathcal{T}_n(a)) &= \mu_n(a)|\tau_{[0,n]}| = |\tau_{[0,n]}| \sum_{\ell} M_{\tau_n}(a, \ell) \mu_{n+1}(\ell) \geq f(n)|\tau_{[0,n]}| \sum_{\ell} \mu_{n+1}(\ell) \\ &= f(n)|\tau_{[0,n]}| \sum_{\ell} \frac{\mu(\mathcal{T}_{n+1}(\ell))}{|\tau_{[0,n+1]}|} = \frac{f(n)}{2f(n)+4} \geq \frac{1}{6}. \end{aligned}$$

A similar computation shows  $\mu(\mathcal{T}_n(b)) \geq 1/6$ . On the other hand, we have

$$\begin{aligned} \sum_{n \geq 0} \mu(\mathcal{T}_n(c)) &= \sum_{n \geq 0} \mu_n(c)|\tau_{[0,n]}| = \sum_{n \geq 0} |\tau_{[0,n]}| \sum_{\ell} M_{\tau_n}(c, \ell) \mu_{n+1}(\ell) \\ &\leq \sum_{n \geq 0} 4|\tau_{[0,n]}| \sum_{\ell} \mu_{n+1}(\ell) = \sum_{n \geq 0} 4|\tau_{[0,n]}| \sum_{\ell} \frac{\mu(\mathcal{T}_{n+1}(\ell))}{|\tau_{[0,n+1]}|} \\ &\leq \sum_{n \geq 0} \frac{4}{2f(n)+4} < +\infty \end{aligned}$$

and thus conditions (11) and (12) are satisfied.

Finally, for all  $n \geq 0$  we have

$$\lim_{N \rightarrow +\infty} \frac{|\text{coinc}_{\mu}(\tau_{[n,N]})|}{|\tau_{[n,N]}|} = 1.$$

Indeed, it is easy to check that

$$1 - \frac{|\text{coinc}_{\mu}(\tau_{[n,N+1]})|}{|\tau_{[n,N+1]}|} \geq \frac{1}{2} \left(1 - \frac{|\text{coinc}(\tau_{[n,N]})|}{|\tau_{[n,N]}|}\right), \quad N \geq n$$

and hence

$$1 - \frac{|\text{coinc}(\tau_{[n,N]})|}{|\tau_{[n,N]}|} = \frac{1}{2^{N-n-1}} \left(1 - \frac{|\text{coinc}(\tau_n)|}{|\tau_n|}\right) \rightarrow 0 \text{ as } N \rightarrow +\infty.$$

From Theorem 2 we conclude that  $\pi_{\text{meq}}$  defines a measure theoretical isomorphism between  $(X_{\tau}, S, \mu)$  and its maximal equicontinuous topological factor.

6.5. **Example 4.** Let  $\mathcal{A} = \{a, b, c\}$ ,  $m : \mathbb{N} \rightarrow \mathbb{N}$  the function such that  $2m(n) + 3 = 3^n$  for all  $n \geq 0$  and consider the directive sequence  $\boldsymbol{\tau} = (\tau_n : \mathcal{A}^* \rightarrow \mathcal{A}^*)_{n \geq 0}$  such that for all  $n \geq 0$  the morphism  $\tau_n : \mathcal{A}^* \rightarrow \mathcal{A}^*$  is defined by

$$\begin{aligned} a &\mapsto (ab)^{m(n)}abc \\ b &\mapsto a(ab)^{m(n)}ac \\ c &\mapsto (ab)^{m(n)}cbc. \end{aligned}$$

Observe that  $\boldsymbol{\tau}$  is a constant length and primitive directive sequence with coincidences. Moreover, the sequence  $\boldsymbol{\tau}$  is recognizable since the composition matrix of the morphism  $\tau_n$  for each  $n \geq 0$  is

$$M_{\tau_n} = \begin{pmatrix} m(n) + 1 & m(n) + 2 & m(n) \\ m(n) + 1 & m(n) & m(n) + 1 \\ 1 & 1 & 2 \end{pmatrix}$$

which is invertible [BSTY19, Theorem 4.6]. Thus  $(X_{\boldsymbol{\tau}}, S)$  is a Toeplitz subshift with maximal equicontinuous topological factor  $(\mathbb{Z}_3, +1)$ . Moreover, from Theorem 11 it has a unique ergodic measure  $\mu$ . Nevertheless  $\pi_{\text{meq}}$  is not a measure theoretical isomorphism but  $(X_{\boldsymbol{\tau}}, S, \mu)$  has discrete spectrum. Indeed, an easy computation similar to the previous examples which is left to the reader shows that for  $\mathcal{A}_{\mu} = \{a, b\}$  we have  $\boldsymbol{\tau}$  is  $(\mathcal{A}_{\mu})$ -adapted to  $\mu$  and

$$1 - \frac{|\text{coinc}_{\mu}(\tau_{[0, N]})|}{|\tau_{[0, N]}|} = \prod_{k=1}^{N-1} \left(1 - \frac{2}{3^k}\right) \left(1 - \frac{|\text{coinc}_{\mu}(\tau_0)|}{|\tau_0|}\right),$$

which does not converge to 0 as  $N \rightarrow +\infty$  and from Theorem 2 we obtain that  $\pi_{\text{meq}}$  is not a measure theoretical isomorphism.

Let us show it has discrete spectrum. Let  $P$  be the subset of  $X_{\boldsymbol{\tau}}$  given by

$$P = \liminf_{n \rightarrow +\infty} (\mathcal{T}_n(a) \cup \mathcal{T}_n(b)).$$

The following arguments mainly come from [BDM10]. The Borel–Cantelli lemma implies that  $\mu(P) = 1$ . For  $n \geq 0$  define

$$f_n(x) = \begin{cases} 0 & \text{if } x \in S^j B_n(a), \text{ where } 0 \leq j < |\tau_{[0, n]}| \text{ is even} \\ 1 & \text{if } x \in S^j B_n(b) \cup S^j B_n(c), \text{ where } 0 \leq j < |\tau_{[0, n]}| \text{ is odd.} \end{cases}$$

Let  $A_n = \{x \in X_{\boldsymbol{\tau}} : f_n(x) \neq f_{n+1}(x)\}$ . A quick computation by cases shows that

$$A_n \subseteq \left( \bigcup_{0 \leq j < |\tau_{[0, n]}|} S^j B_{n+1}(b) \right) \cup \mathcal{T}_n(c) \cup \mathcal{T}_{n+1}(c).$$

It is easy to check that

$$\mu(\mathcal{T}_n(c)) \leq \frac{2}{3^n}, \quad \mu \left( \bigcup_{0 \leq j < |\tau_{[0, n]}|} S^j B_{n+1}(b) \right) \leq \frac{1}{3^n}$$

and consequently  $\sum \mu(A_n)$  converges. The Borel–Cantelli lemma again implies that the sequence  $(f_n)_{n \geq 0}$  converges  $\mu$ -almost everywhere to some measurable function  $f : X_{\boldsymbol{\tau}} \rightarrow \mathbb{Z}/2\mathbb{Z}$  that satisfies  $f(Sx) = f(x) + 1 \pmod{2}$  for  $\mu$ -almost every  $x$ .

Denote by  $(\mathbb{Z}_3 \times \mathbb{Z}/2\mathbb{Z}, +(1, 1))$  the product system of  $(\mathbb{Z}_3, +1)$  and  $(\mathbb{Z}/2\mathbb{Z}, +1)$  and let  $\nu$  be the Haar measure on this product. Consider the measurable factor map  $F : (X_\tau, S, \mu) \rightarrow (\mathbb{Z}_3 \times \mathbb{Z}/2\mathbb{Z}, +(1, 1), \nu)$  defined by

$$\begin{aligned} F : X_\tau &\rightarrow \mathbb{Z}_3 \times \mathbb{Z}/2\mathbb{Z} \\ x &\mapsto (\pi_{\text{meq}}(x), f(x)) \end{aligned}$$

We claim that  $F$  defines a measure theoretical isomorphism. Indeed, it is sufficient to prove that the map  $F$  is injective on  $P$ . If  $F(x) = F(y) = (z, w)$  for some  $x, y$  in  $P$ , we have that for all large values of  $n$  we have both  $x$  and  $y$  belonging to  $S^{z_n} \mathcal{T}_n(w)$ , where  $z = (z_n)_{n \geq 0}$ . Hence, since  $\tau$  is proper, the atoms of the sequence of partitions  $(\mathcal{T}_n)_{n \geq 0}$  generate the topology of  $(X_\tau, S)$  (see Section 2.6) and we obtain  $x = y$ .

**6.6. Example 5.** Let  $\mathcal{A} = \{1, 2, a, b, c\}$ ,  $s : \mathbb{N} \rightarrow \mathbb{N}$  a nonnegative function such that  $2s(n) + 5 = 3^n$  for  $n \geq 0$  and consider the directive sequence  $\tau = (\tau_n : \mathcal{A}^* \rightarrow \mathcal{A}^*)_{n \geq 0}$  such that for all  $n \geq 0$  the morphism  $\tau_n : \mathcal{A}^* \rightarrow \mathcal{A}^*$  is defined by

$$\begin{aligned} 1 &\mapsto a(12)^{s(n)}12bc, & a &\mapsto (ab)^{s(n)}ab12c \\ 2 &\mapsto a(12)^{s(n)}21bc, & b &\mapsto a(ab)^{s(n)}b12c \\ & & c &\mapsto (ab)^{s(n)}ab12c. \end{aligned}$$

Observe that  $\tau$  is a constant length, recognizable and primitive directive sequence with coincidences. Thus  $(X_\tau, S)$  is a Toeplitz subshift with maximal equicontinuous topological factor  $(\mathbb{Z}_3, +1)$ .

The composition matrix of the morphism  $\tau_n$  for each  $n \geq 0$  is

$$M_{\tau_n} = \begin{pmatrix} s(n) + 1 & s(n) + 1 & 1 & 1 & 1 \\ s(n) + 1 & s(n) + 1 & 1 & 1 & 1 \\ 1 & 1 & s(n) + 1 & s(n) + 1 & s(n) + 1 \\ 1 & 1 & s(n) + 1 & s(n) + 1 & s(n) + 1 \\ 1 & 1 & 1 & 1 & 1 \end{pmatrix}$$

The subshift  $(X_\tau, S)$  has two ergodic measures  $\mu_{a,b}$  and  $\mu_{1,2}$ . Indeed, since the rank of each matrix  $M_{\tau_n}$  is two,  $(X_\tau, S)$  has at most two different ergodic measures. Moreover, the system is not uniquely ergodic thanks to Theorem 11: a simple yet tedious computation shows that for any increasing sequence  $(n_k)_{k \geq 0}$  the vectors  $(v_k)_{k \geq 0}$  satisfy  $|v_k| = \prod_{n_k < j \leq n_{k+1}} (2s(j) + 3)$  and so

$$\sum_{k \geq 0} \frac{|v_k|}{|\tau'_k|} = \frac{1}{2s(n_k) + 5} < +\infty.$$

where  $\tau'_k = \tau'_{[n_k, n_{k+1}]}$ ,  $k \geq 0$ .

Let  $\mathcal{A}_{\mu_{a,b}} = \{a, b\}$  and  $\mathcal{A}_{\mu_{1,2}} = \{1, 2\}$ . Observe that, from the form of the composition matrix  $M_{\tau_n}$ , for any ergodic measure  $\mu$  of  $(X_\tau, S)$  we have

$$\mu(\mathcal{T}_n(c)) \rightarrow 0, \quad \mu(\mathcal{T}_n(a)) = \mu(\mathcal{T}_n(b)) \quad \text{and} \quad \mu(\mathcal{T}_n(1)) = \mu(\mathcal{T}_n(2)).$$

Thus, without loss of generality,  $\tau$  is  $(\mathcal{A}_{\mu_{a,b}})$ -adapted to  $\mu_{a,b}$  and  $(\mathcal{A}_{\mu_{1,2}})$ -adapted to  $\mu_{1,2}$ . We left the details to the reader.

The map  $\pi_{\text{meq}}$  is a measure theoretical isomorphism with respect to  $\mu_{1,2}$  whereas it is not with respect to  $\mu_{a,b}$  as it has  $-1$  as an eigenvalue. Moreover, as in the previous example,  $(X_\tau, S, \mu_{a,b})$  has discrete spectrum: it is measure theoretically isomorphic to  $(\mathbb{Z}_3 \times \mathbb{Z}/2\mathbb{Z}, +(1, 1), \nu)$  (here  $\nu$  is the Haar measure on  $\mathbb{Z}_3 \times \mathbb{Z}/2\mathbb{Z}$ ).

We leave it to the reader to verify these statements in the light of the calculations made earlier.

**6.7. Example 6.** Above we gave example of Toeplitz subshifts with discrete spectrum. There are of course many Toeplitz subshifts that has a non discrete spectrum such that those with positive entropy. All examples we gave has a finite topological rank and are thus of entropy zero. It is interesting to give a finite topological rank Toeplitz shift that does not have a discrete spectrum.

Let  $\mathcal{A} = \{1, 2\}$ ,  $s : \mathbb{N} \rightarrow \mathbb{N}$  a nonnegative function such that  $3s(n) + 1 = 5^{2n}$  for  $n \geq 0$  and consider the directive sequence  $\tau = (\tau_n : \mathcal{A}^* \rightarrow \mathcal{A}^*)_{n \geq 0}$  such that for all  $n \geq 0$  the morphism  $\tau_n : \mathcal{A}^* \rightarrow \mathcal{A}^*$  is defined by

$$1 \mapsto (121)^{s(n)}2, \quad 2 \mapsto 1(121)^{s(n)}.$$

The directive sequence  $\tau$  is clearly recognizable. Let us show that  $(X_\tau, S)$  does not have non continuous eigenvalues. Suppose that  $\lambda = \exp(2i\pi\alpha)$  is such an eigenvalue. Then, from [BDM10, Proposition 28] one can suppose  $\alpha = 1/2$ . But from [DFM19, Corollary 16] and some computations one can show  $\lambda = -1$  cannot be an eigenvalue.

Thus, the maximal equicontinuous measurable factor is  $(\mathbb{Z}_5, +1)$  and it cannot be measure theoretically isomorphic to  $(X_\tau, S)$  as  $(\mathbb{Z}_5, +1)$  is coalescent [HP68]. Indeed, otherwise  $\pi_{\text{meq}}$  would be a measure theoretical isomorphism which is not possible because  $\tau$  does not fulfill Item 2 of Theorem 2.

**6.8. Example 7.** We finish this section with the description of a non Toeplitz subshift generated by a constant length directive sequence that has positive topological entropy and where the conclusion of Theorem 2 holds.

Let  $(\ell_n)_{n \geq 0}$  be a sequence of nonnegative numbers such that

$$\ell_{n+1} = (\ell_n - 1)! + 1, \quad n \geq 0.$$

For  $n \geq 0$  let  $\mathcal{A}_n = \{a_n, b_n(1), \dots, b_n(\ell_n - 1)\}$ . The choice of  $\ell_{n+1}$  implies that there exists a bijection  $\pi_n$  between  $\{b_{n+1}(1), \dots, b_{n+1}(\ell_{n+1} - 1)\}$  and the set of all words in  $\mathcal{A}_n^*$ , which can be defined using each letter in  $\{b_n(1), \dots, b_n(\ell_n - 1)\}$  exactly once. Let  $\pi'_n$  be an onto map between  $\{b_{n+1}(1), \dots, b_{n+1}(\ell_{n+1} - 1)\}$  and  $\mathcal{A}_n^2$ . Consider the directive sequence  $\tau = (\tau_n : \mathcal{A}_{n+1}^* \rightarrow \mathcal{A}_n^*)_{n \geq 0}$  such that for all  $n \geq 0$  the morphism  $\tau_n : \mathcal{A}_{n+1}^* \rightarrow \mathcal{A}_n^*$  is defined by

$$\begin{aligned} \tau_n(b_{n+1}(i)) &= \pi_n(b_{n+1}(i))\pi'_n(b_{n+1}(i))a_n, \quad 1 \leq i < \ell_{n+1} \\ \tau_n(a_{n+1}) &= a_n^{\ell_{n+1}}b_n(1). \end{aligned}$$

The fact that  $\tau$  is a constant length directive sequence that is primitive, injective on letters, recognizable and without coincidences is left to the reader. Since every word in  $\mathcal{A}_n^2$  occurs in the images of  $\tau_n$ , it can be checked that the maximal equicontinuous topological factor of  $(X_\tau, S)$  corresponds to the odometer  $(\mathbb{Z}_{(|\tau_{[0, n])}|)_{n \geq 0}}, +1)$ . Thus, since  $\tau$  does not have coincidences  $(X_\tau, S)$  is a minimal non Toeplitz subshift.

Stirling's approximation implies that

$$\log \ell_{n+1} > (\ell_n - 1) \log(\ell_n - 1) - (\ell_n - 1).$$

Since  $\ell_k - 1 \geq 2^k$  for  $k \geq 0$ , inductively we obtain

$$\begin{aligned} \frac{\log \ell_{n+1}}{(\ell_0 + 2) \dots (\ell_n + 2)} &> \prod_{k=0}^n \left(1 - \frac{3}{\ell_k + 2}\right) \log(\ell_0 - 1) - \sum_{k=0}^n \frac{(\ell_k - 1) \dots (\ell_n - 1)}{(\ell_0 + 2) \dots (\ell_n + 2)} \\ &> \prod_{k=0}^n \left(1 - \frac{3}{2^k}\right) \log(\ell_0 - 1) - 2. \end{aligned}$$

From [BH94, Lemma 2.6] the topological entropy of  $(X_\tau, S)$  is

$$\lim_{n \rightarrow +\infty} \frac{\log |\mathcal{A}_n|}{|\tau_{[0, n]}|} = \lim_{n \rightarrow +\infty} \frac{\log \ell_n}{(\ell_0 + 2) \dots (\ell_{n-1} + 2)} > 0$$

if we choose  $\ell_0$  such that  $\prod_{k=0}^{\infty} \left(1 - \frac{3}{2^k}\right) \log(\ell_0 - 1) > 2$ . In particular  $(X_\tau, S)$  is not of finite topological rank.

Finally if  $\mathcal{A}'_n = \{a_n\}$  for  $n \geq 0$  then there exists an ergodic measure  $\mu$  such that  $\tau$  is  $(\mathcal{A}'_n)_{n \geq 0}$ -adapted to  $\mu$ . Theorem 2 implies that  $(X_\tau, S, \mu)$  is isomorphic to  $(\mathbb{Z}_{(|\tau_{[0, n]}|)_{n \geq 0}}, +1, \nu)$ .

#### REFERENCES

- [ABKK17] M. Adamska, S. Bezuglyi, O. Karpel, and J. Kwiatkowski. Subdiagrams and invariant measures on Bratteli diagrams. *Ergodic Theory Dynam. Systems*, 37:2417–2452, 2017.
- [AD23] Felipe Arbulú and Fabien Durand. Dynamical properties of minimal ferenczi subshifts. *Ergodic Theory Dynam. Systems*, pages 1–48, 2023.
- [BBK06] Veronica Baker, Marcy Barge, and Jaroslaw Kwapisz. Geometric realization and coincidence for reducible non-unimodular Pisot tiling spaces with an application to  $\beta$ -shifts. *Ann. Inst. Fourier (Grenoble)*, 56:2213–2248, 2006. Numération, pavages, substitutions.
- [BCBD<sup>+</sup>21] V. Berthé, P. Cecchi Bernales, F. Durand, J. Leroy, D. Perrin, and S. Petite. On the dimension group of unimodular  $\mathcal{S}$ -adic subshifts. *Monatshefte für Mathematik*, 2021.
- [BD02] Marcy Barge and Beverly Diamond. Coincidence for substitutions of Pisot type. *Bull. Soc. Math. France*, 130:619–626, 2002.
- [BD14] Valérie Berthé and Vincent Delecroix. Beyond substitutive dynamical systems:  $\mathcal{S}$ -adic expansions. In *Numeration and substitution 2012*, RIMS Kōkyūroku Bessatsu, B46, pages 81–123. Res. Inst. Math. Sci. (RIMS), Kyoto, 2014.
- [BDM10] Xavier Bressaud, Fabien Durand, and Alejandro Maass. On the eigenvalues of finite rank Bratteli-Vershik dynamical systems. *Ergodic Theory Dynam. Systems*, 30:639–664, 2010.
- [BH94] Mike Boyle and David Handelman. Entropy versus orbit equivalence for minimal homeomorphisms. *Pacific J. Math.*, 164:1–13, 1994.
- [BK06] Marcy Barge and Jaroslaw Kwapisz. Geometric theory of unimodular Pisot substitutions. *Amer. J. Math.*, 128:1219–1282, 2006.
- [BKMS13] S. Bezuglyi, J. Kwiatkowski, K. Medynets, and B. Solomyak. Finite rank Bratteli diagrams: structure of invariant measures. *Trans. Amer. Math. Soc.*, 365:2637–2679, 2013.
- [BPR23] Marie-Pierre Béal, Dominique Perrin, and Antonio Restivo. Recognizability of morphisms. *Ergodic Theory Dynam. Systems*, pages 1–25, 2023.
- [BPRS23] Marie-Pierre Béal, Dominique Perrin, Antonio Restivo, and Wolfgang Steiner. Recognizability in  $\mathcal{S}$ -adic shifts, 2023.
- [BSTY19] Valérie Berthé, Wolfgang Steiner, Jörg M. Thuswaldner, and Reem Yassawi. Recognizability for sequences of morphisms. *Ergodic Theory Dynam. Systems*, 39:2896–2931, 2019.
- [CS01] Vincent Canterini and Anne Siegel. Geometric representation of substitutions of Pisot type. *Trans. Amer. Math. Soc.*, 353:5121–5144, 2001.

- [DDMP21] Sebastián Donoso, Fabien Durand, Alejandro Maass, and Samuel Petite. Interplay between finite topological rank minimal Cantor systems,  $\mathcal{S}$ -adic subshifts and their complexity. *Trans. Amer. Math. Soc.*, 374:3453–3489, 2021.
- [Dek78] F. M. Dekking. The spectrum of dynamical systems arising from substitutions of constant length. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 41:221–239, 1978.
- [DFM19] Fabien Durand, Alexander Frank, and Alejandro Maass. Eigenvalues of minimal Cantor systems. *J. Eur. Math. Soc. (JEMS)*, 21:727–775, 2019.
- [DG16] Tomasz Downarowicz and Eli Glasner. Isomorphic extensions and applications. *Topol. Methods Nonlinear Anal.*, 48:321–338, 2016.
- [DK15] Tomasz Downarowicz and Stanisław Kasjan. Odometers and Toeplitz systems revisited in the context of Sarnak’s conjecture. *Studia Math.*, 229:45–72, 2015.
- [DL12] Fabien Durand and Julien Leroy.  $\mathcal{S}$ -adic conjecture and Bratteli diagrams. *C. R. Math. Acad. Sci. Paris*, 350:979–983, 2012.
- [Dow05] Tomasz Downarowicz. Survey of odometers and Toeplitz flows. In *Algebraic and topological dynamics*, volume 385 of *Contemp. Math.*, pages 7–37. Amer. Math. Soc., Providence, RI, 2005.
- [DP22] Fabien Durand and Dominique Perrin. *Dimension groups and dynamical systems — Substitutions, Bratteli diagrams and Cantor systems*, volume 196 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2022.
- [EG60] Robert Ellis and W. H. Gottschalk. Homomorphisms of transformation groups. *Trans. Amer. Math. Soc.*, 94:258–271, 1960.
- [EM22] Bastián Espinoza and Alejandro Maass. On the automorphism group of minimal  $\mathcal{S}$ -adic subshifts of finite alphabet rank. *Ergodic Theory Dynam. Systems*, 42(9):2800–2822, 2022.
- [Esp22] Bastián Espinoza. Symbolic factors of  $\mathcal{S}$ -adic subshifts of finite alphabet rank. *Ergodic Theory Dynam. Systems*, pages 1–37, 2022.
- [Esp23] Bastián Espinoza. The structure of low complexity subshifts, 2023.
- [Fer96] Sébastien Ferenczi. Rank and symbolic complexity. *Ergodic Theory Dynam. Systems*, 16:663–682, 1996.
- [Fer97] Sébastien Ferenczi. Systems of finite rank. *Colloq. Math.*, 73:35–65, 1997.
- [FFT09] Sébastien Ferenczi, Albert M. Fisher, and Marina Talet. Minimality and unique ergodicity for adic transformations. *J. Anal. Math.*, 109:1–31, 2009.
- [Fog02] N. Pytheas Fogg. *Substitutions in dynamics, arithmetics and combinatorics*, volume 1794 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2002. Edited by V. Berthé, S. Ferenczi, C. Mauduit and A. Siegel.
- [Fur60] Harry Furstenberg. *Stationary processes and prediction theory*. Annals of Mathematics Studies, No. 44. Princeton University Press, Princeton, N.J., 1960.
- [GJ00] Richard Gjerde and O. Johansen. Bratteli-Vershik models for Cantor minimal systems: applications to Toeplitz flows. *Ergodic Theory Dynam. Systems*, 20:1687–1710, 2000.
- [GLL22] France Gheeraert, Marie Lejeune, and Julien Leroy.  $\mathcal{S}$ -adic characterization of minimal ternary dendric shifts. *Ergodic Theory Dynam. Systems*, 42:3393–3432, 2022.
- [GR17] Felipe García-Ramos. Weak forms of topological and measure-theoretical equicontinuity: relationships with discrete spectrum and sequence entropy. *Ergodic Theory Dynam. Systems*, 37:1211–1237, 2017.
- [GRJY21] Felipe García-Ramos, Tobias Jäger, and Xiangdong Ye. Mean equicontinuity, almost automorphy and regularity. *Israel J. Math.*, 243:155–183, 2021.
- [HLSY21] Wen Huang, Zhengxing Lian, Song Shao, and Xiangdong Ye. Minimal systems with finitely many ergodic measures. *Journal of Functional Analysis*, 280:109000, 2021.
- [Hos86] B. Host. Valeurs propres des systèmes dynamiques définis par des substitutions de longueur variable. *Ergodic Theory Dynam. Systems*, 6:529–540, 1986.
- [HP68] Frank Hahn and William Parry. Some characteristic properties of dynamical systems with quasi-discrete spectra. *Math. Systems Theory*, 2:179–190, 1968.
- [HS03] Michael Hollander and Boris Solomyak. Two-symbol Pisot substitutions have pure discrete spectrum. *Ergodic Theory Dynam. Systems*, 23:533–540, 2003.
- [JK69] Konrad Jacobs and Michael Keane. 0–1-sequences of Toeplitz type. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 13:123–131, 1969.



- [Liv87] A. N. Livshits. On the spectra of adic transformations of Markov compact sets. *Uspekhi Mat. Nauk*, 42:189–190, 1987.
- [LV92] A. N. Livshits and A. M. Vershik. Adic models of ergodic transformations, spectral theory, substitutions, and related topics. In *Representation theory and dynamical systems*, volume 9 of *Adv. Soviet Math.*, pages 185–204. Amer. Math. Soc., Providence, RI, 1992.
- [Mar75] Nelson G. Markley. Substitution-like minimal sets. *Israel J. Math.*, 22:332–353, 1975.
- [Mos92] Brigitte Mossé. Puissances de mots et reconnaissabilité des points fixes d’une substitution. *Theoret. Comput. Sci.*, 99:327–334, 1992.
- [Mos96] Brigitte Mossé. Reconnaissabilité des substitutions et complexité des suites automatiques. *Bull. Soc. Math. France*, 124:329–346, 1996.
- [Que87] Martine Queffélec. *Substitution dynamical systems — Spectral Analysis*, volume 1294 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1987.
- [Sen81] E. Seneta. *Nonnegative matrices and Markov chains*. Springer Series in Statistics. Springer-Verlag, New York, second edition, 1981.
- [Wil84] Susan Williams. Toeplitz minimal flows which are not uniquely ergodic. *Z. Wahrsch. Verw. Gebiete*, 67:95–107, 1984.

LABORATOIRE AMIÉNOIS DE MATHÉMATIQUE FONDAMENTALE ET APLIQUÉE, CNRS-UMR 7352, UNIVERSITÉ DE PICARDIE JULES VERNE, 33 RUE SAINT LEU, 80039 AMIENS CEDEX 1, FRANCE.

*Email address:* felipe.arbulu@u-picardie.fr

LABORATOIRE AMIÉNOIS DE MATHÉMATIQUE FONDAMENTALE ET APLIQUÉE, CNRS-UMR 7352, UNIVERSITÉ DE PICARDIE JULES VERNE, 33 RUE SAINT LEU, 80039 AMIENS CEDEX 1, FRANCE.

*Email address:* fabien.durand@u-picardie.fr

LABORATOIRE AMIÉNOIS DE MATHÉMATIQUE FONDAMENTALE ET APLIQUÉE, CNRS-UMR 7352, UNIVERSITÉ DE PICARDIE JULES VERNE, 33 RUE SAINT LEU, 80039 AMIENS CEDEX 1, FRANCE.

*Email address:* bespinoza@dim.uchile.cl