

Is AD Leo entering a polarity reversal? Long-term monitoring of the large-scale magnetic field with ESPaDOnS, NARVAL and SPIRou

Stefano Bellotti, Julien Morin, Lisa Lehmann, Pascal Petit, Gaitee Hussain

► To cite this version:

Stefano Bellotti, Julien Morin, Lisa Lehmann, Pascal Petit, Gaitee Hussain. Is AD Leo entering a polarity reversal? Long-term monitoring of the large-scale magnetic field with ESPaDOnS, NARVAL and SPIRou. Brun, Allan Sacha; Bouvier, Jérôme; Petit, Pascal. The 21st Cambridge Workshop on Cool Stars, Stellar Systems, and the Sun (CS21), Jul 2022, Toulouse, France. , 2022, 10.5281/zen-odo.7323944. hal-04169395

HAL Id: hal-04169395 https://hal.science/hal-04169395

Submitted on 24 Jul 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Is AD Leo entering a polarity reversal? Long-term monitoring of the large-scale magnetic field with **ESPaDOnS, NARVAL and SPIRou**

Stefano Bellotti^{1,3}, Julien Morin², Lisa Lehmann¹, Pascal Petit¹, Gaitee Hussain³ and the SLS consortium 1: IRAP, Toulouse (FR), 2: LUPM, Montpellier (FR), 3: ESA/ESTEC, Noordwijk (NL)









European Space Agency



See electronic version

stefano.bellotti@irap.omp.eu

Stellar magnetic cycles

Target star: AD Leo

With a long-term monitoring of the large-scale magnetic field, we can search for temporal evolution of the topology and uncover stellar dynamo theories. The Sun is an important benchmark: the magnetic cycle is characterised by a 11-yr variation in sunspot number, size and latitude of emergence, and magnetic polarity reversal [1,2]. Spectropolarimetric observations revealed analogous polarity flips also on other stars [3,4].

At the lower end of the main sequence, M dwarfs represent excellent laboratories to study non-solar dynamo mechanisms underlying the generation and preservation of magnetic fields.

The magnetic field generation occurs under different interior physical conditions, as the transition between partly - and fully-convective occurs at spectral type M3. Observational constraints for M dwarfs have been set by high-resolution spectroscopy field measurements in unpolarised light [5] and by tomographic inversion techniques (Zeeman-Doppler Imaging, [6]), revealing a dichotomy in the full-convective regime: strong, dipolar axisymmetric and weak, complex non-axisymmetric fields [7,8]. To explain this phenomenon, either dynamo bistability [9] or magnetic cycles [10] are invoked, but no definite conclusion has been reached so far.

```
Spectral type = M3.5
P_{rot} = 2.23 \pm 0.01 d
v_{eq}sin(i) = 3 \text{ km/s}
Inclination = 20^{\circ}
Mass = 0.42 M_{\odot}
Main topology: dipolar and
axisymmetric [7,11]
Dipole strength: ~1 kG
```

Longitudinal magnetic field

- Disk-integrated, line-of-sight component of the magnetic field (B₁)
- Stellar north pole is dominated by negative dipole all the time
- Global decrease in field strength over 14 years
- Reduced axisymmetry indicated by varying amplitude in the phase-folded time series



FWHM of Stokes /

- FWHM useful activity tracer sensitive to Zeeman effect \rightarrow high Landé factor (g_{eff}) lines can reveal activity signal
- Full vs high-g_{eff} mask [12]: all FWHM values are larger in the latter case
- Constant vs sine fit: the variations of the 2020b epoch are better described by sinusoid fit (22% decrease in residual RMS)



Fig 1. Evolution of B₁. *Left*: optical (circles) and near-infrared (squares) time series between 2006 and 2020. *Right*: near-infrared time series phase-folded at the stellar rotation period.

Principal Component Analysis

• Apply PCA to mean-subtracted near-infrared Stokes V (2019-2020) to infer details on non-axisymmetric field [13, see also poster of L. T. Lehmann]



- First three eigenvectors have signals \rightarrow on average field is non-axisymmetric
- Variations in the coefficients are more sinusoidal and amplified \rightarrow large-scale field is dipolar and its axisymmetry decreases over time

Fig 3. Evolution of PCA coefficients for the first three eigenvectors over the near-infrared epochs: 2019a (top), 2019b2020a (middle) and 2020b (bottom).

- Increased FWHM variations imply lower degree of axisymmetry
 - FWHM phase variations are reflected by ZDI maps



Fig 2. Phase-folded FWHM curves using full and high-g_{eff} masks for near-infrared epochs: 2019a (top), 2019b2020a (middle), 2020b (bottom)

Zeeman-Doppler Imaging

• Reconstruction of the large-scale magnetic field topology by means of a maximum-entropy algorithm



[size]

 G^2)

- poloidal The geometry is (magnetic fraction energy >90%) and dipolar (>70%), as in 2006-2016 observations
- Axisymmetry decreases from >85% to 48% in 2020b and tilt angle increases from 15° to 56° accordingly
- Fig 4. Evolution of ZDI map over near-infrared epochs: 2019a (left), 2019b2020a (middle) and 2020b (right). The magnetic maps are in flat polar



view, with the north pole at the center and the equator marked by a solid line

Conclusions

- Decreasing axisymmetry of the large-scale field confirmed by different techniques
- Global evolution of the large-scale field of AD Leo possibly suggesting a magnetic cycle, in the strong aligned dipole category
- Difference of longitudinal and ZDI field strength between near-infrared and optical domains, to be further investigated



Fig 5. Evolution of the magnetic field topology over 14 years. While the field remains poloidal and dipolar, its secular variation is manifested by a reduced axisymmetry and changing field strength

Bibliography

[1] Hathaway et al., 2010, Living Reviews in Solar Physics, 7, 1 [2] Hale, 1919, ApJ, 49, 153 [3] Petit et al., 2009, A&A, 508, L9 [4] Jeffers et al., 2022, arXiv e-prints, arXiv:2201.07530 [5] Shulyak et al., 2019, A&A, 626, A86 [6] Donati et al., 1997, MNRAS, 291, 658 [7] Morin et al., 2008, MNRAS, 390, 567 [8] Morin et al., 2010, MNRAS, 407, 2269 [9] Gastine et al., 2013, A&A, 549, L5 [10] Kitchiatnikov et al., 2014, MNRAS, 442, L1 [11] Lavail et al., 2018, MNRAS, 479, 4836 [12] Klein et al., 2021, MNRAS, 502, 188 [13] Lehman & Donati, 2022, arXiv e-prints, arXiv:2205.15647

Acknowledgements: Based on observations obtained at the Canada-France-Hawaii Telescope (CFHT) which is operated by the National Research Council (NRC) of Canada, the Institut National des Sciences de l'Univers of the Centre National de la Recherche Scientifique (CNRS) of France, and the University of Hawaii. The observations at the CFHT were performed with care and respect from the summit of Maunakea which is a significant cultural and historic site. We gratefully acknowledge the CFHT QSO observers who made this project possible. We acknowledge funding from ANR of France (SPIaSH), with PI Xavier Delfosse.