



HAL
open science

A Linear Algorithm for Radio k -Coloring Powers of Paths Having Small Diameter

Dipayan Chakraborty, Soumen Nandi, Sagnik Sen, D. Supraja

► **To cite this version:**

Dipayan Chakraborty, Soumen Nandi, Sagnik Sen, D. Supraja. A Linear Algorithm for Radio k -Coloring Powers of Paths Having Small Diameter. 34th International Workshop on Combinatorial Algorithms, Jun 2023, Tainan, Taiwan. pp.148-159, 10.1007/978-3-031-34347-6_13 . hal-04169997

HAL Id: hal-04169997

<https://hal.science/hal-04169997v1>

Submitted on 25 Jul 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

A linear algorithm for radio k -coloring powers of paths having small diameter

Dipayan Chakraborty^{1,*}, Soumen Nandi², Sagnik Sen³, and D K Supraja^{2,3,**}

¹ Université Clermont-Auvergne, CNRS, Mines de Saint-Étienne,
Clermont-Auvergne-INP, LIMOS, 63000 Clermont-Ferrand, France
dipayan.chakraborty@uca.fr

² Netaji Subhas Open University, India
soumen2004@gmail.com

³ Indian Institute of Technology Dharwad, India
sen007isi@gmail.com, dksupraja95@gmail.com

Abstract. The radio k -chromatic number $rc_k(G)$ of a graph G is the minimum integer λ such that there exists a function $\phi : V(G) \rightarrow \{0, 1, \dots, \lambda\}$ satisfying $|\phi(u) - \phi(v)| \geq k + 1 - d(u, v)$, where $d(u, v)$ denotes the distance between u and v . To date, several upper and lower bounds of $rc_k(\cdot)$ is established for different graph families. One of the most notable works in this domain is due to Liu and Zhu [SIAM Journal on Discrete Mathematics 2005] whose main results were computing the exact values of $rc_k(\cdot)$ for paths and cycles for the specific case when k is equal to the diameter.

In this article, we find the exact values of $rc_k(G)$ for powers of paths where the diameter of the graph is strictly less than k . Our proof readily provides a linear time algorithm for providing such labeling. Furthermore, our proof technique is a potential tool for solving the same problem for other graph classes with “small” diameter.

Keywords: radio coloring · radio k -chromatic number · Channel Assignment Problem · power of paths.

1 Introduction and main results

The theory of radio coloring and its variations are popular and well-known mathematical models of the Channel Assignment Problem (CAP) in wireless networks [1,2]. The connection between the real-life problem and the theoretical model has been explored in different bodies of works. In this article, we focus on the theoretical aspects of a particular variant, namely, the radio k -coloring. All the graphs considered in this article are undirected simple graphs and we refer to the book “Introduction to graph theory” by West [14] for all standard notations and terminologies used.

* Corresponding auhtor

** Corresponding auhtor

A λ -radio k -coloring of a graph G is a function $\phi : V(G) \rightarrow \{0, 1, \dots, \lambda\}$ satisfying $|\phi(u) - \phi(v)| \geq k + 1 - d(u, v)$. For every $u \in V(G)$, the value $\phi(u)$ is generally referred to as the *color* of u under ϕ . Usually, we pick λ in such a way that it has a preimage under ϕ , and then, we call λ to be the span of ϕ , denoting it by $\text{span}(\phi)$. The *radio k -chromatic number*⁴ $rc_k(G)$ is the minimum $\text{span}(\phi)$, where ϕ varies over all radio k -colorings of G .

In particular, the radio 2-chromatic number is the most well-studied restriction of the parameter (apart from the radio 1-chromatic number, which is equivalent to studying the chromatic number of graphs). There is a famous conjecture by Griggs and Yeh [6] that claims $rc_2(G) \leq \Delta^2$ where Δ is the maximum degree of G . The conjecture has been resolved for all $\Delta \geq 10^{69}$ by Havet, Reed and Sereni [7].

As one may expect, finding the exact values of $rc_k(G)$ for a general graph is an NP-complete problem [6]. Therefore, finding the exact value of $rc_k(G)$ for a given graph (usually belonging to a particular graph family) offers a huge number of interesting problems. Unfortunately, due to a lack of general techniques for solving these problems, not many exact values are known till date. One of the best contributions in this front remains the work of Liu and Zhu [12] who computed the exact value of $rc_k(G)$ where G is a path or a cycle and $k = \text{diam}(G)$.

As our work focuses on finding radio k -chromatic number of powers of paths, let us briefly recall the relevant related works. For a detailed overview of the topic, we encourage the reader to consult Chapter 7.5 of the dynamic survey on this topic maintained in the Electronic Journal of Combinatorics by Gallian [5] and the survey by Panigrahi [13]. For small paths P_n , that is, with $\text{diam}(P_n) < k$, Kchikech et al. [8] had established an exact formula for $rc_k(P_n)$; whereas, recall that, for paths of diameter equal to $k \geq 2$, Liu and Zhu [12] gave an exact formula for the radio number $rc_k(P_k)$. Moreover, a number of studies on the parameter $rc_k(P_n)$ depending on how k is related to $\text{diam}(P_n)$, or n alternatively, have been done by various authors [8,9,10,3]. So far as works on powers of paths are concerned, the only notable work we know is an exact formula for the radio number $rn(P_n^2)$ of the square of a path P_n by Liu and Xie [11]. Hence the natural question to ask is whether the results for the paths can be extended to paths of a general power m , where $1 \leq m \leq n$.

Progressing along the same line, in this article we concentrate on powers of paths having “small diameters”, that is, $\text{diam}(P_n^m) < k$ and compute the exact value of $rc_k(P_n^m)$, where P_n^m denotes the m -th power graph of a path P_n on $(n + 1)$ vertices. In other words, the graph P_n^m is obtained by adding edges between the vertices of P_n that are at most m distance apart, where $m \leq n$. Notice that, the so-obtained graph is, in particular, an interval graph. Let us now state our main theorem.

Theorem 1. *For all $k > \text{diam}(P_n^m)$ and $m \leq n$, we have*

⁴ In the case that $\text{diam}(G) = k, k + 1$ or $k + 2$, the radio k -chromatic number is alternatively known as the *radio number* denoted by $rn(G)$, the *radio antipodal number* denoted by $ac(G)$ and the *nearly antipodal number* denoted by $ac'(G)$, respectively.

$$rc_k(P_n^m) = \begin{cases} nk - \frac{n^2 - m^2}{2m} & \text{if } \lceil \frac{n}{m} \rceil \text{ is odd and } m|n, \\ nk - \frac{n^2 - s^2}{2m} + 1 & \text{if } \lceil \frac{n}{m} \rceil \text{ is odd and } m \nmid n, \\ nk - \frac{n^2}{2m} + 1 & \text{if } \lceil \frac{n}{m} \rceil \text{ is even and } m|n, \\ nk - \frac{n^2 - (m-s)^2}{2m} + 1 & \text{if } \lceil \frac{n}{m} \rceil \text{ is even and } m \nmid n, \end{cases}$$

where $s \equiv n \pmod{m}$ and $1 \leq s < m$.

In this article, we develop a robust graph theoretic tool for the proof. Even though the tool is specifically used to prove our result, it can be adapted to prove bounds for other classes of graphs. Thus, we would like to remark that, the main contribution of this work is not only in proving an important result that captures a significant number of problems with a unified proof, but also in devising a proof technique that has the potential of becoming a standard technique to attack similar problems. We will prove the theorem in the next section.

Moreover, our proof of the upper bound is by giving a prescribed radio k -coloring of the concerned graph, and then proving its validity, while the lower bound proof establishes its optimality. Therefore, as a corollary to Theorem 1, we can say that our proof provides a linear time algorithm radio k -color powers of paths, optimally.

Theorem 2. *For all $k > \text{diam}(P_n^m)$ and $m \leq n$, one can provide an optimal radio k -coloring of the graph P_n^m in $O(n)$ time.*

We prove Theorem 1 in the next section.

2 Proofs of Theorems 1 and 2

This section is entirely dedicated to the proofs of Theorems 1 and 2. The proofs use specific notations and terminologies developed for making it easier for the reader to follow. The proof is contained in several observations and lemmas and uses a modified and improved version of the DGNS formula [4].

As seen from the theorem statement, the graph P_n^m that we work on is the m^{th} power of the path on $(n+1)$ vertices. One crucial aspect of this proof is the naming of the vertices of P_n^m . In fact, for convenience, we shall assign *two* names to each of the vertices of the graph and use them as required depending on the context. Such a naming convention will depend on the parity of the diameter of P_n^m .

Observation 1. *The diameter of the graph P_n^m is $\text{diam}(P_n^m) = \lceil \frac{n}{m} \rceil$.*

For the rest of this section, let $q = \lfloor \frac{\text{diam}(P_n^m)}{2} \rfloor$.

2.1 The naming conventions

We are now ready to present the first naming convention for the vertices of P_n^m . For convenience, let us suppose that the vertices of P_n^m are placed (embedded) on the X -axis having co-ordinates $(i, 0)$ where $i \in \{0, 1, \dots, n\}$ and two (distinct) vertices are adjacent if and only if their Euclidean distance is at most m .

We start by selecting the layer L_0 consisting of the vertex, named c_0 , say, positioned at $(qm, 0)$ for even values of $\text{diam}(P_n^m)$. On the other hand, for odd values of $\text{diam}(P_n^m)$, the layer L_0 consists of the vertices c_0, c_1, \dots, c_m , say, positioned at $(qm, 0), (qm + 1, 0), \dots, (qm + m, 0)$, respectively, and inducing a maximal clique of size $(m + 1)$. The vertices of L_0 are called the *central vertices*, and those positioned to the left and the right side of the central vertices are naturally called the *left vertices* and the *right vertices*, respectively.

After this, we define the layer L_i as the set of vertices that are at a distance i from L_0 . Observe that the layer L_i is non-empty for all $i \in \{0, 1, \dots, q\}$. Moreover, notice that, for all $i \in \{1, 2, \dots, q\}$, L_i consists of both left and right vertices. In particular, for $i \geq 1$, the left vertices of L_i are named $l_{i1}, l_{i2}, \dots, l_{im}$, sorted according to the increasing order of their Euclidean distances from L_0 . Similarly, for $i \in \{1, 2, \dots, q - 1\}$, the right vertices of L_i are named $r_{i1}, r_{i2}, \dots, r_{im}$, sorted according to the increasing order of their Euclidean distance from L_0 . However, the right vertices of L_q are $r_{q1}, r_{q2}, \dots, r_{qs}$, where $s = (n + 1) - (2q - 1)m - |L_0|$ (observe that this s is the same as the s mentioned in the statement of Theorem 1), again sorted according to the increasing order of their Euclidean distances from L_0 . That is, if $m \nmid n$, then there are $s = (n + 1) - (2q - 1)m - |L_0|$ right vertices in L_q . Besides L_q , every layer L_i , for $i \in \{1, 2, \dots, q - 1\}$, has exactly m left vertices and m right vertices. This completes our first naming convention.

Now, we move to the second naming convention. This depends on yet another observation.

Observation 2. For $k \geq \text{diam}(P_n^m)$, let ϕ be a radio k -coloring of P_n^m . Then $\phi(x) \neq \phi(y)$ for all distinct $x, y \in V(P_n^m)$.

Let ϕ be a radio k -coloring of P_n^m . Thus, due to Observation 2, it is possible to sort the vertices of P_n^m according to the increasing order of their colors. That is, our second naming convention which names the vertices of P_n^m as v_0, v_1, \dots, v_n satisfying $\phi(v_0) < \phi(v_1) < \dots < \phi(v_n)$. Clearly, the second naming convention depends only on the coloring ϕ , which, for the rest of this section, will play the role of any arbitrary radio k -coloring of P_n^m .

2.2 The lower bound

Next, we shall proceed to establish the lower bound of Theorem 1 by showing it to be a lower bound of $\text{span}(\phi)$. To do so, however, we need to introduce yet another notation. Let $f : V(P_n^m) \rightarrow \{0, 1, \dots, q\}$ be the function which indicates the layer of a vertex, that is, $f(x) = i$ if $x \in L_i$. With this notation, we initiate the lower bound proof with the following result.

Lemma 1. For any $i \in \{0, 1, \dots, n-1\}$, we have

$$\phi(v_{i+1}) - \phi(v_i) \geq \begin{cases} k - f(v_i) - f(v_{i+1}) + 1 & \text{if } \text{diam}(P_n^m) \text{ is even,} \\ k - f(v_i) - f(v_{i+1}) & \text{if } \text{diam}(P_n^m) \text{ is odd.} \end{cases}$$

Proof. If $\text{diam}(P_n^m)$ is even, then L_0 consists of the single vertex c_0 . Observe that, as v_i is in $L_{f(v_i)}$, it is at a distance $f(v_i)$ from c_0 . Similarly, v_{i+1} is at a distance $f(v_{i+1})$ from c_0 . Hence, by the triangle inequality, we have

$$d(v_i, v_{i+1}) \leq d(v_i, c_0) + d(c_0, v_{i+1}) = f(v_i) + f(v_{i+1}).$$

Therefore, by the definition of radio k -coloring,

$$\phi(v_{i+1}) - \phi(v_i) \geq k - f(v_i) - f(v_{i+1}) + 1.$$

If $\text{diam}(P_n^m)$ is odd, then L_0 is a clique. Thus, by the definition of layers and the function f , there exist vertices c_j and $c_{j'}$ ($j \neq j'$) in L_0 satisfying $d(v_i, c_j) = f(v_i)$ and $d(v_{i+1}, c_{j'}) = f(v_{i+1})$. Hence, by triangle inequality again, we have

$$d(v_i, v_{i+1}) \leq d(v_i, c_j) + d(c_j, c_{j'}) + d(c_{j'}, v_{i+1}) = f(v_i) + 1 + f(v_{i+1}).$$

Therefore, by the definition of radio k -coloring,

$$\phi(v_{i+1}) - \phi(v_i) \geq k - f(v_i) - f(v_{i+1}).$$

Hence we are done. \square

Notice that it is not possible to improve the lower bound of the inequality presented in Lemma 1. Motivated by this fact, whenever we have

$$\phi(v_{i+1}) - \phi(v_i) = \begin{cases} k - f(v_i) - f(v_{i+1}) + 1 & \text{if } \text{diam}(P_n^m) \text{ is even,} \\ k - f(v_i) - f(v_{i+1}) & \text{if } \text{diam}(P_n^m) \text{ is odd.} \end{cases}$$

for some $i \in \{0, 1, \dots, n-1\}$, we say that the pair (v_i, v_{i+1}) is *optimally colored* by ϕ . Moreover, we can naturally extend this definition to a sequence of vertices of the type $(v_i, v_{i+1}, \dots, v_{i+i'})$ by calling it an *optimally colored sequence* by ϕ if (v_{i+j}, v_{i+j+1}) is optimally colored by ϕ for all $j \in \{0, 1, \dots, i'-1\}$. Furthermore, a *loosely colored sequence* $(v_i, v_{i+1}, v_{i+2}, \dots, v_{i+i'})$ is a sequence that does not contain any optimally colored sequence as a subsequence.

An important thing to notice is that the sequence of vertices (v_0, v_1, \dots, v_n) can be written as a concatenation of maximal optimally colored sequences and loosely colored sequences. That is, it is possible to write

$$(v_0, v_1, \dots, v_n) = Y_0 X_1 Y_1 X_2 \cdots X_t Y_t$$

where Y_i s are loosely colored sequences and X_j s are maximal optimally colored sequences. Here, we allow the Y_i s to be empty sequences as well. In fact, for

$1 \leq i \leq t-1$, a Y_i is empty if and only if there exist two consecutive vertices $v_{s'}$ and $v_{s'+1}$ of P_n^m in the second naming convention such that $(v_{s'}, v_{s'+1})$ is loosely colored and that $X_i = (v_s, v_{s+1}, \dots, v_{s'})$ and $X_{i+1} = (v_{s'+1}, v_{s'+2}, \dots, v_{s''})$ for some $s \leq s' < s''$. Moreover, Y_0 (resp. Y_t) is empty if and only if the pair (v_0, v_1) (resp. (v_{n-1}, v_n)) is optimally colored. By convention, empty sequences are always loosely colored and a sequence having a singleton vertex is always optimally colored. From now onward, whenever we mention a radio k -coloring ϕ of P_n^m , we shall also suppose an associated concatenated sequence using the same notation as mentioned above.

Let us now prove a result which plays an instrumental role in the proof of the lower bound.

Lemma 2. *Let ϕ be a radio- k coloring of P_n^m such that*

$$(v_0, v_1, \dots, v_n) = Y_0 X_1 Y_1 X_2 \dots X_t Y_t.$$

Then, for even values of $\text{diam}(P_n^m)$, we have

$$\text{span}(\phi) \geq \left[n(k+1) - 2 \sum_{i=1}^q i |L_i| \right] + \left[f(v_0) + f(v_n) + \sum_{i=0}^t |Y_i| + t - 1 \right]$$

and, for odd values of $\text{diam}(P_n^m)$, we have

$$\text{span}(\phi) \geq \left[nk - 2 \sum_{i=1}^q i |L_i| \right] + \left[f(v_0) + f(v_n) + \sum_{i=0}^t |Y_i| + t - 1 \right],$$

where $|Y_i|$ denotes the length of the sequence Y_i .

As we shall calculate the two additive components of Lemma 2 separately, we introduce short-hand notations for them for the convenience of reference. So, let

$$\alpha_1 = \begin{cases} n(k+1) - 2 \sum_{i=1}^q i |L_i| & \text{if } \text{diam}(P_n^m) \text{ is even,} \\ nk - 2 \sum_{i=1}^q i |L_i| & \text{if } \text{diam}(P_n^m) \text{ is odd,} \end{cases}$$

and

$$\alpha_2(\phi) = f(v_0) + f(v_n) + \sum_{i=0}^t |Y_i| + t - 1.$$

Observe that α_1 and α_2 are functions of a number of variables and factors such as, n, m, k, ϕ , etc. However, to avoid clumsy and lengthy formulations, we have avoided writing α_1 and α_2 as multivariate functions, as their definitions are not ambiguous in the current context. Furthermore, as k and P_n^m are assumed to be fixed in the current context and, as α_1 does not depend on ϕ (follows from its definition), it is treated and expressed as a constant as a whole. On the other hand, α_2 is expressed as a function of ϕ .

Now we shall establish lower bounds for α_1 and $\alpha_2(\phi)$ separately to prove the lower bound of Theorem 1. Let us start with α_1 first.

Lemma 3. *We have*

$$\alpha_1 = \begin{cases} nk - \frac{n^2+m^2-s^2}{2m} & \text{if } \text{diam}(P_n^m) \text{ is even,} \\ nk - \frac{n^2-s^2}{2m} & \text{if } \text{diam}(P_n^m) \text{ is odd,} \end{cases}$$

where $s = (n+1) - (2q-1)m - |L_0|$.

Next, we focus on $\alpha_2(\phi)$. We shall handle the cases with odd $\text{diam}(P_n^m)$ first.

Lemma 4. *We have*

$$\alpha_2(\phi) \geq \begin{cases} 0 & \text{if } \text{diam}(P_n^m) \text{ is odd and } m|n, \\ 1 & \text{if } \text{diam}(P_n^m) \text{ is odd and } m \nmid n. \end{cases}$$

Next, we consider the cases with even $\text{diam}(P_n^m)$. Before starting with it though, we are going to introduce some terminologies to be used during the proofs. So, let X_i be an optimally colored sequence. As X_i cannot have two consecutive left (resp., right) vertices as elements, the number of left vertices can be at most one more than the number of right vertices and the central vertex, the latter two combined together.

Lemma 5. *We have*

$$\alpha_2(\phi) \geq \begin{cases} 1 & \text{if } \text{diam}(P_n^m) \text{ is even and } m|n, \\ m-s+1 & \text{if } \text{diam}(P_n^m) \text{ is even and } m \nmid n, \end{cases}$$

where $s \equiv n \pmod{m}$.

Combining Lemmas 2, 3, 4 and 5, therefore, we have the following lower bound for the parameter $rc_k(P_n^m)$.

Lemma 6. *For all $k \geq \text{diam}(P_n^m)$ and $m \leq n$, we have*

$$rc_k(P_n^m) \geq \begin{cases} nk - \frac{n^2-m^2}{2m} & \text{if } \lceil \frac{n}{m} \rceil \text{ is odd and } m|n, \\ nk - \frac{n^2-s^2}{2m} + 1 & \text{if } \lceil \frac{n}{m} \rceil \text{ is odd and } m \nmid n, \\ nk - \frac{n^2}{2m} + 1 & \text{if } \lceil \frac{n}{m} \rceil \text{ is even and } m|n, \\ nk - \frac{n^2-(m-s)^2}{2m} + 1 & \text{if } \lceil \frac{n}{m} \rceil \text{ is even and } m \nmid n, \end{cases}$$

where $s \equiv n \pmod{m}$ and $1 \leq s < m$.

Remark 1. Our lower bound technique can be applied to a graph G of diameter more than k also. This can be achieved by taking a subgraph H of G induced on $\bigcup_{i=0}^q L_i$, where $q = \lfloor \frac{k}{2} \rfloor$ and $\text{diam}(H) \leq k$. Thus, a lower bound for H serves as a lower bound for G as well.

2.3 The upper bound

Now let us prove the upper bound. We shall provide a radio k -coloring ψ of P_n^m and show that its span is the same as the value of $rc_k(P_n^m)$ stated in Theorem 1. To define ψ , we shall use both naming conventions. That is, we shall express the ordering (v_0, v_1, \dots, v_n) of the vertices of P_n^m with respect to ψ in terms of the first naming convention.

Let us define some ordering for the right (and similarly for the left) vertices:

- (1) $r_{ij} \prec_1 r_{i'j'}$ if either (i) $j < j'$ or (ii) $j = j'$ and $(-1)^{j-1}i < (-1)^{j'-1}i'$;
- (2) $r_{ij} \prec_2 r_{i'j'}$ if either (i) $j < j'$ or (ii) $j = j'$ and $(-1)^{m-j}i < (-1)^{m-j'}i'$;
- (3) $r_{ij} \prec_3 r_{i'j'}$ if either (i) $j < j'$ or (ii) $j = j'$ and $i > i'$; and
- (4) $r_{ij} \prec_4 r_{i'j'}$ if either (i) $j < j'$ or (ii) $j = j'$ and $(-1)^ji < (-1)^{j'}i'$.

Observe that, the orderings are based on comparing the second co-ordinate of the indices of the right (resp., left) vertices, and if they happen to be equal, then comparing the first co-ordinate of the indices with conditions on their parities. Moreover, all the above four orderings define total orders on the set of all right (resp., left) vertices. Thus, there is a unique increasing (resp., decreasing) sequence of right (or the left) vertices with respect to \prec_1 , \prec_2 , \prec_3 , and \prec_4 . Based on these orderings, we are going to construct a sequence of vertices of the graph and then greedily color the vertices to provide our labeling.

The sequences of the vertices are given as follows:

- (1) An *alternating chain* as a sequence of vertices of the form $(a_1, b_1, a_2, b_2, \dots, a_p, b_p)$ such that (a_1, a_2, \dots, a_p) is the increasing sequence of right vertices with respect to \prec_1 and (b_1, b_2, \dots, b_p) is the decreasing sequence of left vertices with respect to \prec_2 .
- (2) A *canonical chain* as a sequence of vertices of the form $(a_1, b_1, a_2, b_2, \dots, a_p, b_p)$ such that (a_1, a_2, \dots, a_p) is the increasing sequence of right vertices with respect to \prec_3 and (b_1, b_2, \dots, b_p) is the decreasing sequence of left vertices with respect to \prec_3 ;
- (3) A *special alternating chain* as a sequence of vertices of the form $(a_1, b_1, a_2, b_2, \dots, a_p, b_p)$ such that (a_1, a_2, \dots, a_p) is the increasing sequence of right vertices with respect to \prec_2 and (b_1, b_2, \dots, b_p) is the decreasing sequence of left vertices with respect to \prec_1 ; and
- (4) A *special canonical chain* as a sequence of vertices of the form $(a_1, b_1, a_2, b_2, \dots, a_p, b_p)$ such that (a_1, a_2, \dots, a_p) is the increasing sequence of right vertices with respect to \prec_4 and (b_1, b_2, \dots, b_p) is the decreasing sequence of left vertices with respect to \prec_4 .

Notice that the special alternating chains, the reverse alternating chain and the canonical chains can exist only when the number of right and left vertices are equal. Of course, when $m|n$, both the chains exist. Otherwise, we shall modify the names of the vertices a little to make them exist.

We are now ready to express the sequence (v_0, v_1, \dots, v_n) by splitting it into different cases which are depicted in Figures 1, 2, 3 and 4 for example. In the

figures, the both naming conventions for each of the vertices are given so that the reader may cross verify the correctness for that particular instance for each case. For convenience, also recall that $q = \lfloor \frac{\text{diam}(P_n^m)}{2} \rfloor$.

Case 1: when $\text{diam}(P_n^m)$ is even, $m|n$ and $k > \text{diam}(P_n^m)$. First of all, $(v_0, v_1, \dots, v_{2qm-1})$ is the alternating chain. Moreover, $v_n = c_0$.




Fig. 1: *Case 1.* $n = 16$, $m = 4$, $\text{diam}(P_{24}^4) = 4$, $k = 6$.

Case 2: when $\text{diam}(P_n^m)$ is odd, $m|n$ and $k > \text{diam}(P_n^m)$. Let the ordering of the vertices be $(v_0, v_1, \dots, v_{2qm+m})$. Now, $v_{j(2q+1)} = c_j$ for all $0 \leq j \leq m$. The remaining vertices follow the canonical chain.




Fig. 2: *Case 2.* $n = 20$, $m = 4$, $\text{diam}(P_{20}^4) = 5$, $k = 7$.

Case 3: when $\text{diam}(P_n^m)$ is odd, $m \nmid n$ and $k > \text{diam}(P_n^m)$. For any set A , let A^* represent an ordered sequence of the elements of A . Let $G \cong P_n^m$ and $S = V(G) = \{v_0, v_1, v_2, \dots, v_{2qm+s}\}$. Then S^* is defined as described. First, define

$$T = \{v_t : 0 \leq t \leq s(2q+1)\} - \{v_{j(2q+1)} : 0 \leq j \leq s\}.$$

Order T^* as a canonical chain. Also, define $v_{j(2q+1)} = c_j$ for all $0 \leq j \leq s$. Assume G' to be the subgraph of G induced by the subset $S - \{r_{q1}, r_{q2}, \dots, r_{qs}\}$ of S . Then $G' \cong P_{n'}^m$, $m|n'$ and $\text{diam}(G') = \frac{n'}{m}$ is even, where $n' = n - s$. Define

$$v_n = l_{11} \text{ and } U = \{v_t : s(2q+1) + 1 \leq t < n\}.$$

Note that $U \subset V(G')$. Order U^* (as vertices of G') by the following.

- (i) Special alternating chain when m and s have the same parity.
- (ii) Alternating chain when m is even and s is odd.
- (iii) Special canonical chain when m is odd and s is even.




Fig. 3: Case 3. $n = 19$, $m = 4$, $\text{diam}(P_{19}^4) = 5$, $k = 7$, $s = 3$.

Case 4: when $\text{diam}(P_n^m)$ is even, $m \nmid n$ and $k > \text{diam}(P_n^m)$. Notice that, in this case, the left vertices are $(m - s)$ more than the right vertices. Also, L_0 has only one vertex in this case. We shall discard some vertices from the set of left vertices, and then present the ordering. To be specific, we disregard the subset $\{l_{11}, l_{12}, \dots, l_{1(m-s)}\}$, temporarily, from the set of left vertices and consider the alternating chain. First of all, $(v_0, v_1, \dots, v_{2qm-2m+2s-1})$ is the alternating chain. Additionally, $(v_{2qm-2m+2s}, v_{2qm-2m+2s+1}, v_{2qm-2m+2s+2}, \dots, v_{2qm-m+s}) = (c_0, l_{11}, l_{12}, \dots, l_{1(m-s)})$.




Fig. 4: Case 4. $n = 14$, $m = 4$, $\text{diam}(P_{22}^4) = 4$, $k = 6$, $s = 2$.

Thus, we have obtained a sequence (v_0, v_1, \dots, v_n) in each case under consideration. Now, we define, $\psi(v_0) = 0$ and $\psi(v_{i+1}) = \psi(v_i) + k + 1 - d(v_i, v_{i+1})$, recursively, for all $i \in \{0, 1, 2, \dots, n - 1\}$. Next, we note that ψ is a radio k -coloring.

Lemma 7. *The function ψ is a radio k -coloring of P_n^m .*

This brings us to the upper bound for $rc_k(P_n^m)$.

Lemma 8. *For all $k > \text{diam}(P_n^m)$ and $m \leq n$, we have*

$$rc_k(P_n^m) \leq \begin{cases} nk - \frac{n^2 - m^2}{2m} & \text{if } \lceil \frac{n}{m} \rceil \text{ is odd and } m|n, \\ nk - \frac{n^2 - s^2}{2m} + 1 & \text{if } \lceil \frac{n}{m} \rceil \text{ is odd and } m \nmid n, \\ nk - \frac{n^2}{2m} + 1 & \text{if } \lceil \frac{n}{m} \rceil \text{ is even and } m|n, \\ nk - \frac{n^2 - (m-s)^2}{2m} + 1 & \text{if } \lceil \frac{n}{m} \rceil \text{ is even and } m \nmid n, \end{cases}$$

where $s \equiv n \pmod{m}$ and $1 \leq s < m$.

Proof. Observe that, $rc_k(P_n^m) \leq span(\psi)$. So, to prove the upper bound, it is enough to show that for all $k > diam(P_n^m)$ and $s \equiv n \pmod{m}$,

$$span(\psi) = \begin{cases} nk - \frac{n^2 - m^2}{2m} & \text{if } \lceil \frac{n}{m} \rceil \text{ is odd and } m|n, \\ nk - \frac{n^2 - s^2}{2m} + 1 & \text{if } \lceil \frac{n}{m} \rceil \text{ is odd and } m \nmid n, \\ nk - \frac{n^2}{2m} + 1 & \text{if } \lceil \frac{n}{m} \rceil \text{ is even and } m|n, \\ nk - \frac{n^2 - (m-s)^2}{2m} + 1 & \text{if } \lceil \frac{n}{m} \rceil \text{ is even and } m \nmid n. \end{cases}$$

As ψ is explicitly known, it is possible to calculate it and prove the above. However, we omit the rest of the proof due to space constraint. \square

2.4 The proofs

Finally we are ready to conclude the proofs.

Proof of Theorem 1 The proof follows directly from Lemmas 6 and 8. \square

Proof of Theorem 2 Notice that the proof of the upper bound for Theorem 1 is given by prescribing an algorithm (implicitly). The algorithm requires ordering the vertices of the input graph, and then providing the coloring based on the ordering. Each step runs in linear order of the number of vertices in the input graph. Moreover, we have theoretically proved the tightness of the upper bound. Thus, we are done. \square

For the full version of the paper, please go to https://homepages.iitdh.ac.in/~sen/Supraja_IWCA.pdf.

Acknowledgements: This work is partially supported by the following projects: “MA/IFCAM/18/39”, “SRG/2020/001575”, “MTR/2021/000858”, and “NBHM/RP-8 (2020)/Fresh”. Research by the first author is partially sponsored by a public grant overseen by the French National Research Agency as part of the “Investissements d’Avenir” through the IMobS3 Laboratory of Excellence (ANR-10-LABX-0016) and the IDEX-ISITE initiative CAP 20-25 (ANR-16-IDEX-0001).

References

1. G. Chartrand, D. Erwin, F. Harary, and P. Zhang. Radio labelings of graphs. *Bulletin of the Institute of Combinatorics and its Applications*, 33:77–85, 2001.
2. G. Chartrand, D. Erwin, and P. Zhang. A graph labeling problem suggested by FM channel restrictions. *Bulletin of the Institute of Combinatorics and its Applications*, 43:43–57, 2005.
3. G. Chartrand, L. Nebeskỳ, and P. Zhang. Radio k -colorings of paths. *Discussiones Mathematicae Graph Theory*, 24(1):5–21, 2004.
4. S. Das, S. C. Ghosh, S. Nandi, and S. Sen. A lower bound technique for radio k -coloring. *Discrete Mathematics*, 340(5):855–861, 2017.

5. J. A. Gallian. A dynamic survey of graph labeling. *Electronic Journal of combinatorics*, 1(DynamicSurveys):DS6, 2018.
6. J. R. Griggs and R. K. Yeh. Labelling graphs with a condition at distance 2. *SIAM Journal on Discrete Mathematics*, 5(4):586–595, 1992.
7. F. Havet, B. Reed, and J.-S. Sereni. L(2,1)-labelling of graphs. In *ACM-SIAM symposium on Discrete algorithms (SODA 2008)*, pages 621–630, 2008.
8. M. Kchikech, R. Khennoufa, and O. Togni. Linear and cyclic radio k -labelings of trees. *Discussiones Mathematicae Graph Theory*, 27(1):105–123, 2007.
9. R. Khennoufa and O. Togni. A note on radio antipodal colourings of paths. *Mathematica Bohemica*, 130(3):277–282, 2005.
10. S. R. Kola and P. Panigrahi. Nearly antipodal chromatic number $ac'(P_n)$ of the path P_n . *Mathematica Bohemica*, 134(1):77–86, 2009.
11. D. D. F. Liu and M. Xie. Radio number for square paths. *Ars Combinatoria*, 90:307–319, 2009.
12. D. D. F. Liu and X. Zhu. Multilevel distance labelings for paths and cycles. *SIAM Journal on Discrete Mathematics*, 19(3):610–621, 2005.
13. P. Panigrahi. A survey on radio k -colorings of graphs. *AKCE International Journal of Graphs and Combinatorics*, 6(1):161–169, 2009.
14. D. B. West. *Introduction to Graph Theory (2nd Edition)*. Prentice Hall, 2001.