



HAL
open science

Oriented embedding functors of tori as homogeneous spaces

Philippe Gille, Ting-Yu Lee

► **To cite this version:**

Philippe Gille, Ting-Yu Lee. Oriented embedding functors of tori as homogeneous spaces. 2023. hal-04171950v1

HAL Id: hal-04171950

<https://hal.science/hal-04171950v1>

Preprint submitted on 27 Jul 2023 (v1), last revised 10 Jul 2024 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ORIENTED EMBEDDING FUNCTORS OF TORI AS HOMOGENEOUS SPACES

PHILIPPE GILLE AND TING-YU LEE

ABSTRACT. We provide a characterization of homogeneous spaces under a reductive group scheme such that the geometric stabilizers are maximal tori. The quasi-split case over a semilocal base is of special interest and permits to answer a question raised by Marc Levine on homogeneous SL_n -spaces. At the end, we provide an application to the local-global principles for embeddings of étale algebras with involution into central simple algebras with involution.

Keywords: Reductive group schemes, tori, torsors, homogeneous spaces, embeddings, quasi-split groups, local-global principles, étale algebras, central simple algebras.

MSC 2000: 14L15, 20G35

1. INTRODUCTION

Let k be a field and let k_s be a separable closure of k ; we denote by $\Gamma_k = \mathrm{Gal}(k_s/k)$ the absolute Galois group of k . Let X be an affine k -variety which is a SL_n -homogeneous space such that the stabilizer of a geometric point is a maximal torus. Marc Levine asked whether X is G -isomorphic to SL_n/T where T is a maximal torus of SL_n .

Our main classification result (Theorem 3.4) tells us that X is the variety¹ of embeddings of a suitable oriented torus T in SL_n and we explain now the meaning of $X(k)$. Denoting by $\mathbb{G}_m^{n,1}$ the standard split torus of SL_n , we remind the reader that the automorphism group of the root system $A_{n-1} = \Phi(\mathrm{SL}_n, \mathbb{G}_m^{n,1})$ is $S_n \times \mathbb{Z}/2\mathbb{Z}$ if $n \geq 3$ and $\mathbb{Z}/2\mathbb{Z}$ for $n = 2$. We have that $T = R_{A/k}^1(\mathbb{G}_m)$ for an étale k -algebra A (see [14, §7.5]) and that $X(k)$ corresponds to the set of embeddings $f : T \rightarrow \mathrm{SL}_n$ such that the class of the $S_n \times \mathbb{Z}/2\mathbb{Z}$ -torsor $\mathrm{Isom}\left(\Phi(\mathrm{SL}_{n,k_s}, \mathbb{G}_{m,k_s}^{n,1}), \Phi(\mathrm{SL}_{n,k_s}, f(T)_{k_s})\right)$ is $[A] \times 1$ if $n \geq 3$ and simply $[A]$ if $n = 2$. We used implicitly there that the Galois cohomology set $H^1(k, S_n)$ classifies isomorphism classes of étale algebras of degree n and that the oriented type of T is the class of the above torsor.

Date: July 27, 2023.

¹Those varieties (and their generalizations over a base) have been defined by the second author [15].

On the other hand, if we see A as a k -vector space of dimension n , we have an isomorphism $\mathrm{SL}(A) \xrightarrow{\sim} \mathrm{SL}_n$ and the natural embedding $T = R_{A/k}^1(\mathbb{G}_m) \hookrightarrow \mathrm{SL}_1(A)$ defines then an embedding $T \rightarrow \mathrm{SL}_n$ of oriented type $[A] \times 1$. In other words, the SL_n -homogeneous space X carries a k -point whose stabilizer is $T = R_{A/k}^1(\mathbb{G}_m)$. Thus X is SL_n -isomorphic to SL_n/T as desired.

This statement can be strengthened in two ways. First we can replace SL_n by any quasi-split reductive k -group, the presence of k -points on a such homogeneous space being the Gille-Raghunathan's theorem [12, 19].

With more effort, we can replace further the base field k by an arbitrary semilocal ring (with infinite residue fields). This is Corollary 4.7 which involves a generalization of the existence of maximal tori of any orientation in such group schemes (see Theorem 4.6). As in Steinberg's section theorem, the case of type A_{2n} requires an additional argument. Also this generalization needs the notion of versal torsor in that setting (Reichstein-Tossici [21]); we introduced a variation of this technique involving algebraic spaces (see §4.2).

Finally we provide an arithmetic application to the embeddings of étale algebras with involution into central simple algebras with involution. Let (E, σ) be an étale algebra E over k with involution σ and (A, τ) be a central simple algebra A over k with involution τ . We want to know when (E, σ) can be embedded into (A, τ) under some constraints on the dimensions of (E, σ) and (A, τ) (see [18], [2], [3]). In Theorem 4.4.1, we show that when k is a number field and the unitary group associated to (A, τ) is quasi-split, the local-global principle always holds for the embeddings of (E, σ) into (A, τ) .

Acknowledgments. We thank Marc Levine for useful discussions and the Camille Jordan Institute for inviting the second author.

Notation. We use mainly the terminology and notations of Grothendieck-Dieudonné [10, §9.4 and 9.6] which agree with that of Demazure-Grothendieck used in [24, Exp. I.4]

(a) Let S be a scheme and let \mathcal{E} be a quasi-coherent sheaf over S . For each morphism $f : T \rightarrow S$, we denote by $\mathcal{E}_T = f^*(\mathcal{E})$ the inverse image of \mathcal{E} by the morphism f . We denote by $\mathbf{V}(\mathcal{E})$ the affine S -scheme defined by $\mathbf{V}(\mathcal{E}) = \mathrm{Spec}(\mathrm{Sym}^\bullet(\mathcal{E}))$; it is affine over S and represents the S -functor $Y \mapsto \mathrm{Hom}_{\mathcal{O}_Y}(\mathcal{E}_Y, \mathcal{O}_Y)$ [10, 9.4.9].

(b) We assume now that \mathcal{E} is locally free of finite rank and denote by \mathcal{E}^\vee its dual. In this case the affine S -scheme $\mathbf{V}(\mathcal{E})$ is of finite presentation (ibid, 9.4.11); also the S -functor $Y \mapsto H^0(Y, \mathcal{E}_Y) = \mathrm{Hom}_{\mathcal{O}_Y}(\mathcal{O}_Y, \mathcal{E}_Y)$ is representable by the affine S -scheme $\mathbf{V}(\mathcal{E}^\vee)$ which is also denoted by $\mathbf{W}(\mathcal{E})$ [24, I.4.6].

It applies to the locally free coherent sheaf $\mathcal{E}nd(\mathcal{E}) = \mathcal{E}^\vee \otimes_{\mathcal{O}_S} \mathcal{E}$ over S so that we can consider the affine S -scheme $\mathbf{V}(\mathcal{E}nd(\mathcal{E}))$ which is an S -functor in associative commutative and unital algebras [10, 9.6.2]. Now we consider the S -functor $Y \mapsto$

$\text{Aut}_{\mathcal{O}_Y}(\mathcal{E}_Y)$. It is representable by an open S -subscheme of $\mathbf{V}(\text{End}(\mathcal{E}))$ which is denoted by $\text{GL}(\mathcal{E})$ (*loc. cit.*, 9.6.4).

2. GENERALITIES ON HOMOGENEOUS SPACES

Let S be a scheme and let \bullet be the final object of the category of fppf sheaves in groups, that is $\bullet(T) = \text{Hom}_{S\text{-sch}}(T, S) = h_S(T)$ for each S -scheme T .

Let \underline{G} be a fppf S -sheaf in groups. Let \underline{X} be a fppf S -sheaf equipped with a left action of \underline{G} . We say that \underline{X} is homogeneous (resp. principal homogeneous) under \underline{G} if the map $\underline{X} \rightarrow \bullet$ is an epimorphism of fppf S -sheaves and if the action map $\underline{G} \times_S \underline{X} \rightarrow \underline{X} \times_S \underline{X}$, $(g, x) \rightarrow (x, g.x)$ is an epimorphism (resp. an isomorphism) of fppf S -sheaves. Similarly we have the notion of right homogeneous (resp. principal homogeneous) spaces.

Let \underline{E} be a right principal homogeneous \underline{G} -space and denote by \underline{G}' the twist of \underline{G} by \underline{E} through inner automorphisms. Then twisting by \underline{E} gives rise to an equivalence of categories between the category of left homogeneous \underline{G} -spaces and that of left homogeneous \underline{G}' -spaces.

Remark 2.1. For the discussion of the same material in the representable case, see [20, VI.1].

Lemma 2.2. *Let \underline{H} be a fppf group subsheaf of \underline{G} and \underline{X} be the \underline{G} -homogeneous space $\underline{G}/\underline{H}$. Consider the S -sheaf in groups $\underline{A} = \underline{\text{Aut}}_{\underline{G}}(\underline{X})$ which acts on the left of \underline{X} .*

(1) *Let $\underline{N} = \underline{N}_{\underline{G}}(\underline{H})$ be the normalizer of \underline{H} in \underline{G} (as in [13, §2.3]). Then the map $\underline{G} \times_S \underline{N} \rightarrow \underline{G}$, $(g, n) \mapsto gn$ induces an isomorphism $\underline{N}/\underline{H} \xrightarrow{\sim} \underline{A}^{op}$.*

(2) *The action of \underline{A} on \underline{X} is free and the map $\underline{X} \rightarrow \underline{A} \backslash \underline{X}$ is a (left) principal \underline{A} -homogeneous space.*

(3) *Let \underline{F} be a right \underline{A} -torsor and let \underline{X}' be the twist of \underline{X} by \underline{F} . Then \underline{X}' is a \underline{G} -homogeneous space and there is a canonical isomorphism $\underline{A}' \backslash \underline{X}' \xrightarrow{\sim} \underline{A} \backslash \underline{X}$, where $\underline{A}' = \underline{\text{Aut}}_{\underline{G}}(\underline{X}')$.*

(4) *The S -forms of \underline{X} (as \underline{G} -spaces) are classified by $H_{right}^1(S, \underline{A})$.*

Proof. We put $\underline{W} = \underline{N}/\underline{H}$.

(1) The map $\underline{G} \times_S \underline{N} \rightarrow \underline{G}$ induces a map $\underline{X} \times \underline{W} \rightarrow \underline{X}$ hence a group homomorphism $\phi : \underline{W} \rightarrow \underline{A}^{op}$ of S -sheaves in groups.

ϕ is a monomorphism. Let T be a S -scheme and let $w \in \underline{W}(T)$ be an element such that $\phi(w) = 1$. Up to localize we can assume that w arises from an element $n \in \underline{N}(T)$ which satisfies $1 \times n \in \underline{H}(T)$. Thus $w = 1$.

ϕ is an epimorphism. Let T be an S -scheme and let $f \in \underline{A}(T)$. We put $x = f([1]) \in \underline{X}(T)$ and up to localize we can assume that x arises from some $g \in \underline{G}(T)$. We claim that $g \in \underline{N}(T)$. For each T -scheme T' and each $g' \in \underline{G}(T')$ we have $f(g' \cdot [1]) = g' \cdot g_{T'} \cdot [1]$. In particular for each $h' \in \underline{H}(T')$, we have $h' g_{T'} \in g_{T'} \underline{H}(T')$ so

that $g_{T'}$ normalizes $\underline{H}(T')$. The claim is proven and it follows that $f = \phi(w)$ where w is the image of $g \in \underline{N}(T)$ in $\underline{W}(T)$.

(2) By (1), it is equivalent to prove that \underline{W} acts freely on \underline{X} . Let T be a scheme and $x \in \underline{X}(T)$. After localization we can assume that x arises from an element $g \in \underline{G}(T)$. Consider the T -sheaf in groups $\underline{\text{Stab}}_{\underline{W}}(x)$. For each T -scheme T' and $w \in \underline{\text{Stab}}_{\underline{W}}(x)(T')$, up to localization we can assume that w comes from an element $n \in \underline{N}(T')$. Then $[g] = [g] \cdot w = [gn]$, which implies $n \in \underline{H}(T')$ and w is the identity in $\underline{W}(T')$. Hence \underline{W} acts freely on \underline{X} and $\underline{X} \rightarrow \underline{A} \backslash \underline{X}$ is a right torsor of \underline{W} .

(3) Let p be the projection from \underline{X} to the quotient sheaf $\underline{A} \backslash \underline{X}$. Consider the map $\pi : \underline{F} \times \underline{X} \rightarrow \underline{A} \backslash \underline{X}$, which projects $\underline{F} \times \underline{X}$ to \underline{X} and then to the quotient sheaf $\underline{A} \backslash \underline{X}$. Then clearly π induces a map from \underline{X}' to $\underline{A} \backslash \underline{X}$, which we still denote by π .

Note that \underline{G} acts on the \underline{X} -factor of $\underline{F} \times \underline{X}$. As the G -action and the A -action commutes on X , this defines an \underline{G} -action on \underline{X}' . Choose a *fppf*-cover $\{S_i\}$ of S which trivializes the \underline{A} -torsor \underline{F} . For each S_i , there is an isomorphism of right \underline{A} -torsors between \underline{F}_{S_i} and \underline{A}_{S_i} . Hence $\underline{X}'_{S_i} \simeq \underline{X}_{S_i}$ and $\underline{A}'_{S_i} \simeq \underline{A}_{S_i}$.

Fix an isomorphism of \underline{A}_{S_i} -torsors $\iota_i : \underline{F}_{S_i} \rightarrow \underline{A}_{S_i}$. The map ι_i induces an isomorphism between \underline{X}'_{S_i} and \underline{X}_{S_i} which sends $[f, x]$ to $\iota_i(f)(x)$ for $x \in \underline{X}(T_i)$, $f \in \underline{F}(T_i)$ and for arbitrary S_i -scheme T_i . We denote this induced isomorphism still by ι_i . The map ι_i induces an isomorphism between \underline{A}'_{S_i} and \underline{A}_{S_i} by sending a' to $\iota_i \circ a' \circ \iota_i^{-1}$ for $a' \in \underline{A}'(T_i)$ and T_i an S_i -scheme. Then clearly ι_i induces an isomorphism $\bar{\iota}_i : (\underline{A}' \backslash \underline{X}')_{S_i} \rightarrow (\underline{A} \backslash \underline{X})_{S_i}$.

One checks that $p \circ \iota_i = \pi_{S_i}$. Hence $\bar{\iota}_i \circ p' = \pi_{S_i}$ where p' is the projection from \underline{X}' to $\underline{A}' \backslash \underline{X}'$. This implies that $\pi(x') = \pi(a' \cdot x')$ for $x' \in \underline{X}'(T)$, $a' \in \underline{A}'(T)$ and for arbitrary S -scheme T . Hence π induces a map $\bar{\pi}$ from $\underline{A}' \backslash \underline{X}'$ to $\underline{A} \backslash \underline{X}$. On each S_i , we have $\bar{\pi}_{S_i} = \bar{\iota}_i$. As $\bar{\iota}_i$ is an isomorphism, $\bar{\pi}$ is an isomorphism between $\underline{A}' \backslash \underline{X}'$ and $\underline{A} \backslash \underline{X}$.

(4) Let X' be an S -form of X as \underline{G} -spaces. Then the sheaf of isomorphisms $\underline{\text{Isom}}_{\underline{G}}(X, X')$ is a right \underline{A} -torsor and $\underline{\text{Isom}}_{\underline{G}}(\underline{X}, \underline{X}') \wedge^{\underline{A}} \underline{X}$ is canonically isomorphic to \underline{X}' .

Conversely given a right \underline{A} -torsor \underline{F} , let \underline{X}' be the twist of \underline{X} by \underline{F} . According to the proof of (3), the twist \underline{X}' is a G -homogeneous space and is isomorphic to \underline{X} *fppf*-locally. Hence \underline{X}' is an S -form of \underline{X} . We claim that the sheaf of isomorphisms $\underline{\text{Isom}}(\underline{X}, \underline{X}')$ is isomorphic to \underline{F} .

Note that $\underline{A} \wedge^{\underline{A}} \underline{X}$ is canonically isomorphic to \underline{X} and the \underline{A} -action on \underline{X} corresponds to the \underline{A} -action on the \underline{A} -factor of $\underline{A} \wedge^{\underline{A}} \underline{X}$. Regard \underline{A} as a right \underline{A} -torsor. Then there is a natural map ι between $\underline{\text{Isom}}_{\underline{A}}(\underline{A}, \underline{F})$ and $\underline{\text{Isom}}_{\underline{G}}(\underline{X}, \underline{X}') = \underline{\text{Isom}}_{\underline{G}}(\underline{A} \wedge^{\underline{A}} \underline{X}, \underline{F} \wedge^{\underline{A}} \underline{X}')$. To be precise, for an S -scheme S' and $\phi \in \underline{\text{Isom}}_{\underline{A}}(\underline{A}, \underline{F})(S')$, $\iota(\phi)([(a, x)]) = [(\phi(a), x)]$ for any S' -scheme S'' and $[(a, x)] \in (\underline{A} \wedge^{\underline{A}} \underline{X})(S'')$.

The sheaf of automorphisms of \underline{A} (as a right \underline{A} -torsor) is \underline{A} itself which acts on the left of \underline{A} . As $\underline{\text{Isom}}_{\underline{A}}(\underline{A}, \underline{F})$ is a right $\underline{\text{Aut}}_{\underline{A}}(\underline{A})$ -torsor, it is a right \underline{A} -torsor. It is easy

to check that ι is compatible with the \underline{A} -action. Since $\underline{\text{Isom}}_{\underline{A}}(\underline{A}, \underline{F})$ and $\underline{\text{Isom}}_{\underline{G}}(\underline{X}, \underline{X}')$ are both right \underline{A} -torsors, and ι is compatible with the \underline{A} -action, ι is an isomorphism.

There is a natural map i from $\underline{\text{Isom}}_{\underline{A}}(\underline{A}, \underline{F})$ to \underline{F} which sends $\phi \in \underline{\text{Isom}}_{\underline{A}}(\underline{A}, \underline{F})(S')$ to $\phi(1_{\underline{A}}) \in \underline{F}(S')$ for all S -schemes S' . Since i is compatible with the right \underline{A} -action, i is an isomorphism between right \underline{A} -torsors, and $i \circ \iota^{-1}$ gives the desired isomorphism between $\underline{\text{Isom}}_{\underline{G}}(\underline{X}, \underline{X}')$ and \underline{F} . Therefore the map which sends an S -form \underline{X}' to the right \underline{A} -torsor $\underline{\text{Isom}}_{\underline{G}}(\underline{X}, \underline{X}')$ is a bijection. \square

3. EMBEDDING FUNCTORS

Let S be a scheme. For a point $s \in S$, let $\kappa(s)$ be the residue field of s and $\overline{\kappa(s)}$ be the algebraic closure of $\kappa(s)$. Let \bar{s} be the scheme $\text{Spec}(\overline{\kappa(s)})$.

3.1. Twisted root data and Weyl groups. Let M be a \mathbf{Z} -lattice and M^\vee be its dual lattice. Let R and R^\vee be finite subsets of M and M^\vee respectively. Suppose that (M, M^\vee, R, R^\vee) satisfy the root data axioms.

Denote by \underline{M}_S (resp. \underline{R}_S) the constant sheaf of S associated to M (resp. R). Let T be an S -torus and denote by \mathcal{M} (resp. \mathcal{M}^\vee) the fppf sheaf of characters (resp. cocharacters) of T . Let $\Psi = (\mathcal{M}, \mathcal{M}^\vee, \mathcal{R}, \mathcal{R}^\vee)$ be a twisted root datum. (c.f. [24] Exp. XXI, 1.1). We say that the twisted root datum Ψ is of *type* (M, M^\vee, R, R^\vee) at $s \in S$ if $\Psi_{\bar{s}} \simeq (\underline{M}_{\bar{s}}, \underline{M}_{\bar{s}}^\vee, \underline{R}_{\bar{s}}, \underline{R}_{\bar{s}}^\vee)$.

We denote by $W(\Psi)$ the Weyl group of the twisted root datum Ψ . If Ψ is split of constant type, Ψ can be written as $(\underline{M}_S, \underline{M}_S^\vee, \underline{R}_S, \underline{R}_S^\vee)$ for some root datum (M, M^\vee, R, R^\vee) . In this case, the Weyl group $W(\Psi)$ is a finite group generated by the reflections s_α defined by

$$s_\alpha(x) = x - (\alpha^\vee, x)\alpha, \text{ for } \alpha \in R \text{ and } x \in M.$$

For $\alpha \in R$, the reflection s_α induces an automorphism \tilde{s}_α on T by

$$\tilde{s}_\alpha(t) = t(\alpha^\vee(\alpha(t)))^{-1} \text{ for any } S\text{-scheme } S', \alpha \in R \text{ and } t \in T(S').$$

(See [24] Exp. XXII, 3.3.) We denote by $W(T)$ the group generated by \tilde{s}_α . The map sending s_α to \tilde{s}_α defines an isomorphism from $W(\Psi)^{op}$ to $W(T)$. For $w \in W(\Psi)$, we denote by \tilde{w} its image in $W(T)$. In general, $W(\Psi)$ (resp. $W(T)$) is a finite étale group scheme over S , which is regarded as a subgroup scheme of $\underline{\text{Aut}}(\mathcal{M})$ (resp. $\underline{\text{Aut}}(T)$).

Let G be a reductive group scheme over S . Suppose that G has a maximal torus T over S . We denote by $\Phi(G, T)$ the twisted root datum of G with respect to T . Let $\underline{N}_G(T)$ the normalizer of T in G . Then $\underline{N}_G(T)/T$ is isomorphic to $W(T) \simeq W(\Phi(G, T))^{op}$. (cf. [24] Exp. XXII, 3.1-3.4)

3.2. The Orientation. Let Ψ_1, Ψ_2 be two twisted root data. Suppose that Ψ_2 and Ψ_1 are of the same type at each $s \in S$. Let $\underline{\text{Isom}}(\Psi_1, \Psi_2)$ be the sheaf of isomorphisms between Ψ_1 and Ψ_2 . Then $\underline{\text{Isom}}(\Psi_1, \Psi_2)$ is a right principal homogeneous space of $\underline{\text{Aut}}(\Psi_1)$ and a left principal homogeneous of $\underline{\text{Aut}}(\Psi_2)$. Define

$$\underline{\text{Isomext}}(\Psi_1, \Psi_2) = W(\Psi_2) \backslash \underline{\text{Isom}}(\Psi_1, \Psi_2).$$

Suppose $\underline{\text{Isomext}}(\Psi_1, \Psi_2)(S)$ is nonempty. For $u \in \underline{\text{Isomext}}(S)$, we define $\underline{\text{Isomint}}_u(\Psi_1, \Psi_2)$ to be the fiber of $\underline{\text{Isom}}(\Psi_1, \Psi_2) \rightarrow \underline{\text{Isomext}}(\Psi_1, \Psi_2)$ at u .

Let G be a reductive group scheme over S . The *type* of G at s is the type of the root datum of $G_{\bar{s}}$ with respect to its maximal torus. (ref. [24], Exp. 22, Def. 2.6.1, 2.7).

Suppose that G has a maximal torus T . We define $\underline{\text{Isomext}}(G, \Psi)$ to be $\underline{\text{Isomext}}(\Phi(G, T), \Psi)$. For a different choice of maximal tori T' , there is a canonical isomorphism between $\underline{\text{Isomext}}(\Phi(G, T), \Psi)$ and $\underline{\text{Isomext}}(\Phi(G, T'), \Psi)$ (ref. [15, §1.2.1]). Hence $\underline{\text{Isomext}}(G, \Psi)$ is well-defined.

We define $\underline{\text{Isomext}}(\Psi, G)$ in a similar way. As there is a canonical isomorphism between $\underline{\text{Isom}}(\Phi(G, T), \Psi)$ and $\underline{\text{Isom}}(\Psi, \Phi(G, T))$, which sends $f \in \underline{\text{Isom}}(\Phi(G, T), \Psi)(S')$ to $f^{-1} \in \underline{\text{Isom}}(\Psi, \Phi(G, T))(S')$, the two functors $\underline{\text{Isomext}}(G, \Psi)$ and $\underline{\text{Isomext}}(\Psi, G)$ are canonically isomorphic.

An *orientation* of G with respect to Ψ is an element of $\underline{\text{Isomext}}(G, \Psi)(S)$.

A twisted root datum Ψ is *admissible* for G if at each $s \in S$, the type of $G_{\bar{s}}$ is the same as the type of $\Psi_{\bar{s}}$, and $\underline{\text{Isomext}}(G, \Psi)(S) \neq \emptyset$.

Let G' be an S -form of G . Let $\underline{\text{Isom}}(G, G')$ be the sheaf of group isomorphisms between G and G' . Note that G acts on itself by conjugation. Thus we can define the right quotient of $\underline{\text{Isom}}(G, G')$ by the adjoint quotient G_{ad} of G and denote $\underline{\text{Isom}}(G, G')/G_{ad}$ by $\underline{\text{Isomext}}(G, G')$.

Let Ψ_1, Ψ_2 and Ψ_3 be twisted root data over S . Suppose that all of them are of the same type at each geometric fibre of S . Then we have the following map :

$$(3.1) \quad \underline{\text{Isomext}}(\Psi_1, \Psi_2) \times \underline{\text{Isomext}}(\Psi_2, \Psi_3) \rightarrow \underline{\text{Isomext}}(\Psi_1, \Psi_3),$$

which comes from the composition of isomorphisms

$$\underline{\text{Isom}}(\Psi_1, \Psi_2) \times \underline{\text{Isom}}(\Psi_2, \Psi_3) \rightarrow \underline{\text{Isom}}(\Psi_1, \Psi_3).$$

Suppose G and Ψ_1 are of the same type at each geometric fiber of S . Similarly we have the following pairings coming from the composition of isomorphisms between root data.

Proposition 3.2.1. *We have the following pairings*

- (1) $\underline{\text{Isomext}}(G, \Psi_1) \times \underline{\text{Isomext}}(\Psi_1, \Psi_2) \rightarrow \underline{\text{Isomext}}(G, \Psi_2)$.
- (2) $\underline{\text{Isomext}}(G, \Psi_1) \times \underline{\text{Isomext}}(G, \Psi_2) \rightarrow \underline{\text{Isomext}}(\Psi_1, \Psi_2)$.
- (3) $\underline{\text{Isomext}}(G, \Psi_1) \times \underline{\text{Isomext}}(G', \Psi_1) \rightarrow \underline{\text{Isomext}}(G, G')$.
- (4) $\underline{\text{Isomext}}(G', G) \times \underline{\text{Isomext}}(G, \Psi) \rightarrow \underline{\text{Isomext}}(G', \Psi)$.

Proof. The first two assertions come from the definition of $\underline{\text{Isomext}}(G, \Psi)$ and the composition of isomorphisms between root data. For (3) and (4), we notice that if G and G' have maximal tori T and T' respectively, then $\underline{\text{Isomext}}(\Phi(G, T), \Phi(G', T'))$ is representable and isomorphic to $\underline{\text{Isomext}}(G, G')$. As reductive groups have maximal tori étale locally, assertions (3) and (4) can be deduced from descent. We refer to [15, Prop. 1.4] for more details. \square

Denote by $G//G$ the adjoint quotient of G (ref [16]). Suppose that G has a maximal torus T . Let $T/W(T)$ be the quotient sheaf of T by the action of Weyl group $W(T)$. Then $T/W(T)$ is isomorphic to $G//G$ ([16] Thm. 4.1).

Consider the quotient map (of fppf-sheaves) $\pi : T \rightarrow T/W(T)$. Let $x \in (G//G)(S)$. As $T/W(T)$ and $G//G$ are isomorphic, we can regard x as an element in $(T/W(T))(S)$. Suppose that there is a semisimple regular element $t_1 \in G(S)$ whose image in $(G//G)(S)$ is x . Let T_1 be the centralizer $\underline{C}_G(t_1)$, which is also a maximal torus. Let Ψ (resp. Ψ_1) be the root datum associated to T (resp. T_1). Then the natural inclusion from T to G gives an orientation $v \in \underline{\text{Isomext}}(G, \Psi)(S)$. Similarly, we get an orientation $v_1 \in \underline{\text{Isomext}}(G, \Psi_1)(S)$ from the natural inclusion of T_1 in G . Then by 3.2.1 (2) we get an orientation $u \in \underline{\text{Isomext}}(\Psi, \Psi_1)(S)$ coming from $v \cdot v_1$. Then we have the following proposition.

Proposition 3.2.2. *The left $W(T)$ -torsor $\pi^{-1}(x)$ is isomorphic to $\underline{\text{Isomint}}_u(\Psi, \Psi_1)$.*

Proof. For an S -scheme S' and $t \in \pi^{-1}(x)(S')$, we denote by $\underline{\text{Transp}}(t, t_1)$ the transporter from t to t_1 . Both t and t_1 are mapped to the same element in $(G//G)(S)$, and hence $\underline{\text{Transp}}(t, t_1) \rightarrow \bullet$ is surjective. Since t_1 is a semisimple regular element, so is t . As t lies in T , the centralizer $\underline{C}_G(t)$ of t is the torus $T_{S'}$ and $\underline{\text{Transp}}(t, t_1)$ is a right $T_{S'}$ -torsor.

The conjugation action of T on $\Psi = \Phi(G, T)$ is trivial. Therefore the canonical morphism

$$\underline{\text{Transp}}(t, t_1) \wedge^{T_{S'}} \Psi_{S'} \rightarrow \Psi_{1, S'}$$

defines an element in $\underline{\text{Isomint}}_u(\Psi, \Psi_1)(S')$. In this way, we get a map

$$i : \pi^{-1}(x) \rightarrow \underline{\text{Isomint}}_u(\Psi, \Psi_1).$$

To be precise, write the root datum Ψ as $(\mathcal{M}, \mathcal{M}^\vee, \mathcal{R}, \mathcal{R}^\vee)$. For an S' -scheme S'' and $h \in \underline{\text{Transp}}(t, t_1)(S'')$, $i(t)(m) = m \circ \text{int}(h^{-1})$ for all $m \in \mathcal{M}_{S''} = \text{Hom}_{S''}(T_{S''}, \mathbb{G}_{m, S''})$. Note that $i(t)$ is independant of the choice of h since T acts trivially on \mathcal{M} .

As $\underline{\text{Isomint}}_u(\Psi, \Psi_1)$ is a right $W(\Psi)$ -torsor and $W(T) \simeq W(\Psi)^{op}$, we can regard it as left $W(T)$ -torsor. We write down explicitly this action. For $w \in W(T)_{S'}$, let n_w be a lift of w in $N(T)(S'')$ for some S' -scheme S'' . As T acts on \mathcal{M} trivially, $\text{int}(n_w^{-1})$ induces an automorphism on $\mathcal{M}_{S'}$. For $f \in \underline{\text{Isomint}}_u(\Psi, \Psi_1)(S')$, $(w \cdot f)(m) := f(m \circ \text{int}(n_w))$ for $m \in \mathcal{M}_{S'} = \text{Hom}_{S'-grp}(T_{S'}, \mathbb{G}_{m, S'})$.

Since $\pi^{-1}(x)$ and $\underline{\text{Isomint}}_u(\Psi, \Psi_1)$ are both $W(T)$ -torsors, to check i is an isomorphism, it suffices to check the map i is $W(T)$ -equivariant. Let $w \in W(T)_{S'}$ and

n_w be a lift of w in $N(T)(S'')$ for some S' -scheme S'' . For $h \in \underline{\text{Transp}}(t, t_1)(S'')$, $hn_w^{-1} \in \underline{\text{Transp}}(w \cdot t, t_1)(S'')$. Thus

$$\begin{aligned}
 (3.2) \quad i(w \cdot t)(m) &= m \circ \text{int}((hn_w^{-1})^{-1}) \\
 &= m \circ \text{int}(n_w) \circ \text{int}(h^{-1}) \\
 &= i(t)(m \circ \text{int}(n_w)) \\
 &= (w \cdot i(t))(m).
 \end{aligned}$$

Hence i is an isomorphism between $W(T)$ -torsors. \square

3.3. Embedding functors. Let G be a reductive group scheme over S and $\Psi = (\mathcal{M}, \mathcal{M}^\vee, \mathcal{R}, \mathcal{R}^\vee)$ be an admissible root datum for G . Let T be the S -torus with sheaf of characters \mathcal{M} .

Suppose there is an embedding f of T into G as algebraic group schemes. Then f induces an isomorphism from the character group of $f(T)$ to \mathcal{M} . We call this map $f^\#$. If $f^\#$ induces an isomorphism between $\Phi(G, f(T))$ and Ψ , then we say f is an embedding with respect to the twisted root datum Ψ .

Define the *embedding functor* $\mathfrak{E}(G, \Psi)$ as follows: for each S -scheme S' ,

$$\mathfrak{E}(G, \Psi)(S') = \left\{ f : T_{S'} \hookrightarrow G_{S'} \left| \begin{array}{l} f \text{ is both a closed immersion and a group} \\ \text{homomorphism which induces an isomorphism} \\ f^\# : \Phi(G_{S'}, f(T_{S'})) \xrightarrow{\sim} \Psi_{S'} \end{array} \right. \right\}$$

Suppose $\underline{\text{Isomext}}(G, \Psi)(S)$ is nonempty. Fix an orientation $v \in \underline{\text{Isomext}}(G, \Psi)(S)$. The *oriented embedding functor* $\mathfrak{E}(G, \Psi, v)$ over S is defined by

$$\mathfrak{E}(G, \Psi, v)(S') = \left\{ f : T_{S'} \hookrightarrow G_{S'} \left| \begin{array}{l} f \in \mathfrak{E}(G, \Psi)(S'), \text{ and the image of } f^\# \\ \text{in } \underline{\text{Isomext}}(G, \Psi)(S') \text{ is } v. \end{array} \right. \right\}$$

The oriented embedding functor $\mathfrak{E}(G, \Psi, v)$ is representable by an affine S -scheme which is a left homogeneous G -space ([15, 1.6]). For an S -scheme S' and an element $f \in \mathfrak{E}(G, \Psi, v)(S')$, the stabilizer of f in $G_{S'}$ is $f(T_{S'})$, which is a maximal torus of $G_{S'}$. It admits a right action of $W(T)$ and the quotient sheaf $\mathfrak{E}(G, \Psi, v)/W(T)$ identifies with the S -scheme \mathcal{T}_G of maximal tori of G . (See [15, §1.1 and §1.2] for more details.)

Remark 3.1. In [15, §1.2.2] we define the oriented embedding functor $\mathfrak{E}(G, \Psi, v')$ by orientation $v' \in \underline{\text{Isomext}}(\Psi, G)(S)$ instead of $v \in \underline{\text{Isomext}}(G, \Psi)(S)$ here. However when v' is the image of v under the canonical isomorphism $\underline{\text{Isomext}}(G, \Psi) \rightarrow \underline{\text{Isomext}}(\Psi, G)$, these two definitions are clearly equivalent.

Lemma 3.2. Keep the notation as above.

- (1) We have an isomorphism $W(\Psi) \xrightarrow{\sim} \underline{\text{Aut}}_G(\mathfrak{E}(G, \Psi, v))$ as group schemes.
- (2) Let Ψ' be another admissible root datum for G , and $v' \in \underline{\text{Isomext}}(G, \Psi')(S)$. Let $u \in \underline{\text{Isomext}}(\Psi, \Psi')(S)$ be the orientation $v' \cdot v$ by the pairing (3.2.1). Under the

identification of $W(\Psi)$ and $\underline{\text{Aut}}_G(\mathfrak{E}(G, \Psi, v))$ in (1), there is a natural isomorphism between $\underline{\text{Isomint}}_u(\Psi, \Psi')$ and $\underline{\text{Isom}}(\mathfrak{E}(G, \Psi, v), \mathfrak{E}(G, \Psi', v'))$ as right $W(\Psi)$ -torsors.

(3) Let F be a right $W(\Psi)$ -torsor and $\Psi' = F \wedge^{W(\Psi)} \Psi$. Let c be the canonical element of $\underline{\text{Isomext}}(\Psi, \Psi')(S)$, and let $v_1 \in \underline{\text{Isomext}}(G, \Psi')$ be $c \cdot v$ by the pairing (3.2.1). By identifying $W(\Psi)$ with $\underline{\text{Aut}}_G(\mathfrak{E}(G, \Psi, v))$, $F \wedge^{W(\Psi)} \mathfrak{E}(G, \Psi, v)$ is isomorphic to $\mathfrak{E}(G, \Psi', v_1)$.

(4) Let E be a right G -torsor and denote by G' the twist of G by E via inner automorphisms. Let c be the canonical element of $\underline{\text{Isomext}}(G', G)(S)$. Then the homogeneous G' -space $E \wedge^G \mathfrak{E}(G, \Psi, v)$ is isomorphic to $\mathfrak{E}(G', \Psi, v')$ where $v' \in \underline{\text{Isomext}}(G', \Psi)(S)$ is $c \cdot v$ by the pairing (3.2.1).

Proof. (1) For an S -scheme S' and $w \in W(\Psi)(S')$, we denote by \tilde{w} the image of w in $W(T)(S')$. Define a morphism $\iota : W(\Psi) \rightarrow \underline{\text{Aut}}_G(\mathfrak{E}(G, \Psi, v))$ as the following. For an S -scheme S' and $w \in W(\Psi)(S')$,

$$\iota(w)(x) := x \circ \tilde{w} \text{ for all } S'\text{-schemes } S'', \text{ and } x \in \mathfrak{E}(G, \Psi, v)(S'').$$

For w_1 and $w_2 \in W(\Psi)(S')$,

$$\begin{aligned} \iota(w_1 \circ w_2)(x) &= x \circ \widetilde{(w_1 \circ w_2)} \\ (3.3) \quad &= x \circ \tilde{w}_2 \circ \tilde{w}_1 \\ &= (\iota(w_1) \circ \iota(w_2))(x). \end{aligned}$$

Hence ι is a group homomorphism.

Clearly ι is monomorphism. To see that ι is an isomorphism, it suffices to check ι is an isomorphism of fppf-sheaves. Let $\{S_i\}$ be a fppf-cover of S such that G_{S_i} and Ψ_{S_i} are both split. Then $\mathfrak{E}(G, \Psi, v)(S_i)$ is not empty. Choose $x_i \in \mathfrak{E}(G, \Psi, v)(S_i)$. By Lemma 2.2, $\underline{\text{Aut}}_G(\mathfrak{E}(G, \Psi, v))(S_i)$ is isomorphic to $(\underline{N}_G(x_i(T))/x_i(T))^{op}$, where $\underline{N}_G(x_i(T))$ is the normalizer of $x_i(T)$ in G . Since $\underline{N}_G(x_i(T))/x_i(T) \simeq W(\Phi(G_{S_i}, x_i(T_{S_i})))^{op}$, the group $\underline{N}_G(x_i(T))/x_i(T)$ is isomorphic to $W(\Psi_{S_i})^{op}$. Thus $\underline{\text{Aut}}_G(\mathfrak{E}(G, \Psi, v))_{S_i} \simeq W(\Psi_{S_i})$. As ι is a monomorphism and $\underline{\text{Aut}}_G(\mathfrak{E}(G, \Psi, v))_{S_i}$ is isomorphic to $W(\Psi_{S_i})$, ι is an isomorphism.

(2) Write $\Psi' = (\mathcal{M}', (\mathcal{M}')^\vee, \mathcal{R}', (\mathcal{R}')^\vee)$ and denote by T' the torus with sheaf of characters \mathcal{M}' . For an S -scheme S' and $f \in \underline{\text{Isomint}}_u(\Psi, \Psi')(S')$, we denote by \tilde{f} the isomorphism from $T'_{S'}$ to $T_{S'}$ induced by f . Define $\eta : \underline{\text{Isomint}}_u(\Psi, \Psi') \rightarrow \underline{\text{Isom}}(\mathfrak{E}(G, \Psi, v), \mathfrak{E}(G, \Psi', v'))$ by $\eta(f)(x) = x \circ \tilde{f}$, for all S' -scheme S'' , and $x \in \mathfrak{E}(G, \Psi, v)(S'')$.

Next we show that η is compatible with the $W(\Psi)$ -action under the identification of $W(\Psi)$ and $\underline{\text{Aut}}_G(\mathfrak{E}(G, \Psi, v))$. Let $w \in W(\Psi)(S')$. Then

$$\begin{aligned}
\eta(f \circ w)(x) &= x \circ (\widetilde{f \circ w}) \\
&= x \circ \tilde{w} \circ \tilde{f} \\
(3.4) \quad &= \eta(f)(x \circ \tilde{w}) \\
&= \eta(f)(\iota(w)(x)) \\
&= (\eta(f) \circ \iota(w))(x).
\end{aligned}$$

This shows that η is compatible with the $W(\Psi)$ -action. Since $\underline{\text{Isomint}}_u(\Psi, \Psi')$ and $\underline{\text{Isom}}(\mathfrak{E}(G, \Psi, v), \mathfrak{E}(G, \Psi', v'))$ are both right $W(\Psi)$ -torsors, this implies that η is an isomorphism.

(3) Write Ψ' as $(\mathcal{M}', (\mathcal{M}')^\vee, \mathcal{R}', (\mathcal{R}')^\vee)$ and let T' be the torus with character group \mathcal{M}' . For an S -scheme S' and $f \in F(S')$, we define $\varphi_f^\sharp : \mathcal{M}_{S'} \rightarrow \mathcal{M}'_{S'}$ as $\varphi_f^\sharp(m) = (f, m)$ for all S' -scheme S'' and $m \in \mathcal{M}_{S'}(S'')$. Denote by φ_f the group homomorphism from T' to T defined by φ_f^\sharp .

Define $\varsigma : F \times \mathfrak{E}(G, \Psi, v) \rightarrow \mathfrak{E}(G, \Psi', v')$ as follows. For an S -scheme S' and $(f, x) \in (F \times \mathfrak{E}(G, \Psi, v))(S')$, let $\varsigma(f, x) = x \circ \varphi_f$. Clearly $x \circ \varphi_f$ is an embedding of T' in G with orientation v' . Note that for $w \in W(\Psi)(S')$, we have

$$\begin{aligned}
\varphi_{fw}^\sharp(m) &= (fw, m) \\
(3.5) \quad &= (f, w^{-1}m) \\
&= \varphi_f^\sharp(w^{-1}m).
\end{aligned}$$

Therefore $\varphi_{fw}^\sharp = \varphi_f^\sharp \circ w^{-1}$ and $\varphi_{fw} = \tilde{w}^{-1} \circ \varphi_f$. It follows that

$$\begin{aligned}
\varsigma(fw, x) &= x \circ \varphi_{fw} \\
(3.6) \quad &= x \circ \tilde{w}^{-1} \circ \varphi_f \\
&= (w^{-1} \cdot x) \circ \varphi_f \\
&= \varsigma(f, w \cdot x).
\end{aligned}$$

Hence ς induces a morphism from $F \wedge^{W(\Psi)} \mathfrak{E}(G, \Psi, v)$ to $\mathfrak{E}(G, \Psi', v')$, which is clearly an isomorphism.

(4) For an S -scheme S' and $e \in E(S')$, we define $f_e : G_{S'} \rightarrow G'_{S'}$ by $f_e(g) = [e, g]$ for all S' -scheme S'' and $g \in G(S'')$. Then f_e is an isomorphism between $G_{S'}$ and $G'_{S'}$.

For $y = (e, x) \in (E \times_S \mathfrak{E}(G, \Psi, v))(S')$, we define

$$\iota : E \times_S \mathfrak{E}(G, \Psi, v) \rightarrow \mathfrak{E}(G', \Psi, v')$$

as $\iota(y) = f_e \circ x$. It is clear that $\iota(y) \in \mathfrak{E}(G', \Psi)(S')$. As x is an embedding of orientation v , by the definition of canonical orientation, the embedding $\iota(y)$ is of

orientation v' . Since $\iota(e, x) = \iota(e \cdot g, g^{-1} \cdot x)$ for all $g \in G(S')$, ι induces an isomorphism from $E \wedge^G \mathfrak{E}(G, \Psi, v)$ to $\mathfrak{E}(G', \Psi, v')$. \square

Remark 3.3. Let G^{sc} be the simply connected cover of G and let G_{ad} be its adjoint quotient. On the other hand let Ψ^{sc} (resp. Ψ_{ad}) the simply connected (resp. adjoint) root datum associated to Ψ . Then we can attach to (Ψ, v) an oriented root datum (Ψ^{sc}, v^{sc}) (resp. (Ψ_{ad}, v_{ad})) and isomorphisms

$$\mathfrak{E}(G^{sc}, \Psi^{sc}, v^{sc}) \xrightarrow{\sim} \mathfrak{E}(G, \Psi, v) \xrightarrow{\sim} \mathfrak{E}(G_{ad}, \Psi_{ad}, v_{ad}).$$

It implies that we can deal in practice with semisimple simply connected (resp. adjoint) group schemes. Another advantage is that we have isomorphisms of finite étale group schemes

$$\underline{\text{Isomext}}(G^{sc}, \Psi^{sc}) \xrightarrow{\sim} \underline{\text{Isomext}}(G_{ad}, \Psi_{ad}) \xrightarrow{\sim} \underline{\text{Isomext}}(\underline{\text{Dyn}}(G), \underline{\text{Dyn}}(\Psi)).$$

(See [15, Cor. 1.7].)

3.4. The main result.

Theorem 3.4. *Let X be a separated S -scheme satisfying one of the following conditions:*

- (i) S is locally noetherian;
- (ii) X is locally of finite type over S .

We assume that X is a left homogeneous space under G and that for each point $s \in S$, the stabilizers of $X_{\bar{s}}$ are maximal tori of $G_{\bar{s}}$.

(1) *There exists a root datum $\Psi = (\mathcal{M}, \mathcal{M}^\vee, \mathcal{R}, \mathcal{R}^\vee)$ which is admissible for G and an orientation $v \in \underline{\text{Isomext}}(G, \Psi)(S)$ such that X is isomorphic to $\mathfrak{E}(G, \Psi, v)$ as left G -homogeneous spaces.*

(2) *The oriented root datum (Ψ, v) in (1) is unique up to isomorphism.*

Proof. We shall establish first that the local stabilizers of X are maximal tori. Since G is smooth over S with connected fibers, X is smooth of finite presentation over S according to [20, prop. VI.1.2].

(1) Let $(S_i)_{i \in I}$ be a fppf cover of S such that X_{S_i} is G_{S_i} -isomorphic to G_{S_i}/H_i where H_i is a S_i -subgroup scheme of G_{S_i} (see [24, IV 6.7.3]). Since H_i is the stabilizer of a point of $X(S_i)$ (which is S_i -separated), H_i is a closed S_i -subgroup scheme of G_{S_i} and is in particular affine.

Again by [20, prop. VI.1.2], H_i is flat locally of finite presentation over S_i so is affine flat of finite presentation over S_i . As the geometric fibers of H_i are tori, it follows that H_i is a S_i -torus [24, X.4.9]. Therefore H_i is a maximal S_i -torus of G_{S_i} for each $i \in I$.

In particular, $C(G)$ acts trivially on X , so X is a G_{ad} -homogeneous space whose local stabilizers are maximal tori. By Remark 3.3 we reduce to the case that G is an adjoint semisimple S -group scheme. We denote by G^q a quasi-split S -form of G . Since

G is an inner form of G^q , there is a canonical orientation $c \in \underline{\text{Isomext}}(G, G^q)(S)$. Let T^q be a maximal torus of G^q , Ψ^q be the root datum $\Phi(G^q, T^q)$ and $v^q \in \underline{\text{Isomext}}(G^q, \Psi^q)(S)$ be the orientation induced by the inclusion of T^q to G^q . Let $u = c \cdot v^q \in \underline{\text{Isomext}}(G, \Psi^q)(S)$ (by 3.2.1) and consider the homogeneous G -space $X^q = \mathfrak{E}(G, \Psi^q, u)$. This is a left G homogeneous space whose local stabilizers are maximal tori.

Claim. X is an S -form of X^q .

Denote by $\underline{\text{Isom}}_G(X^q, X)$ the sheaf of G -space isomorphisms over S . Since $\Phi(G_{S_i}, H_i)$ is fppf-locally isomorphic to $\Psi_{S_i}^q$ over S_i , we can find a fppf-cover S_{ij} of S_i such that there is $f \in X^q(S_{ij})$ with $f(T_{S_{ij}}^q) = H_{i, S_{ij}}$. As the stabilizer $\underline{\text{Stab}}_{G_{S_{ij}}}(f)$ is $H_{i, S_{ij}}$, we have $X_{S_{ij}}^q \simeq G_{S_{ij}}/H_{i, S_{ij}}$ as $G_{S_{ij}}$ -spaces. Hence there is an isomorphism between X^q and X over S_{ij} which preserves the G -structure. This proves that $\underline{\text{Isom}}_G(X^q, X)$ is nonempty. The canonical isomorphism $\underline{\text{Isom}}_G(X^q, X) \wedge^{\underline{\text{Aut}}_G(X^q)} X^q \rightarrow X$ then shows that X is an S -form of X^q . Our claim is established.

By identifying $W(\Psi^q)$ with $\underline{\text{Aut}}_G(X^q)$ as in Lemma 3.2 (1), we denote by Ψ the root datum $\underline{\text{Isom}}_G(X^q, X) \wedge^{\underline{\text{Aut}}_G(X^q)} \Psi^q$. Let $c' \in \underline{\text{Isomext}}(\Psi^q, \Psi)(S)$ be the canonical orientation, and v be the orientation $u \cdot c'$ by pairing (3.2.1). We conclude that X is isomorphic to the embedding functor $\mathfrak{E}(G, \Psi, v)$ by Lemma 3.2(3).

(2) It follows from Lemma 3.2(2). □

4. THE CASE OF A SEMILOCAL RING

4.1. Schematically dominant morphisms. We recall that a morphism of schemes $f : Y \rightarrow X$ is schematically dominant if $\mathcal{O}_X \rightarrow f_*\mathcal{O}_Y$ is injective; if f is an immersion we say that X is schematically dense in Y [11, 11.10.2]. If $g : X' \rightarrow X$ is flat, and f is schematically dominant and quasi-compact, then $f' : Y \times_X X' \rightarrow X'$ is schematically dominant (*loc. cit.*, 11.10.5).

If $f : X \rightarrow Y$ is an S -morphism of schemes, we say that f is schematically S -dominant (or universally schematically dominant with respect to S) if for each S -scheme S' , $f_{S'} : X_{S'} \rightarrow Y_{S'}$ is schematically dominant; similarly if f is an immersion, we say that Y is schematically S -dense in X . Under mild assumptions, this property can be checked fiberwise (*ibid*, 11.10.9). The base change property above extends mechanically: If $g : X' \rightarrow X$ is flat and $f : Y \rightarrow X$ is schematically S -dominant and quasi-compact, then $f' : Y \times_X X' \rightarrow X'$ is schematically S -dominant.

Rydh extended that to morphism of algebraic spaces [22, §7.5]. A morphism of algebraic spaces $f : X \rightarrow Y$ is schematically dominant if the morphism $\mathcal{O}_Y \rightarrow f_*\mathcal{O}_X$ is injective in the small étale site; if f is an immersion, we say that X is schematically dense in Y .

As in [23, §3.5] the base change property extends in that framework: given a flat morphism $g : X' \rightarrow X$ of algebraic spaces assuming that $f : Y \rightarrow X$ is schematically dominant and quasi-compact, then $f' : Y \times_X X' \rightarrow X'$ is schematically dominant.

If S is a scheme and $f : Y \rightarrow X$ is a morphism of S -algebraic spaces, we say that $f : X \rightarrow Y$ is schematically S -dominant if for each S -scheme S' , $f_{S'} : X_{S'} \rightarrow Y_{S'}$ is schematically dominant; similarly if f is an immersion, we say that Y is schematically S -dense in X . One last time, given a flat morphism $g : X' \rightarrow X$ and assuming that $f : Y \rightarrow X$ is schematically S -dominant and quasi-compact, then $f' : Y \times_X X' \rightarrow X'$ is schematically S -dominant.

4.2. (4)-Versal torsors. Our goal is to adapt the framework of versal torsors [9, §5] from fields to the semilocal ring setting by using algebraic spaces for which we use Olsson's book as reference [O] as well with the Stacks project [26]. It is close of Reichstein-Tossici's definition of (3)-versality for algebraic groups [21, def. 1.3].

Let $S = \text{Spec}(R)$ be a semilocal scheme and let G be an affine flat S -group scheme of finite presentation.

Definition 4.1. *Let $f : E \rightarrow X$ be a G -torsor where E is a qcqs S -scheme and X is a qcqs S -algebraic space. We say that $E \rightarrow X$ is (4)-versal if for any open retrocompact subspace U schematically S -dense in X then for each semilocal R -algebra B whose residue fields of maximal ideals are infinite, and for each G -torsor F over B , there exists $x \in U(B)$ such that $E_x \cong F$ as G -torsors.*

Remarks 4.2. (a) We observe that if $f : E \rightarrow X$ is a (4)-versal G -torsor so is $f^{-1}(V) \rightarrow V$ for each schematically S -dense open retrocompact subspace V of X .

(b) There exists a dense open subspace X' of X which is a scheme [26, Tag 06NH]. If X' can be chosen furthermore schematically S -dense retrocompact, we see then that the open dense retrocompact subscheme $U' = U \times_X X'$ can be chosen in the definition. This is the case if S is noetherian of dimension ≤ 1 and X is reduced separated according to [26, Tag 0ADD].

Lemma 4.3. *Let $G \rightarrow \text{GL}(\mathcal{E})$ be a faithful linear representation of G where \mathcal{E} is a locally free R -module of finite rank.*

(1) *The fppf quotient $X = \text{GL}(\mathcal{E})/G$ is representable by an algebraic space and the quotient map $\text{GL}(\mathcal{E}) \rightarrow X$ is a (4)-versal G -torsor with right G -action.*

(2) *Assume that $\mathbf{V}(\mathcal{E})$ admits an open universally dense S -subscheme V which is G -stable and such that G acts freely on V . Let $Y = V/G$ be the quotient algebraic space. Then $V \rightarrow Y$ is a (4)-versal G -torsor.*

Proof. (1) If R is noetherian, then the fppf quotient $X = \text{GL}(\mathcal{E})/G$ is represented by an S -algebraic space [1, th. 3.1.1]. The usual noetherian reduction trick shows that it is the case in general. Let B be a semilocal ring whose residue fields of maximal ideals are infinite and F be a G -torsor over B . Denote by κ_i the residue fields of maximal ideals \mathfrak{M}_i of B .

We first consider the case that B is indecomposable, i.e. $\text{Spec } B$ is connected. In this case $\mathcal{E} \otimes_R B$ becomes a free module over B of rank n , and we can regard $\text{GL}(\mathcal{E})_B$

as $\mathrm{GL}_{n,B}$. Consider the exact sequence of pointed sets:

$$1 \rightarrow G(B) \rightarrow \mathrm{GL}_n(B) \rightarrow X(B) \xrightarrow{\delta} H_{fppf}^1(B, G) \rightarrow H_{fppf}^1(B, \mathrm{GL}_n).$$

As B is semilocal, we have $H_{fppf}^1(B, \mathrm{GL}_n) = 1$ since all locally free B -modules of rank r are trivial. Hence the torsor F is a pull-back of $\mathrm{GL}(\mathcal{E}) \rightarrow X$ at some element $x \in X(B)$. Denote by \bar{x}_i the image of x in $X(\kappa_i)$.

Denote by X_{κ_i} (resp. X_B) the functor obtained from X restricted to the category of κ_i -schemes (resp. B -schemes). Then X_{κ_i} is an algebraic space over $\mathrm{Spec} \kappa_i$. Note that we have a natural left $\mathrm{GL}(\mathcal{E})$ -action on X . Hence x defines a morphism f_x from $\mathrm{GL}_{n,B}$ to X_B as follows:

$$f_x(g) = g \cdot x,$$

for all B -algebra C and all $g \in \mathrm{GL}_{n,B}(C)$. This induces a morphism $f_{\bar{x}_i} : \mathrm{GL}_{n,\kappa_i} \rightarrow X_{\kappa_i}$.

Let U be an open retrocompact subspace of X which is schematically S -dense in X . We have the following commutative diagram

$$\begin{array}{ccc} U \times_{X, f_x} \mathrm{GL}_{n,B} & \longrightarrow & \mathrm{GL}_{n,B} \\ \downarrow & & \downarrow f_x \\ U_B & \longrightarrow & X_B \end{array}$$

The algebraic space $U \times_{X, f_x} \mathrm{GL}_{n,B}$ is representable by an open retrocompact subscheme V_x of $\mathrm{GL}_{n,B}$ which is S -dense according to base change property listed in 4.1. Since κ_i is infinite, $V_x(\kappa_i)$ is nonempty. Hence for each i , there is a point $g_i \in V_x(\kappa_i)$ such that $g_i \cdot x_i$ is in the image of $U(\kappa_i)$. By Chinese remainder theorem, there is an element $\bar{g} \in \mathrm{GL}_{n,B}(B/\prod \mathfrak{M}_i)$ which lifts g_i for all i . As B is semilocal, we can lift \bar{g} to an element $g \in \mathrm{GL}_{n,B}(B)$. By our construction g is in $V_x(B)$. Hence $g \cdot x$ is in the image of $U(B)$. Since $\delta(x) = \delta(g \cdot x)$, F is a pull back of $\mathrm{GL}(\mathcal{E}) \rightarrow X$ at $g \cdot x \in U(B)$.

As every semilocal ring B can be decomposed into a finite product of indecomposable semilocal rings B_i , the first assertion follows.

(2) Let B be a semilocal ring whose residue fields of maximal ideals are infinite and F be a G -torsor over B . Denote by κ_i the residue fields of maximal ideals \mathfrak{M}_i of B . Let U be an open retrocompact subspace of V/G which is S -schematically dense. We consider the G -torsor $P = V \times_{V/G} U$ over U and want to show that it is (4)-versal, that is, for a given G -torsor F over B to find $u \in U(B)$ such that $P_u \cong F$. The Chinese Remainder theorem implies that $V(B) \neq \emptyset$. We pick $v \in V(B)$, we can define a right G -equivariant morphism f_v from $\mathrm{GL}(\mathcal{E})$ to $\mathbf{V}(\mathcal{E})$ by

$$f_v(\sigma) = v \circ \sigma,$$

where σ is an element of $\mathrm{GL}(\mathcal{E})(C)$ and C is a B -algebra. We have then a commutative diagram of G -torsors

$$\begin{array}{ccc} \mathrm{GL}(\mathcal{E})_B & \xrightarrow{f_v} & V_B \\ q \downarrow & & q' \downarrow \\ \mathrm{GL}(\mathcal{E})_B/G_B & \xrightarrow{\bar{f}_v} & (V/G)_B. \end{array}$$

The point is that $\bar{f}_v^{-1}(U_B)$ is retrocompact and schematically B -dense in $X_B = \mathrm{GL}(\mathcal{E})_B/G_B$ according to the preliminaries of 4.1.

According to (1) applied to B , the G -torsor $\mathrm{GL}(\mathcal{E}) \rightarrow X$ is a (4)-versal torsor. There exists $x \in \bar{f}_v^{-1}(U_B)$ such that $q^{-1}(x) \cong F$. We put $u = \bar{f}_v(x)$ and conclude that $P_u = q'^{-1}(u) \cong F$.

Thus $V \rightarrow V/G$ is also a (4)-versal torsor. □

In this paper we shall use only the next special case.

Lemma 4.4. *Let G be a finite locally free S -scheme. Let \mathcal{E} be a locally free G - R -module of finite rank. Let $U \subset \mathbf{V}(\mathcal{E})$ be a G -stable retrocompact S -dense open subscheme such that G acts freely on U . Then U/G is representable by an S -scheme X and the quotient map $U \rightarrow X$ is a versal (4)-torsor.*

Proof. According to [8, III.2.6.1, Corollaire], U/G is representable by an S -scheme X and $U \rightarrow X$ is a G -torsor. Lemma 4.3 shows that $U \rightarrow X$ is a versal (4)-torsor. □

Remark 4.5. In this setting, it is not clear whether (4)-versal torsors always exist.

4.3. Embeddings in quasi-split groups. In this section, we assume that S is a semilocal scheme which is a finite field or whose residue fields (of maximal ideals) are infinite. Let G be a quasi-split reductive S -group scheme.

Let (B, T) be a Killing couple of G . We identify the Weyl group $W(T)$ with $N_G(T)/T$ and denote by $\mathbf{Dyn}(G)$ the Dynkin scheme (which is finite étale over S). We have a canonical isomorphism $\mathbf{Dyn}(\Phi(G, T)) \xrightarrow{\sim} \mathbf{Dyn}(G)$.

According to [24, XXIV.3.13], if G is simply connected, we have an isomorphism $T \xrightarrow{\sim} R_{\mathbf{Dyn}(G)/S}(\mathbb{G}_{m, \mathbf{Dyn}(G)})$. We denote by T^{reg} the open subscheme of regular elements of T . Then $W(T)$ stabilizes T^{reg} and acts freely on it. According to [8, III.2.6.1, Corollaire], the fppf sheaf $T^{reg}/W(T)$ is representable and the map $\pi : T^{reg} \rightarrow T^{reg}/W(T)$ is a $W(T)$ -torsor.

Denote by $G//G$ the adjoint quotient. We use now Chevalley's theorem [16] in this setting $u : T/W(T) \xrightarrow{\sim} G//G$. Away of type A_{2n} , we can deal with the Steinberg cross section $c : G//G \rightarrow G$ of the quotient map $G \rightarrow G//G$ [16, th. 5.7]; in that case we can associate to an element $x \in (T^{reg}/W(T))(S)$ the semisimple regular element $g = c(u(x)) \in G(S)$ giving rise to the maximal S -subtorus $T_x = C_G(g)$ of G . The following generalizes to the semilocal setting a result of Gille-Raghunathan [12, 19].

Theorem 4.6. *Let S be a semilocal scheme which is a finite field or whose residue fields (at maximal ideals) are infinite.*

Let G be a quasi-split reductive S -group scheme. Let Ψ be a root data which is admissible for G and let $v \in \underline{\text{Isomext}}(G, \Psi)(S)$ be an orientation. Then $\mathfrak{E}(G, \Psi, v)(S) \neq \emptyset$.

Proof. The finite field is obvious since all homogeneous G -spaces have a k -rational point [25, III.2.4]. We assume that $S = \text{Spec}(R)$ where R is a semilocal ring whose residue fields (of maximal ideals) $\kappa_1, \dots, \kappa_c$ are infinite.

Without loss of generality, we can assume that S is connected and G is semisimple simply connected. Furthermore using the decomposition [24, XXIV.5.5] and the usual Weil restriction argument, we can assume that the geometric fibers of G are almost simple. Let (B, T) be a Killing couple of G . We denote by $W(T) = N_G(T)/T$ the Weyl group and by $\mathbf{Dyn}(G)$ the Dynkin scheme (which is finite étale).

Let c be the orientation coming from the natural inclusion $T \hookrightarrow G$, and denote by $u \in \underline{\text{Isomext}}(\Phi(G, T), \Psi)(S)$ the image of (c, v) under the morphism in Prop. 3.2.1. We consider the left W -torsor $J = \underline{\text{Isomint}}_u(\Phi(G, T), \Psi)$. According to Lemma 3.2, we have $\mathfrak{E}(G, \Psi, v) = J \wedge^W \mathfrak{E}(G, \Phi(G, T), c)$.

By Lemma 4.4, $\pi : T^{reg} \rightarrow T^{reg}/W$ is a (4)-versal (left) $W(T)$ -torsor. In particular there exists $x \in (T^{reg}/W)(S)$ such that $\pi^{-1}(x) \cong J$ (as left $W(T)$ -torsors).

Case 1. G is without A_{2n} -components. By Steinberg cross section, there is a semisimple regular element $t \in G(S)$ mapped to x . The centralizer of t is a maximal torus T' of G over S . Let $c' \in \underline{\text{Isomext}}(G, \Phi(G, T'))(S)$ be the canonical orientation induced by the natural inclusion from T' to G . Denote by $u' \in \underline{\text{Isomext}}(\Phi(G, T), \Phi(G, T'))(S)$ be the image of (c, c') under the pairing in Prop. 3.2.1. By Prop. 3.2.2 the left $W(T)$ -torsors $\pi^{-1}(x)$ and $\underline{\text{Isomint}}_{u'}(\Phi(G, T), \Phi(G, T'))$ are isomorphic. It follows that $J \wedge^W \mathfrak{E}(G, \Phi(G, T), c) = \mathfrak{E}(G, \Psi, v)$ is isomorphic to $\pi^{-1}(x) \wedge^W \mathfrak{E}(G, \Phi(G, T), c) \cong \mathfrak{E}(G, \Phi(G, T'), c')$. As $\mathfrak{E}(G, \Phi(G, T'), c')(S) \neq \emptyset$, we conclude that $\mathfrak{E}(G, \Psi, v)(S)$ is not empty.

Case 2. G is of A_{2n} -type. First note that for type A , we have a Cartan involution defined over S , hence if we we have an embedding with respect to one orientation, we have an embedding with respect to the other orientation. Hence it suffices to show that $\mathfrak{E}(G, \Psi) \neq \emptyset$. The key point is to reduce to the case that G is of type A_{2n+1} .

Let R/S be an étale quadratic algebra which splits G . We can assume that G is the unitary group $U(V, h)$ for some non-degenerate hermitian form (V, h) over R/S , where V is a free R -module V of rank $2n + 1$ and h is of Witt index n . (See [15, Cor. 1.7].)

Let \mathfrak{S}_i be the symmetric group of the set $\{1, \dots, i\}$. We have a natural injection from \mathfrak{S}_{2n+1} to \mathfrak{S}_{2n+2} and hence an injective homomorphism ι from $\mathfrak{S}_{2n+1} \times \mathbb{Z}/2\mathbb{Z}$ to $\mathfrak{S}_{2n+1} \times \mathbb{Z}/2\mathbb{Z}$ which maps $\mathbb{Z}/2\mathbb{Z}$ isomorphically to itself. The map ι induces a map $\iota_* : H_{\text{ét}}^1(S, \mathfrak{S}_{2n+1} \times \mathbb{Z}/2\mathbb{Z}) \rightarrow H_{\text{ét}}^1(S, \mathfrak{S}_{2n+2} \times \mathbb{Z}/2\mathbb{Z})$. Note that $\mathfrak{S}_{2n+1} \times \mathbb{Z}/2\mathbb{Z}$ is isomorphic

to the Weyl group of the split root datum of GL_{2n} . Hence $H_{\acute{e}t}^1(S, \mathfrak{S}_{2n+1} \times \mathbb{Z}/2\mathbb{Z})$ classifies those root data which are étale locally isomorphic to the split root datum of $\mathrm{GL}_{2n+1, S}$.

Thus $\Psi = (\mathcal{M}, \mathcal{M}^\vee, \mathcal{R}, \mathcal{R}^\vee)$ corresponds to some $[\alpha] \in H_{\acute{e}t}^1(S, \mathfrak{S}_{2n+1} \times \mathbb{Z}/2\mathbb{Z})$. We twist the split root datum of $\mathrm{GL}_{2n+2, S}$ by $\iota_*([\alpha])$ and denote it by $\Psi' = (\mathcal{M}', \mathcal{M}'^\vee, \mathcal{R}', \mathcal{R}'^\vee)$. Let T and T' be the tori determined by \mathcal{M} and \mathcal{M}' respectively. Note that by our construction \mathcal{M}'^\vee can be written as $\mathcal{M}^\vee \oplus \mathcal{E}^\vee$ with \mathcal{E} is locally free of rank 1 in such a way that R^\vee is injectively sent to a subset of R'^\vee . Fix such an isomorphism $\mathcal{M}'^\vee \simeq \mathcal{M}^\vee \oplus \mathcal{E}^\vee$. This in turns gives an isomorphism $T' \simeq T \times \mathbf{R}_{R/S}^{(1)}(\mathbf{G}_m)$. We denote by T_0 the subtorus $1 \times \mathbf{R}_{R/S}^{(1)}(\mathbf{G}_m) \subseteq T'$ (under the above isomorphism).

Write the Witt decomposition of V as $\langle a \rangle \perp \mathbb{H}_n$, where \mathbb{H}_n is the maximal hyperbolic Hermitian subspace of V . Let $(V', h') = (V, h) \perp \langle -a \rangle$. Let $G' = U(V', h')$. By our assumption, there is an orientation between Ψ and G . Hence Ψ and G are both become inner forms after base change to R . By our construction of Ψ' and G' , they are also both of inner form after base change to R . Hence there is an orientation v' between Ψ' and G' . Since G' is quasi-split of type A_{2n+1} , by Case 1 we have $\mathfrak{E}(G', \Psi', v')(S) \neq \emptyset$.

Choose $f \in \mathfrak{E}(G', \Psi', v')(S)$. Consider the subgroup $f(T_0)$ in G' . Let C be the centralizer $\underline{\mathrm{Centr}}_{G'}(f(T_0))$, which is a Levi subgroup of G' . Clearly $f(T') \subseteq C$. Let V_0 be the maximal submodule of V fixed by $f(T_0)$. Namely, V_0 is the maximal submodule on which $f(T_0)$ acts trivially. Note that as $f(T_0)$ is a torus, V is étale locally decomposed into eigenspace of $f(T_0)$. Thus the rank of V_0 is $2n + 1$, which can be checked étale locally. As h' is a hyperbolic Hermitian form on V' , V_0 must contains a totally isotropic submodule of rank n . Let h_0 be the restriction of h' on V_0 . As (V_0, h_0) and (V, h) are R -module of the same rank and with maximal Witt index, we have $U(V_0, h_0) \simeq U(V, h) = G$.

Clearly C stabilizes V' and hence induces an action of $C/f(T_0)$ on (V', h') . This gives a group homomorphism of $C/f(T_0)$ to $U(V_0, h_0)$, which is an isomorphism at each geometric point \bar{s} of S . Therefore $C/f(T_0) \simeq U(V_0, h_0)$ and f gives an embedding of $T \simeq T'/T_0$ in G with respect to Ψ . \square

Corollary 4.7. *Let S be a semilocal scheme which is the spectrum of a finite field or the spectrum of a semilocal rings whose residue fields at the maximal ideals are infinite. Let G be a quasi-split reductive S -group scheme. Let X be an S -scheme satisfying one of the following conditions:*

- (i) S is noetherian;
- (ii) X is locally of finite type over S .

We assume that X is a left homogeneous under G and that for each point $s \in S$, the stabilizers of $X_{\bar{s}}$ are maximal tori of $G_{\bar{s}}$. Then $X \cong G/T$ where T is a maximal S -torus of G .

Proof. It follows immediately from Theorem 3.4 and Theorem 4.6. \square

4.4. An application to the local-global principle for embeddings. In this section we assume that k is a global field with characteristic not 2. Let K be a field, A be a central simple algebra over K with involution τ , and E be an étale algebra over K with involution σ . Suppose that $\tau|_K = \sigma|_K$ and that k is the field of invariants K^τ .

Let $\dim_L A = n^2$. Assume that $\dim_K E = n$, and that if $K \neq k$, then $\dim_k E^\sigma = n$. If $K = k$, then we assume that $\dim_k E^\sigma = \lfloor \frac{n+1}{2} \rfloor$.

The local-global principle for embeddings of K -algebras with involution (E, σ) into (A, τ) is first considered in [18]. One can associate a root datum Ψ to (E, σ) and a reductive group $U(A, \tau)^\circ$ to (A, τ) , where $U(A, \tau)^\circ$ is the connected component of the unitary group $U(A, \tau)$. (See [15] §1.3.2.) In [15] §1.3.2 and [2] §1.2 and 1.3, it is shown that there is a bijection between the embeddings of algebras with involutions over k and the embeddings of corresponding root data into reductive groups.

Combining with Theorem 4.6, we have the following theorem.

Theorem 4.4.1. *Let (E, σ) and (A, τ) be as above. If $U(A, \tau)^\circ$ is quasi-split, then the local-global principle holds for the K -algebra embeddings of (E, σ) into (A, τ) .*

Proof. Let Ψ be the root datum associated to (E, σ) ([15] §1.3.2). Let $G = U(A, \tau)^\circ$. By Theorem 4.6, it suffices to show that $\text{Isomext}(G, \Psi)(k) \neq \emptyset$.

First we consider the case where $K \neq k$. In this case, we can always fix an orientation by [15] Theorem 1.15 (2) and Remark 1.16.

Next we consider the case where $K = k$. If τ is symplectic involution or τ is orthogonal with n odd, then G is of type $C_{n/2}$ or $B_{(n+1)/2}$. Then there is always an orientation between Ψ and G .

Consider the case that τ is orthogonal with n even. Suppose there is an embedding η_v of $(E \times_k k_v, \sigma \otimes \text{id}_{k_v})$ into $(A \times_k k_v, \sigma \otimes \text{id}_{k_v})$ for all places v of k . Then η_v gives an isomorphism between the discriminant $\Delta(E)$ of E and the center $Z(A, \tau)$ of the Clifford algebra of (A, τ) over k_v (see [3] §2.3). As they are both quadratic étale algebras over k , this means that $\Delta(E)$ is isomorphic to $Z(A, \tau)$ over k . By [2] Proposition 1.3.1, an isomorphism between $\Delta(E)$ and $Z(A, \tau)$ gives an orientation between Ψ and G .

Hence for $K = k$, we can always fix an orientation between Ψ and G . \square

REFERENCES

- [1] S. Anantharaman, *Schémas en groupes, espaces homogènes et espaces algébriques sur une base de dimension 1*, Mémoires de la Société Mathématique de France **33** (1973), 5-79.
- [2] E. Bayer-Fluckiger, T.-Y. Lee, and R. Parimala, *Embedding functor for classical groups and Brauer–Manin obstruction*, Pac. J. Math. **279** (2015), 87-100.
- [3] E. Bayer-Fluckiger, T.-Y. Lee, and R. Parimala, *Embeddings of maximal tori in classical groups and explicit Brauer–Manin obstruction*, J. Eur. Math. Soc. **20** (2018), 137-163.
- [4] A. Borel, *Linear algebraic groups*, Graduate Texts in Mathematics **126** (2nd ed.), Berlin, New York: Springer-Verlag.

- [5] S. Bosch, W. Lütkebohmert, M. Raynaud, *Néron models*, Ergebnisse der Mathematik und ihrer Grenzgebiete **21** (1990), Springer.
- [6] F. Bruhat, J. Tits, *Groupes réductifs sur un corps local : II. Schémas en groupes. Existence d'une donnée radicielle valuée*, Publications Mathématiques de l'IHÉS **60** (1984), 5-184.
- [7] B. Calmès, J. Fasel, *Groupes classiques*, Autour des schémas en groupes, vol II, Panoramas et Synthèses **46** (2015), 1-133.
- [8] M. Demazure et P. Gabriel, *Groupes algébriques*, Masson (1970).
- [9] S. Garibaldi, A. Merkurjev et J.-P. Serre, *Cohomological invariants in Galois cohomology*, University Lecture Series **28** (2003), American Mathematical Society.
- [10] A. Grothendieck, J.-A. Dieudonné, *Eléments de géométrie algébrique. I*, Grundlehren der Mathematischen Wissenschaften 166; Springer-Verlag, Berlin, 1971.
- [11] A. Grothendieck (avec la collaboration de J. Dieudonné), *Eléments de Géométrie Algébrique IV*, Publications mathématiques de l'I.H.É.S. no 20, 24, 28 and 32 (1964 - 1967).
- [12] P. Gille, *Type des tores maximaux des groupes semi-simples*, J. Ramanujan Math. Soc. **19** (2004), 213-230.
- [13] P. Gille, *Sur la classification des schémas en groupes semi-simples*, "Autour des schémas en groupes, III", Panoramas et Synthèses **47** (2015), 39-110.
- [14] A. Grothendieck, *Le groupe de Brauer. I. Algèbres d'Azumaya et interprétations diverses*, Dix exposés sur la cohomologie des schémas, 46-66, Adv. Stud. Pure Math., 3, North-Holland, Amsterdam, 1968.
- [15] T.-Y. Lee, *Embedding functors and their arithmetic properties*, Comment. Math. Helv. **89** (2014), 671-717.
- [16] T.-Y. Lee, *Adjoint quotients of reductive groups*, Panoramas et Synthèses **47** (2015), 131-145.
- [17] B. Margaux, *Vanishing of Hochschild cohomology for affine group schemes and rigidity of homomorphisms between algebraic groups*, Doc. Math. **14** (2009), 653-672.
- [O] M. Olsson, *Algebraic spaces and stacks*, American Mathematical Society Colloquium Publications, 62. American Mathematical Society, Providence, RI, 2016.
- [18] G. Prasad and A. S. Rapinchuk, *Local-Global principles for embedding of fields with involution into simple algebras with involution*, Comment. Math. Helv. **85** (2010), 583-645.
- [19] M. S. Raghunathan, *Tori in quasi-split-groups*, J. Ramanujan Math. Soc. **19** (2004), 281-287.
- [20] M. Raynaud, *Faisceaux amples sur les schémas en groupes et les espaces homogènes*, Lecture Notes in Mathematics **119** (1970), Springer-Verlag, Berlin-New York.
- [21] Z. Reichstein, D. Tossici, *Special groups, versality and the Grothendieck-Serre conjecture*, preprint (2019), arXiv:1912.08109
- [22] D. Rydh, *Submersions and effective descent of étale morphism*, Bull. Soc. math. France **138**(2010) 181-230.
- [23] D. Rydh, *Approximation of Sheaves on Algebraic Stacks*, International Mathematics Research Notices (2016), 717-737.
- [24] *Séminaire de Géométrie algébrique de l'I.H.É.S., 1963-1964, Schémas en groupes, dirigé par M. Demazure et A. Grothendieck*, Lecture Notes in Math. 151-153. Springer (1970).
- [25] J.P. Serre, *Cohomologie galoisienne*, 5-ième version révisée, Lecture Notes in Mathematics **5**, Springer.
- [26] Stacks project, <https://stacks.math.columbia.edu>

P. GILLE, INSTITUT CAMILLE JORDAN - UNIVERSITÉ CLAUDE BERNARD LYON 1 43 BOULEVARD DU 11 NOVEMBRE 1918, 69622 VILLEURBANNE CEDEX - FRANCE

Email address: `gille@math.univ-lyon1.fr`

T.-Y. LEE, ASTRONOMY MATHEMATICS BUILDING 5F, NO. 1, SEC. 4, ROOSEVELT RD., TAIPEI 10617, TAIWAN.

Email address: `tingyulee@ntu.edu.tw`