

#### Comparison of metabolic and immune biomarkers between primiparous and multiparous dairy cows during the peripartum period

Angélique Corset, Anne Boudon, Sabrina Philau, Ophelie Dhumez, Aude Remot, Pierre Germon, Marion Boutinaud

#### ▶ To cite this version:

Angélique Corset, Anne Boudon, Sabrina Philau, Ophelie Dhumez, Aude Remot, et al.. Comparison of metabolic and immune biomarkers between primiparous and multiparous dairy cows during the peripartum period. ADSA annual meeting 2023, Jun 2023, Ottawa, Canada. , Journal of Dairy Science, 106 (supplement 1), pp.231-232, 2023, Abstracts of the 2023 American Dairy Science Association® Annual Meeting. hal-04172030

HAL Id: hal-04172030

https://hal.science/hal-04172030

Submitted on 27 Jul 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





Égalité

Fraternité







#### Comparison of metabolic and biomarkers between immune primiparous and multiparous dairy cows during the peripartum period

A. Corset<sup>1,2</sup>, A. Boudon<sup>1</sup>, S. Philau<sup>1</sup>, O. Dhumez<sup>1</sup>, A. Remot<sup>3</sup>, P. Germon<sup>3</sup>, M. Boutinaud<sup>1</sup>

1 INRAE-Institut Agro Rennes Angers, UMR 1348 PEGASE, Saint-Gilles, France;


- 2 Biodevas laboratoires, Savigné-l'Évêque, France;
- 3 INRAE-Université de Tours, UMR 1282 ISP, Nouzilly, France.

### Introduction

- The peripartum period, before calving and the first month of lactation, is a critical period for dairy cows
- The onset of lactation: Increased milk production and decreased immune capacity due to inflammation
- Objective of this study: Investigate the influence of parity during this period of risk on metabolic and immune biomarkers (chemokines, pro-inflammatory and anti-inflammatory cytokines ) in the dairy cows


# Material & Methods

### Sampling kinetics




Blood incubation without or with Echerischia coli in monovette ® for 24 hours at 38°C

Fluorescence Median Intensities cytokines (MFI) of 15 plasma measured by Luminex® technology 1



- 4 chemokines or chemoattractant cytokines: CCL2, CCL3, CCL4, CXCL10
- 8 pro-inflammatory cytokines: IFNγ, IL17A, IL1α, IL1b, IL2, IL6, IL8, TNFα
- 3 anti-inflammatory cytokines: IL10, IL1Rα, IL4



Statistical

Type III ANOVA mixed linear model with the parity, weeks, their interaction, and cow as a random effect

† p < 0.1 \* p < 0.05 P value: NS p > 0.1 \*\* D < 0.01


## Results

8 multiparous

7 primiparous

### Metabolic biomarkers

As described in the literature 2, at the beginning of lactation multiparous dairy cows produce more milk and their metabolism is more sollicitated.



NEFA = Non Esterified Fatty Acid BHB =  $\beta$ -hydroxybutyrate

## Plasma cytokines


#### Without *E. Coli* stimulation:


- chemoattractant & proinflammatory cytokines primiparous > multiparous
- anti-inflammatory cytokines =
- total average of the 15 cytokines tend to have primiparous > multiparous
- differences independent week of lactation



#### With *E. Coli* stimulation:

- chemoattractant & proinflammatory cytokines primiparous > multiparous
- anti-inflammatory cytokines = (not showed)
- average the cytokines tend primiparous > multiparous in week -3 and 8 (not showed)





# Conclusion

Our study shows that compared with primiparous, multiparous cows have:

- a greater metabolic solicitation

- less excessive inflammation

These results suggests that multiparous cows may have acquired a better regulation of inflammation by decreasing the pro-inflammatory cytokine production compared with primiparous cows.

**References** 1. Lesueur, J. et al., 2022 Standardized Whole Blood Assay and Bead-Based Cytokine Profiling Reveal Commonalities and Diversity of the Response to Bacteria and TLR Ligands in Cattle. Front immunol.

2. Wathes, D.C. et al., 2007 Differences between primiparous and multiparous dairy cows in the inter-relationships between metabolic traits, milk yield and body condition score in the periparturient period. Domest Anim Endocrinol

angelique.corset@inrae.fr