



**HAL**  
open science

## **A marine protected area network does not confer community structure resilience to a marine heatwave across coastal ecosystems**

Joshua Smith, Christopher Free, Cori Lopazanski, Julien Brun, Clarissa Anderson, Mark Carr, Joachim Claudet, Jenifer Dugan, Jacob Eurich, Tessa Francis, et al.

### ► To cite this version:

Joshua Smith, Christopher Free, Cori Lopazanski, Julien Brun, Clarissa Anderson, et al.. A marine protected area network does not confer community structure resilience to a marine heatwave across coastal ecosystems. *Global Change Biology*, In press, 10.1111/gcb.16862 . hal-04173111

**HAL Id: hal-04173111**

**<https://hal.science/hal-04173111>**

Submitted on 28 Jul 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 A marine protected area network does not confer  
2 community structure resilience to a marine heatwave  
3 across coastal ecosystems

4  
5 Joshua G. Smith<sup>1,2\*</sup>, Christopher M. Free<sup>3,4</sup>, Cori Lopazanski<sup>1,3</sup>, Julien Brun<sup>1</sup>, Clarissa R.  
6 Anderson<sup>5</sup>, Mark H. Carr<sup>6</sup>, Joachim Claudet<sup>7</sup>, Jenifer E. Dugan<sup>4</sup>, Jacob G. Eurich<sup>8,1</sup>, Tessa B.  
7 Francis<sup>9</sup>, Scott L. Hamilton<sup>10</sup>, David Mouillot<sup>11,12</sup>, Peter T. Raimondi<sup>6</sup>, Richard M. Starr<sup>10</sup>, Shelby  
8 L. Ziegler<sup>13</sup>, Kerry J. Nickols<sup>14,15</sup>, Jennifer E. Caselle<sup>4</sup>

9 <sup>1</sup> National Center for Ecological Analysis and Synthesis, University of California, Santa Barbara, Santa  
10 Barbara, CA, USA

11 <sup>2</sup> Conservation and Science Division, Monterey Bay Aquarium, CA, USA

12 <sup>3</sup> Bren School of Environmental Science and Management, University of California, Santa Barbara, Santa  
13 Barbara, CA, USA

14 <sup>4</sup> Marine Science Institute, University of California, Santa Barbara, Santa Barbara, CA, USA

15 <sup>5</sup> Scripps Institution of Oceanography/Southern California Coastal Ocean Observing System, University of  
16 California, San Diego, Scripps Institution of Oceanography, La Jolla, CA 92037, USA

17 <sup>6</sup> Department of Ecology and Evolutionary Biology, University of California Santa Cruz, Santa Cruz, CA,  
18 USA

19 <sup>7</sup> National Center for Scientific Research, PSL Université Paris, CRIOBE, CNRS-EPHE-UPVD, Maison de  
20 l'Océan, Paris, France

21 <sup>8</sup> Environmental Defense Fund, Santa Barbara, CA, USA

22 <sup>9</sup> Puget Sound Institute, University of Washington, Tacoma, WA, USA

23 <sup>10</sup> Moss Landing Marine Laboratories, San Jose State University, Moss Landing, CA, USA

24 <sup>11</sup> MARBEC, University of Montpellier, CNRS, IFREMER, IRD, Montpellier, France

25 <sup>12</sup> Institut Universitaire de France, IUF, Paris, France

26 <sup>13</sup> Odum School of Ecology, University of Georgia, Athens, GA, USA

27 <sup>14</sup> Department of Biology, California State University Northridge, Northridge, CA, USA

28 <sup>15</sup> Current affiliation: Ocean Visions, Leesburg, VA, USA

29

30 \* **Corresponding author:** National Center for Ecological Analysis and Synthesis, University of California,  
31 Santa Barbara, Santa Barbara, CA, USA; [joshsmith@nceas.ucsb.edu](mailto:joshsmith@nceas.ucsb.edu)

32 **Running Title:** MPAs do not confer climate resilience

33

34

35

36

## 37 Abstract

38 Marine protected areas (MPAs) have gained attention as a conservation tool for enhancing  
39 ecosystem resilience to climate change. However, empirical evidence explicitly linking MPAs to  
40 enhanced ecological resilience is limited and mixed. To better understand whether MPAs can  
41 buffer climate impacts, we tested the resistance and recovery of marine communities to the  
42 2014-2016 Northeast Pacific heatwave in the largest scientifically designed MPA network in the  
43 world off the coast of California, USA. The network consists of 124 MPAs (48 no-take, 76  
44 partial-take) implemented at different times, with full implementation completed in 2014. We  
45 compared fish, benthic invertebrate, and macroalgal community structure inside and outside of  
46 seven no-take MPAs across rocky intertidal, kelp forest, shallow reef, and deep reef nearshore  
47 habitats in California's Central Coast region from 2007-2020. We also explored whether MPA  
48 features, including age, size, depth, proportion rock, historic fishing pressure, habitat diversity  
49 and richness, connectivity, and fish biomass response ratios (proxy for ecological performance),  
50 conferred climate resilience for kelp forest and rocky intertidal habitats spanning 28 MPAs  
51 across the full network. Ecological communities dramatically shifted due to the marine heatwave  
52 across all four nearshore habitats, and MPAs did not facilitate habitat-wide resistance or  
53 recovery. Only in protected rocky intertidal habitats did community structure significantly resist  
54 marine heatwave impacts. Community shifts were associated with a pronounced decline in the  
55 relative proportion of cold-water species and an increase in warm-water species. MPA features  
56 did not explain resistance or recovery to the marine heatwave. Collectively, our findings suggest  
57 that MPAs have limited ability to mitigate the impacts of marine heatwaves on community  
58 structure. Given that mechanisms of resilience to climate perturbations are complex, there is a  
59 clear need to expand assessments of ecosystem-wide consequences resulting from acute  
60 climate-driven perturbations, and the potential role of regulatory protection in mitigating  
61 community structure changes.

62 **Keywords:** California, marine protected area networks, marine heatwaves, community  
63 composition, community structure, resilience, climate change

## 64 1. Introduction

65 Climate change can rapidly reshape the distribution of species and the composition of  
66 ecological communities (Smale et al., 2019; Pörtner et al., 2023), imperiling nature's  
67 contributions to people. In particular, episodic periods of anomalous ocean warming, hereafter  
68 'marine heatwaves', are driving pronounced shifts in species distributions across marine  
69 ecosystems (Azzurro & D'Amen, 2022; Olsen et al., 2022), with direct implications for ecological  
70 processes and associated human benefits (Smale et al., 2019; Cheung et al., 2021; Payne et  
71 al., 2021; Cinner et al., 2022; Smith et al., 2023). While the urgency to plan for adaptation to  
72 climate change is clear, as marine heatwaves increase in frequency and severity (Holbrook et  
73 al., 2019), pathways to enhance ecosystem resilience are mixed. Therefore, understanding how  
74 ecological communities resist, recover from, or are transformed by climate perturbations, such  
75 as marine heatwaves, represents one of the most pressing challenges for building ecosystem  
76 resilience capacity (Mason et al., 2022).

77 Marine protected areas (MPAs) are an important conservation strategy for preserving  
78 biodiversity and nature's contributions to people (Gronrud-Colvert et al., 2021). MPAs may  
79 provide network-scale population connectivity that can enhance spillover of individuals and the  
80 replenishment of populations in both protected and fished areas (Harrison et al., 2012;  
81 Williamson et al., 2016; Baetscher et al., 2019; Di Lorenzo et al., 2020; Goetze et al., 2021).  
82 Although most MPAs were initially designed to reduce the effects of overfishing and habitat loss,  
83 they are frequently hypothesized to provide long-term protection against climate impacts  
84 (Roberts et al., 2017; Hofmann et al., 2021). For example, networks of MPAs may provide  
85 refugia as species redistribute in response to climate change, owing to lower anthropogenic  
86 stressors and higher population sizes resulting from reduced harvest (McLeod et al., 2009; Carr  
87 et al., 2017). MPA features such as habitat diversity, historic fishing pressure, age, and size  
88 may also influence the capacity for MPAs to provide ecological resilience (Jacquemont et al.,  
89 2022). As such, MPAs and MPA networks are being increasingly highlighted as a key tool for  
90 enhancing climate resilience (IUCN-WCPA, 2008; Jacquemont et al., 2022).

91 Despite the growing number of studies examining MPAs as a tool for mitigating climate  
92 impacts, the effectiveness of MPAs (both individually and as networks) for enhancing the  
93 resilience (i.e., resistance to and recovery from disturbance, Bates et al., 2019) of marine  
94 communities to climate change remains contested (Johnson et al., 2022, but see Jacquemont et  
95 al. 2022). Climate change stressors (e.g., ocean acidification, sea level rise, hypoxia, warming)

96 may have effects on populations and communities that occur regardless of regulatory protection  
97 (Bates et al., 2019; Bruno et al., 2019). Marine heatwaves can also reduce connectivity by  
98 changing prevailing current patterns – a key design attribute for many MPA networks (Lima et  
99 al., 2021). Mixed evidence surrounding the efficacy of MPAs in providing climate resilience may  
100 be explained in part by the single-habitat (e.g., coral reef, seagrass, kelp forest, etc.) focus of  
101 many studies, and also in part by the ways in which assemblages of species are partitioned.

102 Our understanding of whether and how MPAs confer ecological resilience to climate  
103 change can be improved by synthesizing the effects of regulatory protection across multiple  
104 taxa, habitats, ecosystems, and protection levels. Potential mechanisms producing ecological  
105 shifts in response to climate change may include altered mortality due to physiological  
106 environmental tolerances, changes in species interactions (e.g., competition, predation,  
107 disease, facilitation), adult movement across habitats and along the coast, ontogenetic shifts,  
108 and changes in recruitment success (Harley et al., 2006). Moreover, the ways in which species  
109 responses are evaluated (e.g., trait-based, functional groups, feeding guilds, evolutionary  
110 lineages, etc.) can influence detection of climate-driven outcomes. Therefore, long-term  
111 monitoring across multiple habitats subjected to similar (or the same) perturbations is needed to  
112 thoroughly examine the impacts of climate change on marine ecosystems, to test whether (and  
113 which) MPAs confer climate resilience, and to assess the relative resilience among ecosystems  
114 and taxa. Such cross-ecosystem studies are rare, especially those that include monitoring  
115 across networks of MPAs before, during, and after extreme climate change-driven  
116 perturbations.

117 In 1999, California passed the Marine Life Protection Act (MLPA), which expanded its  
118 system of MPAs to function as a coherent ecological network and to address six goals aimed at  
119 conservation, fisheries, and other human benefits (Marine Life Protection Act, 1999; Gleason et  
120 al., 2013). Guided by these goals, California established a network of 124 MPAs (48 no-take, 76  
121 partial-take) distributed along the state's entire 1,300-kilometer coastline that protects 16% of  
122 state waters. The network protects hard- and soft-bottom habitats ranging in depth from the  
123 intertidal to depths of 1,000 m. However, with few exceptions, ecological monitoring studies  
124 focused on hard-bottom habitats. Leading up to and following MPA implementation, an  
125 extensive ecological monitoring effort of these habitats began to support adaptive management  
126 of the network (Botsford et al., 2014). During the course of California's MPA monitoring, a major  
127 marine heatwave occurred that was the consequence of two environmental anomalies: a 2014-  
128 2015 warming event known as "the Blob," and a major El Niño event in 2015-2016 (Bond et al.,

129 2015; Di Lorenzo & Mantua, 2016; Gentemann et al., 2017). This pronounced climate  
130 perturbation was the largest marine heatwave on record in California (Laufkötter et al., 2020).

131 This study leveraged over a decade of monitoring across the rocky intertidal, kelp forest,  
132 shallow reef, and deep reef habitats to evaluate whether a network of MPAs confers community  
133 structure resilience to climate disturbances. Specifically, we tested the following hypotheses: (1)  
134 the 2014-16 marine heatwave event resulted in the reorganization of community structure  
135 across multiple nearshore habitats and taxa, (2) MPAs enhanced the ability for ecological  
136 communities to resist and recover from the impacts of climate perturbations, and (3) species  
137 traits (thermal affinities) explained differential responses to the marine heatwave in relation to  
138 regulatory protection (MPAs). After testing these hypotheses, we explored whether MPA  
139 features (size, age, historic fishing pressure, habitat diversity, and connectivity) enhanced  
140 ecological community resilience across habitats to inform the design of MPA networks that are  
141 resilient to climate change.

## 142 2. Methods

### 143 2.1. MPA sampling and data collection

144 As part of the State of California's long-term MPA evaluation and monitoring program,  
145 several habitat-specific research groups conduct annual surveys designed to monitor ecological  
146 changes over time inside MPAs and at areas of comparable habitat outside MPAs (**Figure 1A;**  
147 **Figure S1**, Supplementary Appendix *Figures*). Conceptually, while MPAs could be considered  
148 the "treatment" or "control" in our analyses (i.e., fishing is considered either the experiment or  
149 the control), we take a social-ecological system perspective and consider human activities,  
150 including fishing, to be integral system components (Kandel et al., 2022). Thus, in our analytical  
151 framework, MPAs are considered the experimental treatment where fishing is removed (e.g.,  
152 Pacoureaux et al., 2023).

153 Our analyses focused on four habitats that have extensive long-term monitoring data:  
154 rocky intertidal, kelp forest, shallow reef, and deep reef (**Figure 1A; Figure S1**, Supplementary  
155 Appendix *Figures*). Across the four habitats included in our analyses, three organismal groups  
156 were sampled: macroalgae (rocky intertidal and kelp forest), conspicuous mobile and sessile  
157 invertebrates (rocky intertidal and kelp forest), and fishes (kelp forest, shallow reef, and deep  
158 reef). We focused our community structure analyses on MPAs located along the temperate

159 Central Coast of California (**Figure 1A**) because this area was the most comprehensively  
160 sampled region of the MPA network and had sufficient pre-marine heatwave data to evaluate  
161 baseline community structure (**Figure S1**, Supplementary Appendix *Figures*). To evaluate MPA  
162 features as potential drivers of ecological resilience, we included MPAs across all of California  
163 for two of the habitats (rocky intertidal and kelp forest), where more extensive spatial and  
164 temporal coverage exists (see section 2.6).

165 Two general types of data were used in our analyses: species counts (kelp forest,  
166 shallow reef, deep reef), and the proportional cover of invertebrates and macroalgae (rocky  
167 intertidal and kelp forest invertebrates and algae). While biomass is often used as a common  
168 measurable response across taxa (Duffy et al., 2017), it is not suitable for our analyses because  
169 of the large number of taxa for which it is difficult to accurately measure biomass (e.g.,  
170 macroalgae and invertebrates). Therefore, we elected to use a taxonomic abundance-based  
171 approach to compare changes in community structure. All analyses were conducted using  
172 published data for each habitat (Brooks et al., 2022; Cieri et al., 2022; Malone et al., 2022;  
173 MARINe et al., 2022).

## 174 2.2. Changes in community structure associated with the marine heatwave

175 We used non-metric multidimensional scaling (nMDS) to visualize community structure  
176 before (2007-13), during (2014-16), and after (2017-20) the 2014-16 marine heatwave. Prior to  
177 ordination, a base similarity matrix was constructed for each habitat type using Bray-Curtis  
178 dissimilarity on Hellinger-transformed counts or on percent cover of species. The Hellinger  
179 transformation converts absolute counts to the square root of proportional counts, which  
180 reduces the disproportionate contribution of highly abundant species (Legendre & Gallagher,  
181 2001). We elected to use the transformation to relative species abundance to improve the  
182 robustness, comparability, and ecological interpretation of our community structure analyses.  
183 Each site sampled inside or outside of a MPA surveyed in a single year represents a single unit  
184 of replication. To visualize the state of each ecological community, we plotted centroids and  
185 95% confidence ellipses that represent the generalized position of the community in ordinated  
186 2-dimensional nMDS space before, during, and after the marine heatwave event.

187 We used a pairwise permutational analysis of variance (PERMANOVA) to test three  
188 hypotheses surrounding community structure changes (**Figure 2A**). First, if MPAs do not  
189 mitigate marine heatwave impacts on community structure, then communities inside MPAs

190 should respond similarly (i.e., comparable change in multivariate distance) to those outside.  
191 Second, if MPAs confer resistance, then communities inside MPAs should remain unchanged  
192 while those outside shift. Finally, if MPAs enhance recovery, then communities inside MPAs  
193 should shift during the marine heatwave, but return to or move back in the direction of their  
194 previous state following the disturbance. To test these hypotheses, a single PERMANOVA was  
195 performed for each habitat and site type (inside or outside a MPA) using pairwise comparisons  
196 between marine heatwave periods (before vs. during, before vs. after) using the *pairwiseAdonis*  
197 wrapper function (Martinez Arbizu, 2019) in the *vegan* package (Oksanen et al., 2022) in R (R  
198 Core Team, 2021).

### 199 2.3. Effect of MPAs on community structure resistance and recovery

200 We used a multivariate distance-based approach to test whether MPAs conferred  
201 ecological community resistance to or recovery from the marine heatwave. We refer to this  
202 measurement as ‘multivariate distance’ to distinguish it from ‘geographic distance,’ which is  
203 often a measure of climate change impacts on marine species distributions. We define  
204 resistance as community structure (i.e., relative abundance of species) that remained  
205 unchanged during the marine heatwave (low change in multivariate distance-based centroids),  
206 and recovery as a community that returned to a similar structure post-heatwave. Resistance  
207 was evaluated by calculating the multivariate vector distance (in high dimensional space using  
208 the Bray-Curtis dissimilarity matrix) of the centroid of the ecological community between the  
209 periods before (2007-2013) and during (2014-2016) the marine heatwave (e.g., high resistance  
210 is indicated by a smaller change in multivariate distance-based centroids).

211 Recovery was evaluated by calculating the multivariate distance between the before  
212 (2007-2013) and after period (2017-2020) centroids. We calculated the change in multivariate  
213 distance between centroids using the *betadisper* function in the *vegan* package (Oksanen et al.,  
214 2022). The *betadisper* function returns the principal coordinates of centroids, which we used to  
215 calculate multivariate distance after ensuring positive-definite eigenvalues. Finally, statistical  
216 significance was tested using a PERMANOVA on community structure resistance (before vs.  
217 during) and recovery (before vs. after).

218 To explore whether the timing of community shifts coincided with temporal changes in  
219 oceanographic variables, we used a Granger causality test on Bray-Curtis dissimilarity. For this  
220 analysis, annual dissimilarity was calculated for each site (inside or outside a MPA) and habitat



221 relative to 2007. Dissimilarity was then offset (lagged) against oceanographic variables (see  
222 section 2.5) in one-year increments for a maximum of three years (maximum lag based on  
223 length of the time series). We used 2007 as the baseline year because it preceded the marine  
224 heatwave and because we were interested in lag effects specifically related to the marine  
225 heatwave, rather than gradual environmental changes over time. However, 2008 was used as  
226 the baseline year for the deep reef habitat because surveys were not extensively conducted in  
227 2007. It is important to note that the Granger test only examines lagged effects between the  
228 time series of community change and the environmental variables, it does not examine  
229 correlations, which are explored using other diagnostics described in section 2.5.

## 230 2.4. Species associated with community changes

231 To evaluate which species were associated with community change, we used two  
232 approaches. First, we used a Similarity Percentage Analysis (SIMPER) to decompose  
233 community structure using Bray-Curtis dissimilarity and to examine the percent contribution of  
234 each species to the before versus after marine heatwave communities. However, because  
235 SIMPER is known to confound the mean between group variation and dispersion (Warton et al.,  
236 2012), we cross-validated the SIMPER output using the *mvabund* package (Wang et al., 2022)  
237 in R. The *mvabund* package uses fitted generalized linear models to account for nonlinear  
238 mean-to-variance relationships of each species. Both of these analyses were conducted on the  
239 absolute abundance of species (rather than relative abundance) because they are not sensitive  
240 to common or rare taxa. This combination of approaches allowed us to determine the  
241 contribution of individual taxa to observed community structure differences. Additionally,  
242 because the PERMANOVA analyses revealed that community change was similar inside and  
243 outside of MPAs, we analyzed compositional changes overall rather than within each site by  
244 protection status. Finally, we constructed an analysis of deviance table using the multivariate  
245 generalized linear model fits to test whether species structure changed as a result of the marine  
246 heatwave (before vs. after). Test statistics and p-values for each species were generated using  
247 the PIT-trap (probability integral transform) resampling method.

## 248 2.5. Environmental correlates and species traits

249 We explored temperature and oceanographic conditions before, during, and after the  
250 marine heatwave to evaluate whether community structure shifts were explained by  
251 environmental changes (**Figure 1B-F**, **Figure S2** Supplementary Appendix *Figures*). For these

252 environmental analyses, we used air temperature (AT, °C) measured *in situ* at rocky intertidal  
253 long-term monitoring sites (as an environmental correlate for rocky intertidal communities only,  
254 Supplementary Appendix *Methods*); sea surface temperature (SST, °C) at 1 km daily resolution  
255 from MURSST (Chin et al., 2017), sea bottom temperature (SBT, °C) at 8 km daily resolution  
256 from GLORYS (Jean-Michel et al. 2021), the Biologically Effective Upwelling Transport Index  
257 (BEUTI,  $\text{mmol m}^{-1} \text{s}^{-1}$ ; Jacox et al., 2018) calculated at 1 degree latitude bins, and the  
258 Multivariate Oceanographic Climate Index (MOCI), which is a long-term (30 year) indicator of  
259 several oceanographic and atmospheric conditions (García-Reyes & Sydeman, 2017)  
260 calculated at the regional level (Central California). We selected these environmental indicators  
261 and associated products because they provide biologically meaningful climatology at the best  
262 available spatial resolution for our study area.

263 To process the environmental data, we first calculated the monthly mean AT, SST, SBT,  
264 BEUTI, and quarterly MOCI values at each site. We then calculated monthly anomalies for AT,  
265 SST, SBT, and BEUTI as the difference between the observed monthly mean and the baseline  
266 average (long-term average for each month, 2000-2012 for AT, 1988-2012 for BEUTI, 1993-  
267 2012 for SBT, and 2002-2012 for SST; start year of the historical climatology reflects the first  
268 year with data). Baseline averages were calculated through the end of 2012 to accommodate a  
269 buffer before the onset of the marine heatwave in 2014. For MOCI, we calculated the annual  
270 mean (2000-2020) at each site as a standard index. To visualize and pair the environmental  
271 data with the long-term biological monitoring data, we calculated the mean anomalies across all  
272 calendar months for each year (**Figure 1B-F**). Finally, to determine whether observed shifts in  
273 community structure were associated with changes in oceanographic conditions, we used the  
274 *envfit* function in the *vegan* package (Oksanen et al., 2022) to overlay the four environmental  
275 variables as vectors on the ordinated community data. The *envfit* function computes the rank-  
276 based correlation coefficient and significance is evaluated through permutation-based  
277 resampling.

278 We used a trait-based fourth-corner model to evaluate effects of environmental variables  
279 and species thermal affinities on the relative abundance of all recorded taxa. Species thermal  
280 affinities were obtained from literature reported classifications and expert judgment  
281 (Supplementary Appendix, *Methods*). The fourth-corner model relates the interaction between  
282 the environment and species traits (thermal affinities) on variation in the abundance of taxa  
283 (Brown et al., 2014). Briefly, we used the *mvabund* package to fit simultaneous generalized  
284 linear models for counts of all species at each site as a function of the environmental and

285 oceanographic conditions (AT, SST, SBT, BEUTI, MOCI), the relative representation of thermal  
286 affinities, and their interaction.

## 287 2.6. Drivers of resistance and recovery

288 The ecological and design context of California's network is well documented, allowing  
289 us to explore whether certain MPA features conferred community structure resilience to the  
290 marine heatwave. These analyses were spatially expanded to include a subset of MPAs  
291 covering the entire network from northern to southern California, thereby providing a greater  
292 scope of MPA feature diversity (**Figure S3**, Supplementary Appendix *Figures*). However, only  
293 the kelp forest and rocky intertidal habitats contained sufficient pre-heatwave monitoring data  
294 with consistent annual surveys in all bioregions of the state to appropriately examine drivers of  
295 ecological resilience among MPAs at the network level (**Figure 1A**; **Figure S1**, Supplementary  
296 Appendix *Figures*). MPA features included habitat richness and habitat diversity (calculated  
297 using Shannon-Wiener on depth-stratified hard and soft bottom, proportion of rock, and the  
298 extent of kelp forest canopy, rocky intertidal, sandy beach, coastal marsh, tidal flats, and  
299 armored shore), historic fishing pressure, MPA age and size, connectivity, and kelp forest fish  
300 biomass response ratios (as a proxy for MPA ecological performance, see Supplementary  
301 Appendix *Methods*).

302 We developed a two-stage multivariate model to examine the effect of MPA features on  
303 community resistance and recovery. First, we used a PERMANOVA to identify MPAs that  
304 significantly resisted or recovered from the marine heatwave (building on the methods outlined  
305 in section 2.3). We then used logistic regression to evaluate the probability that specific MPA  
306 features enhanced resistance or resilience. For the logistic regression, any MPA community that  
307 significantly resisted or recovered from the marine heatwave was assigned a target level of '1',  
308 and any MPA community that did not resist or recover from the marine heatwave a '0.'

## 309 3. Results

### 310 3.1. Community structure resistance and recovery

311 Ecological community change was widespread across all habitats, and three habitats  
312 (kelp forest, shallow reef, deep reef) showed no clear differences inside and outside of MPAs in  
313 the magnitude of community change (**Figure 2B**). Prior to the marine heatwave, community

314 structure was similar inside and outside of MPAs in all habitats except the rocky intertidal  
315 (indicated by the non-overlapping ellipses in **Figure 2B**). However, ecological community  
316 structure dramatically shifted as a result of the marine heatwave event across all measured  
317 nearshore habitats (n=4), regardless of whether the communities (n=5) were inside or outside of  
318 MPAs (**Figures 2B and S8, Table 1**).

319 Community structure changes coincided with oceanographic conditions associated with  
320 the marine heatwave (**Figure 1, Tables 1 and 2**). At the onset of the marine heatwave in 2014,  
321 sea surface temperature was anomalously warm by as much as 2°C, and sea bottom  
322 temperature was also 1°C above the baseline average. MOCI and BEUTI experienced  
323 precipitous changes during the marine heatwave, reflecting reduced upwelling and productivity,  
324 and these anomalies persisted until at least mid-2016 (**Figure 1 E-F**). Although each  
325 oceanographic anomaly aligned well with community shifts in nMDS space (**Figure 2B**),  
326 significant correlations were habitat-specific (**Table 2**). In the rocky intertidal, warmer air  
327 temperature was the strongest environmental correlate with community structure changes,  
328 although warmer SST and declines in upwelling (BEUTI) were also correlated with community  
329 change. For shallow reef, kelp forest fishes, and kelp forest invertebrates and algae, SST, SBT,  
330 and MOCI were significantly associated with community change, such that communities shifted  
331 with increases in surface and bottom water temperature and positive values of the MOCI index.  
332 In the deep reef habitat (fishes only), the only significant oceanographic correlate was BEUTI  
333 (**Table 2**), where the community shifted in response to reduced upwelling. Finally, there were no  
334 significant temporal lags determined except for one interaction between the shallow reef and  
335 BEUTI (*lag* = 3 years for sites inside and outside of MPAs, **Table S4**).

336 For all habitats except the rocky intertidal, MPAs did not impart increased resistance to  
337 or recovery from marine heatwave-driven community changes compared to sites outside of  
338 MPAs (**Figure 3**). The rocky intertidal and deep reef were the only two habitats that resisted the  
339 marine heatwave (based on non-significant PERMANOVA result, **Figure 3A**). The rocky  
340 intertidal was the only habitat where community structure inside MPAs was not significantly  
341 different either during or post-marine heatwave (**Figure 3B**). Kelp forest invertebrates, algae,  
342 and fishes all substantially shifted during the marine heatwave, and these changes persisted for  
343 several years later with no apparent recovery. The shallow reef habitat responded similarly with  
344 pronounced shifts in community structure during the marine heatwave, although the trajectory of  
345 the shallow reef habitat started to move towards the pre-heatwave state beginning in the year  
346 2018, with the biggest shifts towards recovery occurring inside MPAs (**Figures S5 and S6**,

347 Supplementary Appendix *Figures*). In the deep reef habitat, year-to-year changes were more  
348 variable, resulting in a larger multivariate distance relative to the other habitats, but this  
349 community change was not significantly different from the baseline year (**Figures 3A** and **S9**  
350 Supplementary Appendix *Figures*). However, recovery was more variable than resistance. The  
351 shallow reef habitat exhibited relatively greater recovery (i.e., less shift in multivariate distance  
352 compared to the pre-heatwave community state) than the kelp forest and deep reef habitats.

### 353 3.2. Species responses

354 The multivariate analyses revealed several species that explained differences between  
355 the pre- and post-heatwave periods. Among the three habitat types with monitoring of fish  
356 species (kelp forest, shallow reef, deep reef), the blue and deacon rockfish complex (*Sebastes*  
357 *mystinus* and *S. diaconus*) was positively correlated with the post-heatwave period (**Table S5**,  
358 Supplementary Appendix, *Tables*). Fish species that declined and were found within the top  
359 80% contribution of pre- vs. post-heatwave community structure included: *Sebastes serranoides*  
360 (kelp forest), *S. chrysomelas* and *carinatus* (kelp forest), *S. miniatus* (kelp forest, shallow reef),  
361 *Brachyistius frenatus* (kelp forest), *S. semicinctus* (deep reef), and *S. hopkinsi* (deep reef). For  
362 invertebrates and algae, the multivariate analyses revealed an increase in the abundance of  
363 purple sea urchins (*Strongylocentrotus purpuratus*) and a decline in macroalgae in both the  
364 rocky intertidal and kelp forest habitats.

### 365 3.3. Community structure and thermal traits

366 The relative proportional representation of thermal affinities significantly changed during  
367 the marine heatwave for kelp forest invertebrates and algae, kelp forest fishes, and shallow reef  
368 fishes ( $p < 0.001$ ,  $F_{6,708} = 62.24$ ;  $p < 0.05$ ,  $F_{6,700} = 2.18$ ;  $p < 0.001$ ,  $F_{4,315} = 37.41$ , respectively).  
369 Cold-temperate species significantly declined during the marine heatwave for these habitats  
370 (**Figure 4A**). During the marine heatwave, there was a slight increase in cosmopolitan species  
371 for the rocky intertidal and kelp forest (invertebrates and algae), and an even more pronounced  
372 increase in warm-temperate and subtropical fish species in the kelp forest, shallow reef, and  
373 deep reef habitats. Importantly, these changes in community composition that occurred during  
374 the marine heatwave (2014-2016) persisted into the following years (2017 and beyond), which  
375 partially explains the lack of observed recovery to the pre-heatwave community structure.

376 Variation in species abundance was explained by significant interactions between  
377 thermal affinities and oceanographic variables for all habitats from the fourth corner model  
378 (**Figure 4B**, **Figure S7** Supplementary Appendix, *Figures*). In general, species thermal affinities  
379 responded similarly across habitats. The abundance of cold-temperate species declined with  
380 increased sea surface temperature anomalies and reduced upwelling (BEUTI), while the  
381 abundance of warm-temperate, subtropical, and tropical species increased with the  
382 oceanographic conditions that were associated with the onset of the marine heatwave, such as  
383 MOCI and warming water temperature. Sea bottom temperature had the most variable  
384 interactive effect across habitats and thermal affinities. Interestingly, thermal associations for the  
385 deep reef habitat had an opposite sign for cold-temperate and warm-temperate interactions,  
386 relative to the other habitats.

### 387 3.4. MPA features and ecological stability

388 Community structure responses were highly variable across the state-level network for  
389 MPAs. In general, rocky intertidal communities showed higher resistance than kelp forest  
390 communities, as indicated by smaller shifts in multivariate distance (**Figure 5C**; smaller points).  
391 While shifts in community structure were generally larger in the kelp forest (**Figure 5C**; larger  
392 points), in some cases, the magnitude of shifts inside MPAs was less relative to shifts outside of  
393 MPAs (**Figure 5A**). In general, MPA protection fostered more resistance than recovery (**Figure**  
394 **5C**; i.e., more blue points in the resistance column and more red points in the recovery column),  
395 although these results were not statistically significant. For example, kelp forest communities in  
396 the Point Buchon, South Point, and Harris Point MPAs resisted and moved towards recovery  
397 from the heatwave more than communities outside these MPAs (**Figure 5C**; blue points).  
398 However, kelp forest communities in the Scorpion, Anacapa Island, and Campus Point MPAs  
399 were less successful at resisting shifts than communities outside these MPAs (**Figure 5C-left**;  
400 red points). Similarly, kelp forest communities in the Point Lobos, Santa Barbara, Farnsworth,  
401 and Blue Cavern MPAs were less successful at recovering from shifts than communities outside  
402 these MPAs (**Figure 5C-right**; red points). None of the evaluated MPA features (age, size,  
403 historic fishing pressure, habitat diversity, connectivity, biomass response ratios) were  
404 statistically significant drivers of this variability (**Figure S11**, Supplementary Appendix *Figures*).

## 405 4. Discussion

406 Our findings suggest that the 2014-16 Northeast Pacific marine heatwave impacted  
407 ecological community structure across four nearshore habitats, and that MPAs did not confer  
408 widespread resistance or recovery. Marine protected areas have gained increased attention as  
409 a conservation strategy for mitigating the effects of climate change (Wilson et al., 2018; Duncan  
410 et al., 2019), but evidence of their efficacy in providing ecosystem-wide resilience to climate  
411 disturbances remains mixed (Roberts et al., 2017; Bates et al., 2019; Jacquemont et al., 2022).  
412 Critically, international efforts to conserve 30% of marine habitats via MPA implementation by  
413 the year 2030 (IUCN, 2014) highlight the need for planning that considers the effect of large  
414 climate-driven perturbations on local ecological processes. Therefore, while MPAs may be  
415 implemented with specific conservation targets, our study suggests that extreme climate  
416 perturbations, such as marine heatwaves, can overwhelm intended climate benefits of MPAs  
417 over the short-term (< 5 years following a marine heatwave). These results highlight the  
418 ecosystem-wide consequences of acute climate-driven perturbations despite regulatory  
419 protection.

420 Several ecological processes may explain the pronounced shifts observed across  
421 habitats, such as altered adult movement, changes to recruitment regimes, tropicalization,  
422 deborealization, and shifts in species interactions such as herbivory, competition, predation, and  
423 disease (Vergés et al., 2014; García Molinos et al., 2016; Wernberg et al., 2016; Free et al.,  
424 2019; Mignot et al., 2022). First, our study found that the relative proportion of warm-temperate  
425 and subtropical species increased during and after the marine heatwave. The increased  
426 proportional representation of species with warm-water thermal affinities may be explained by  
427 adult movement into the study area, changes in recruitment patterns, or by a decline in the  
428 relative abundance of cold-temperate species (Sanford et al., 2019; Walker et al., 2020;  
429 Fredston et al., 2021). For example, the large increase in blue rockfish in kelp forest, shallow  
430 reef, and deep reef habitats was associated with strong recruitment and pelagic young-of-the-  
431 year abundance of rockfishes in midwater trawl surveys at the start of the marine heatwave  
432 period (Field et al. 2021). Large increases in the subtropical wrasse, seniorita (*Oxyjulis*  
433 *californica*), were also observed in multiple habitats, contributing to community change. Second,  
434 during the marine heatwave, a large-scale outbreak of herbivorous sea urchins occurred  
435 throughout the study region that coincided with precipitous declines in kelp (and other  
436 macroalgae) and the loss of an important benthic mesopredator, the sunflower sea star

437 (*Pycnopodia helianthoides*), due to a large-scale marine disease outbreak (Harvell et al., 2019;  
438 Smith et al., 2021).

439 Species life history traits in relation to the marine heatwave event may explain why the  
440 rocky intertidal was the only habitat where community structure exhibited resilience. Variation in  
441 stress tolerance to air temperature is a fundamental driver of rocky intertidal community  
442 structure. Although 2014 was the warmest year on record in California, the sustained sea  
443 surface temperature anomalies associated with ocean warming during the marine heatwave  
444 may have impacted the rocky intertidal less than the subtidal habitats. Resilience in rocky  
445 intertidal MPAs may be explained by increased propagule delivery to MPAs, particularly by long-  
446 lived foundation species that stabilize the community (Raimondi & Smith, 2022), or through  
447 regulatory protection that prohibits the harvest of habitat-forming mussels (*Mytilus spp.*) in  
448 MPAs. Additionally, the rocky intertidal is the only habitat that used a fixed-plot design  
449 (*Supplementary Appendix, Methods*) and this methodological form of sampling could have  
450 resulted in smaller propagated spatial variance, which could be more biased towards  
451 conclusions of enhanced resilience relative to the sampling used in the other habitats. More  
452 research is needed to distill the mechanistic processes associated with MPA resilience in the  
453 rocky intertidal.

454 The primary regulation associated with MPAs in California involves a restriction of fishing  
455 activities; thus, we were interested in understanding whether ecological communities inside no-  
456 take MPAs are more resilient to marine heatwaves than our control social-ecological system.  
457 We consider human activities, such as fishing, as integral parts of the social-ecological system,  
458 since fishing occurred decades before MPA establishment and the marine heatwave, with the  
459 restriction of harvest through regulatory protection as the primary conservation mechanism.  
460 However, there are multiple viewpoints pertaining to interactions between humans and the  
461 oceans, and similarly there are many approaches to evaluating the effects of fishing and  
462 regulatory protection. Alternative frameworks might consider areas inside MPAs as “reference”  
463 locations for places where harvest occurs (Costello 2014). While we believe human and natural  
464 systems are now so integrated in coastal areas that there are virtually no pristine ecosystems,  
465 the framing of considering MPAs as controls and fished systems as experimental treatments  
466 would not impact our results.

467 It is not surprising that community shifts occurred inside and outside MPAs  
468 simultaneously for most habitats, since many of the species exhibiting the biggest changes are



469 not directly targeted by fishing activities (e.g., invertebrates and algae). In cases where MPAs  
470 do not confer ecological resilience to marine heatwaves, it is important to note that the capacity  
471 of MPAs to harbor higher biomass than unprotected sites can still be maintained (Frid et al.,  
472 2023). Other studies have documented increases in targeted species biomass inside of MPAs  
473 within this system, which is a focal MPA conservation objective (Hamilton et al., 2010; Caselle  
474 et al., 2015; Ziegler et al., 2022). Similar responses to the marine heatwave in southern  
475 California fish communities were reported by Freedman et al. (2020), who found that non-  
476 targeted subtropical species were most responsible for community reorganization. Ziegler et al.  
477 (2023) also reported that fish species diversity recovered more quickly after the heatwave inside  
478 of MPAs.

479         Although the anomalously warm environmental conditions subsided after 2016, changes  
480 in community structure persisted, especially in the kelp forest, shallow reef, and deep reef  
481 habitats. These results are particularly interesting in the context of ecosystem stability and  
482 transition dynamics. Given the highly variable recruitment of the dominant fishes in the study  
483 region (Schroeder et al., 2019; Field et al., 2021), it is possible that sufficient time has not  
484 elapsed for the communities to return to a pre-perturbed state. Lagged effects due to  
485 ontogenetic shifts and methodological sampling that disproportionately select for adults may  
486 also contribute to the observed lack of recovery. For example, declines in vermilion rockfish  
487 (*Sebastes miniatus*) in the shallow reef habitat coincided with increases in that species in the  
488 deep reef habitat, likely reflecting ontogenetic movements from shallow to deep habitats (Love  
489 et al., 2002; Love, 2011). In addition, the shallow reef habitat monitoring program used hook  
490 and line sampling, which disproportionately selects individuals that are older than 2-3 years,  
491 which is why lagged effects of oceanography were more likely to be detected in this habitat  
492 (Ziegler et al., 2023). Moreover, many species in the California Current are slow-growing, late to  
493 mature, long-lived, and relatively sedentary with small home ranges (Love et al., 2002; Love,  
494 2011). These life history traits may create inertia to resistance or recovery from change.  
495 However, the persistence of novel community structure configurations observed in this study,  
496 despite the return of pre-heatwave environmental conditions, highlights the need for further  
497 assessments of community transition dynamics over ecologically meaningful timescales.

498         Our finding of no effect of MPA features on dampening marine heatwave-driven  
499 responses in the rocky intertidal and kelp forest habitats is particularly interesting given that  
500 habitat mosaics, historic fishing effort, and connectivity have been linked to MPA conservation  
501 capacity (Bastari et al., 2016). The widespread effects of marine heatwaves and other climate

502 perturbations may overwhelm local stressors (e.g., fishing, marine pollution) that MPAs were  
503 initially successful at mitigating (Bruno et al., 2019). Indeed, even biodiversity rich spots or deep  
504 water communities are susceptible to the impacts of climate change (Kocsis et al., 2021;  
505 Emblemsvåg et al., 2022). Our results highlight the need for management frameworks that  
506 include climate adaptation strategies, the protection of range-shift corridors, and consideration  
507 of the compounding effects of both fishing and marine heatwaves on resilience (Burrows et al.,  
508 2014). More investigation is needed to better understand the drivers that can help MPAs to  
509 maintain higher fish biomass than unprotected sites under climate change (Frid et al., 2023).

510           Increasing our understanding of the pathways through which marine heatwaves  
511 restructure ecological communities is central to developing adaptive management solutions. In  
512 our study, we used a taxonomic (i.e., species) abundance-based approach to evaluate changes  
513 in community structure, but functional diversity may also be impacted by warming events and  
514 should be explored (McLean et al., 2019; Murgier et al., 2021). Prolonged warming events may  
515 differentially impact groups and guilds of species with similar traits, such as dispersal ability,  
516 thermal tolerance, metabolic rate, and mobility (Duncan et al., 2019; Harvey et al., 2021).  
517 Moreover, to understand the ability of MPA networks to resist and recover from future marine  
518 heatwaves, it is critical to have sufficient monitoring across multiple habitats, taxa, and regions.  
519 Because pre-perturbation data are essential for establishing baselines and comparatively  
520 evaluating ecological responses, consistent monitoring is fundamental to capture the effect of  
521 marine heatwave events. Although the timing of marine heatwaves is unpredictable, evidence  
522 suggests that the frequency and magnitude of abrupt warming events are expected to increase  
523 (Frölicher et al., 2018; Holbrook et al., 2020).

524           Evidence of MPAs as a tool for mitigating the effects of climate change remains  
525 controversial, especially since most MPAs were initially designed for other conservation  
526 priorities. Our analysis of ecological community change across a network of MPAs highlights  
527 that widespread effects of climate-driven stressors such as marine heatwaves can dramatically  
528 restructure ecological communities, regardless of regulatory protection. Ultimately, improved  
529 resilience capacity of MPA networks will require integrating adaptive management with careful  
530 consideration of how abrupt climate change-driven perturbations may inhibit intended  
531 conservation outcomes.

## 532 Acknowledgements

533 We thank the hundreds of individuals who contributed to data collection for each of the long-  
534 term monitoring programs included in this study. We also thank R. Brooks, I. Brunjes, K. Cieri,  
535 R. Gaddam, D. Malone, S. Worden, and M. Yeager for assistance with data curation. This  
536 research was funded in part by the California Ocean Protection Council (OPC, #C0752005) as  
537 part of the Decadal Review of California's Marine Protected Areas. The Ocean Protection  
538 Council also provided support for long-term habitat monitoring (#C0302700, #C0752003). Other  
539 sources of support for habitat monitoring include California Sea Grant (R/MPA-43, R/MPA-44,  
540 R/MPA-45, R/MPA-46, R/MPA-48) and the David and Lucile Packard Foundation. CMF was  
541 funded through the Arnhold UC Santa Barbara-Conservation International Climate Solutions  
542 Collaborative. JC was funded through BiodivERSA (METRODIVER and MOVE) and Fondation  
543 de France (MultiNet) projects. JED is grateful for support from the Santa Barbara Coastal Long  
544 Term Ecological Research project (National Science Foundation OCE-1831937). JEC and MHC  
545 were supported in part by the David and Lucile Packard Foundation through the Partnership for  
546 Interdisciplinary Studies of Coastal Oceans (PISCO).

## 547 Author Contributions Statement

548 JGS, CMF, CL, SH, RS, DM, KN, and JC conceived the ideas and designed methodology; JGS,  
549 CMF, CL, JB, MC, SH, PR, SZ, and JC collected the data; JGS, CMF, CL, JB, and CA analyzed  
550 the data; JGS led the writing of the manuscript. All authors contributed substantially to revisions.  
551 The authors have agreed to be listed and approve the manuscript for submission.

## 552 Conflict of Interests Statement

553 The authors have no conflicts of interest to declare.

## 554 Data Availability Statement

555 All source code and data are available on GitHub: <https://github.com/NCEAS/ca-mpa>

556 **References**

- 557 Azzurro, E., & D'Amen, M. (2022). Climate change paves the way for a new inter-ocean fish  
558 interchange. *Frontiers in Ecology and the Environment*, 20(10), 558–563.  
559 <https://doi.org/10.1002/fee.2459>
- 560 Baetscher, D. S., Anderson, E. C., Gilbert-Horvath, E. A., Malone, D. P., Saarman, E. T., Carr,  
561 M. H., & Garza, J. C. (2019). Dispersal of a nearshore marine fish connects marine  
562 reserves and adjacent fished areas along an open coast. *Molecular Ecology*, 28(7),  
563 1611–1623. <https://doi.org/10.1111/mec.15044>
- 564 Bastari, A., Micheli, F., Ferretti, F., Pusceddu, A., & Cerrano, C. (2016). Large marine protected  
565 areas (LMPAs) in the Mediterranean Sea: The opportunity of the Adriatic Sea. *Marine*  
566 *Policy*, 68, 165–177. <https://doi.org/10.1016/j.marpol.2016.03.010>
- 567 Bates, A. E., Cooke, R. S. C., Duncan, M. I., Edgar, G. J., Bruno, J. F., Benedetti-Cecchi, L.,  
568 Côté, I. M., Lefcheck, J. S., Costello, M. J., Barrett, N., Bird, T. J., Fenberg, P. B., &  
569 Stuart-Smith, R. D. (2019). Climate resilience in marine protected areas and the  
570 'Protection Paradox.' *Biological Conservation*, 236, 305–314.  
571 <https://doi.org/10.1016/j.biocon.2019.05.005>
- 572 Bond, N. A., Cronin, M. F., Freeland, H., & Mantua, N. (2015). Causes and impacts of the 2014  
573 warm anomaly in the NE Pacific. *Geophysical Research Letters*, 42(9), 3414–3420.  
574 <https://doi.org/10.1002/2015GL063306>
- 575 Botsford, L. W., White, J. W., Carr, M. H., & Caselle, J. E. (2014). Chapter Six—Marine  
576 Protected Area Networks in California, USA. In M. L. Johnson & J. Sandell (Eds.),  
577 *Advances in Marine Biology* (Vol. 69, pp. 205–251). Academic Press.  
578 <https://doi.org/10.1016/B978-0-12-800214-8.00006-2>
- 579 Brooks, R., Ziegler, S., Starr, R., Wendt, D., Ruttenberg, B., Bellquist, L., Caselle, J., Morgan,  
580 S., Mulligan, T., Semmens, B., Tyburczy, J., & Hamilton, S. (2022). *Nearshore Fishes*

- 581 *Abundance and Distribution Data, California Collaborative Fisheries Research Program*  
582 *(CCFRP)*. [https://search.dataone.org/view/urn%3Auuid%3Af843f110-e691-4d26-bf12-](https://search.dataone.org/view/urn%3Auuid%3Af843f110-e691-4d26-bf12-3854a4b641cd)  
583 [3854a4b641cd](https://search.dataone.org/view/urn%3Auuid%3Af843f110-e691-4d26-bf12-3854a4b641cd)
- 584 Brown, A. M., Warton, D. I., Andrew, N. R., Binns, M., Cassis, G., & Gibb, H. (2014). The fourth-  
585 corner solution – using predictive models to understand how species traits interact with  
586 the environment. *Methods in Ecology and Evolution*, *5*(4), 344–352.  
587 <https://doi.org/10.1111/2041-210X.12163>
- 588 Bruno, J. F., Côté, I. M., & Toth, L. T. (2019). Climate Change, Coral Loss, and the Curious  
589 Case of the Parrotfish Paradigm: Why Don't Marine Protected Areas Improve Reef  
590 Resilience? *Annual Review of Marine Science*, *11*(1), 307–334.  
591 <https://doi.org/10.1146/annurev-marine-010318-095300>
- 592 Burrows, M. T., Schoeman, D. S., Richardson, A. J., Molinos, J. G., Hoffmann, A., Buckley, L.  
593 B., Moore, P. J., Brown, C. J., Bruno, J. F., Duarte, C. M., Halpern, B. S., Hoegh-  
594 Guldborg, O., Kappel, C. V., Kiessling, W., O'Connor, M. I., Pandolfi, J. M., Parmesan,  
595 C., Sydeman, W. J., Ferrier, S., ... Poloczanska, E. S. (2014). Geographical limits to  
596 species-range shifts are suggested by climate velocity. *Nature*, *507*(7493), Article 7493.  
597 <https://doi.org/10.1038/nature12976>
- 598 Carr, M. H., Robinson, S. P., Wahle, C., Davis, G., Kroll, S., Murray, S., Schumacker, E. J., &  
599 Williams, M. (2017). The central importance of ecological spatial connectivity to effective  
600 coastal marine protected areas and to meeting the challenges of climate change in the  
601 marine environment. *Aquatic Conservation: Marine and Freshwater Ecosystems*, *27*(S1),  
602 6–29. <https://doi.org/10.1002/aqc.2800>
- 603 Caselle, J. E., Rassweiler, A., Hamilton, S. L., & Warner, R. R. (2015). Recovery trajectories of  
604 kelp forest animals are rapid yet spatially variable across a network of temperate marine  
605 protected areas. *Scientific Reports*, *5*(1), Article 1. <https://doi.org/10.1038/srep14102>
- 606 Cheung, W. W. L., Frölicher, T. L., Lam, V. W. Y., Oyinlola, M. A., Reygondeau, G., Sumaila, U.

- 607 R., Tai, T. C., Teh, L. C. L., & Wabnitz, C. C. C. (2021). Marine high temperature  
608 extremes amplify the impacts of climate change on fish and fisheries. *Science*  
609 *Advances*, 7(40), eabh0895. <https://doi.org/10.1126/sciadv.abh0895>
- 610 Chin, T. M., Vazquez-Cuervo, J., & Armstrong, E. M. (2017). A multi-scale high-resolution  
611 analysis of global sea surface temperature. *Remote Sensing of Environment*, 200, 154–  
612 169. <https://doi.org/10.1016/j.rse.2017.07.029>
- 613 Cieri, K., Starr, R., Caselle, J., Kahn, A., Lauermann, A., Lindholm, J., Tissot, B., Ziegler, S.,  
614 Bretz, C., Carlson, P., Hoeke, J., Jainese, C., Martel, G., McDermott, S., Mohay, J., &  
615 Salinas-Ruiz, P. (2022). *Monitoring and Evaluation of Mid-Depth Rock Ecosystems in*  
616 *the California MLPA Marine Protected Area Network: ROV Dataset*.  
617 <https://opc.dataone.org/view/urn%3Auuid%3A7353f779-b722-4064-8074-3e9c651ed38e>
- 618 Cinner, J. E., Caldwell, I. R., Thiault, L., Ben, J., Blanchard, J. L., Coll, M., Diedrich, A., Eddy, T.  
619 D., Everett, J. D., Folberth, C., Gascuel, D., Guiet, J., Gurney, G. G., Heneghan, R. F.,  
620 Jägermeyr, J., Jiddawi, N., Lahari, R., Kuange, J., Liu, W., ... Pollnac, R. (2022).  
621 Potential impacts of climate change on agriculture and fisheries production in 72 tropical  
622 coastal communities. *Nature Communications*, 13(1), Article 1.  
623 <https://doi.org/10.1038/s41467-022-30991-4>
- 624 Marine Life Protection Act, Chapter 10.5, California State Legislature, Fish and Game Code  
625 (FGC) (1999).  
626 [https://leginfo.legislature.ca.gov/faces/codes\\_displayText.xhtml?lawCode=FGC&division](https://leginfo.legislature.ca.gov/faces/codes_displayText.xhtml?lawCode=FGC&division=3.&title=&part=&chapter=10.5.&article=)  
627 [=3.&title=&part=&chapter=10.5.&article=](https://leginfo.legislature.ca.gov/faces/codes_displayText.xhtml?lawCode=FGC&division=3.&title=&part=&chapter=10.5.&article=)
- 628 Di Lorenzo, E., & Mantua, N. (2016). Multi-year persistence of the 2014/15 North Pacific marine  
629 heatwave. *Nature Climate Change*, 6(11), Article 11.  
630 <https://doi.org/10.1038/nclimate3082>
- 631 Di Lorenzo, M., Guidetti, P., Di Franco, A., Calò, A., & Claudet, J. (2020). Assessing spillover  
632 from marine protected areas and its drivers: A meta-analytical approach. *Fish and*

- 633 *Fisheries*, 21(5), 906–915. <https://doi.org/10.1111/faf.12469>
- 634 Duffy, J. E., Godwin, C. M., & Cardinale, B. J. (2017). Biodiversity effects in the wild are  
635 common and as strong as key drivers of productivity. *Nature*, 549(7671), Article 7671.  
636 <https://doi.org/10.1038/nature23886>
- 637 Duncan, M. I., Bates, A. E., James, N. C., & Potts, W. M. (2019). Exploitation may influence the  
638 climate resilience of fish populations through removing high performance metabolic  
639 phenotypes. *Scientific Reports*, 9(1), Article 1. [https://doi.org/10.1038/s41598-019-](https://doi.org/10.1038/s41598-019-47395-y)  
640 [47395-y](https://doi.org/10.1038/s41598-019-47395-y)
- 641 Emblemsvåg, M., Werner, K. M., Núñez-Riboni, I., Frelat, R., Torp Christensen, H., Fock, H. O.,  
642 & Primicerio, R. (2022). Deep demersal fish communities respond rapidly to warming in  
643 a frontal region between Arctic and Atlantic waters. *Global Change Biology*, 28(9), 2979–  
644 2990. <https://doi.org/10.1111/gcb.16113>
- 645 Field, J. C., Miller, R. R., Santora, J. A., Tolimieri, N., Haltuch, M. A., Brodeur, R. D., Auth, T. D.,  
646 Dick, E. J., Monk, M. H., Sakuma, K. M., & Wells, B. K. (2021). Spatiotemporal patterns  
647 of variability in the abundance and distribution of winter-spawned pelagic juvenile  
648 rockfish in the California Current. *PLOS ONE*, 16(5), e0251638.  
649 <https://doi.org/10.1371/journal.pone.0251638>
- 650 Fredston, A., Pinsky, M., Selden, R. L., Szuwalski, C., Thorson, J. T., Gaines, S. D., & Halpern,  
651 B. S. (2021). Range edges of North American marine species are tracking temperature  
652 over decades. *Global Change Biology*, 27(13), 3145–3156.  
653 <https://doi.org/10.1111/gcb.15614>
- 654 Free, C. M., Thorson, J. T., Pinsky, M. L., Oken, K. L., Wiedenmann, J., & Jensen, O. P. (2019).  
655 Impacts of historical warming on marine fisheries production. *Science*, 363(6430), 979–  
656 983. <https://doi.org/10.1126/science.aau1758>
- 657 Frid, O., Malamud, S., Di Franco, A., Guidetti, P., Azzurro, E., Claudet, J., Micheli, F., Yahel, R.,  
658 Sala, E., & Belmaker, J. (2023). Marine protected areas' positive effect on fish biomass

- 659 persists across the steep climatic gradient of the Mediterranean Sea. *Journal of Applied*  
660 *Ecology*, 60(4), 638–649. <https://doi.org/10.1111/1365-2664.14352>
- 661 Frölicher, T. L., Fischer, E. M., & Gruber, N. (2018). Marine heatwaves under global warming.  
662 *Nature*, 560(7718), Article 7718. <https://doi.org/10.1038/s41586-018-0383-9>
- 663 García Molinos, J., Halpern, B. S., Schoeman, D. S., Brown, C. J., Kiessling, W., Moore, P. J.,  
664 Pandolfi, J. M., Poloczanska, E. S., Richardson, A. J., & Burrows, M. T. (2016). Climate  
665 velocity and the future global redistribution of marine biodiversity. *Nature Climate*  
666 *Change*, 6(1), Article 1. <https://doi.org/10.1038/nclimate2769>
- 667 García-Reyes, M., & Sydeman, W. J. (2017). California Multivariate Ocean Climate Indicator  
668 (MOCI) and marine ecosystem dynamics. *Ecological Indicators*, 72, 521–529.  
669 <https://doi.org/10.1016/j.ecolind.2016.08.045>
- 670 Gentemann, C. L., Fewings, M. R., & García-Reyes, M. (2017). Satellite sea surface  
671 temperatures along the West Coast of the United States during the 2014–2016 northeast  
672 Pacific marine heat wave. *Geophysical Research Letters*, 44(1), 312–319.  
673 <https://doi.org/10.1002/2016GL071039>
- 674 Gleason, M., Fox, E., Ashcraft, S., Vasques, J., Whiteman, E., Serpa, P., Saarman, E.,  
675 Caldwell, M., Frimodig, A., Miller-Henson, M., Kirlin, J., Ota, B., Pope, E., Weber, M., &  
676 Wiseman, K. (2013). Designing a network of marine protected areas in California:  
677 Achievements, costs, lessons learned, and challenges ahead. *Ocean & Coastal*  
678 *Management*, 74, 90–101. <https://doi.org/10.1016/j.ocecoaman.2012.08.013>
- 679 Goetze, J. S., Wilson, S., Radford, B., Fisher, R., Langlois, T. J., Monk, J., Knott, N. A.,  
680 Malcolm, H., Currey-Randall, L. M., Ierodiaconou, D., Harasti, D., Barrett, N., Babcock,  
681 R. C., Bosch, N. E., Brock, D., Claudet, J., Clough, J., Fairclough, D. V., Heupel, M. R.,  
682 ... Harvey, E. S. (2021). Increased connectivity and depth improve the effectiveness of  
683 marine reserves. *Global Change Biology*, 27(15), 3432–3447.  
684 <https://doi.org/10.1111/gcb.15635>



- 685 Grorud-Colvert, K., Sullivan-Stack, J., Roberts, C., Constant, V., Pike, E. P., Kingston, N.,  
686 Laffoley, D., Sala, E., Claudet, J., Friedlander, A. M., Gill, D. A., Lester, S. E., Day, J. C.,  
687 Gonçalves, E. J., Ahmadi, G. N., Rand, M., Villagomez, A., Ban, N. C., Gurney, G. G.,  
688 ... Lubchenco, J. (2021). The MPA Guide: A framework to achieve global goals for the  
689 ocean. *Science*, 373, eabf0861.
- 690 Hamilton, S. L., Caselle, J. E., Malone, D. P., & Carr, M. H. (2010). Incorporating biogeography  
691 into evaluations of the Channel Islands marine reserve network. *Proceedings of the*  
692 *National Academy of Sciences*, 107(43), 18272–18277.  
693 <https://doi.org/10.1073/pnas.0908091107>
- 694 Harley, C. D. G., Randall Hughes, A., Hultgren, K. M., Miner, B. G., Sorte, C. J. B., Thornber, C.  
695 S., Rodriguez, L. F., Tomanek, L., & Williams, S. L. (2006). The impacts of climate  
696 change in coastal marine systems. *Ecology Letters*, 9(2), 228–241.  
697 <https://doi.org/10.1111/j.1461-0248.2005.00871.x>
- 698 Harrison, H. B., Williamson, D. H., Evans, R. D., Almany, G. R., Thorrold, S. R., Russ, G. R.,  
699 Feldheim, K. A., van Herwerden, L., Planes, S., Srinivasan, M., Berumen, M. L., &  
700 Jones, G. P. (2012). Larval Export from Marine Reserves and the Recruitment Benefit  
701 for Fish and Fisheries. *Current Biology*, 22(11), 1023–1028.  
702 <https://doi.org/10.1016/j.cub.2012.04.008>
- 703 Harvell, C. D., Montecino-Latorre, D., Caldwell, J. M., Burt, J. M., Bosley, K., Keller, A., Heron,  
704 S. F., Salomon, A. K., Lee, L., Pontier, O., Pattengill-Semmens, C., & Gaydos, J. K.  
705 (2019). Disease epidemic and a marine heat wave are associated with the continental-  
706 scale collapse of a pivotal predator (*Pycnopodia helianthoides*). *Science Advances*, 5(1),  
707 eaau7042. <https://doi.org/10.1126/sciadv.aau7042>
- 708 Harvey, B. P., Marshall, K. E., Harley, C. D. G., & Russell, B. D. (2021). Predicting responses to  
709 marine heatwaves using functional traits. *Trends in Ecology & Evolution*.  
710 <https://doi.org/10.1016/j.tree.2021.09.003>

- 711 Hofmann, G., Hazen, E., Ambrose, R., Aseltine-Beilson, D., Caselle, J., Chan, F., Levine, A.,  
712 Micheli, F., Sunday, J., & White, W. (2021). *Climate Resilience and California's Marine*  
713 *Protected Area Network: A Report by the Ocean Protection Council Science Advisory*  
714 *Team Working Group and California Ocean Science Trust.*
- 715 Holbrook, N. J., Scannell, H. A., Sen Gupta, A., Benthuyesen, J. A., Feng, M., Oliver, E. C. J.,  
716 Alexander, L. V., Burrows, M. T., Donat, M. G., Hobday, A. J., Moore, P. J., Perkins-  
717 Kirkpatrick, S. E., Smale, D. A., Straub, S. C., & Wernberg, T. (2019). A global  
718 assessment of marine heatwaves and their drivers. *Nature Communications*, 10(1),  
719 Article 1. <https://doi.org/10.1038/s41467-019-10206-z>
- 720 Holbrook, N. J., Sen Gupta, A., Oliver, E. C. J., Hobday, A. J., Benthuyesen, J. A., Scannell, H.  
721 A., Smale, D. A., & Wernberg, T. (2020). Keeping pace with marine heatwaves. *Nature*  
722 *Reviews Earth & Environment*. <https://doi.org/10.1038/s43017-020-0068-4>
- 723 IUCN-WCPA. (2008). *Establishing resilient marine protected area networks - making it happen:*  
724 *Full technical version, including ecological, social and governance considerations, as*  
725 *well as case studies*. IUCN World Commission on Protected Areas, National Oceanic  
726 and Atmospheric Administration, and The Nature Conservancy.  
727 <https://portals.iucn.org/library/node/9284>
- 728 Jacox, M. G., Edwards, C. A., Hazen, E. L., & Bograd, S. J. (2018). Coastal Upwelling  
729 Revisited: Ekman, Bakun, and Improved Upwelling Indices for the U.S. West Coast.  
730 *Journal of Geophysical Research: Oceans*, 123(10), 7332–7350.  
731 <https://doi.org/10.1029/2018JC014187>
- 732 Jacquemont, J., Blasiak, R., Le Cam, C., Le Gouellec, M., & Claudet, J. (2022). Ocean  
733 conservation boosts climate change mitigation and adaptation. *One Earth*, 5(10), 1126–  
734 1138. <https://doi.org/10.1016/j.oneear.2022.09.002>
- 735 Johnson, J. V., Dick, J. T. A., & Pincheira-Donoso, D. (2022). Marine protected areas do not  
736 buffer corals from bleaching under global warming. *BMC Ecology and Evolution*, 22(1),

- 737           58. <https://doi.org/10.1186/s12862-022-02011-y>
- 738 Kandel, P., Pandit, R., White, B., & Polyakov, M. (2022). Do protected areas increase  
739 household income? Evidence from a Meta-Analysis. *World Development*, 159, 106024.  
740 <https://doi.org/10.1016/j.worlddev.2022.106024>
- 741 Kocsis, Á. T., Zhao, Q., Costello, M. J., & Kiessling, W. (2021). Not all biodiversity rich spots are  
742 climate refugia. *Biogeosciences*, 18(24), 6567–6578. [https://doi.org/10.5194/bg-18-](https://doi.org/10.5194/bg-18-6567-2021)  
743 6567-2021
- 744 Laufkötter, C., Zscheischler, J., & Frölicher, T. L. (2020). High-impact marine heatwaves  
745 attributable to human-induced global warming. *Science*, 369(6511), 1621–1625.  
746 <https://doi.org/10.1126/science.aba0690>
- 747 Legendre, P., & Gallagher, E. D. (2001). Ecologically meaningful transformations for ordination  
748 of species data. *Oecologia*, 129(2), 271–280. <https://doi.org/10.1007/s004420100716>
- 749 Lima, L. S., Gherardi, D. F. M., Pezzi, L. P., Passos, L. G. dos, Endo, C. A. K., & Quimbayo, J.  
750 P. (2021). Potential changes in the connectivity of marine protected areas driven by  
751 extreme ocean warming. *Scientific Reports*, 11(1), Article 1.  
752 <https://doi.org/10.1038/s41598-021-89192-6>
- 753 Love, M. S. (2011). *Certainly More Than You Want to Know About the Fishes of the Pacific*  
754 *Coast*. Really Big Press.
- 755 Love, M. S., Yoklavich, M., & Thorsteinson, L. K. (2002). *The Rockfishes of the Northeast*  
756 *Pacific*. University of California Press.
- 757 Malone, D. P., Davis, K., Lonhart, S. I., Parsons-Field, A., Caselle, J. E., & Carr, M. H. (2022).  
758 Large-scale, multidecade monitoring data from kelp forest ecosystems in California and  
759 Oregon (USA). *Ecology*, 103(5), e3630. <https://doi.org/10.1002/ecy.3630>
- 760 MARINe, PISCO, & Raimondi, P. (2022). *MARINe/PISCO: Intertidal: MARINe Coastal*  
761 *Biodiversity Surveys: Point Contact Surveys Summarized*.  
762 [https://doi.org/10.6085/AA/marine\\_cbs.5.5](https://doi.org/10.6085/AA/marine_cbs.5.5)

- 763 Martinez Arbizu, P. (2020). pairwiseAdonis: Pairwise multilevel comparison using adonis. R  
764 package version 0.4
- 765 Mason, J. G., Eurich, J. G., Lau, J. D., Battista, W., Free, C. M., Mills, K. E., Tokunaga, K.,  
766 Zhao, L. Z., Dickey-Collas, M., Valle, M., Pecl, G. T., Cinner, J. E., McClanahan, T. R.,  
767 Allison, E. H., Friedman, W. R., Silva, C., Yáñez, E., Barbieri, M. Á., & Kleisner, K. M.  
768 (2022). Attributes of climate resilience in fisheries: From theory to practice. *Fish and*  
769 *Fisheries*, 23(3), 522–544. <https://doi.org/10.1111/faf.12630>
- 770 McLean, M., Mouillot, D., Lindegren, M., Villéger, S., Engelhard, G., Murgier, J., & Auber, A.  
771 (2019). Fish communities diverge in species but converge in traits over three decades of  
772 warming. *Global Change Biology*, 25(11), 3972–3984. <https://doi.org/10.1111/gcb.14785>
- 773 McLeod, E., Salm, R., Green, A., & Almany, J. (2009). Designing marine protected area  
774 networks to address the impacts of climate change. *Frontiers in Ecology and the*  
775 *Environment*, 7(7), 362–370. <https://doi.org/10.1890/070211>
- 776 Mignot, A., von Schuckmann, K., Landschützer, P., Gasparin, F., van Gennip, S., Perruche, C.,  
777 Lamouroux, J., & Amm, T. (2022). Decrease in air-sea CO<sub>2</sub> fluxes caused by persistent  
778 marine heatwaves. *Nature Communications*, 13(1), Article 1.  
779 <https://doi.org/10.1038/s41467-022-31983-0>
- 780 Murgier, J., McLean, M., Maire, A., Mouillot, D., Loiseau, N., Munoz, F., Violle, C., & Auber, A.  
781 (2021). Rebound in functional distinctiveness following warming and reduced fishing in  
782 the North Sea. *Proceedings of the Royal Society B: Biological Sciences*, 288(1942),  
783 20201600. <https://doi.org/10.1098/rspb.2020.1600>
- 784 Oksanen, J., Simpson, G. L., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., Friendly,  
785 M., McGlinn, D., O'Hara, R. B., Solymos, P., Stevens, M. H. H., Szoecs, E., & Wagner,  
786 H. (2022). *vegan: Community Ecology Package* (R package version 2.6-2).  
787 <https://CRAN.R-project.org/package=vegan>
- 788 Olsen, A. Y., Larson, S., Padilla-Gamiño, J. L., & Klinger, T. (2022). Changes in fish

- 789 assemblages after marine heatwave events in West Hawai'i Island. *Marine Ecology*  
790 *Progress Series*, 698, 95–109. <https://doi.org/10.3354/meps14156>
- 791 Pacoureau, N., Carlson, J. K., Kindsvater, H. K., Rigby, C. L., Winker, H., Simpfendorfer, C. A.,  
792 Charvet, P., Pollom, R. A., Barreto, R., Sherman, C. S., Talwar, B. S., Skerritt, D. J.,  
793 Sumaila, U. R., Matsushiba, J. H., VanderWright, W. J., Yan, H. F., & Dulvy, N. K.  
794 (2023). Conservation successes and challenges for wide-ranging sharks and rays.  
795 *Proceedings of the National Academy of Sciences*, 120(5), e2216891120.  
796 <https://doi.org/10.1073/pnas.2216891120>
- 797 Payne, M. R., Kudahl, M., Engelhard, G. H., Peck, M. A., & Pinnegar, J. K. (2021). Climate risk  
798 to European fisheries and coastal communities. *Proceedings of the National Academy of*  
799 *Sciences*, 118(40), e2018086118. <https://doi.org/10.1073/pnas.2018086118>
- 800 Pörtner, H.-O., Scholes, R. J., Arneth, A., Barnes, D. K. A., Burrows, M. T., Diamond, S. E.,  
801 Duarte, C. M., Kiessling, W., Leadley, P., Managi, S., McElwee, P., Midgley, G., Ngo, H.  
802 T., Obura, D., Pascual, U., Sankaran, M., Shin, Y. J., & Val, A. L. (2023). Overcoming  
803 the coupled climate and biodiversity crises and their societal impacts. *Science*,  
804 380(6642), eabl4881. <https://doi.org/10.1126/science.abl4881>
- 805 R Core Team. (2021). *R: A Language and Environment for Statistical Computing*. R Foundation  
806 for Statistical Computing. <http://www.R-project.org/>
- 807 Raimondi, P., & Smith, J. (2022). *Assessment of Rocky Intertidal habitats for the California*  
808 *Marine Protected Area Decadal report* [A report to the Ocean Protection Council and UC  
809 Sea Grant]. [https://www.opc.ca.gov/webmaster/\\_media\\_library/2022/05/MPA\\_RO1.pdf](https://www.opc.ca.gov/webmaster/_media_library/2022/05/MPA_RO1.pdf)
- 810 Roberts, C. M., O'Leary, B. C., McCauley, D. J., Cury, P. M., Duarte, C. M., Lubchenco, J.,  
811 Pauly, D., Sáenz-Arroyo, A., Sumaila, U. R., Wilson, R. W., Worm, B., & Castilla, J. C.  
812 (2017). Marine reserves can mitigate and promote adaptation to climate change.  
813 *Proceedings of the National Academy of Sciences*, 114(24), 6167–6175.  
814 <https://doi.org/10.1073/pnas.1701262114>

- 815 Sanford, E., Sones, J. L., García-Reyes, M., Goddard, J. H. R., & Largier, J. L. (2019).  
816 Widespread shifts in the coastal biota of northern California during the 2014–2016  
817 marine heatwaves. *Scientific Reports*, 9(1), Article 1. [https://doi.org/10.1038/s41598-](https://doi.org/10.1038/s41598-019-40784-3)  
818 019-40784-3
- 819 Schroeder, I. D., Santora, J. A., Bograd, S. J., Hazen, E. L., Sakuma, K. M., Moore, A. M.,  
820 Edwards, C. A., Wells, B. K., & Field, J. C. (2019). Source water variability as a driver of  
821 rockfish recruitment in the California Current Ecosystem: Implications for climate change  
822 and fisheries management. *Canadian Journal of Fisheries and Aquatic Sciences*, 76(6),  
823 950–960. <https://doi.org/10.1139/cjfas-2017-0480>
- 824 Smale, D. A., Wernberg, T., Oliver, E. C. J., Thomsen, M., Harvey, B. P., Straub, S. C.,  
825 Burrows, M. T., Alexander, L. V., Benthuisen, J. A., Donat, M. G., Feng, M., Hobday, A.  
826 J., Holbrook, N. J., Perkins-Kirkpatrick, S. E., Scannell, H. A., Sen Gupta, A., Payne, B.  
827 L., & Moore, P. J. (2019). Marine heatwaves threaten global biodiversity and the  
828 provision of ecosystem services. *Nature Climate Change*, 9(4), Article 4.  
829 <https://doi.org/10.1038/s41558-019-0412-1>
- 830 Smith, J. G., Tomoleoni, J., Staedler, M., Lyon, S., Fujii, J., & Tinker, M. T. (2021). Behavioral  
831 responses across a mosaic of ecosystem states restructure a sea otter–urchin trophic  
832 cascade. *Proceedings of the National Academy of Sciences*, 118(11), e2012493118.  
833 <https://doi.org/10.1073/pnas.2012493118>
- 834 Smith, K. E., Burrows, M. T., Hobday, A. J., King, N. G., Moore, P. J., Sen Gupta, A., Thomsen,  
835 M. S., Wernberg, T., & Smale, D. A. (2023). Biological Impacts of Marine Heatwaves.  
836 *Annual Review of Marine Science*, 15(1), 119–145. [https://doi.org/10.1146/annurev-](https://doi.org/10.1146/annurev-marine-032122-121437)  
837 marine-032122-121437
- 838 Vergés, A., Steinberg, P. D., Hay, M. E., Poore, A. G. B., Campbell, A. H., Ballesteros, E., Heck,  
839 K. L., Booth, D. J., Coleman, M. A., Feary, D. A., Figueira, W., Langlois, T., Marzinelli, E.  
840 M., Mizerek, T., Mumby, P. J., Nakamura, Y., Roughan, M., van Sebille, E., Gupta, A. S.,

- 841 ... Wilson, S. K. (2014). The tropicalization of temperate marine ecosystems: Climate-  
842 mediated changes in herbivory and community phase shifts. *Proceedings of the Royal*  
843 *Society B: Biological Sciences*, 281(1789), 20140846.  
844 <https://doi.org/10.1098/rspb.2014.0846>
- 845 Walker, H. J., Hastings, P. A., Hyde, J. R., Lea, R. N., Snodgrass, O. E., & Bellquist, L. F.  
846 (2020). Unusual occurrences of fishes in the Southern California Current System during  
847 the warm water period of 2014–2018. *Estuarine, Coastal and Shelf Science*, 236,  
848 106634. <https://doi.org/10.1016/j.ecss.2020.106634>
- 849 Wang, Y., Naumann, U., Eddelbuettel, D., Wilshire, J., & Warton, D. (2022). *mvabund*:  
850 *Statistical Methods for Analysing Multivariate Abundance Data* (R package version  
851 4.2.1.). <https://CRAN.R-project.org/package=mvabund>
- 852 Warton, D. I., Wright, S. T., & Wang, Y. (2012). Distance-based multivariate analyses confound  
853 location and dispersion effects. *Methods in Ecology and Evolution*, 3(1), 89–101.  
854 <https://doi.org/10.1111/j.2041-210X.2011.00127.x>
- 855 Wernberg, T., Bennett, S., Babcock, R. C., de Bettignies, T., Cure, K., Depczynski, M., Dufois,  
856 F., Fromont, J., Fulton, C. J., Hovey, R. K., Harvey, E. S., Holmes, T. H., Kendrick, G.  
857 A., Radford, B., Santana-Garcon, J., Saunders, B. J., Smale, D. A., Thomsen, M. S.,  
858 Tuckett, C. A., ... Wilson, S. (2016). Climate-driven regime shift of a temperate marine  
859 ecosystem. *Science*, 353(6295), 169–172. <https://doi.org/10.1126/science.aad8745>
- 860 Williamson, D. H., Harrison, H. B., Almany, G. R., Berumen, M. L., Bode, M., Bonin, M. C.,  
861 Choukroun, S., Doherty, P. J., Frisch, A. J., Saenz-Agudelo, P., & Jones, G. P. (2016).  
862 Large-scale, multidirectional larval connectivity among coral reef fish populations in the  
863 Great Barrier Reef Marine Park. *Molecular Ecology*, 25(24), 6039–6054.  
864 <https://doi.org/10.1111/mec.13908>
- 865 Wilson, J. R., Lomonico, S., Bradley, D., Sievanen, L., Dempsey, T., Bell, M., McAfee, S.,  
866 Costello, C., Szuwalski, C., McGonigal, H., Fitzgerald, S., & Gleason, M. (2018).

867 Adaptive comanagement to achieve climate-ready fisheries. *Conservation Letters*, 11(6),  
868 e12452. <https://doi.org/10.1111/conl.12452>

869 Ziegler, S. L., Brooks, R. O., Hamilton, S. L., Ruttenberg, B. I., Chiu, J. A., Fields, R. T., Waltz,  
870 G. T., Shen, C., Wendt, D. E., & Starr, R. M. (2022). External fishing effort regulates  
871 positive effects of no-take marine protected areas. *Biological Conservation*, 269,  
872 109546. <https://doi.org/10.1016/j.biocon.2022.109546>

873 Ziegler, S. L., Johnson, J. M., Brooks, R. O., Johnston, E. M., Mohay, J. L., Ruttenberg, B. I.,  
874 Starr, R. M., Waltz, G. T., Wendt, D. E., & Hamilton, S. L. (2023). Marine protected  
875 areas, marine heatwaves, and the resilience of nearshore fish communities. *Scientific*  
876 *Reports*, 13(1), Article 1. <https://doi.org/10.1038/s41598-023-28507-1>



## Tables and Figures

**Table 1.** Results from a series of pairwise permutational analysis of variance (PERMANOVA) tests on community resistance (before-during) and recovery (before-after). PERMANOVAs were performed separately for each habitat and calculated on Bray-Curtis dissimilarity matrices. Pseudo p-values were generated using 999 permutations of residuals. Asterisks denote that community structure was significantly different for pairwise comparisons (Resistance = before vs. during, Recovery = before vs. after).

Habitat	MPA type	Transition	B-C distance	Sum of squares	R-squared	Psuedo F	P(perm)
Rocky intertidal	Outside	Resistance	0.059	0.155	0.02	1.883	0.071
		Recovery	0.053	0.219	0.025	2.635	<b>0.006*</b>
	Inside	Resistance	0.064	0.154	0.026	1.824	0.114
		Recovery	0.063	0.174	0.026	1.946	0.091
Kelp forest inverts and algae	Outside	Resistance	0.085	0.49	0.061	4.381	<b>0.001*</b>
		Recovery	0.125	0.921	0.099	8.211	<b>0.001*</b>
	Inside	Resistance	0.08	0.333	0.047	2.823	<b>0.003*</b>
		Recovery	0.142	0.849	0.102	7.189	<b>0.001*</b>
Kelp forest fishes	Outside	Resistance	0.081	0.249	0.047	3.22	<b>0.007*</b>
		Recovery	0.1	0.64	0.109	9.098	<b>0.001*</b>
	Inside	Resistance	0.066	0.21	0.05	2.944	<b>0.015*</b>
		Recovery	0.124	0.505	0.109	7.569	<b>0.001*</b>
Shallow reef	Outside	Resistance	0.073	0.146	0.061	2.349	0.064
		Recovery	0.071	0.586	0.188	9.515	<b>0.001*</b>
	Inside	Resistance	0.066	0.159	0.093	3.676	<b>0.002*</b>
		Recovery	0.112	0.486	0.225	11.882	<b>0.001*</b>
Deep reef	Outside	Resistance	0.136	0.198	0.124	1.839	0.064
		Recovery	0.167	0.681	0.308	6.687	<b>0.001*</b>
	Inside	Resistance	0.104	0.121	0.082	1.163	0.296
		Recovery	0.107	0.337	0.163	2.53	<b>0.015*</b>

**Table 2.** EnvFit scores indicating the significance level of correlations between environmental conditions (air temperature °C, AT; sea surface temperature °C, SST; sea bottom temperature °C, SBT, biologically effective upwelling transport index  $\text{mmol m}^{-1}\text{s}^{-1}$ , BEUTI; multivariate oceanographic climate index, MOCI) and community change in non-metric multidimensional scaling (nMDS) space.

Habitat	Environmental variable	Type	R square	p-value
Rocky intertidal	AT	Anomaly	0.07	<b>0.043*</b>
		Absolute	0.33	<b>0.001*</b>
	SST	Anomaly	0.04	0.149
		Absolute	0.09	<b>0.012*</b>
	BEUTI	Anomaly	0.04	0.121
		Absolute	0.09	<b>0.011*</b>
MOCI	Absolute	0.02	0.324	
Kelp forest inverts and algae	SST	Anomaly	0.10	<b>0.004*</b>
		Absolute	0.04	0.077
	BEUTI	Anomaly	0.03	0.128
		Absolute	0.02	0.323
	MOCI	Absolute	0.07	<b>0.009*</b>
	SBT	Anomaly	0.17	<b>0.001*</b>
Absolute		0.14	<b>0.001*</b>	
Kelp forest fishes	SST	Anomaly	0.06	<b>0.016*</b>
		Absolute	0.04	0.066
	BEUTI	Anomaly	0.05	<b>0.029*</b>
		Absolute	0.03	0.17
	MOCI	Absolute	0.12	<b>0.002*</b>
	SBT	Anomaly	0.07	<b>0.012*</b>
Absolute		0.01	0.635	
Rocky reef fishes	SST	Anomaly	0.17	<b>0.002*</b>
		Absolute	0.18	<b>0.002*</b>
	BEUTI	Anomaly	0.02	0.401
		Absolute	0.03	0.257
	MOCI	Absolute	0.29	<b>0.001*</b>
	SBT	Anomaly	0.16	<b>0.002*</b>
Absolute		0.12	<b>0.009*</b>	
Deep reef fishes	SST	Anomaly	0.05	0.524
		Absolute	0.08	0.292
	BEUTI	Anomaly	0.25	<b>0.014*</b>
		Absolute	0.27	<b>0.016*</b>
	MOCI	Absolute	0.02	0.726
	SBT	Anomaly	0.03	0.699
Absolute		0.15	0.099	




Figure 1. The (A) coverage of ecological monitoring in marine protected areas (MPAs) in California's Central Coast region and (B-F) exposure of these MPAs to five indicators of environmental conditions. In (A), points indicate which habitats were monitored both inside and outside of MPAs (i.e., points indicate data availability, not the location of sampling sites). Red polygons indicate no-take MPAs ( $n = 12$ , called State Marine Reserves), and blue polygons indicate partial-take MPAs ( $n = 10$ , called State Marine Conservation Areas; see table S1 for list of partial-take MPAs that are de facto no-take MPAs by habitat). The dark horizontal lines delineate the Central Coast region, and the thin gray line indicates state waters (3 nautical miles offshore and all of Monterey Bay). Environmental indicators include: (B) air temperature (AT), (C) sea surface temperature (SST); (D) sea bottom temperature (SBT); (E) the multivariate oceanographic climate index (MOCI); and (F) the biologically effective upwelling index (BEUTI). Lines indicate the median and shading indicates the 95% confidence interval. The 2014-2016 marine heatwave (MHW) is indicated by the vertical red rectangle.

387x484mm (236 x 236 DPI)






Figure 2. (A) Potential and (B) observed shifts in community structure from 2007 to 2020. Panel A illustrates a typology of potential community structure shifts: (i) “No impact”, where community structure does not shift, and “impact”, where community structure significantly shifts. Non-overlapping ellipses indicate significant differences in community structure. In this example, pre-heatwave community structure was significantly different inside (solid lines ellipses around points) vs. outside (dashed lines around triangles) MPAs; (ii) “No MPA benefit”, where communities inside MPAs respond similarly to those outside; (iii) “MPA Resistance”, where communities inside MPAs remain stable while those outside shift; and (iv) “Recovery”, where communities inside MPAs shift during the heatwave, but return to their previous state following the disturbance while communities outside of MPAs do not return. Panel B shows observed shifts in community structure before (2007-2013), during (2014-2016), and after (2017-2020) the marine heatwave inside (circles) and outside (triangles) MPAs in the Central Coast region of California. Each sub-panel represents a habitat and each point depicts the centroid position with 95% confidence ellipses. Also included are vectors for each environmental anomaly: air temperature (AT, rocky intertidal only), sea surface temperature (SST), sea bottom temperature (SBT), the multivariate ocean climate index (MOCI), and the biologically effective upwelling transport index (BEUTI). The trajectory of each vector reflects its correlation with community structure (Table 2). Therefore, indicators that are highly correlated with changes in community structure are aligned with the centroids (points).

452x484mm (236 x 236 DPI)




Figure 3. Community resistance (before-to-during marine heatwave comparison) and recovery (before-to-after marine heatwave comparison) as measured by multivariate distance between centroids. Each point depicts the multivariate distance between the pre-heatwave centroid (before) and the during-heatwave or the post-heatwave centroid inside (red) and outside (blue) MPAs. Therefore, higher values indicate less resistance or recovery. Error bars depict the pooled standard error between centroids, and the asterisks denote significant differences in community structure between marine heatwave periods, as derived from a pairwise permutational analysis of variance (PERMANOVA). Points without asterisks indicate resistance or recovery (as measured by non-significant PERMANOVA, Table 1). Note that some communities have inherently greater interannual variability, potentially resulting from variation in sampling methodology, which can lead to larger mean multivariate distances that are not significantly different (Supplementary Appendix, Figure S9).

452x258mm (236 x 236 DPI)




Figure 4. The (A) proportional community composition by thermal affinity groups from 2007-2020 and the (B) interactive effect of environmental anomalies (air temperature, AT, rocky intertidal only; sea surface temperature, SST; sea bottom temperature, SBT; multivariate oceanographic climate index, MOCI; biologically effective upwelling transport index, BEUTI) and species traits on variance in the abundance of species over the same time period. In (A), vertical lines bound the 2014-2016 marine heatwave years. In (B), colors indicate the standardized coefficients for all trait-oceanographic indicator interaction terms, based on multiple generalized linear-LASSO models. Coefficients were scaled to unit variance to make them visually comparable. Darker squares indicate stronger associations; positive associations are red and negative associations are blue. Black x's mark interaction terms dropped in model selection. White squares without x's indicate traits that were never observed. Dark grey squares indicate non-applicable indicators.

419x516mm (236 x 236 DPI)






Figure 5. Conceptual schematics illustrating scenarios where (A) community structure shifts less inside a MPA than outside, thereby reducing community structure shifts, and (B) community structure shifts more inside a MPA than outside, thereby exacerbating community structure shifts relative to the outside sites. Panel C shows resistance (before-to-during marine heatwave comparison) and recovery (before-to-after marine heatwave comparison) of community structure to the 2014-2016 marine heatwave by marine protected area (MPA) and habitat type along a latitudinal gradient. Point size indicates the Bray-Curtis multivariate distance for resistance or recovery. Therefore, smaller points indicate MPAs that exhibited greater resistance or recovery (i.e., communities inside the MPA shifted smaller multivariate distances). Point color indicates the magnitude of the shift of community structure inside a given MPA relative to its paired outside site. Red shades indicate MPAs where the change in multivariate distance was greater than the paired outside site (i.e., shift exacerbated inside MPAs relative to outside). Blue shades indicate MPAs where the change in multivariate distance was greater in the paired outside site than inside the MPA (i.e., community structure changes were less in the MPA). Note that the rocky intertidal does not use a paired inside-outside sampling design (outside site selection is described in Supplementary Appendix, Methods) and therefore the grey points indicate MPAs without the necessary data from adjacent outside sites. Missing points indicate insufficient data to evaluate changes. SMR = State Marine Reserve (no-take MPA); SMCA = State Marine Conservation Area (partial-take MPA, with a few official and habitat-specific de facto exceptions).