



**HAL**  
open science

## An updated list of the Culicoides (Diptera, Ceratopogonidae) fauna from Ecuador

Juan D Mosquera, Sonia Zapata, Gustavo Spinelli, Moises Gualapuro, Renato León, Denis Augot

### ► To cite this version:

Juan D Mosquera, Sonia Zapata, Gustavo Spinelli, Moises Gualapuro, Renato León, et al.. An updated list of the Culicoides (Diptera, Ceratopogonidae) fauna from Ecuador. *Parasite*, 2022, 29, pp.63. 10.1051/parasite/2022061 . hal-04178171

**HAL Id: hal-04178171**

**<https://hal.science/hal-04178171>**

Submitted on 7 Aug 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## An updated list of the *Culicoides* (Diptera, Ceratopogonidae) fauna from Ecuador

Juan D. Mosquera<sup>1</sup>, Sonia Zapata<sup>1,2,\*</sup> , Gustavo Spinelli<sup>3,4</sup> , Moises Gualapuro<sup>1</sup> , Renato León<sup>1</sup> , and Denis Augot<sup>5,6,\*</sup> 

<sup>1</sup> Instituto de Microbiología, Colegio de Ciencias Biológicas y Ambientales, Universidad San Francisco de Quito, Diego de Robles and Pampite, 170901 Quito, Ecuador

<sup>2</sup> Estación de Biodiversidad Tiputini, Colegio de Ciencias Biológicas y Ambientales, Universidad San Francisco de Quito USFQ, Quito, Ecuador

<sup>3</sup> División Entomología, Museo de La Plata, Paseo del Bosque, 1900 La Plata, Argentina

<sup>4</sup> Instituto de Limnología “Dr. Raúl A. Ringuelet”, Consejo Nacional de Investigaciones Científicas y Técnicas de la Argentina, Universidad Nacional de La Plata, Boulevard 120 y 62, 1900 La Plata, Buenos Aires, Argentina

<sup>5</sup> UscVecpar, ANSES-LSA-EA7510, SFR Cap Santé, Université de Reims Champagne-Ardenne, 51 rue Cognacq-Jay, 51096 Reims Cedex, France

<sup>6</sup> Anses, INRAe, ENVA, UMR-BIPAR, Laboratoire de Santé Animale, 14 rue Pierre et Marie Curie, 94701 Maisons-Alfort Cedex, France

Received 23 July 2022, Accepted 2 December 2022, Published online 23 December 2022

**Abstract** – An updated list of biting midges of the genus *Culicoides* inhabiting Ecuador is provided. Entomological investigations were carried out from July 2010 to May 2019 using CDC light traps in three Ecuadorian regions (Amazon basin, Andean foothills and highlands) and Pacific Coast). A total of 12,073 *Culicoides* specimens from seven subgenera and nine species groups were collected. More species and higher variation were found in the Amazon basin than in either of the Andes regions or coastal sites. A total of 53 species were identified. Of these, 15 are herein reported as new species records for Ecuador: *Culicoides acotylus* Lutz, *C. aitkeni* Wirth & Blanton, *C. benarrochi* Ortiz & Mirsa, *C. carvalhoi* Wirth & Blanton, *C. freitasi* Wirth & Blanton, *C. ginesi* Ortíz, *C. lopesi* Barretto, *C. lyrinotatus* Wirth & Blanton, *C. profundus* Santarém, Felipe-Bauer & Trindade, *C. pseudoreticulatus* Santarém, Felipe-Bauer & Castellón, *C. quasiparaensis* Clastrier, *C. vernoni* Wirth & Blanton, *C. youngi* Wirth & Barreto and two new species. Our results show that the updated list of the Ecuadorian *Culicoides* fauna comprises 70 species. This inventory highlights the presence of species that have been incriminated as vectors of disease elsewhere in animals and humans, mainly *C. insignis* and *C. paraensis*.

**Key words:** Biting midges, New records, Diversity, *Culicoides*, Ecuador.

**Résumé** – Une liste actualisée de la faune des *Culicoides* (Diptera, Ceratopogonidae) de l'Équateur. Une liste actualisée des *Culicoides* vivant en Équateur est fournie. Des enquêtes entomologiques ont été menées de juillet 2010 à mai 2019 à l'aide de pièges lumineux CDC dans trois régions équatoriennes (le bassin amazonien, les Andes (piémont et hautes terres) et la côte Pacifique). Au total, 12 073 spécimens de *Culicoides* de sept sous-genres et neuf groupes d'espèces ont été collectés. Plus d'espèces et une variation plus élevée ont été trouvées dans le bassin amazonien que dans les régions des Andes ou les sites côtiers. Au total, 53 espèces ont été identifiées. Parmi celles-ci, 15 sont signalées ici comme nouvelles espèces pour l'Équateur : *Culicoides acotylus* Lutz, *C. aitkeni* Wirth & Blanton, *C. benarrochi* Ortiz & Mirsa, *C. carvalhoi* Wirth & Blanton, *C. freitasi* Wirth & Blanton, *C. ginesi* Ortíz, *C. lopesi* Barretto, *C. lyrinotatus* Wirth & Blanton, *C. profundus* Santarém, Felipe-Bauer & Trindade, *C. pseudoreticulatus* Santarém, Felipe-Bauer & Castellón, *C. quasiparaensis* Clastrier, *C. vernoni* Wirth & Blanton, *C. youngi* Wirth & Barreto et deux nouvelles espèces. Nos résultats montrent que la liste actualisée de la faune culicoïde équatorienne comprend 70 espèces. Cet inventaire met en évidence la présence d'espèces qui ont été incriminées ailleurs comme vecteurs de maladies chez l'animal et l'homme, principalement *C. insignis* et *C. paraensis*.

Edited by Jean-Lou Justine.

\*Corresponding authors: [szapata@usfq.edu.ec](mailto:szapata@usfq.edu.ec); [denis.augot@anses.fr](mailto:denis.augot@anses.fr)

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Introduction

Biting midges of the genus *Culicoides* Latreille (Diptera: Ceratopogonidae) are the smallest (1–3 mm) of the hematophagous flies and are broadly distributed throughout the world [6]. These insects are vectors of livestock viruses, including blue-tongue virus (BTV), African horse sickness virus (AHSV), Akabane virus (AKAV), Epizootic hemorrhagic disease virus (EHDV) [28], Schmallenberg virus [22], as well as Oropouche virus (OROV), which causes clinical symptoms such as fever, headache, muscle and joint pain, skin rash, meningitis and/or encephalitis in humans [36].

Both BTV and OROV are encountered in South America [28]. Based on vector distribution and environmental conditions, the epidemiology of BTV can be categorized into three zones: endemic (South America north of the Tropic of Capricorn), epidemic (south of the Tropic of Capricorn to Uruguay) and incursion zone (South of Uruguay) [37]. In the Neotropical region, *Culicoides insignis* Lutz and to a lesser extent *C. filarifer* Hoffman, *C. pusillus* Lutz and *C. trilineatus* Fox [24] are suspected to transmit BTV, while *C. paraensis* (Goeldi) is a proven vector of OROV [28].

In Ecuador, OROV has been reported in the Pastaza province in the Ecuadorian Amazon basin [26] and in the Esmeraldas province in the coastal region [58]. Similarly, BTV has been reported: in cattle in El Oro Province in southern Ecuador [23], as well as in Pichincha, Napo, Esmeraldas and Manabi Provinces [11, 50]; in sheep in West of Ecuador [27] and in Pichincha Province [30]. EHDV is present in Pichincha, Napo, Esmeraldas Provinces [50]. In contrast, BTV and epizootic hemorrhagic disease virus (EHDV) are not present in cattle in the Galapagos Islands [51].

Inventories of *Culicoides* have been carried out in the Neotropical region since the middle of the 20th century. The studies carried out by O.P. Forattini and W.W. Wirth and collaborators, and the standardization of morphological characters used to describe *Culicoides* species led to the production of identification aids and dichotomous keys [17, 21, 35, 43, 53, 56].

Moreover, in the list of the biting midges of the World [7], more than 300 new species were described from the Neotropical region between 1853 and 2016.

Ortíz and León [34] first investigated the genus in Ecuador, with the description of five new species (*C. balsapambensis* Ortíz & León, *C. camposi* Ortíz & León, *C. contubernalis* Ortíz & León, *C. insinuatus* Ortíz & León and *C. limonensis* Ortíz & León). Subsequently, a series of publications by several authors contributed to the rise in the number of species in Ecuador to 55 (Table 1) [1, 15, 17, 47, 49, 52, 55, 57].

The number of *Culicoides* species reported in South America varies by country: 44 species have been reported in Argentina [2], 18 in Bolivia [25], 151 in Brazil [45], 114 in Colombia [56], 31 in Peru [32], and 81 in Venezuela [8].

In Ecuador, biting midges are not monitored as part of a national vector surveillance programme in contrast to European countries that have implemented extensive control strategies in response to BTV emergence (Regulation Commission (EC) No.

1266/2007). Nevertheless, in the last decade, a few local academic studies have contributed to better knowledge of biting midges (particularly in taxonomy and distribution) [19, 30, 31]. We aimed to revise the current inventory of *Culicoides* species present in Ecuador in the light of the recent detection of BTV and OROV. The purpose of this study was to produce a comprehensive list of *Culicoides* species in Ecuador, to provide information that may contribute to implement adequate surveillance and vector control strategies in the future.

## Materials and methods

Biting midges were captured using CDC light traps between July 2010 and October 2013 and in May 2019 in nine localities from three regions (Fig. 1): The Amazon basin, from Yasuní-Tiputini region (Orellana Province, July 2010 and October 2013) and Cofán-Dureno (Sucumbíos Province, July 2011); the Andean highlands of Urkusiki region (Imbabura Province, September 2012); the northern Andean foothills of Pacto region (Pichincha Province, March 2011), Paraiso Escondido (Pichincha Province, November 2012), Santo Domingo de los Colorados (Santo Domingo de los Tsachilas Province, August 2013), the southern Andean foothills of Caluma and Echeandia (Bolívar Province, March 2013); and the coastal region of Junin (Manabi Province, March 2013) and Santo Domingo de Onzole (Esmeraldas Province, February 2013 and May 2019). This last collection of 2019 was carried out due to the recent report of Oropouche cases in this coastal province. In Orellana Province, research was conducted at Tiputini Biodiversity Station (TBS). TBS is located on the north bank of the Tiputini River, bordering Yasuní National Park and within Yasuní Biosphere Reserve, one of the most diverse regions of the world [4]. Most of the collections were carried out during the dry season (from July to February) since these sites are more accessible.

The collection sites were selected based on the known ubiquitous capture sites of *Culicoides* spp., like animal shelters, forests, and mud rich sites with organic matter [6, 52]. Seven-night catches per site were performed monthly. Traps were set approximately 1 h before sunset until 1 h after sunrise under favorable climatic conditions (absence of heavy rain and/or wind) [1]. All these specimens were stored in 70–100% ethanol. In the laboratory, *Culicoides* specimens were separated from other arthropods using a stereo microscope (Olympus SZ51), placed in 1.5 mL Eppendorf® tubes containing 70% ethanol and stored at –20 °C. Specimens were identified using different morphological keys [13, 15–17, 44, 47, 52, 54, 56, 57]. The *Culicoides* species were classified into subgenera, species groups according to Borkent and Dominiak [7].

Specimen identification was performed, using a microscope, after mounting the head, wings, genitalia and spermathecae on microscope slides with Gum Chloral [1] or Euparal medium [20]. Digital images of the wings were obtained using an Olympus (BX41 or BX53) microscope equipped with an Olympus SC100 camera and software (cellSens or stream). All voucher specimens were deposited in the entomological collections of San Francisco de Quito University, Ecuador.

**Table 1.** *Culicoides* species list for Ecuador.

Subgenus	Species (synonym)	Trap locality	References
<i>Anilomyia</i> Vargas, 1960	<i>C. efferus</i> Fox, 1952	(Ti)	[56]
	<i>C. metagonatus</i> Wirth & Blanton, 1956	(Pa, Ti2)	[49, 56]
<i>Avaritia</i> Fox, 1955	<i>C. pusillus</i> Lutz, 1913	(PE, SD, Oz)	[17, 52, 54]
<i>Diphaomyia</i> Vargas, 1960	<i>C. freitasi</i> Wirth & Blanton 1973	(Ti)	This study
<i>Haematomyidium</i> Goeldi, 1905	<i>C. debilipalpis</i> Lutz, 1913 ( <i>khalafi</i> Beck, 1957; <i>ichesi</i> Ronderos & Spinelli, 1995)		[17, 52]
	<i>C. equatoriensis</i> Barbosa, 1952 (as a variety of <i>debilipalpis</i> )	(Ti2)	[49, 56]
	<i>C. ginesi</i> Ortíz, 1951	(Ti, CD, Ti2)	This study
	<i>C. glabrior</i> Macfie, 1940 (as a variety of <i>debilipalpis</i> ) ( <i>grahambelli</i> , Forattini, 1956)	(Ti, CD, Ti2)	[49, 56]
	<i>C. insinuatus</i> Ortíz & León, 1955	(CD)	[17, 49, 54, 56]
	<i>C. limonensis</i> Ortíz & León, 1955	(Ti2)	[17, 49, 54, 56]
	<i>C. neoparaensis</i> Tavares & Souza, 1978	(Ti)	[13, 49]
	<i>C. paraensis</i> (Goeldi), 1905 ( <i>undecimpunctatus</i> Kieffer, 1917)	(Ti, CD, Ti2)	[13, 17, 57]
	<i>C. quasiparaensis</i> Clastrier, 1971	(Ti2)	This study
	<i>C. youngi</i> Wirth & Barreto, 1978	(Pa)	This study
<i>Hoffmania</i> Fox, 1948	<i>C. aikeni</i> Wirth & Blanton, 1968	(Ti)	This study
	<i>C. batesi</i> Wirth & Blanton, 1973 ( <i>sanmartini</i> Wirth & Barreto, 1978)	(PE, Oz, Ti, Ti2)	[47, 49, 56]
	<i>C. contubernalis</i> Ortíz & León, 1955 (as a variety of <i>rozeboomi</i> )	(Ti)	[16, 34]
	<i>C. diabolicus</i> Hoffman, 1925	(Pa, Ti, PE, Oz, CE, SD, Ti2, Oz2)	[47, 49, 52]
	<i>C. filariferus</i> Hoffman, 1939		[54, 56]
	<i>C. foxi</i> Ortíz, 1950	(Oz, Ti, Oz2)	[47]
	<i>C. fusipalpis</i> Wirth & Blanton, 1973	(Ti2)	[47, 49, 54, 56]
	<i>C. guttatus</i> (Coquillett), 1904	(Ti, PE, Oz, Ti2, Oz2)	[17]
	<i>C. heliconiae</i> Fox & Hoffman, 1944 ( <i>rozeboomi</i> Barbosa, 1947)	(Pa, Ti, PE, Ju)	[16, 49, 56]
	<i>C. hylas</i> Macfie, 1940	(Pa, Ti, CD, PE, Ju, Ti2)	[1]
	<i>C. insignis</i> Lutz, 1913	(Pa, CD, CE, Oz2)	[47]
	<i>C. lutzi</i> Costa Lima, 1937		[17, 47]
	<i>C. ocumarensis</i> Ortíz, 1950	(Oz, SD)	[47, 56]
	<i>C. pseudodiabolicus</i> Fox, 1946	(Oz, Oz2)	[47, 54, 56]
	<i>C. pseudoheliconiae</i> FelipeBauer, 2008	(Pa, Ti, Oz)	[1]
	<i>C. tidwelli</i> Spinelli, 1993		[47, 49]
	<i>C. trinidadensis</i> Hoffman, 1925 ( <i>oliveri</i> Fox & Hoffman, 1944; <i>wokei</i> Barbosa, 1947 (preoccupied by <i>Culicoides wokei</i> Fox, 1947); <i>diminutus</i> Barbosa, 1951)		[47, 49]
	<i>C. verecundus</i> Macfie, 1948	(Pa)	[17, 49, 52, 54, 56]
<i>Macfiella</i> Fox, 1955	<i>C. phlebotomus</i> (Williston), 1896 ( <i>amazonius</i> Macfie, 1935)		[49, 52, 56]
<i>Mataemyia</i> Vargas, 1960	<i>C. bricenoi</i> Ortíz, 1951	(Ti, Ti2)	[49, 54, 56]
	<i>C. dicrourus</i> Wirth & Blanton, 1955		[49, 56]
<i>Oecacta</i> Poey, 1853	<i>C. alahialinus</i> Barbosa, 1952		[17, 49, 52, 56]
	<i>C. barbosai</i> Wirth & Blanton, 1956		[49, 52, 56]
	<i>C. furens</i> (Poey), 1853 ( <i>maculithorax</i> (Williston), 1896; <i>dovei</i> Hall, 1932; <i>birabeni</i> Cavalieri, 1966)		[17, 49, 52, 56]


(Continued on next page)

**Table 1.** (Continued)

Subgenus	Species (synonym)	Trap locality	References
<i>Psychophaena</i> Philippi, 1865	<i>C. venezuelensis</i> Ortíz & Mirsa, 1950 ( <i>pictipennis</i> (Philippi), 1865; <i>ortizi</i> Fox, 1952)	(Ur, CD)	[56]
<b>Species group</b>			
Acotylus group	<i>C. acotylus</i> Lutz, 1913 ( <i>panamericanus</i> Fox, 1947)	(Ti)	This study
Carpenteri group	<i>C. belemensis</i> Wirth & Blanton, 1973	(Ti, Ti2)	[1]
	<i>C. camposi</i> Ortíz & León, 1955 ( <i>fairchildi</i> Wirth & Blanton, 1955)	(Ti)	[17, 49, 52, 54, 56]
	<i>C. carpenteri</i> Wirth & Blanton, 1953		[49, 56]
Dasyophrus group	<i>C. dasyophrus</i> Macfie, 1940	(Ti2)	[49, 56]
Eublepharus group	<i>C. eublepharus</i> Macfie, 1948 ( <i>transferrans</i> Ortíz, 1953)	(Ti)	[17, 49, 52, 54, 56]
	<i>C. propriipennis</i> Macfie, 1948		[49, 54, 56]
	<i>C. rangeli</i> Ortíz & Mirsa, 1952 ( <i>donajii</i> Vargas, 1954; <i>patulipalpis</i> Wirth and Blanton, 1959)	(Ti2)	[17, 49, 54, 56]
Fluvialis group	<i>C. balsapambensis</i> Ortíz & León, 1955 (as a variety of <i>C. pifanoi</i> )	(Ti)	[17, 49, 52, 56]
	<i>C. castillae</i> Fox, 1946 ( <i>gibsoni</i> Wirth, 1952; <i>flochabonnenci</i> Ortíz & Mirsa, 1952)	(Ti, PE, Oz, Ju, CE, SD, Ti2)	[17, 49, 52, 56]
	<i>C. leopoldoi</i> Ortíz, 1951	(Ti, PE, Ti2)	[56]
	<i>C. tetrathyris</i> Wirth & Blanton, 1959	(Ti, PE, Oz, CE, SD, Ti2)	[49, 54, 56]
	<b>C. sp1</b>	(Ti)	This study
Leoni group	<i>C. benarrochi</i> Ortíz & Mirsa, 1952	(Ti)	This study
	<i>C. gabaldoni</i> Ortíz, 1954		[49, 56]
	<i>C. glabellus</i> Wirth & Blanton, 1956	(Ti, PE)	[49, 56]
	<i>C. leoni</i> Barbosa, 1952		[17, 49, 56]
Limai group	<i>C. carvalhoi</i> Wirth & Blanton, 1973	(Ti)	This study
	<i>C. limai</i> Barretto, 1944	(Ti, PE)	[17, 49, 52, 54, 56]
	<i>C. lopesi</i> Barretto, 1944	(Ti)	This study
	<i>C. vernoni</i> Wirth & Blanton, 1973	(Ti)	This study
	<b>C. sp2</b>	(Ti)	This study
Monticola group	<i>C. monticola</i> Wirth & Lee, 1967 ( <i>pichindensis</i> Browne, 1980)		[49, 56]
Pachymerus group	<i>C. pachymerus</i> Lutz, 1914	(Ti)	[17]
Reticulatus group	<i>C. lyrinotatus</i> Wirth & Blanton, 1955	(Ti)	This study
	<i>C. paucienfuscatus</i> Barbosa, 1947	(Ti, Ti2)	[56]
	<i>C. pifanoi</i> Ortíz, 1951 ( <i>tricoloratus</i> Wirth & Blanton, 1953)	(Ti, Oz, Ti2)	[54]
	<i>C. profundus</i> Santarém, Felipe-Bauer & Trindade, 2014	(Ti)	This study
	<i>C. pseudoreticulatus</i> Santarém, Felipe-Bauer & Castellón, 2014	(Ti, Ti2)	This study
Stigmalis group	<i>C. alvarezi</i> Ortíz, 1957		[49, 56]
	<i>C. fluviatilis</i> (Lutz), 1914 ( <i>scorzai</i> Ortíz, 1956)		[49, 54, 56]

*Andean highlands*: (Ur: Urkisiki); *Northern Andean foothills*: (Pa: Pacto), (PE: Paraiso Escondido), (SD: Santo Domingo de los Colorados); *Southern Andean foothills*: (CE: Caluma-Echandia); *Amazon Basin*: (CD: Cofán-Dureno), (Ti, Ti2: Tiputini); *Coast*: (Oz, Oz2: Santo Domingo de Onzole) and (Ju: Junin).

¥ Specimens from Ecuador previously considered to be *C. verecundus* were restored from synonymy by Felipe-Bauer *et al.* [15] and are now known as *C. contubernalis* Ortíz & León.



**Figure 1.** Sites of entomological collections.

## Results


Fifty-three species of the genus *Culicoides* were identified morphologically from a total of 12,073 captured *Culicoides* specimens. The locality with the highest abundance was Yasuni-Tiputini in Orellana Province in both collection periods, with 7,691 specimens collected. We also found 73 specimens in Cofan-Dureno, 323 in Pacto, 243 in Paraiso Escondido, 194 in Santo Domingo de los Colorados, 19 in Urkusiki, 1117 in Caluma-Echeandia, 1533 in Santo Domingo de Onzole, and 880 in Junin. Females represent 71.52% of the collected biting midges, while 28.48% were males. We identified 29 species, belonging to seven different subgenera (*Anilomyia* Vargas, *Avaritia* Fox, *Diphaomyia* Vargas, *Haematomyidium* Goeldi, *Hoffmania* Fox, *Mataemyia* Vargas, *Psychophaena* Philippi) (Table 1, Fig. 2). Moreover, we caught 24 species belonging to nine species groups (acotylus, carpenteri, dasyophrus, eublepharus, fluvialis, leoni, limai, pachymerus and reticulatus) (Table 1, Fig. 2). Nine species were found in three Ecuadorian regions (*C. batesi* Wirth & Blanton, *C. castillae* Fox, *C. diabolicus* Hoffman, *C. guttatus* (Coquillett), *C. heliconiae* Fox & Hoffman, *C. hylas* Macfie, *C. insignis* Lutz, *C. pseudoheliconiae* Felipe-Bauer, *C. tetrathyris* Wirth & Blanton). We reported two new species collected in TBS (Table 1). *Culicoides pseudodiabolicus* Fox was only found in the coastal region of Santo Domingo de Onzole (Esmeraldas) in both collection periods. Forty-eight species were caught in the Amazon basin, and 45 of them were collected in TBS. Two species (*C. youngi* Wirth & Barreto, *C. verecundus* Macfie) were found only in the Andean foothills. *Culicoides venezuelensis* Ortíz

& Mirsa was the only species found in the Andean highlands. For the distribution of other species, see Table 1.

## Discussion

Fifteen species are herein newly recorded for Ecuador (Table 1 and Fig. 2): *C. aitkeni* Wirth & Blanton, *C. acotylus* Lutz, *C. benarrochi* Ortiz & Mirsa, *C. carvalhoi* Wirth & Blanton, *C. freitasi* Wirth & Blanton, *C. ginesi* Ortíz, *C. lopesi* Barretto, *C. lyrinotatus* Wirth & Blanton, *C. profundus*, Santarém, Felipe-Bauer & Trindade, *C. pseudoreticulatus* Santarém, Felipe-Bauer & Castellón, *C. quasiparaensis* Clastrier, *C. vernoni* Wirth & Blanton, *C. youngi* Wirth & Barreto and two new species; one belongs to the fluvialis group according to DNA barcoding [1] and the other belongs to the limai group according to morphological characters.

Additionally, we detected 13 species that have been reported previously only once in Ecuador (Table 1): *C. efferus* Fox, *C. belemensis* Wirth & Blanton, *C. foxi* Ortiz, *C. guttatus* (Coquillett), *C. hylas* Macfie, *C. insignis*, *C. leopoldo* Ortiz, *C. neoparaensis* Tavares & Souza, *C. pachymerus* Lutz, *C. paucifuscatus* Barbosa, *C. pifanoi* Ortíz, *C. pseudoheliconiae* Felipe-Bauer and *C. venezuelensis* Ortíz & Mirsa. *Culicoides venezuelensis* was found for the first time in the Ecuadorian Amazon basin (Sucumbios province) and Andean highlands (Imbabura province). *Culicoides batesi*, *C. castillae*, *C. diabolicus*, *C. guttatus*, *C. heliconiae*, *C. hylas*, *C. insignis*, *C. pseudoheliconiae* and *C. tetrathyris* were found in three regions (Coast, Andean, and Amazon basin). The highest number of



**Figure 2.** Wing photographs of *Culicoides* species in Ecuador. The new *Culicoides* records are: *C. acotylus*, *C. aitkeni*, *C. benarrochi*, *C. carvalhoi*, *C. freitasi*, *C. ginesi*, *C. lopesi*, *C. lyrinotatus*, *C. profundus*, *C. pseudoreticulatus*, *C. quasiparaensis*, *C. vernoni*, *C. youngi* and two new species (*Culicoides* n. sp. 1 and *Culicoides* n. sp. 2).




Figure 2. Continued.






Figure 2. Continued.




Figure 2. Continued.



**Figure 2.** Continued.

species was found in the Amazon basin, in the Tiputini region. This finding corroborates other studies that suggest that Tiputini harbors immense biodiversity and probably the highest diversity of insects in the world (100,000 species/ha) [4, 59]. Our results increase the number of Ecuadorian *Culicoides* species to 70 (Table 1).

*Culicoides insignis*, *C. pachymerus*, and *C. paraensis* have medical and/or veterinary importance because they have been reported as possible causative agents of dermatozoonosis [42, 46]. Intriguingly, three species identified in this study (*C. foxi*, *C. insignis*, and *C. filarifer*) have been reported to carry DNA from *Leishmania brasiliensis* and *Le. Amazonensis* [39]. Additionally, *C. foxi* is a causative agent of allergic dermatitis and *C. pifanoi* is known as a causative agent of filariasis [8]. Moreover, *C. acotylus*, *C. fluvialis*, and *C. leopoldoi* are suspected to cause allergic dermatitis in humans [8]. Finally, *C. guttatus* may play a role as a vector of mansonellosis, a neglected tropical disease [38] and of BTV [8]. Further investigations are needed to determine the role of these species in dermatological disorders.

According to [6, 10, 45], the reported distribution of the species newly cited herein for Ecuador are: i) *C. acotylus* in Brazil (Amazonas, Pará, Rondônia, Roraima, and Mato Grosso), Honduras, Mexico, Panama, Suriname, Trinidad and Tobago, and Venezuela; ii) *C. aitkeni* in Brazil (Amazonas, Pará); Trinidad and Tobago; iii) *C. benarrochi* in Brazil (Amazonas, Roraima, and Rio de Janeiro), Trinidad and Tobago and Venezuela; iv) *C. carvalhoi* in Brazil (Pará); v) *C. freitasi* in

Brazil (Amazonas and Pará); vi) *C. ginesi* from El Salvador to Panama, northeastern Argentina, Brazil (Pará, Rondônia), Colombia, Trinidad and Tobago and Venezuela; vii) *C. lopesi* in Brazil (Acre, Amazonas, Rio de Janeiro, and São Paulo), Panama and Suriname; viii) *C. lyrinotatus* in Brazil (Pará), Nicaragua and Panama; ix) *C. profundus* in Brazil (Amazonas, Pará and Rondônia); x) *C. pseudoreticulatus* in Brazil (Amazonas, Rondônia and Roraima); xi) *C. quasiparaensis* in Brazil (Acre, Amazonas, Pará, Rondônia, Roraima, and Maranhão), Colombia, El Salvador, French Guiana, Honduras and Peru; xii) *Culicoides vernoni* in Brazil (Amazonas, Rondônia and Pará), Bolivia, Colombia and Costa Rica; and xiii) *C. youngi* in Brazil (Amazonas) and Colombia.

*Culicoides paraensis* is distributed from the USA to Argentina [49, 56], and it is abundant in the humid tropics in Panama [52] and in urban areas with banana crops in the Amazon region and in the southern states of Brazil [14]. It was collected in South Carolina zoos [33]. In Peru, populations of *C. paraensis* seemed to be highest from October to December, with numbers fluctuating depending on the year and collection site [29]. Interestingly, the presence of *C. paraensis*, one of the known vectors of OROV [40], was also confirmed herein in Ecuador. An estimated half-million people have been affected by OROV since it was first isolated in Brazil [9]. In Ecuador, OROV was detected in the Esmeraldas Province (Coast) [58] and there is serological evidence in febrile patients from the Amazon basin (Pastaza Province) [26]. In our study, *C. paraensis* was found

in Orellana province, located north of Pastaza province, suggesting that this vector may be transmitting OROV in Pastaza province where serologic evidence was detected [26]. In contrast, *C. paraensis* was absent in the Coastal region, (Esmeraldas and Manabi Provinces). Future investigations will address the distribution of this species particularly in forested and urban areas in Ecuador. In this study, *C. neoparaensis*, *C. paraensis*, and *C. quasiparaensis* were collected in the same area (Yasuní-Tiputini – Amazon basin). These three species belong to the paraensis group that includes seven species (*C. diversus* Felipe-Bauer, *C. filiductus* Wirth, *C. peruvianus* Felipe-Bauer, and *C. austroparaensis* Spinelli) [13, 48]. It is worth remarking that our identifications were based only on morphological characters; thus, further molecular analysis would be needed to explore the boundaries of these closely related/similar species.

Finally, *C. insignis* is considered the main candidate vector species of BTV in South America [5], an arbovirus that infects livestock worldwide and is responsible for global losses of agriculture of up to US\$3 billion [41]. Bluetongue virus has been detected in blood samples from Ecuadorian cattle [11, 50]; however, its vector species in Ecuador have not been confirmed yet. *Culicoides insignis* is distributed in the USA (Alabama, Georgia, and Florida states), Mexico (Chiapas, Yucatán), Central America and Caribbean to central Argentina [6, 49, 56]. It is abundant in the arid tropics and absent in humid tropics in Panama [52]. Populations of *C. insignis* were highest from July to December at most sites in the Caribbean, and from June to October in Central America [18]. Therefore, in this region, nulliparous and parous females of this species were captured throughout the year [18]. In this study, *C. insignis* was collected in the Amazon basin and in the foothills (from north to south) of the Andean region until 2013, but we also found specimens in 2019 in the coastal region of Esmeraldas province; thus, it is possible that this species has favorable adaptation capacities due to its broad distribution throughout South America [3, 12].

In conclusion, our examination of fresh material revealed 15 new species records for the country, and confirmed 13 species previously reported in Ecuador. Herein, we present an updated list of the Ecuadorian *Culicoides* that comprises 70 species, including possible vectors of human and veterinary interest. This species inventory is a prerequisite for the future development of a barcode library and the construction of an image database of *Culicoides* wings. Both may be useful for further ecological studies and to establish risk maps for *Culicoides*-borne diseases in Ecuador.

**Acknowledgements.** This study was funded by Universidad San Francisco de Quito and Université de Reims Champagne-Ardenne. We are especially grateful to Rosita Bayas who generated the map of entomological collection sites. Also, we thank L. Hadj-Henni and Carolina Molina for their important contributions to this manuscript. We would like to thank Dr. Maria Luiza Felipe-Bauer (reviewer of this article) for her time and constructive comments, which have greatly improved the manuscript.

## Author contributions

DA, SZ and RL designed the study. JM, SZ, DA, MG and RL carried entomological collections, separation, and identifica-

tion of specimens. GS and DA confirmed identification of *Culicoides* species. JM, DA, GS, MG, RL and SZ wrote the paper and approved the final version.

## References

- Augot D, Mathieu B, Hadj-Henni L, Barriel V, Zapata-Mena S, Smolis S, Slama D, Randrianambintsoa FJ, Trueba G, Kaltenbach M, Rahola N, Depaquit J. 2017. Molecular phylogeny of 42 species of *Culicoides* Diptera. Ceratopogonidae from three continents. Parasite, 24, 23.
- Ayala MM, Marino PI. 2018. New records of biting midges of the genus *Culicoides* Latreille Diptera: Ceratopogonidae in Argentina. Revista de la Sociedad Entomológica Argentina, 77, 36–38.
- Aybar CAV, Juri MJD, De Grosso MSL, Spinelli GR. 2011. Spatial and temporal distribution of *Culicoides insignis* and *Culicoides paraensis* in the subtropical mountain forest of Tucumán, northwestern Argentina. Florida Entomologist, 944, 1018–1025.
- Bass MS, Finer M, Jenkins CN, Kreft H, Cisneros-Heredia DF, McCracken SF, Pitman NCA, English PH, Swing K, Villa G, DiFiore A, Voigt CC, Kunz TH. 2010. Global conservation significance of Ecuador's Yasuní National Park. PloS One, 51, e8767.
- Borkent A. 2005. The biting midges, the Ceratopogonidae Diptera, in Biology of Diseases Vectors. Marquardt W, Editor. Academic Press: San Diego, CA. p. 113–126.
- Borkent A, Spinelli GR. 2007. Neotropical Ceratopogonidae (Diptera: Insecta), vol IV, Pensoft Publishers: Sofia, Bulgaria. p. 198.
- Borkent A, Dominiak P. 2020. Catalog of the biting midges of the World (Diptera: Ceratopogonidae). Zootaxa, 4787(1), 1–377.
- Cazorla-Perfetti D. 2014. Catálogo de las especies de ceratopogonidae diptera: nematocera registradas para Venezuela y su importancia agro-ecológica y sanitaria. Saber, 264, 395–408.
- Da Rosa JFT, De Souza WM, De Paula Pinheiro F, Figueiredo ML, Cardoso JF, Acrani GO, Nunes MRT. 2017. Oropouche virus: clinical, epidemiological, and molecular aspects of a neglected Orthobunyavirus. American Journal of Tropical Medicine and Hygiene, 965, 1019–1030.
- De Sousa FE, Almeida JF, Pessoa FAC. 2016. List of *Culicoides* biting midges (Diptera: Ceratopogonidae) from the state of Amazonas, Brazil, including new records. Check List, 12, 2002.
- De la Torre E, Moreira N, Saegerman C, De Clercq K, Salinas M, Maldonado A, Jarrín D, Sol Vaca M, Pachacama S, Espinoza J, Delgado H, Barrera M. 2021. Bluetongue virus infections in cattle herds of Manabí province of Ecuador. Pathogens, 10, 1445.
- Diaz F, Ronderos MM, Spinelli GR. 2005. The immatures of the Neotropical species *Culicoides venezuelensis* Ortiz and Mirsa (Diptera: Ceratopogonidae). Transactions of the American Entomological Society, 131, 375–385.
- Felipe-Bauer ML, Cáceres AG, Silva CS, Valderrama-Bazan W, Gonzales-Perez A. 2003. Two new *Culicoides* of the paraensis species group Diptera: Ceratopogonidae from the Amazonian region of Peru. Memórias do Instituto Oswaldo Cruz, 988, 1051–1058.
- Felipe-Bauer ML, Sternheim US. 2008a. *Culicoides paraensis* (Diptera: Ceratopogonidae) infestations in cities of the Itapocu river valley, Southern Brazil. Entomological News, 119, 185–192.

15. Felipe-Bauer ML, Cáceres AG, Silva CS, Valderrama-Bazan W, Gonzales-Perez A, Costa JM. 2008b. Description of *Culicoides pseudoheliconiae* sp. n. from Peruvian Amazon and revalidation of *Culicoides contubernalis* Ortiz and León (Diptera: Ceratopogonidae). *Memórias do Instituto Oswaldo Cruz*, 1033, 259–262.
16. Felipe-Bauer ML, Damasceno CP, Py-Daniel V, Spinelli GR. 2009. *Culicoides baniwa* sp.nov. from the Brazilian Amazon Region with a synopsis of the *hylas* species group (Diptera: Ceratopogonidae). *Memórias do Instituto Oswaldo Cruz*, 104 (6), 851–857.
17. Forattini OP. 1957. *Culicoides* da região neotropical Diptera. Ceratopogonidae. Arquivos da Faculdade de Higiene e Saúde Pública da Universidade de São Paulo, 11, 161–526.
18. Greiner EC, Mo CL, Homan EJ, Gonzalez J, Oviedo MT, Thompson LH, Gibbs EP. 1993. Epidemiology of bluetongue in Central America and the Caribbean: initial entomological findings. *Medical Veterinary Entomology*, 7, 309–315.
19. Gualapuro MR. 2013. Contribución al estudio de la fauna de *Culicoides* (Diptera: Ceratopogonidae) en la zona norte del Ecuador. USFQ, Ecuador: Quito.
20. Hadj-Henni L, Djerada Z, Millot C, Augot D. 2021. Comprehensive characterisation of *Culicoides clastri* and *C. festivipennis* (Diptera: Ceratopogonidae) according to morphological and morphometric characters using a multivariate approach and DNA barcode. *Scientific Reports*, 11, 521.
21. Huerta H, Benítez-Alva JI, Concha-Suarez J, Ibáñez-Piñón CR, Manrique-Saïde P. 2020. New records of genus *Culicoides* Latreille from Oaxaca, Mexico (Diptera: Ceratopogonidae). *Acta Zoológica Mexicana (nueva serie)*, 36, 1–27.
22. Koenraadt CJ, Balenghien T, Carpenter S, Ducheyne E, Elbers AR, Fife M, Garros C, Ibáñez-Justicia A, Kampen H, Kormelink RJ, Losson B, Van der Poel W, De Regge N, Van Rijn P, Sanders C, Schaffner F, Van Oldruitenborgh-Oosterbaan M, Takken W, Werner D, Seelig F. 2014. Bluetongue, Schmallenberg-what is next? *Culicoides*-borne viral diseases in the 21st Century. *BMC Veterinary Research*, 101, 1–7.
23. Lopez W, Nicoletti P, Gibbs E. 1985. Antibody to bluetongue virus in cattle in Ecuador. *Tropical Animal Health Production*, 17, 82.
24. Maclachlan NJ, Guthrie AJ. 2010. Re-emergence of bluetongue, African horse sickness, and other orbivirus diseases. *Veterinary Research*, 416, 35.
25. Mancini JMD, Aybar CAV, Fuenzalida AD, De Grosso MSL, Quintana MG. 2016. Ceratopogonidae Diptera: Nematocera of the piedmont of the Yungas forests of Tucumán: ecology and distribution. *PeerJ*, 4, e2655.
26. Manock SR, Jacobsen KH, De Bravo NB, Russell KL, Negrete M, Olson JG, Sanchez JL, Blair PJ, Smalligan RD, Quist BK, Espín JF, Espinoza WR, MacCormick F, Fleming LC, Kochel T. 2009. Etiology of acute undifferentiated febrile illness in the Amazon basin of Ecuador. *American Journal of Tropical Medicine & Hygiene*, 811, 146–151.
27. Mantilla A. 1998. Anticuerpos del virus de la LA en ovinos de cria e. ANCO: Instituto nacional de Higiene.
28. Mellor P, Boorman J, Baylis M. 2000. *Culicoides* biting midges: their role as arbovirus vectors. *Annual Review of Entomology*, 451, 307–340.
29. Mercer DR, Castillo-Pizango MJ. 2005. Changes in relative species compositions of biting midges (Diptera: Ceratopogonidae) and an outbreak of Oropouche virus in Iquitos, Peru. *Journal of Medical Entomology*, 42, 554–558.
30. Merino Mena CX. 2011. Identificación de anticuerpos precipitantes para el virus de la lengua azul en suero de ovinos de la parte alta y baja de la provincia de Pichincha. Facultad De Ingeniería Y Ciencias Agropecuarias: Ecuador.
31. Mosquera JD. 2017. Research about virus of medical and veterinary importance transmitted by *Culicoides* – Investigación del virus de importancia médica y veterinaria transmitidos por culicoides. Master's Thesis. Universidad San Francisco de Quito (USFQ): Quito, Ecuador.
32. Navarro D, Rojas M, Jurado J, Manchego A, Ramírez M, Castillo A, Rivera H. 2019. Detección molecular del virus de Lengua Azul en *Culicoides insignis* y en ovinos de Pucallpa, Perú. *Revista de Investigaciones Veterinarias del Peru*, 301, 465–476.
33. Nelder MPD, Swanson A, Adler PH, Grogan WL. 2010. Biting midges of the genus *Culicoides* in South Carolina zoos. *Journal of Insect Science*, 10, 55.
34. Ortíz I, León LA. 1955. Los *Culicoides* (Diptera: Ceratopogonidae) de la Republica del Ecuador. *Boletín de Informaciones Científicas Nacionales*, 67, 564–590.
35. Perruolo GJ. 2009. Keys of species of *Culicoides* (Diptera: Ceratopogonidae) associated with cattle operations in Neotropical Region. *Revista Científica (Maracaibo)*, 19(2), 124–133.
36. Pinheiro FP, Travassos da Rosa APA, Travassos da Rosa JFS, Ishak R, Freitas RB, Gomes RMLC, LeDuc JW, Oliva OFP. 1981. Oropouche virus. *American Journal of Tropical Medicine & Hygiene*, 30, 149–160.
37. Portela Lobato ZI, Maldonado Coelho Guedes MI, Diniz Matos AC. 2015. Bluetongue and other orbiviruses in South America: gaps and challenges. *Veterinaria Italiana*, 51(4), 253–262.
38. Racourt C. 2017. Review of *Mansonella ozzardi* and its vectors in the Neotropical region with emphasis on the current situation in Haiti. *Journal of Bacteriology and Parasitology*, 8, 2.
39. Rebêlo JMM, Rodrigues BL, Abreu-Bandeira MDC, Moraes JLP, Fonteles RS, Pereira SRF. 2016. Detection of *Leishmania amazonensis* and *Leishmania braziliensis* in *Culicoides* (Diptera, Ceratopogonidae) in an endemic area of cutaneous leishmaniasis in the Brazilian Amazonia. *Journal of Vector Ecology*, 41, 303–308.
40. Roberts DR, Hoch AL, Dixon KE, Llewellyn CH. 1981. Oropouche virus. *American Journal of Tropical Medicine and Hygiene*, 301, 165–171.
41. Rushton J, Lyons N. 2015. Economic impact of Bluetongue: a review of the effects on production. *Veterinaria Italiana*, 51, 401–406.
42. Santamaría E, Cabrera OL, Zipa Y, Ferro C, Ahumada ML, Pardo RH. 2008. Preliminary evaluation of the *Culicoides* biting nuisance (Diptera: Ceratopogonidae) in the province of Boyaca, Colombia. *Biomedica*, 28, 497–509.
43. Santarém MC, Da Trindade RL, Da Silva Tdo N, Castellón EG, Patiu CA, Felipe-Bauer ML. 2014. New Neotropical *Culicoides* and redescription of *Culicoides reticulatus* Lutz (Diptera: Ceratopogonidae). *Zootaxa*, 3795, 255–274.
44. Santarém MC, Farias E, Felipe-Bauer ML. 2015. *Culicoides castelloni* sp.nov. from the Brazilian Amazon Region with a revision of the *reticulatus* species group (Diptera, Ceratopogonidae). *Annals of the Brazilian Academy of Sciences*, 87, 955–972.
45. Santarém MC, Felipe-Bauer ML. 2022. Brazilian Species of Biting Midges (Diptera: Ceratopogonidae) [Espécies de Maruins do Brasil]. *Coleção de Ceratopogonidae. CCER, Fiocruz, Brazil*. 68 pp. Available online from <https://portal.fiocruz.br/documento/especies-maruins-do-brasil>
46. Sherlock IA, Guitton N. 1965. Dermatozoonosis by *Culicoides'* bite Diptera: Ceratopogonidae in Salvador, State of Bahia, Brazil. III. Epidemiological aspects. *Memórias do Instituto Oswaldo Cruz*, 63, 1–12.
47. Spinelli GR, Greiner E, Wirth W. 1993. The Neotropical bloodsucking midges of the *Culicoides guttatus* group of the subgenus *Hoffmania* Diptera: Ceratopogonidae. *Contributions of the American Entomological Institute*, 27, 1–91.

48. Spinelli GR, Ronderos MM, Díaz F, Marino PI. 2005. The bloodsucking biting midges of Argentina (Diptera: Ceratopogonidae). *Memórias do Instituto Oswaldo Cruz*, 100(2), 137–150.
49. Spinelli GR, Santamaría E, Cabrera OL, Ronderos MM, Suárez MF. 2009. Five new species of *Culicoides* Latreille described from Colombia, yielding a new species list and country records Diptera: Ceratopogonidae. *Memórias do Instituto Oswaldo Cruz*, 104(1), 81–92.
50. Verdezoto J, Breard E, Viarouge C, Quenault H, Lucas P, Sailleau C, Zientara S, Augot D, Zapata S. 2018. Novel serotype of bluetongue virus in South America and first report of epizootic haemorrhagic disease virus in Ecuador. *Transboundary and Emerging Diseases*, 65(1), 244–247.
51. Vinuesa RL, Cruz M, Bréard E, Viarouge C, Zanella G. 2018. Bluetongue virus and epizootic hemorrhagic disease virus survey in cattle of the Galapagos Islands. *Journal of Veterinary Diagnostic Investigation*, 31(2), 271–275.
52. Wirth W, Blanton FS. 1959. Biting midges of the genus *Culicoides* from Panama Diptera: Heleidae. *Proceedings of the United States National Museum*, 109, 237–482.
53. Wirth W, Blanton FS. 1968. A revision of the Neotropical biting midges of the *Hylas* group of *Culicoides* (Diptera, Ceratopogonidae). *Florida Entomologist*, 51, 201–215.
54. Wirth W, Blanton FS. 1973. A review of the maruins or biting midges of the genus *Culicoides* Diptera: Ceratopogonidae in the Amazon Basin. *Amazoniana: Limnologia et Oecologia Regionalis Systematis Fluminis Amazonas*, 4, 405–470.
55. Wirth W, Dyce AL, Peterson BV, Roper I. 1984. An atlas of wing photographs, with a summary of the numerical characters of the Nearctic species of *Culicoides* (Diptera: Ceratopogonidae). *Contributions of the American Entomological Institute*, 22, 1–46.
56. Wirth W, Dyce AL, Spinelli GR. 1988. An atlas of wing photographs, with a summary of the numerical characters of the Neotropical species of *Culicoides* (Diptera: Ceratopogonidae). *Contributions of the American Entomological Institute*, 25, 1–72.
57. Wirth W, Felipe-Bauer ML. 1989. The Neotropical biting midges related to *Culicoides paraensis* Diptera: Ceratopogonidae. *Memórias do Instituto Oswaldo Cruz*, 84, 551–565.
58. Wise EL, Pullan ST, Márquez S, Paz V, Mosquera JD, Zapata S, Jackson SK, Fejer G, Trueba G, Logue CH. 2018. Isolation of Oropouche virus from febrile patient, Ecuador. *Emerging Infectious Diseases*, 24, 935–937.
59. Zook D. 2010. Tropical rainforests as dynamic symbiospheres of life. *Symbiosis*, 51, 27–36.

**Cite this article as:** Mosquera JD, Zapata S, Spinelli G, Gualapuro M, León R & Augot D. 2022. An updated list of the *Culicoides* (Diptera, Ceratopogonidae) fauna from Ecuador. *Parasite* 29, 63.



An international open-access, peer-reviewed, online journal publishing high quality papers on all aspects of human and animal parasitology

Reviews, articles and short notes may be submitted. Fields include, but are not limited to: general, medical and veterinary parasitology; morphology, including ultrastructure; parasite systematics, including entomology, acarology, helminthology and protistology, and molecular analyses; molecular biology and biochemistry; immunology of parasitic diseases; host-parasite relationships; ecology and life history of parasites; epidemiology; therapeutics; new diagnostic tools.

All papers in Parasite are published in English. Manuscripts should have a broad interest and must not have been published or submitted elsewhere. No limit is imposed on the length of manuscripts.

**Parasite** (open-access) continues **Parasite** (print and online editions, 1994-2012) and **Annales de Parasitologie Humaine et Comparée** (1923-1993) and is the official journal of the Société Française de Parasitologie.

Editor-in-Chief:  
Jean-Lou Justine, Paris

Submit your manuscript at  
<http://parasite.edmgr.com/>