

Link marine restoration to marine spatial planning through ecosystem-based management to maximize ocean regeneration

Elisabetta Manea, Tundi Agardy, Lucia Bongiorni

▶ To cite this version:

Elisabetta Manea, Tundi Agardy, Lucia Bongiorni. Link marine restoration to marine spatial planning through ecosystem-based management to maximize ocean regeneration. Aquatic Conservation: Marine and Freshwater Ecosystems, In press, 10.1002/aqc.3999. hal-04178937

HAL Id: hal-04178937 https://hal.science/hal-04178937v1

Submitted on 8 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1	Link marine restoration to marine spatial planning through ecosystem based management to		
2	maximize ocean regeneration		
3	Elisabetta Manea ^{1,2*} , Tundi Agardy ^{3*} , Lucia Bongiorni ^{2*}		
4			
5	¹ Laboratoire d'Ecogeochimie des Environnements Benthiques (LECOB), Sorbonne Université,		
6	Université Pierre-et-Marie-Curie, Centre National de la Recherche Scientifique (CNRS),		
7	Observatoire Oceanologique, F-66650 Banyuls sur Mer, France		
8	² Institute of Marine Sciences, National Research Council, CNR-ISMAR, Arsenale, Tesa 104,		
9	Castello 2737/F, 30122, Venice, Italy		
10	³ Sound Seas, 26 Van Nuys Road, Colrain, MA 01340, USA		
11			
12	Main Corresponding author details		
13	Elisabetta Manea		
14	Current email address: elisabetta.manea@obs-banyuls.fr		
15			
16	Abstract		
17	1. The speed at which marine and coastal ecosystems are being degraded due to cumulative		
18	impacts limits the effectiveness of conservation strategies. To abate ocean degradation and		
19	allow ocean regeneration, conservation planning needs to be improved and ecological		
20	restoration will be needed.		
21	2. This study explores the potential of incorporating restoration into marine spatial planning		
22	(MSP) anchored to ecosystem based management (EBM), termed EB-MSP, for maximizing		
23	ocean regeneration. Our perspective explicitly brings both passive and active restoration into		
24	EB-MSP in a broad and holistic framework for achieving the recovery of ocean ecosystems,		
25	their functions, and their valuable services.		
26	3. By proposing a restoration-focused EB-MSP framework we highlight the co-benefits of		
27	interlinking MSP and marine restoration through the EBM core principles. Such benefits		
28	include: a scaling up of restoration effectiveness, a greater guarantee that sustainability and		

29 conservation goals will be met, and improvements in MSP as an integrated planning tool with

the potential to address climate change. Together this will promote ocean regeneration
 alongside management for sustainable use, to prevent further degradation and to allow much needed ecological recovery.

33

Keywords: addressing climate change, blue economy, ecological principles, ecosystem services,
MSP, ocean sustainability, restoration planning, restoring marine ecosystems

36

37 **1. Introduction**

38 Achieving sustainability in the use of ocean space and resources requires improving the ecological status of ecosystems and securing the interconnections between them. Many marine and coastal 39 ecosystems have been degraded by cumulative impacts that strain their resilience and exert pressure 40 on the ability of the ocean to provide ecosystem services (Halpern et al., 2019). Systematic 41 42 conservation planning can identify marine areas to conserve, but as local anthropogenic and climate change pressures intensify, degraded environmental conditions are inevitable (Saeedi et al., 2019). 43 Restoration science has matured to be able to offer practical guidance on how to assist the recovery 44 and the re-establishment of many degraded marine habitats such as seagrass beds, mangroves, oyster 45 beds and coral reefs. 46

47 Restoration, rehabilitation and remediation (see Table 1 for terminology) are a set of interconnected 48 approaches for environmental recovery, the so-called "restorative continuum" (Gann et al., 2019; 49 Chazdon et al., 2021). Collectively these approaches can lead to ocean regeneration, a term we use to 50 depict the recovery of ocean ecosystems, their functions, and their valuable services. Restoration and 51 conservation strategies are synergic, and must draw on their complementary strengths to achieve their 52 goals (Wiens & Hobbs, 2015).

Marine restoration has been featured in numerous international environmental commitments
(Abelson et al., 2020). The need to restore degraded ecosystems has been recognized for years by the

Convention on Biological Diversity (CBD) initiatives and agreements (Miller, 1999; Maes et al., 55 2008; De Groot et al., 2013) and by diverse European policy instruments (e.g., Habitats Directive, 56 EEC, 1992; Marine Strategy Framework Directive, EC, 2008); quantitative targets for restoration 57 now figure prominently in the CBD's Global Biodiversity Framework agreed in late 2022 (CBD 58 2022). Regeneration of aquatic ecosystems is now one of the main missions of the EU Biodiversity 59 Strategy 2030 which supports the Green Deal - the European agreement set to address the climate 60 crisis by protecting and restoring natural ecosystems and related capital (EU, 2019). Moreover, 61 marine ecosystem restoration is recognized as necessary to achieve many United Nations Sustainable 62 Development Goals (SDGs), especially SDG14 'Life Below Water' (Diz et al., 2018) and is 63 prominent in the UN Decade of Ecosystem Restoration (Aronson et al., 2020). Recently, the 64 European Commission has committed to proposing a legal framework for nature restoration 65 (https://environment.ec.europa.eu/publications/nature-restoration-law en), a unique opportunity for 66 67 a significant improvement in the ecological quality of restoration outcomes, including its implementation at large scale (Shumway et al., 2021; Cliquet et al., 2022). As such, ecological 68 69 restoration is broadly being recognized as a main pillar of ocean management in aiming to reverse 70 degradation trajectories of nature in peril (Coleman et al., 2020).

Marine spatial planning (MSP), a focus of numerous international marine policies and agreements 71 (e.g., UK Marine Policy Statement, 2011; European Directive of Maritime Spatial Planning, 2014; 72 73 Marine and Coastal Act of the Parliament of Victoria, Australia, 2018), is a public process aiming to allocate maritime activities within the marine space by minimizing conflicts and maximizing 74 75 sustainability (Ehler & Douvere, 2009; Agardy, 2010; Frazão Santos et al., 2019). MSP is under development or on track to be implemented in 75 countries around the world (half of the world 76 77 countries with territorial waters; Ehler, 2021) and has the potential to expand even further under the EU and UNESCO IOC commitment (EC, 2014; https://ioc.unesco.org/news/european-commission-78 and-unesco-renew-their-joint-efforts-advance-marine-spatial-planning). It is a key tool in the UN 79

B0 Decade of Ocean Science for Sustainable Development (Heymans et al., 2020), supporting the
achievement of sustainability goals around the world.

MSP has roots in marine conservation, with early spatial planning focused on marine protected area 82 site selection and zoning (Agardy, 2010; Vaughan & Agardy, 2020). Yet in the last fifteen years 83 MSP has gone beyond conservation, aspiring to become a multi-objective approach that balances 84 85 ecological and socio-economic goals while delivering conservation outcomes (Shabtay et al., 2019; 86 Gissi et al., 2022). When MSP recognizes ecological systems as multiple interacting elements, it rests on the principles of ecosystem based management (EBM). Such MSP, called 'ecosystem based 87 marine spatial planning' or EB-MSP, focuses management strategies on "how the ecosystem works 88 89 and functions" (Curtin & Prellezo, 2010; UNEP, 2011) by reducing threats and by explicitly supporting the recovery of marine ecosystem functioning (Arkema et al., 2006; Foley et al., 2010). 90 In practice, however, most MSP processes fail to implement EBM and often marginalize conservation 91 92 objectives (Frazão Santos et al., 2021; Fabbrizzi et al., 2023; Reimer et al., 2023). And the explicit strategic inclusion of restoration processes as an objective of MSP is even more rare, accentuating 93 the risk that conserving and restoring marine ecosystems remains an aspiration rather than an 94 95 outcome.

The intersection of MSP, in general terms, with marine restoration strategies has been recommended 96 97 previously (Lester et al., 2020; Fraschetti et al., 2021; Fabrizzi et al., 2023), but no systematic framework exists for achieving this. We contend that MSP that steadfastly stays true to the EBM 98 approach and principles can boost marine restoration. Conversely, explicitly bridging the worlds of 99 MSP and marine restoration - whereby planners guide investment in marine restoration, represents a 100 significant opportunity for MSP to augment its potential for promoting ocean regeneration and the 101 102 continuous delivery of ecosystem services to obtain both conservation and long-lasting socioeconomic outcomes. 103

- 104
- 105

106 2. Envisioning a restoration-focused EB-MSP framework through EBM principles

107 We present a rationale for the proposed perspective and define why and how a marine restorationfocused EB-MSP framework can simultaneously catalyse restoration projects and achieve both 108 109 conservation and blue economy objectives (Gilliland & Laffoley, 2008). An ecosystem based approach can have many dimensions, however we reduce these to the five core principles of EBM 110 pointed by UNEP in its EBM Manual (UNEP, 2011). In shorthand these are: 1) recognizing 111 connections; 2) taking an ecosystem services approach; 3) addressing cumulative impacts; 4) 112 managing for multiple uses; and 5) embracing change, learning, and adapting. Figure 1 summarizes 113 how these five EBM core principles can provide an overarching guide for inserting marine restoration 114 115 in MSP; in other words, we provide tips for an EB-MSP + restoration approach to operationalize the restoration-focused EB-MSP framework. The principles and how they provide insertion points for 116 restoration in marine planning are described in detail in the following sections. 117

118

119 2.1 Recognizing connections

Much of restoration planning as currently practiced is small in scope, and because it is undertaken in 120 a geographically limited scale often fails to acknowledge connectivity, which takes place at broader 121 122 scales. We refer to 'small scale' restoration that focuses on a single habitat and localized human uses 123 in contrast to 'large scale' restoration that covers Large Marine Ecosystems (Sherman & Alexander, 1986) or national waters, inherently encompassing multiple ecosystems (Collie et al., 2013). 124 Connectivity, defined as "the degree to which landscapes and seascapes enable species to move freely 125 126 and ecological processes to function unimpeded" (UNEP, 2019; Balbar & Metaxas, 2019), is central to many ecological processes, including migration of adults and dispersal of juveniles, nutrient fluxes, 127 gene flow, demographic recovery, and movement (Treml & Halpin, 2012; Roberts et al., 2021). EBM 128 recognizes the need to consider ecological connectivity in order to effectively conserve and restore 129 marine ecosystems; when connectivity is not considered, ecological processes can be disrupted 130 131 leading to negative cascading effects on multiple ecosystem components (UNEP, 2011; Laffoley et

5

al., 2019). Some areas are more critical for maintaining ecological connectivity than others because
they differ in their functions as food subsidies, refuges from weather or predators, accessibility to
dispersal pathways, and in numerous other ecological properties that help to shape individual fitness,
population demography, and assemblage composition (Fobert et al., 2019).

Addressing ecological connectivity, by identifying and prioritizing connected sites when planning 136 restoration actions, has been identified as the most promising strategy when the aim is to restore 137 138 species populations (Gilby et al., 2018; Fraschetti et al., 2021; Fabbrizzi et al., 2023). For example, artificial reefs or structures deployed underwater to enhance repopulation of both pelagic and benthic 139 species (e.g., fish, oysters, corals), have been shown to be more effective when networks of these 140 141 sites are designed to optimize the maintenance of connections between areas and species populations, and by considering distance, dispersal mechanisms, currents, etc. (Blouet et al., 2022; Paxton et al., 142 2022; Swam et al., 2022). Similarly, coral reef restoration is most successful when corals are 143 144 strategically outplanted in areas where their larvae dispersal targets a greater number of surrounding reefs, which increases bet-hedging and contributes to the replenishment of the ecosystem beyond the 145 146 outplant site (Frys et al., 2020). Considering connectivity in reef restoration planning can also allow inclusion of other ecosystems that support reef functioning, such as seagrass beds or estuarine 147 ecosystems. A focus on connections can also highlight potential synergies between land and reef 148 149 restoration projects, as is the case of reforestation actions that improve water quality benefiting coral reefs (Suárez-Castro et al., 2021). 150

Maintaining the links between diverse habitats across wide seascapes is critical for the population dynamic of many mobile species (McMahon et al., 2012). Biophysical models able to predict larval dispersal, for instance, combine different environmental datasets in order to describe connectivity and predict areas most likely connected (Jonsson et al., 2020; Sciascia et al., 2020; Swam et al., 2022). EB-MSP relies heavily on such modelling approaches and can also guide allocation of restoration initiatives. Moreover, planning that takes into account connectivity can allow restoration initiatives to operate at large scales, which is essential to increase the chance of ocean regeneration success (Duarte et al., 2015; Fabbrizzi et al., 2023). An example is provided in the large restoration program being undertaken in the Solent (UK), where active restoration of oyster reefs (Collins, 2022) is coupled to passive restoration provided by restrictions on bottom trawling and dredging, along with restoration of riparian habitats that result in improved water quality in estuarine / nearshore waters. A melded EB-MSP and restoration framework thus makes it possible to optimize connectivity by considering the big picture and the restoration opportunities at the landscape / seascape scale.

It has been showed that where multiple interconnected habitats are co-restored, their positive interactions mutually benefit each other to stabilize and even accelerate ecosystem recovery (McAfee et al., 2022a). This has been well documented in case-studies from South Australia, which showed how constructed boulder reefs provide opportunities to co-restore shellfish and kelp forests, while stabilizing sediment for seagrass recovery. Other case describes re-introducing tidal flows into tidallyrestricted areas (e.g., via tidal gates, sea walls), providing the opportunity to restore mosaics of connected intertidal seagrass, mangrove and saltmarsh habitats (McAfee et al., 2021).

Resting on the connectivity principle, a restoration focused-EB-MSP framework approach could 171 172 guide a multi-habitat approach and also the prioritization of restoration interventions to be explicitly included in zoning of marine areas, i.e the allocation of ocean space for various uses, protection modes 173 and managing objects (Agardy, 2010). EB-MSP can direct active, assisted, or reconstructive 174 restoration (Atkinson & Bonser, 2020) (see Table 1 for terminology) that enhances connectivity 175 among populations and habitats. Active restoration includes engineered replanting, shoreline or reef 176 stabilization, pollution controls, species reintroductions, removal of non-native invasive species, and 177 178 other deliberate actions by managers meant to either restore habitats that were previously present, or enhance degraded habitats to make them more resilient to human and climate change pressures. Many 179 180 options are available to planners, however, choosing particular passive or active restoration measures requires a case-by-case cost-benefit analysis, considering the trade-offs of the two approaches and 181 their direct and indirect costs (e.g., longer recovery time and vigilance costs in natural restoration 182 strategies and material and labour costs in active restoration strategies; Zahawi et al., 2014). With 183

7

restoration-focused EB-MSP, passive and active restoration which restores connectivity can lead to
long-lasting outcomes even over large geographic scales (Diefenderfer et al., 2021).

Marine plans that incorporate restoration measures in particular zones at various scales (within, for 186 instance, protected areas, at the scale of subnational regions, or even at wider national scales) can 187 make them more effective. Additionally, implicitly linking local restoration interventions with 188 broader scale restoration policies through a spatially nested and coordinated restoration strategy is 189 recommended for improved delivery of both ecological and socio-economic benefits (Gilby et al., 190 2021). Indeed, this would prioritize localized, small scale and interconnected restoration actions 191 within a wide area – e.g., by identifying interconnected sites to facilitate the simultaneous restoration 192 193 of oyster reefs, seagrasses and mangroves. This can optimize restoration investments to obtain the greatest ecological and socio-economic benefits across a wide area - e.g., improved habitat that 194 supports fish populations and fisheries at the scale of multiple estuaries (Gilby et al., 2021). 195

196

197 **2.2 Taking an ecosystem services approach**

EBM that aims to maintain the delivery of ecosystem services (ES), or the goods and services 198 provided by the diversity of species and their functions (for instance food provision, shoreline 199 stabilization and buffering of land from storms, hydrological balances, pest control, and carbon 200 201 storage; Normile, 2010), can create public support for management actions. A prerequisite for ensuring the capacity of ecosystems to deliver ES is the maintenance of their ecological processes 202 and properties. Under this perspective, an ecosystem can be considered recovered when the 203 204 biodiversity it accomodated before degradation is restored and it reacquires its capability of delivering 205 ES (Orth et al., 2020).

Although there may be considerable uncertainty about how quickly full recovery of ES can be accomplished, ES studies can be used within a restoration-focused EB-MSP framework to guide the achievement of conservation and related human wellbeing. When spatial multi-criteria decision frameworks are applied to prioritize and select sites to be restored, planning interventions have the potential to enhance ES delivery and catalyse positive biodiversity and socio-economic outcomes(Pittman et al., 2022).

Operationalizing the ES approach to serve restoration-focused EB-MSP may require targeted ES 212 assessments to ascertain conditions prior to ecosystem degradation, in order to provide baseline 213 information. ES assessment helps set thresholds, identify baseline conditions, and guide restoration 214 to meet established targets. Such assessments can feed decisions based on explicitly identified trade-215 offs, to pinpoint the most beneficial planning solution (White et al., 2012). Information coming from 216 such assessments should be considered during the initial phase of MSP, which is focused on creating 217 the knowledge framework that will guide the building of the spatial plan. Subsequently, once 218 219 restoration has been initiated, the rates at which ES are delivered should be monitored for long enough to provide meaningful information regarding restoration success (Gómez-Baggethun et al., 2019). 220 Knowing which areas previously delivered important ES, or would in the future if restored, can help 221 222 prioritize site selection for spatial zoning and make explicit the benefits of restoration. Thus taking an ES approach not only promotes more ecologically sustainable planning, it also can create impetus 223 224 for restoration and the enhanced ES flows that such restoration would bring.

In restoration-focused EB-MSP, economies of scale can be achieved if single restoration projects are 225 226 strategically integrated within larger MSP programmes and are planned and designed in a way to be 227 physically and functionally linked to underpin ecological connectivity, and thus ES delivery. Thus, although restoration can be costly, especially when performed at large scales, when appropriately 228 planned benefits can outweigh costs (De Groot et al., 2013; Strassburg et al., 2019). Evaluations that 229 230 confirm this include a successful large-scale seagrass restoration project along the mid-Atlantic coast of the US in which the numerous ES recovered, delivering benefits beyond the scale of the restored 231 area (Orth et al., 2020). These benefits include improved water quality and ample habitat recovery 232 with consequent increasing of fish populations to support the fishery and an augmented capacity of 233 nitrogen and carbon sequestration to abate climate change effects. 234

Restoration-focused EB-MSP can also guide investment in restoring blue carbon ecosystems. In some 235 236 cases such restoration combines recovery of mangrove, salt marsh, or seagrass with active restoration involving human-engineered infrastructures, such as submerged breakwaters that mimick coral reef 237 structures to reduce shoreline erosion (Stender et al., 2021). Such restoration has been shown to 238 enhance opportunities for sustainable energy production (Vanderklift et al., 2019; Thiele et al., 2020), 239 240 and at the same time can increase coastal protection in the face of catastrophic climate change effects. 241 Moreover, considering the interconnected multi-habitat approach also gives more chance to increase the connectivity between carbon source and sink habitats (e.g., between adjacent kelp forests and 242 seagrass, respectively), and through enhanced organic carbon transfer and burial, maximize the 243 244 carbon sequestration potential (Smale et al., 2018; McAfee et al., 2022a).

Balancing the trade-offs beween ecological restoration benefits and economic interests is critical. Efforts should be dedicated to analyse the complex interactions among multiple ES and human needs, and set threshold values for ecosystem management (Wang et al., 2022). We suggest that a restoration-focused EB-MSP could drive all these in a unique coordinated framework.

249

250 2.3 Addressing cumulative impacts

Restoration efforts which consider only single stressors and limit single sector / use-impact (e.g., 251 252 limiting pollutant inputs or overfishing), may fail because degradation of an ecosystem is usually caused by multiple activities and pressures (Brown et al., 2013; Gissi et al., 2021). Considering and 253 analysing all sources of pressures in a multi-use context, as EBM does, is essential in identifying 254 255 those that are the main cause of ecosystem degradation trajectory (Merovich & Petty, 2007; Teichert et al., 2016). EB-MSP combines studies on cumulative impacts and ES provisioning hotspots and 256 257 such an approach can allow prioritization of restoration interventions and increase their efficiency 258 and success rates (Allan et al., 2013). Indeed, once cumulative impacts are addressed, the stage is set for natural recovery or restoration actions through integrated and multi-use management (Paschke et 259 al., 2019). One example is provided by Farella and colleagues (2020) who supported the development 260

of MSP in Emilia-Romagna Region (Italy) through the application of an ad-hoc MSP modelling framework. They simultaneously analysed cumulative pressures and their effects on the capability of habitats and species populations inhabiting the territorial waters of the region to deliver ES. The study provided practical guidelines and delinated some areas of conservation to support the restoration of overexploited fish populations and of species of conservation priority.

266 Map-based threats assessments coupled to restoration science can help identify areas of greatest 267 restoration need, guide the prioritization of threats to be addressed, and steer implementation of a portfolio of restoration actions (Allan et al., 2013, Neeson et al., 2016). Sophisticated tools to inform 268 EB-MSP and support decision makers in managing multiple anthropogenic uses and impacts exist 269 270 (e.g., predictive models as habitat suitability models - HSM, Marxan and Bayesian belief networks for scenario analyses, InVEST for ES assessment and trade-off), and can also incorporate knowledge 271 272 on vulnerability and natural recovery capability of marine ecosystems in multi-use contexts (e.g., 273 Cumulative Effects Assessment - CEA – models; Andersen et al., 2013; Menegon et al., 2018; see the above cited example by Farella et al., 2020). Another example that combines different modelling 274 275 approaches is provided by Uusitalo et al. (2016), who explored the effect of diverse scenarios of 276 fishing pressure and nutrients inputs reduction in the Baltic Sea to inform the management of cumulative pressures to allow the recovery of Baltic ecosystems and fish populations. 277

By considering cumulative effects, EB-MSP can drive multi-sector cooperation and ultimately 278 effective management (Guerry et al., 2012; Halpern et al., 2019). That said, knowledge on 279 vulnerability and natural recovery of species and habitats is still limited, complicated by multiple 280 281 factors, including differences in the type and severity of the various impacts (Duarte et al., 2015 and references therein), which increases uncertainty of model outputs. However, field studies and 282 manipulative experiments are filling knowledge gaps and are being incorporated in EB-MSP 283 284 decision-support tools to increase their reliability (see for instance Kotta et al., 2019). Furthermore, advances in technology and transdisciplinary research such as use of large-scale satellite data 285 (Klemas, 2013; Ouellette & Getinet, 2016) and machine learning algorithms can increase the efficacy 286

and efficiency of restoration (Zellmer et al., 2019). Where knowledge gaps remain unfilled, an EBMSP + restoration approach should acknowledge uncertainty and apply the precautionary principle,
common practice for EB-MSP (Manea et al., 2020) and fundamental in restoration practices (Gann
et al., 2019).

291

292 **2.4. Managing for multiple uses**

Managing for multiple uses under EBM implies cross-sectoral coordination. To do this effectively, 293 cumulative impacts should be systematically assessed, along with analysis of capacity for addressing 294 295 these impacts with existing institutions, policies, regulations, and norms (Agardy, 2010). EB-MSP integrates this holistic management to respond to multiple stressors and maximize efficiencies 296 297 through coordinated and cooperative actions. Such an approach not only prevents further 298 environmental degradation, but also sets the stage for natural recovery and restoration. Management 299 that is supported by EB-MSP can (and should, wherever possible) include a blueprint for both passive and active restoration (see Table S1 for possible restoration measures). Marine restoration projects 300 301 should likewise include multi-use management as a goal (Paschke et al., 2019), focusing on managing the pressures that have caused degradation or impede recovery. 302

With all this in mind, the first strategy of an EB-MSP + restoration approach will likely be passive 303 restoration (here used as synonym of natural restoration), in which pressures, wherever their 304 provenance, are reduced to promote the natural recovery of ecosystems and of ecological processes 305 306 (Chazdon et al., 2021). Through management measures, EB-MSP can mitigate anthropogenic pressures in areas where passive recovery will require time, for instance by imposing fishery bans or 307 preventing coastal infrastructure development that hinders species recolonization. Furthermore, EB-308 309 MSP can guide siting and establishment of MPAs by highlighting which areas / habitats are ecologically most critical to protect. With protected areas utilized in concert with effective 310 management of multiple uses, socio-economic development and ecosystem recovery can be in 311 balance (Trouillet & Jay, 2021). 312

Since EB-MSP should be coordinated at diverse governmental and / or institutional levels, restoration 313 314 within an EB-MSP framework might need to extend beyond jurisdictional boundaries, requiring coordination between different political institutions (van Tatenhove et al., 2021). At the same time, 315 316 EB-MSP is multi-sectoral and aims to build sectoral integration and coordination among multiple governance and management levels (Grip & Blomqvist, 2021), thus it can only be effective by 317 actively involving a variety of stakeholders. Planners should involve stakeholders to understand their 318 319 priorities, allow co-creation of plans, and to secure their support for management and restoration policies, regulations, and interventions. 320

McAfee and colleagues (2022b) provided an exemplary case of successful large-scale restoration 321 322 project of the disappeared oyster reefs in Australia. The iniative was triggered by awareness of the numerous economic and ecological benefits that this habitat could have brought if restored. Through 323 the practical and financial support of a multitude of stakeholders (e.g., scientists, restoration 324 325 practicionnaires, policymakers, managers, local communities), multidisciplinary teams, and the coordination of both public and private sectors (e.g., research, aquaculture, recreational fishing) over 326 327 a six year period (2015-2021), a substantial investment of funding has enabled 35 restoration projects. 328 This large-scale restoration initiative is ongoing and anticipates additional local projects supported by the increased motivation of the multiple parties involved. 329

330 Restoration embedded into EB-MSP can thus benefit many stakeholders and can lead to long-lasting positive outcomes that will expand stakeholder support, especially when done at large scales 331 (Bayraktarov et al., 2020). A participatory approach to restoration can emphasize the human-332 333 ecosystem relationship, thereby increasing the willingness of stakeholders to support or even engage in restoration. Volunteers can be fundamental for supporting restoration, especially when active 334 restoration is done across large geographies (Orth et al., 2020). Stakeholders can also be engaged in 335 evaluation of the success of restoration initiatives, by providing data on benefits flows and perceptions 336 about the level to which the ES delivery has been recovered. This is particularly useful for gauging 337 recovery of cultural ES, which are difficult to measure because they are intangible but directly linked 338

to human perception and health (Pouso et al., 2020). Furthermore, involving stakeholders can reduce
the risk that unrealistic expectations are set and then not met, generating mistrust in restoration
initiatives (Kodikara et al., 2017).

342

343 **2.5 Embracing change, learning, and adapting**

Embracing change is imperative for achieving positive outcomes and improve prospects for success of future initiatives (Ellison et al., 2020). Part of embracing change is recognizing where change is happening and why, and learning from the application of management over time.

Detailed information on past and present distribution and condition of ecosystems, causes of 347 348 degradation and related footprints, conditions of the surrounding environments, and changes the area is likely to face in the future is all needed in a restoration-focused EB-MSP framework. Planning is 349 built on baseline inventories of information from diverse knowledge sources, and involves a broad 350 351 range of experts including local knowledge (Lombard et al., 2019). Inclusion of local knowledge can help reconstruct pre-disturbance conditions necessary for setting final goals (Gann et al., 2019). The 352 integration of indigenous and scientific knowledge is fundamental especially where published 353 information is scant. To deal with knowledge limitations planners and restoration experts can also 354 rely on accessible satellite data on distribution of habitats and environmental conditions (e.g., 355 356 temperature conditions, clorophyll concentration; Fingas, 2019). Another strategy can be integrating the existing monitoring frameworks at both local and large scales - e.g., the European Marine Strategy 357 Framework Directive (Maccarrone et al., 2015; Abramic et al., 2020) within restoration-focused EB-358 359 MSP, to optimize multi-scale monitoring efforts and obtain long-term data sets (Manea et al., 2022). The integration of multiple knowledge sources is thus crucial for adapting management and 360 restoration as conditions change. A practical example on the potential of integrating local monitoring 361 with remote sensing data is provided by McClenachan et al. (2020) in which they combined these 362 diverse data sources to assess the restoration success of small-size shoreline and oyster reef 363 restoration projects at the scale of the ecosystem of Mosquito Lagoon, Florida, USA. 364

Adaptive management relies on the lessons learned from monitoring and evaluating the performance 365 of implemented management strategies and measures over time, while accounting for possible 366 environmental shifts due to climate change (Thom, 2000; Ellison et al., 2020). Adaptive restoration 367 similarly relies on systematic and long-lasting monitoring to track pace of recovery and regeneration. 368 Since the time needed for an ecosystem to recover after restoration interventions varies (Kodikara et 369 370 al., 2017; Boström-Einarsson et al., 2020), and because species and habitats present different recovery capabilities (Bekkby et al., 2020), course corrections may be necessary. EB-MSP can and should 371 respond to the highly dynamic nature of marine systems, where environmental conditions are 372 constantly shifting. Climate change drives many of these shifts (Roberts et al., 2020), and discussions 373 374 on climate change mitigation and adaptation in coastal areas are acelerating (Flannery et al., 2020), focused on how to make MSP an effective instrument to minimize climate impacts, support climate 375 adaptation, and allow mitigation actions (Frazão Santos et al., 2019; 2020). Restoration-focused EB-376 377 MSP can build on conservation planning for climate change adaptation; key strategies under this approach includes vulnerability assessments, supporting ecosystem resilience, protecting climate 378 379 refugia and predicting species and ecosystem shifts (Wilson et al., 2020). An EB-MSP + restoration 380 approach could strengthen nature based solutions (Gijsman et al., 2021) that enhance blue carbon, as well as restoration of ecosystems that abate coastal vulnerability, eliminate cumulative pressures in 381 382 climate refugia (i.e., passive restoration), and foster active restoration projects in areas accommodating shifts. 383

384

385 **3. Conclusions**

To date, MSP and restoration have been on separate tracks, ignoring both the potential synergies and co-benefits provided by linking the two. MSP is a powerful tool to support marine conservation (Ehler & Douvere, 2009; Reimer et al. 2023), but neither effective management of marine use nor conservation can match the pace of marine environmental degradation (Coleman et al., 2020). Since

most marine ecosystems are no longer pristine, MSP that does not give space, quite literally, for 390 391 restoration is unlikely to result in desired outcomes of maintaining ES delivery under a blue economy. The elements for a transformative shift in how to achieve ocean regeneration (summaried in Figure 392 2) are all here today, but disparate and unsystematic planning is unable to capitalize on the immense 393 opportunities that exist for this transformation to happen. Across the world today MSP is undertaken 394 at various scales to achieve diverse objectives (Ehler, 2021), mainly targeting blue growth as a goal, 395 396 with only marginal and insignificant attention paid to the environmental and social sustainability of 397 that economic growth (Frazão Santos et al., 2021). MSP processes that do base planning on EBM principles are increasing but rarely strive to enhance ocean health. At the same time, the world is 398 399 witnessing a proliferation of active and passive restoration projects, primarily aimed at rebuilding one habitat type across relatively small geographical scales (Fraschetti et al., 2021; Fabbrizzi et al., 2023). 400 At the moment, these restoration projects tend to be opportunistic rather than being a strategic part of 401 402 geographically large, integrated marine plans.

The enabling conditions that could allow the needed transformation to occur include a solid 403 404 understanding of the ecology of marine and coastal systems, accounting for the identification of 405 ecologically critical areas to protect or restore, and a solid understanding of the connectivity between different elements of the ecosystem. Alongside this approach, communications about the values of a 406 407 healthy ocean (taking an ecosystem services perspective), and institutional integration that allows effective management of all uses of ocean space and resources (considering cumulative impacts and 408 managing for multiple use) are needed. The transformation is further made possible by learning from 409 410 trialing management, requiring a continuous linking of restoration with initial planning and with the 'replanning' that is needed for adaptive management. 411

Restoration-focused EB-MSP can support the restorative continuum, i.e. both active and passive restoration interventions, with the former designed to catalyze the latter. Passive restoration mediated by EB-MSP with a wide scope is fundamental because local restoration projects, even if supported by solid scientific understanding and post-care and monitoring actions, can fail in the face of multiple anthropogenic pressures (Diefenderfer et al., 2021). Thus, the strategic easing of pressures on
ecosystems led by EB-MSP can predispose ecosystems for natural recovery, leading to restored
connectivity and functionalities, and simultaneously fostering the effectiveness of assisted or
reconstructive restoration projects.

Utilising a restoration-focused EB-MSP framework, a potential paradigm shift in marine planning outlined in Figure 2, will ensure that interventions end up creating truly resilient 'systems', instead of the façade of reconstructed habitats. Such successful interventions will build on ecological science, oceanography and hydrology, as well as user knowledge, to create understanding of ecosystem connectivity, vulnerability and recovery rates (Danovaro et al., 2021; Fabbrizzi et al., 2023). Planners that consider all the important elements of marine ecosystems, at various ecological levels and spatial scales, through restoration-focused EB-MSP will see their plan lead to durable outcomes.

427

As a summary, a list of priority main actions that can be taken in an EB-MSP restoration framework:
- put in place passive restoration as a first strategy to abate pressures wherever their provenance to
promote the natural recovery of ecosystems;

- establish active restoration measures as a second strategy following the logic of ecological
connectivity and linking local restoration interventions with broader scale restoration policies and
within spatial plans in concert with MPA designation;

- adopt the ES approach to guide the optimization of conservation and restoration measures and the
setting of thresholds, starting from previous ES assessment and trough the identification of baseline
conditions and trade-offs;

- apply the modelling framework on which MSP relies with a view to identifying restoration measures
to be integrated into spatial plans and integrate empirical scientific knowledge from restoration
ecology into the models;

- coordinate restoration at multiple governmental, and thus geographical, scales;

17

- engage with stakeholders in defining objectives, fund raising strategies, monitoring and evaluating
of restoration performance;

- support and take advantage of existing monitoring frameworks integration also for tracking
restoration success;

- rely on the integration of multiple knowledge sources for building the initial assessment of
environmental conditions and for laying the foundations for marine restoration;

- address climate change by: investing in nature based solutions that favor blue carbon ecosystems
and restoration of ecosystems that abate coastal vulnerability, eliminating cumulative pressures in
climate refugia (i.e., passive restoration), and fostering active restoration projects in areas where
habitats is expected to shift.

451

We suggest that adopting an EB-MSP + restoration approach and demonstrating restoration-focused EB-MSP will spur replication and a wide upscaling of effective marine management for ocean regeneration. MSP that fully incorporates marine restoration through EBM within its goals has the potential to support and boost the recovery of ecological structure, function, and resilience, while enabling enhanced ES delivery that maximizes socio-ecological and economic benefits-sharing. Multiple restorative actions can thus be implemented within EB-MSP to contemporarily boost sustainable economy and ocean regeneration.

In sum, restoration-focused EB-MSP is desirable for numerous reasons: i) it allows scaling-up of 459 marine restoration advancements within large-scale planning mechanisms; ii) it delivers important 460 461 means for sustainable blue economy and helps meet conservation objectives; and iii) it can lead to integrated MSP, able to better manage ecosystems across biomes and the pressures derived from 462 climate change. Recovery of marine life, and broader ocean regeneration, can only be achieved if 463 immediate strategic action is taken (Duarte et al., 2020). Restoration-focused EB-MSP can be the 464 vehicle to accomplish this, allowing marine life to recover from centuries of impact and promoting a 465 466 more sustainable way for humankind to benefit from the global ocean.

18

467

468 **Declaration of interests**

469 The authors declare no competing interests.

470

471 Acknowledgments

472 EM acknowledges the support of the European Maritime and Fisheries Fund of the European Union through the projects SUPREME "Supporting maritime spatial Planning in the Eastern 473 Mediterranean", grant no. EASME/EMFF/2015/ 1.2.1.3/01/S12.742087 and SIMWESTMED 474 "Supporting Implementation of Maritime Spatial Planning in the Western Mediterranean region", 475 476 grant no. EASME/EMFF/2015/ 1.2.1.3, and the Marie Curie Actions through the project RESTORE, project number 101062275; LB was supported by the European Union's Horizon Europe research 477 478 and innovation programme through the project MSP4BIO "Improved Science-Based Maritime Spatial Planning to Safeguard and Restore Biodiversity in a coherent European MPA network", grant 479 no. 963646 and the "National Biodiversity Future Center - NBFC" funded under the National 480 Recovery and Resilience Plan (NRRP), by the European Union - NextGenerationEU, project code 481 CN_00000033. This study reflects only the authors' views and not those of the European Union. 482

483

484 Credit authorship contribution statement

EM, TA and LB: Conceptualization, Methodology, Investigation, Writing - Original Draft, Writing Review & Editing, Visualization, Supervision.

487

488 **References**

-Abelson, A., Reed, D.C., Edgar, G.J., Smith, C.S., Kendrick, G.A., Orth, R.J., et al. (2020).

490 Challenges for restoration of coastal marine ecosystems in the Anthropocene. Frontiers in Marine

491 Science, 7: 892. https://doi.org/10.3389/fmars.2020.544105

- 492 -Abramic, A., Nogueira, N., Sepulveda, P., Cavallo, M., Fernández-Palacios, Y., Andrade, C., et al.
- 493 (2020). Implementation of the marine strategy framework directive in macaronesia and synergies
- 494 with the maritime spatial planning process. Marine Policy, 122: 104273.
- 495 https://doi.org/10.1016/j.marpol.2020.104273

- 496 -Agardy, T. (2010). Ocean Zoning: Making Marine Management More Effective. Earthscan, London.
- Allan, J.D., McIntyre, P.B., Smith, S.D.P., Halpern, B.S., Boyer, G.L., Buchsbaum, A., et al. (2013).
- 498 Joint analysis of stressors and ecosystem services to enhance restoration effectiveness. *Proceedings*
- 499 of the National Academy for Sciences, U.S.A., 110: 372–377.
 500 https://doi.org/10.1073/pnas.1213841110
- Andersen, J.H., Stock, A., Heinänen, S., Mannerla, M. & Vinther, M. (2013). Human uses, pressures
- and impacts in the eastern North Sea. Aarhus University, DCE-Danish Centre for Environment andEnergy, 136.
- -Arkema, K.K., Abramson, S.C., & Dewsbury, B.M. (2006). Marine ecosystem-based management:
- from characterization to implementation. Frontiers in Ecology and the Environment, 4: 525-532.
- 506 https://doi.org/10.1890/1540-9295(2006)4[525:MEMFCT]2.0.CO;2
- -Aronson, J., Floret, C., Le Floc'h, E., Ovalle, C. & Pontanier, R. (1993). Restoration and rehabilitation of degraded ecosystems in arid and semi-arid lands. I. A view from the south.
- 509 *Restoration Ecology*, 1(1), 8-17. https://doi.org/10.1111/j.1526-100X.1993.tb00004.x
- -Aronson, J. & Le Floc'h, E. (1996). Vital landscape attributes: missing tools for restoration ecology.
- 511 Restoration Ecology, 4: 377-387. https://doi.org/10.1111/j.1526-100X.1996.tb00190.x
- -Aronson, J., Goodwin, N., Orlando, L., Eisenberg, C., & Cross, A. T. (2020). A world of
- 513 possibilities: six restoration strategies to support the United Nation's Decade on Ecosystem
- 514 Restoration. *Restoration Ecology*, 28(4), 730-736. https://doi.org/10.1111/rec.13170
- -Atkinson, J., & Bonser, S.P. (2020). "Active" and "passive" ecological restoration strategies in meta-
- analysis. *Restoration Ecology*, 28(5), 1032-1035. https://doi.org/10.1111/rec.13229
- Balbar, A. C., & Metaxas, A. (2019). The current application of ecological connectivity in the design
- 518 of marine protected areas. *Global Ecology and Conservation*, 17: e00569.
- 519 https://doi.org/10.1016/j.gecco.2019.e00569

- Bayraktarov, E., Brisbane, S., Hagger, V., Smith, C. S., Wilson, K. A., Lovelock, C. E., et al. (2020).
- 521 Priorities and motivations of marine coastal restoration research. Frontiers in Marine Sciences, 7:
- 522 484. https://doi.org/10.3389/fmars.2020.00484
- -Bradshaw, A.D. (2002). Introduction and philosophy. In Handbook of Ecological Restoration.
- 524 Principles of Restoration, vol. 1, M.R., Perrow, A.J., Davy, ed. (Cambridge University Press) pp. 3-
- 525 9.
- -Bekkby, T., Papadopoulou, N., Fiorentino, D., McOwen, C.J., Rinde, E., Boström, C., et al. (2020).
- 527 Habitat features and their influence on the restoration potential of marine habitats in Europe. *Frontiers*
- 528 *in Marine Sciences*, 7: 184. https://doi.org/10.3389/fmars.2020.00184
- -Blouet, S., Bramanti, L. & Guizien, K. (2022). Artificial reefs geographical location matters more
- than shape, age and depth for sessile invertebrate colonization in the Gulf of Lion (NorthWestern
- 531 Mediterranean Sea). *Peer Community Journal*, 2: e24. DOI: 10.24072/pcjournal.107
- -Boström-Einarsson, L., Babcock, R.C., Bayraktarov, E., Ceccarelli, D., Cook, N., Ferse, S.C.A., et
- al. (2020). Coral restoration–A systematic review of current methods, successes, failures and future
- directions. *PloS ONE*, 15: e0226631. https://doi.org/10.1371/journal.pone.0226631
- -Brown, C.J., Saunders, M.I., Possingham, H.P. & Richardson, A.J. (2013). Interactions between
- 536 global and local stressors of ecosystems determine management effectiveness in cumulative impact
- mapping. Diversity and Distributions, 20: 538–546. <u>https://doi.org/10.1111/ddi.12159</u>
- 538 -CBD (Convention on Biological Diversity) 2022. Final text of Kunming-Montreal Global
- 539 Biodiversity Framework. CBD/COP/15/L25. Montreal Canada
- -Chazdon, R.L., Falk, D. A., Banin, L.F., Wagner, M., Wilson, S.J., Grabowski, R.C., & Suding, K.N.
- 541 (2021). The intervention continuum in restoration ecology: rethinking the active–passive dichotomy.
- 542 *Restoration Ecology*, e13535. https://doi.org/10.1111/rec.13535
- -Clewell, A.F. & Aronson, J. (2013). Ecological restoration: principles, values, and structure of an
- 544 emerging profession. (Island Press).

- -Cliquet, A., Telesetsky, A., Akhtar-Khavari, A., & Decleer, K. (2022). Upscaling ecological
 restoration: toward a new legal principle and protocol on ecological restoration in international
 law. *Restoration Ecology*, *30*(4), e13560. https://doi.org/10.1111/rec.13560
- -Coleman, M.A., Wood, G., Filbee-Dexter, K., Minne, A.J., Goold, H.D., Verges, A., et al. (2020).
- 549 Restore or redefine: future trajectories for restoration. Frontiers in Marine Science, 7: 237.
- 550 https://doi.org/10.3389/fmars.2020.00237
- -Collie, J. S., Beck, M. W., Craig, B., Essington, T. E., Fluharty, D., Rice, J., & Sanchirico, J. N.
- 552 (2013). Marine spatial planning in practice. Estuarine, Coastal and Shelf Science, 117: 1-11.
- 553 https://doi.org/10.1016/j.ecss.2012.11.010
- -Collins, K.J., Hauton, C., Jensen, A.J., and Mallinson, J.J. 2022. Solent oyster enhancement reef
- concrete colonisation trials. *IOP Conference Series: Materials Science and Engineering*, *1245*:012012. doi:10.1088/1757-899X/1245/1/012012
- -Curtin, R., & Prellezo, R. (2010). Understanding marine ecosystem based management: a literature
 review. *Marine Policy*, *34*: 821-830. https://doi.org/10.1016/j.marpol.2010.01.003
- -Danovaro, R., Aronson, J., Cimino, R., Gambi, C., Snelgrove, P.V. & Van Dover, C. (2021). Marine
 ecosystem restoration in a changing ocean. *Restoration Ecology*, 29: e13432.
 https://doi.org/10.1111/rec.13432
- -De Groot, R.S., Blignaut, J., Van Der Ploeg, S., Aronson, J., Elmqvist, T. & Farley, J. (2013).
- 563 Benefits of investing in ecosystem restoration. Conservation Biology, 27: 1286-1293.
- 564 https://doi.org/10.1111/cobi.12158
- -Diefenderfer, H.L., Steyer, G.D., Harwell, M.C., LoSchiavo, A.J., Neckles, H.A., Burdick, D.M., et
- al. (2021). Applying cumulative effects to strategically advance large-scale ecosystem restoration.
- 567 Frontiers in Ecology and the Environment, 19(2), 108-117. https://doi.org/10.1002/fee.2274
- -Diz, D., Johnson, D., Riddell, M., Rees, S., Battle, J., Gjerde, K., et al. (2018). Mainstreaming marine
- biodiversity into the SDGs: the role of other effective area-based conservation measures (SDG 14.5).
- 570 Marine Policy, 93: 251-261. https://doi.org/10.1016/j.marpol.2017.08.019

- -Duarte, C.M., Borja, A., Carstensen, J., Elliott, M., Krause-Jensen, D. & Marbà, N. (2015).
- 572 Paradigms in the recovery of estuarine and coastal ecosystems. *Estuaries and Coasts, 38*: 1202-1212.
- 573 https://doi.org/10.1007/s12237-013-9750-9
- -Duarte, C.M., Agusti, S., Barbier, E., Britten, G.L., Castilla, J.C., Gattuso, J-P., et al. (2020).
 Rebuilding marine life. *Nature*, *580*: 39-51. https://doi.org/10.1038/s41586-020-2146-7
- -EC, 2008. European Commission. Directive 2008/56/EC of the European Parliament of the Council
- 577 of 17 June 2008 establishing a framework for community action in the field of marine environmental
- 578 policy (Marine Strategy Framework Directive). Official Journal of the European Union, L164 (2008),
- 579 p. 19±40.
- -EC, 2014. European Commission. Directive 2014/89/EC of the European Parliament of the Council
 of 23 July 2014 establishing a framework for maritime spatial planning. Official Journal of the
 European Union.
- -EEC, 1992. European Commission. Council Directive 92/43/EEC of May 21 1992 on the conservation of natural habitats and of wild fauna and flora. Official Journal of the European Communities.
- -Ehler, C. & Douvere, F. (2009). Marine Spatial Planning: a step-by-step approach toward ecosystem-
- 587 based management. Intergovernmental Oceanographic Commission and Man and the Biosphere
- 588 Programme. IOC Manual and Guides No. 53, ICAM Dossier No. 6. Paris: UNESCO.
- -Ehler, C.N. (2021). Two decades of progress in Marine Spatial Planning. Marine Policy, 132:
- 590 104134. https://doi.org/10.1016/j.marpol.2020.104134
- -Elliott, M., Burdon, D., Hemingway, K. L. & Apitz, S. E. (2007). Estuarine, coastal and marine
- 592 ecosystem restoration: confusing management and science–a revision of concepts. *Estuarine, Coastal*
- 593 and Shelf Science, 74: 349-366. https://doi.org/10.1016/j.ecss.2007.05.034
- -Ellison, A.M., Felson, A.J. & Friess, D.A. (2020). Mangrove Rehabilitation and Restoration as
- 595 Experimental Adaptive Management. Frontiers in Marine Science, 7: 327.
- 596 https://doi.org/10.3389/fmars.2020.00327

- -Emu Ltd. (2004). Marine Aggregate Restoration and Enhancement: Strategic Policy Overview.
- 598Report No. 04/J/01/06/0548/0437. Emu Ltd. on behalf of The British Marine Aggregate Producers
- 599 Association, the Crown Estate and English Nature.
- -EU, 2019. COM(2019) 640 final. Communication from the Commission to the European Parliament,
- 601 the European Council, the Council, the European Economic and Social Committee and the Committee
- 602 of the Regions. The European Green Deal.
- -Fabbrizzi, E., Giakoumi, S., De Leo, F., Tamburello, L., Chiarore, A., Colletti, A., et al. (2023). The
 challenge of setting restoration targets for macroalgal forests under climate changes. *Journal of Environmental Management, 326:* 116834. https://doi.org/10.1016/j.jenvman.2022.116834
- -Farella, G., Menegon, S., Fadini, A., Depellegrin, D., Manea, E., Perini, L. & Barbanti, A. (2020).
- 607 Incorporating ecosystem services conservation into a scenario-based MSP framework: An Adriatic
- 608
 case
 study.
 Ocean
 and
 Coastal
 Management,
 193:
 105230.

 609
 https://doi.org/10.1016/j.ocecoaman.2020.105230
 htttps://doi.001
- 610 Fingas, M. (2019). Remote sensing for marine management. In World seas: An environmental
- evaluation (pp. 103-119). Academic Press. https://doi.org/10.1016/B978-0-12-805052-1.00005-X
- -Flannery, W., Toonen, H., Jay, S. & Vince, J. (2020). A critical turn in marine spatial planning.
- 613 *Maritime Studies*, 19: 223-228. https://doi.org/10.1007/s40152-020-00198-8
- -Fobert, E.K., Treml, E.A. & Swearer, S.E. (2019). Dispersal and populationconnectivity are
- 615 phenotype dependent in amarine metapopulation. *Proceedings of the Royal Society B*, 286: 20191104.
- 616 http://dx.doi.org/10.1098/rspb.2019.1104
- -Foley, M.M., Halpern, B.S., Micheli, F., Armsby, M.H., Caldwell, M.R., Crain, C.M., et al. (2010).
- Guiding ecological principles for marine spatial planning. *Marine Policy*, *34*: 955-966.
 https://doi.org/10.1016/j.marpol.2010.02.001
- -Fraschetti, S., McOwen, C., Papa, L., Papadopoulou, N., Bilan, M., Boström, C., et al. (2021). Where
- 621 is more important than how in coastal and marine ecosystems restoration. Frontiers in Marine
- 622 Science, 8: 626843. https://doi.org/10.3389/fmars.2021.626843

- -Frazão Santos, C. F., Ehler, C. N., Agardy, T., Andrade, F., Orbach, M. K., & Crowder, L. B. (2019).
- Marine spatial planning. In World seas: an environmental evaluation, C. Sheppard, ed. (Academic
 Press.), pp. 571-592. https://doi.org/10.1016/B978-0-12-805052-1.00033-4
- -Frazão Santos, C., Agardy, T., Andrade, F., Calado, H., Crowder, L.B., Ehler, C.N., et al. (2020).
- 627 Integrating climate change in ocean planning. *Nature Sustainability*, 3(7), 505-516.
 628 https://doi.org/10.1038/s41893-020-0513-x
- -Frazão Santos, C., Agardy, T., Andrade, F., Crowder, L. B., Ehler, C. N., & Orbach, M. K. (2021).
- Major challenges in developing marine spatial planning. *Marine Policy*. 132: 103248
 https://doi.org/10.1016/j.marpol.2018.08.032
- -Frys, C., Saint-Amand, A., Le Hénaff, M., Figueiredo, J., Kuba, A., Walker, B., et al. (2020). Fine-
- 633 Scale Coral Connectivity Pathways in the Florida Reef Tract: Implications for Conservation and
- Restoration. *Frontiers in Marine Science*, 7: 312. https://doi.org/10.3389/fmars.2020.00312
- -Gann, G.D., McDonald, T., Walder, B., Aronson, J., Nelson, C.R., Jonson, J., et al. (2019).
- 636 International principles and standards for the practice of ecological restoration. Second edition.
- 637 *Restoration Ecology*, 27: S1-S46. https://doi.org/10.1111/rec.13035
- -Gijsman, R., Horstman, E.M., van der Wal, D., Friess, D.A., Swales, A. & Winjberg, K.M. (2021).
- 639 Nature-based engineering: A review on reducing coastal flood risk with mangroves. Frontiers in
- 640 *Marine Science*, 8: 825. https://doi.org/10.3389/fmars.2021.702412
- -Gilby, B., Olds, A. D., Connolly, R. M., Henderson, C. J. & Schlacher, T. A. (2018). Spatial
- 642 restoration ecology: placing restoration in a landscape context. *Bioscience* 68: 1007–1019.
- 643 https://doi.org/10.1093/biosci/biy126
- -Gilby, B.L., Olds, A.D., Brown, C.J., Connolly, R.M., Henderson, C.J., Maxwell, P.S., & Schlacher,
- T.A. (2021). Applying systematic conservation planning to improve the allocation of restoration
- 646 actions at multiple spatial scales. Restoration Ecology, 29(5), e13403.
- 647 https://doi.org/10.1111/rec.13403

- -Gilliland, P.M., & Laffoley, D. (2008). Key elements and steps in the process of developing
 ecosystem-based marine spatial planning. *Marine Policy*, *32:* 787-796.
 https://doi.org/10.1016/j.marpol.2008.03.022
- -Gissi, E., Manea, E., Mazaris, A.D., Fraschetti, S., Almpanidou, V., Bevilacqua, S., et al. (2021). A
 review of the combined effects of climate change and other local human stressors on the marine
 environment. *Science of the Total Environment*, 755: 142564.
 https://doi.org/10.1016/j.scitotenv.2020.142564
- -Gissi, E., Maes, F., Kyriazi, Z., Ruiz-Frau, A., Santos, C. F., Neumann, B., et al. (2022).
- 656 Contributions of marine area-based management tools to the UN sustainable development goals.
- 657 Journal of Cleaner Production, 330: 129910. https://doi.org/10.1016/j.jclepro.2021.129910
- -Gómez-Baggethun, E., Tudor, M., Doroftei, M., Covaliov, S., Năstase, A., Onără, D. F., et al. (2019).
- 659 Changes in ecosystem services from wetland loss and restoration: An ecosystem assessment of the
- 660 Danube Delta (1960–2010). *Ecosystem Services*, 39: 100965.
- 661 https://doi.org/10.1016/j.ecoser.2019.100965
- -Grip, K. & Blomqvist, S. (2021). Marine spatial planning: Coordinating divergent marine interests.
- 663 *Ambio*, 50(6), 1172-1183. https://doi.org/10.1007/s13280-020-01471-0
- -Guerry, A.D., Ruckelshaus, M.H., Arkema, K.K., Bernhardt, J. R., Guannel, G., Kim, C-K., et al.
- 665 (2012). Modeling benefits from nature: using ecosystem services to inform coastal and marine spatial
- 666 planning. International Journal of Biodiversity Science & Management, 8: 107-121.
- 667 https://doi.org/10.1080/21513732.2011.647835
- -Halpern, B.S., Frazier, M., Afflerbach, J., Lowndes, J.S., Micheli, F., O'Hara, C., et al. (2019).
- Recent pace of change in human impact on the world's ocean. *Scientific Report, 9:* 1-8.
 https://doi.org/10.1038/s41598-019-47201-9
- -Heymans, J.J., Bundy, A., Christensen, V., Coll, M., De Mutsert, K., Fulton, E.A., et al. (2020). The
- ocean decade: a true ecosystem modeling challenge. Frontiers in Marine Science, 7: 766.
- 673 https://doi.org/10.3389/fmars.2020.554573

- -Jonsson, P. R., Moksnes, P. O., Corell, H., Bonsdorff, E., & Nilsson Jacobi, M. (2020). Ecological
 coherence of Marine Protected Areas: New tools applied to the Baltic Sea network. *Aquatic*
- 676 *Conservation: Marine and Freshwater Ecosystems, 30*(4), 743-760. https://doi.org/10.1002/aqc.3286
- -Klemas, V. (2013). Using Remote Sensing to Select and Monitor Wetland Restoration Sites: An
- 678 Overview. Journal of Coastal Research, 29: 958–970. https://doi.org/10.2112/JCOASTRES-D-12-
- 679 00170.1
- -Kodikara, K.A.S., Mukherjee, N., Jayatissa, L.P., Dahdouh-Guebas, F. & Koedam, N. (2017). Have
- mangrove restoration projects worked? An in-depth study in Sri Lanka. *Restoration Ecology*, 25: 705-
- 682 716. https://doi.org/10.1111/rec.12492
- -Kotta, J., Vanhatalo, J., Jänes, H., Orav-Kotta, H., Rugiu, L., Jormalainen, V., et al. (2019).
- 684 Integrating experimental and distribution data to predict future species patterns. *Scientific Report*,
- 685 9(1), 1-14. https://doi.org/10.1038/s41598-018-38416-3
- -Laffoley, D., Baxter, J. M., Day, J. C., Wenzel, L., Bueno, P., & Zischka, K. (2019). Marine protected
- areas, in: C. Sheppard (Ed.), World Seas An Environ. Eval. Vol. III Ecol. Issues Environ. Impacts,
- 688 2nd ed., Elsevier, 2019, pp. 549–569. https://doi.org/10.1016/B978-0-12-805052-1.00027-9
- -Lester, S.E., Dubel, A.K., Hernan, G., McHenry, J., & Rassweiler, A. (2020). Spatial Planning
- 690 Principles for Marine Ecosystem Restoration. Frontiers in Marine Science, 7: 328.
- 691 https://doi.org/10.3389/fmars.2020.00328
- -Lombard, A.T., Ban, N.C., Smith, J.L., Lester, S.E., Sink, K. J., Wood, S.A., et al. (2019). Practical
- approaches and advances in spatial tools to achieve multi-objective marine spatial planning. *Frontiers*
- *in Marine Science*, *6*: 166. https://doi.org/10.3389/fmars.2019.00166
- Maes, F. (2008). The international legal framework for marine spatial planning. *Marine Policy*,
 32(5), 797-810. https://doi.org/10.1016/j.marpol.2008.03.013
- 697 McAfee, D., Larkin, C., Connell, S.D. (2021). Multi-species restoration accelerates recovery of
- 698 extinguished oyster reefs. Journal of Applied Ecology, 58(2), 286-294. doi: 10.1111/1365-
- 699 2664.13719

- McAfee, D., Reis-Santos, P., Jones, A.R., Gillanders, B.M., Mellin, C., Nagelkerken, I., et al.
- 701 (2022a). Multi-habitat seascape restoration: optimising marine restoration for coastal repair and social
- benefit. Frontiers in Marine Science, 9. https://doi.org/10.3389/fmars.2022.910467
- McAfee, D., McLeod, I. M., Alleway, H. K., Bishop, M. J., Branigan, S., Connell, S. D., et al.
- (2022b). Turning a lost reef ecosystem into a national restoration program. *Conservation Biology*,
- 705 *36*(6), e13958. https://doi.org/10.1111/cobi.13958
- -Maccarrone, V., Filiciotto, F., de Vincenzi, G., Mazzola, S. & Buscaino, G. (2015). An Italian
 proposal on the monitoring of underwater noise: Relationship between the EU Marine Strategy
 Framework Directive (MSFD) and marine spatial planning directive (MSP). *Ocean and Coastal Management, 118*: 215-224. https://doi.org/10.1016/j.ocecoaman.2015.07.006
- McClenachan, G. M., Donnelly, M. J., Shaffer, M. N., Sacks, P. E., & Walters, L. J. (2020). Does 710 size matter? Quantifying the cumulative impact of small-scale living shoreline and oyster reef 711 712 restoration projects on shoreline erosion. *Restoration* Ecology, 28(6), 1365-1371. https://doi.org/10.1111/rec.13235 713
- -Manea, E., Bianchelli, S., Fanelli, E., Danovaro, R. & Gissi, E. (2020). Towards an ecosystem-based
- 715 marine spatial planning in the deep Mediterranean Sea. Science of the Total Environment, 715:
- 716 136884. https://doi.org/10.1016/j.scitotenv.2020.136884
- Manea, E., Bergami, C., Pugnetti, A., Gianni, F., Oggioni, A., Bandelj, V., Cataletto, B., Pleslic, G.
- 718 & Bongiorni, L. (2022). An ecosystem-based system of variables to enhance marine species and
- habitat monitoring and conservation: The Adriatic Natura 2000 case study. Frontiers in Marine
- 720 Science, 9: 920366. doi: 10.3389/fmars.2022.920366
- -McMahon, K.W., Berumen, M.L. & Thorrold, S.R. (2012). Linking habitat mosaics and connectivity
- in a coral reef seascape. *Proceedings of the National Academy of Sciences*, U.S.A., 109: 15372-15376.
- 723 https://doi.org/10.1073/pnas.1206378109

- -Menegon, S., Depellegrin, D., Farella, G., Gissi, E., Ghezzo, M., Sarretta, A., et al. (2018). A
- modelling framework for MSP-oriented cumulative effects assessment. *Ecological Indicators*, 91:
- 726 171-181. https://doi.org/10.1016/j.ecolind.2018.03.060
- -Merovich, G.T. & Petty, J.T. (2007). Interactive effects of multiple stressors and restoration priorities
- in a mined Appalachian watershed. Hydrobiologia, 575(1), 13-31. https://doi.org/10.1007/s10750-
- 729 006-0277-у
- -Miller, K.R. (1999). Bioregional planning and biodiversity conservation. *Partnership for Protection*,
- 731 New Strategies for Planning and Management for Protected Areas. Stolton, S., and Dudley, N., eds.
- 732 London: Earthscan Publication, 41-49.
- -Neeson, T.M, Smith, S.D.P, Allan, J.D & McIntyre, P.B. (2016). Prioritizing ecological restoration
- among sites in multi-stressor landscapes. *Ecological Applications*, 26: 1785–1796.
 https://doi.org/10.1890/15-0948.1
- -Normile, D. (2010). UN biodiversity summit yields welcome and unexpected progress. *Science*, *330*:
 742-743. DOI: 10.1126/science.330.6005.742
- -NRC, (1992). National Research Council. Restoration of Aquatic Ecosystems—Science,
 Technology and Public Policy. National Academy Press, Washington, DC, 576 pp.
- -Orth, R.J., Lefcheck, J.S., McGlathery, K.S., Aoki, L., Luckenbach, M.W., Moore, K.A., et al.
- (2020). Restoration of seagrass habitat leads to rapid recovery of coastal ecosystem services. *Science*
- 742 *Advances*, *6*(41), eabc6434. DOI: 10.1126/sciadv.abc6434
- -Ouellette, W. & Getinet, W. (2016). Remote sensing for marine spatial planning and integrated
- coastal areas management: Achievements, challenges, opportunities and future prospects. Remote
- 745 Sensing Applications: Society and Environment, 4: 138-157.
 746 https://doi.org/10.1016/j.rsase.2016.07.003
- -Paschke, M.W., Perkins, L.B. & Veblen, K.E. (2019). Restoration for multiple use. Restoration
- 748 *Ecology*, 27: 701-704. https://doi.org/10.1111/rec.12949

- -Paxton, A. B., Steward, D. A. N., Harrison, Z. H. & Taylor, J. C. (2022). Fitting ecological principles 749 750 of artificial reefs into the ocean planning puzzle. Ecosphere, 13(2), e3924. https://doi.org/10.1002/ecs2.3924 751
- Pittman, S.J., Stamoulis. K.A., Antonopoulou, M., Das, H.S., Shahid, M., Delevaux, J.M.S.,
- 753 Wedding, L.M. & Mateos-Molina, D. (2022). Rapid Site Selection to Prioritize Coastal Seascapes for
- Nature-Based Solutions With Multiple Benefits. *Frontiers in Marine Science*, 9:832480. doi:
 10.3389/fmars.2022.832480.
- -Pouso, S., Borja, A. & Uyarra, M.C. (2020). An Interdisciplinary Approach for Valuing Changes
- 757 After Ecological Restoration in Marine Cultural Ecosystem Services. Frontiers in Marine Science, 7:
- 758 715. doi: 10.3389/fmars.2020.00715
- -Reimer, J. M., Devillers, R., Trouillet, B., Ban, N. C., Agardy, T., & Claudet, J. (2023). Conservation
 ready marine spatial planning. *Marine Policy*, 153:105655.
 https://doi.org/10.1016/j.marpol.2023.105655
- -Roberts, C.M., O'Leary, B.C. & Hawkins, J.P. (2020). Climate change mitigation and nature
 conservation both require higher protected area targets. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *375*: 20190121. https://doi.org/10.1098/rstb.2019.0121
- -Roberts, K.E., Cook, C.N., Beher, J. & Treml, E.A. (2021). Assessing the current state of ecological
- connectivity in a large marine protected area system. *Conservation Biology*, 35: 699-710.
 https://doi.org/10.1111/cobi.13580
- -Saeedi, H., Reimer, J.D., Brandt, M.I., Dumais, P.O., Jażdżewska, A.M., Jeffery, N.W., et al. (2019).
- Global marine biodiversity in the context of achieving the Aichi Targets: ways forward and
 addressing data gaps. *PeerJ*, 7: e7221. https://doi.org/10.7717/peerj.7221
- -Sciascia, R., Guizien, K., Padrón, M., Bramanti, L. & Magaldi, M. (2020). Larval buoyancy matters:
- sinking changes larval transport predictions among *Paramuricea clavata* populations. Preprint at
- HAL open science. https://hal.archives-ouvertes.fr/hal-02558272

- -SER, (2004). Society for Ecological Restoration International Science & Policy Working Group.
- 775 The SER International Primer on Ecological Restoration. www.ser.org & Tucson: Society for
- 776 Ecological Restoration International.
- -Shabtay, A., Portman, M. E., Manea, E., & Gissi, E. (2019). Promoting ancillary conservation
- through marine spatial planning. *Science of the Total Environment*, 651: 1753-1763.
- 779 https://doi.org/10.1016/j.scitotenv.2018.10.074
- -Sherman, K. & Alexander, L.M. (1986). Variability and management of large marine ecosystems.
- 781 American Association for the Advancement of Science Selected Symposium 99. Westview,
- 782 Boulder, Colorado, USA.
- -Shumway, N., Bell-James, J., Fitzsimons, J. A., Foster, R., Gillies, C. & Lovelock, C.E. (2021).
- Policy solutions to facilitate restoration in coastal marine environments. *Marine Policy*, *134*: 104789.
- 785 https://doi.org/10.1016/j.marpol.2021.104789
- -Simenstad, C., Reed, D. & Ford, M. (2006). When is restoration not?: Incorporating landscape-scale
 processes to restore self-sustaining ecosystems in coastal wetland restoration. *Ecological Engineering*, 26: 27-39. https://doi.org/10.1016/j.ecoleng.2005.09.007
- Smale, D.A., Moore, P.J., Queirós, A., Higgs, S., Burrows, M.T. (2018). Appreciating
- interconnectivity between habitats is key to blue carbon management. *Frontiers in Ecology and the*
- 791 *Environment*, 16(2), 71–73. doi: 10.1002/fee.1765
- Stender, Y., Foley, M., Rodgers, K.U., Jokiel, P., & Singh, A. (2021). Evaluating the feasibility and
- advantage of a multi-purpose submerged breakwater for harbor protection and benthic habitat
- r94 enhancement at Kahului Commercial Harbor, Hawai'i: case study. SN Applied Science, 3: 167
- 795 https://doi.org/10.1007/s42452-020-04072-4
- -Strassburg, B.B., Beyer, H.L., Crouzeilles, R., Iribarrem, A., Barros, F., de Siqueira, M.F., et al.
- 797 (2019). Strategic approaches to restoring ecosystems can triple conservation gains and halve costs.
- 798 *Nature Ecology and Evolution, 3*(1), 62-70. https://doi.org/10.1038/s41559-018-0743-8

- -Suárez-Castro, A.F., Beyer, H.L., Kuempel, C.D., Linke, S., Borrelli, P. & Hoegh-Guldberg, O.
- 800 (2021). Global forest restoration opportunities to foster coral reef conservation. *Global Change*
- 801 *Biology*, 27(20), 5238-5252. https://doi.org/10.1111/gcb.15811
- -Swam, L. M., Couvillion, B., Callam, B., La Peyre, J. F. & La Peyre, M. K. (2022). Defining oyster
- resource zones across coastal Louisiana for restoration and aquaculture. *Ocean and Coastal Management*, 225: 106178. https://doi.org/10.1016/j.ocecoaman.2022.106178
- ovan Tatenhove, J. P., Ramírez-Monsalve, P., Carballo-Cárdenas, E., Papadopoulou, N., Smith, C. J.,
- Alferink, L., et al. (2021). The governance of marine restoration: insights from three cases in two
- European seas. *Restoration Ecology*, 29: e13288. https://doi.org/10.1111/rec.13288
- -Teichert, N., Borja, A., Chust, G., Uriarte, A. & Lepage, M. (2016). Restoring fish ecological quality
- in estuaries: implication of interactive and cumulative effects among anthropogenic stressors. *Science*
- 810 of the Total Environment, 542: 383-393. https://doi.org/10.1016/j.scitotenv.2015.10.068
- -Thiele, T., Alleng, G., Biermann, A., Corwin, E., Crooks, S., Fieldhouse, P., et al. (2020). Blue
- 812 Infrastructure Finance: A new approach, integrating Nature-based Solutions for coastal resilience.813 IUCN, Gland, Switzerland.
- -Thom, R. M. (2000). Adaptive management of coastal ecosystem restoration projects. *Ecological*
- Engineering, 15(3-4), 365-372. https://doi.org/10.1016/S0925-8574(00)00086-0
- -Treml, E.A. & Halpin, P.N. (2012). Marine population connectivity identifies ecological neighbors
 for conservation planning in the Coral Triangle. *Conservation Letters*, 5: 441-449.
- 818 https://doi.org/10.1111/j.1755-263X.2012.00260.x
- -Trouillet, B., & Jay, S. (2021). The complex relationships between marine protected areas and
 marine spatial planning: Towards an analytical framework. *Marine Policy*, *127*: 104441.
 https://doi.org/10.1016/j.marpol.2021.104441
- -UNEP (2011): Taking Steps toward Marine and Coastal Ecosystem-Based Management An
 Introductory Guide.

- -UNEP (2019). Frontiers 2018/19 Emerging Issues of Environmental Concern. United Nations 824 825 Environment Programme, Nairobi.
- -Uusitalo, L., Korpinen, S., Andersen, J. H., Niiranen, S., Valanko, S., Heiskanen, A. S., & Dickey-826
- Collas, M. (2016). Exploring methods for predicting multiple pressures on ecosystem recovery: a 827
- case study on marine eutrophication and fisheries. Continental Shelf Research, 121: 48-60. 828
- https://doi.org/10.1016/j.csr.2015.11.002 829
- 830 -Vanderklift, M.A., Marcos-Martinez, R., Butler, J.R., Coleman, M., Lawrence, A., Prislan, H., et al.
- (2019). Constraints and opportunities for market-based finance for the restoration and protection of 831
- blue carbon ecosystems. Marine Policy, 107: 103429. https://doi.org/10.1016/j.marpol.2019.02.001 832
- 833 -Vaughan, D., & Agardy, T. (2020). Marine protected areas and marine spatial planning-allocation
- of resource use and environmental protection. In Marine Protected Areas (pp. 13-35). Elsevier. 834 https://doi.org/10.1016/B978-0-08-102698-4.00002-2
- 835
- 836 - Wang, C., Tang, C., Fu, B., Lü, Y., Xiao, S., & Zhang, J. (2022). Determining critical thresholds of
- ecological restoration based on ecosystem service index: a case study in the Pingjiang catchment in 837
- southern China. Journal Environmental Management, 303, 114220. 838 of https://doi.org/10.1016/j.jenvman.2021.114220 839
- -White, C., Halpern, B.S. & Kappel, C.V. (2012). Ecosystem service tradeoff analysis reveals the 840
- value of marine spatial planning for multiple ocean uses. Proceedings of the Royal Society B: 841
- Biological Sciences, 109(12), 4696-4701. https://doi.org/10.1073/pnas.1114215109 842
- -Wiens J.A. & Hobbs R.J. (2015). Integrating conservation and restoration in a changing world. 843 844 BioScience, 65(3), 302-313. https://doi.org/10.1093/biosci/biu235
- -Wilson, K. L., Tittensor, D. P., Worm, B., & Lotze, H. K. (2020). Incorporating climate change 845
- adaptation into marine protected area planning. Global Change Biology, 26(6), 3251-3267. 846
- https://doi.org/10.1111/gcb.15094 847
- -Zahawi, R.A., Reid, J.L., & Holl, K.D. (2014). Hidden costs of passive restoration. Restoration 848
- Ecology, 22(3), 284-287. https://doi.org/10.1111/rec.12098 849

- -Zellmer, A.J., Claisse, J.T., Williams, C.M., Schwab, S. & Pondella, D.J. (2019). Predicting optimal
- 851 sites for ecosystem restoration using stacked-species distribution modeling. Frontiers in Marine
- *Science, 6:* 3. https://doi.org/10.3389/fmars.2019.00003
- 853
- 854 Tables
- 855 **Table 1.** Restoration-related terminologies and their definitions.

Term	Definition
Ecological restoration	The process of assisting the recovery of an ecosystem that has been degraded, damaged, or destroyed (SER, 2004). Returning an ecosystem to a close approximation of its condition prior to disturbance by re-establishing pre disturbance aquatic functions and ecosystem processes, and related physical, chemical and biological characteristics, and allowing reintroduction of indigenous species (NRC, 1992; Aronson & Le Floc'h, 1996; Simenstad et al., 2006).
	functions (contrast with rehabilitation). When full recovery is the goal, an important benchmark is when the ecosystem demonstrates self-organization (Gann et al., 2019). It includes the following components and perspectives (Clewell & Aronson, 2013; Gann et al., 2019):
	Adaptive component: it aims to move a degraded ecosystem to a trajectory of recovery that allows adaptation to local and global changes, as well as persistence and evolution of its component species; Ecological perspective: it is an intentional activity that reinitiates ecological
	processes that were interrupted when an ecosystem was impaired; Conservation perspective: it recovers biodiversity in the face of an unprecedented, human-mediated extinction crisis;
	Socio-economic perspective: ecological restoration recovers ecosystem services (ES) from which people benefit.

"Passive" or "Natural" restoration	Allowing natural or unassisted ecosystem recovery after removing a source of disturbance (Atkinson & Bonser, 2020).
"Active" or "Assisted", and "Reconstructive" restoration	Assisted restoration: Abiotic - e.g. Active remediation of substrate conditions (physical or chemical), habitat creation, reshaping watercourses, reintroduction of environmental water flows, applying artificial disturbance to promote seed germination; Biotic - e.g. Invasive species management, reintroduction of species, augmenting or reinforcing depleted populations of species (Atkinson & Bonser, 2020).
	Reconstructive restoration: A combination of the above strategies with the reintroduction of a major proportion of the desired biota. Possibly mimicking natural successional dynamics (Atkinson & Bonser, 2020).
Rehabilitation	The act of partially or, more rarely, fully replacing structural or functional characteristics of an ecosystem that have been reduced or lost (Elliott et al., 2007). In the short term, management measures favour one group of species or ES (Aronson & Le Floc'h, 1996; Simenstad et al., 2006).
	The goal of rehabilitation projects is not native ecosystem recovery, but rather reinstating a level of ecosystem functioning for renewed and ongoing provision of ES potentially derived from non-native ecosystems as well (Gann et al., 2019).
Reallocation	Literally, this entails changing the way something is allocated for conversion of an ecosystem to a different kind of ecosystem or land use primarily for purposes other than the conservation management of local native ecosystems (Aronson et al., 1993). Reallocation can favor the development of new trajectories that over the long-term produce new ecosystems and uses (Aronson and & Le Floc'h, 1996, Elliott et al., 2007).
Remediation	Action taken, following anthropogenic disturbance, to restore or enhance the ecological value of a site (Emu Ltd., 2004), hence giving emphasis to the action or process rather than the end-point reached (Bradshaw, 2002; Elliott et al., 2007).

Recovery	The capacity or the process of a system to return to pre-disturbance condition, the
	original or reference state, after being in a degraded or disrupted one (Elliott et al.,
	2007). Recovery is the outcome sought or achieved of a restoration action. It can be
	active (human induced) or passive (natural).

Figure legends





Figure 1. EB-MSP and marine restoration benefits sharing. Schematic representation of how an EB-

862 MSP implementing EBM core principles can operatively support marine restoration and the derived

benefits. EBM principles are taken from UNEP (2011).



864

Figure 2. How restoration-focused EB-MSP drives more efficient ocean regeneration. The elements 865 for a transformative shift from an old to a new ocean regeneration paradigm are all here today, but 866 disparate and unsystematic planning is unable to capitalize on the immense opportunities that exist 867 for this transformation to happen. This transformation can be reached by staying focused on the five 868 EBM core principles - recognizing connections (to build a solid understanding of the ecology of 869 marine and coastal systems, and identify critical ecological elements to protect or restore as well as 870 the links that connect those), taking an ecosystem services approach (as a way to communicate the 871 values of healthy oceans), addressing cumulative impacts and managing for multiple uses (to allow 872 institutional integration and effective management of uses, resources, and ocean space), embracing 873 874 change, learning, adaptation (to trial solutions and practise adaptive management, allowing also the incorporation of human development and climate changes). This is what the restoration-focused EB-875 876 MSP frameworks makes possible.