



**HAL**  
open science

# Robust estimator of the Ruin Probability in infinite time for heavy-tailed distributions

Modou Kebe, El-Hadji Deme, Yousri Slaoui, Solym Mawaki Manou-Abi

► **To cite this version:**

Modou Kebe, El-Hadji Deme, Yousri Slaoui, Solym Mawaki Manou-Abi. Robust estimator of the Ruin Probability in infinite time for heavy-tailed distributions. 2023. hal-04183503

**HAL Id: hal-04183503**

**<https://hal.science/hal-04183503v1>**

Preprint submitted on 19 Aug 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# Robust estimator of the Ruin Probability in infinite time for heavy-tailed distributions

Modou Kebe <sup>(1)</sup>, El-Hadji Deme <sup>(1,\*)</sup>,  
Yousri Slaoui <sup>(2)</sup> and Solym M. Manou-Abi <sup>(3)</sup>

(1) LERSTAD, UFR SAT, Université Gaston Berger, BP 234, Saint-Louis, Sénégal.

(2) LaMMASD, Département de Mathématiques, Université de Kara, BP 404, Kara, Togo

(3) Laboratoire de Mathématiques et Applications, UMR CNRS 7348 ,Université de Poitiers, France.

**Abstract.** The probability of ruin of an insurance company is one of the main risk measures considered in risk theory, and the problems of its calculation and approximation have attracted much attention. Statistical estimations have been developed on the ruin probability in infinite time for insurance losses from heavy-tailed distributions. However, these estimation suffer heavily from under-coverage or have a robustness problem, particularly when losses are contaminated by large variations in the arrival of claims. We therefore need another method for estimating the probability of ruin in infinite time for heavy-tailed losses. This is why, in this paper, we introduce a robust estimator of the infinite-time probability of ruin for such distributions. Our methodology is based on extreme value theory, which offers adequate statistical results for such distributions. Our approach is based on a sensitive distribution known as the t-Hill estimator (t-score or score moment estimation) for the index of any tail distribution and introduced in [Fabián and Stehlík (2009)]. We establish their asymptotic normality, and through a simulation study, illustrate their behavior in terms of absolute bias and mean squared error. The simulation results clearly show that our estimators perform well and that they are fairly robust to outliers.

**Keywords:** Ruin Probability, Estimation, Robustness, Tail index, Heavy-tailed, Reinsurance.

**Mathematics Subject Classification:** 62H12· 62H05· 60G70

(1) Corresponding author: [elhadji.deme@ugb.edu.sn](mailto:elhadji.deme@ugb.edu.sn)

# 1 Introduction

The effect of insurance operations is the total or partial transfer of the financial consequences of the risk incurred by the insured to an insurance company. The expenses covered by the company may correspond either to indemnities to be paid to third parties in respect of the insured's liability (civil, professional or other), or to compensation for damage suffered by the insured. But what happens to these insurance companies when they themselves are exposed to risk? One approach to this problem is based on the use of ruin theory (see, e.g., [Panjer and Willmot (1992)]).

In the field of insurance, risk is defined as the probability that an insurance company's reserve, i.e. the difference between the premium amount and the insurance amount, will be reduced of an insurance company, which is the difference between the total premiums received and the total amount of claims paid, becomes negative at some point. At that point, ruin is said to occur, due to a miscalculation of the policyholders' contribution rate, or claims that are too large to cover. Indeed, the probability of such an event is seen as a means of controlling risk behavior. It is also a useful way of controlling the insurer's funds in long-term planning. Let's recall the definition of a standard mathematical model for insurance risk (see, for example, [Čizěk *et al* (2005)], p. 345).

The insurance company's initial capital is denoted  $u$ . The number of claims over the period  $(0, t]$ , denoted  $M_t$ , is described by a Poisson process with a fixed intensity (rate)  $\lambda > 0$ . We also define a sequence of non-negative random variables  $\{X_i\}_{i=1}^{\infty}$  independent and identically distributed (i.i.d. ) with a loss distribution function  $F_1(x) := \mathbb{P}(X_i \leq x)$  representing loss severity, with an unknown finite mean  $\mu_1 = \int_0^{\infty} (1 - F_1(x)) dx$ . Knowledge of this mean value has been of great interest to insurance companies, since it is one of the most commonly used premium calculation principles, known as *the net premium*, and corresponds to the expected amount of claims for a given insurance period.

Let's then assume that  $X_i$ 's are independent of  $M_t$  and that the insurer collects a premium at a constant rate  $c$  per unit time and that the net profit condition is met, i.e.  $c/\lambda > \mu_1$ . The classical risk process  $\{R_t\}_{t>0}$  is given by:

$$R_t := u + ct - \sum_{i=1}^{M_t} X_i, \quad t > 0.$$

The corresponding claim surplus process is defined by

$$S_t := u - R_t = ct - \sum_{i=1}^{M_t} X_i, \quad t > 0.$$

First of all, we're interested in the probability that  $S_t$  exceeds an initial reserve  $u$  at a time  $t$  before or at a horizon  $T$ . Explicitly, this probability can be written as follows:

$$\psi_T(u) := \mathbb{P} \left\{ \sup_{0 < t \leq T} S_t > u \right\}.$$

The ruin probability in infinite time is defined by,

$$\phi(u) := \lim_{T \rightarrow \infty} \psi_T(u). \tag{1.1}$$

In the actuarial field, the costs of large claims require the modeling of rare events, i.e. events with a low probability of occurrence, but with large claims amounts and disastrous effects. The analysis of these

extreme events can be carried out using the extreme value methodology, whose distribution functions  $F_1$  are heavy-tailed and mainly characterized by their index, which indicates the size and frequency of certain extreme phenomena within a given probability distribution (see for example [Vandewalle and Beirlant (2006)]).

The heavy-tailed nature of claims requires particular attention to the analysis of tail distributions. Extreme value theory (EVT) therefore offers suitable statistical tools for modeling these distribution tails, see for example [Matthys *et al.* (2004)], [Vandewalle and Beirlant (2006)] [Reiss and Thomas (2007)], [Necir *et al.* (2010)], [Deme *et al.* (2013a)], [Deme *et al.* (2015)], [Deme *et al.* (2021)] etc... In addition, reinsurance companies have to calculate premiums to cover these excess claims, which are generally very high. EVT has become one of the leading theories in the development of statistical models for high insurance losses.

Now, suppose that  $F_1$  is heavy-tailed, that is:

$$\lim_{x \rightarrow \infty} \exp(\delta x) (1 - F_1) = \infty, \text{ for all } \delta > 0. \quad (1.2)$$

The class of regularly varying functions provides good examples of heavy-tailed models. We can cite the following models: Pareto, Burr, Student, Lévy-stable and log-gamma (see, for example, [Beirlant *et al.* (2001)]). In the remainder of this paper, we restrict ourselves to this class of distributions. In other words, we assume that the survival function  $1 - F_1$  is smoothly varying at infinity with index  $-1/\gamma < 0$ , that

$$\lim_{t \rightarrow \infty} \frac{1 - F_1(tx)}{1 - F_1(t)} = x^{-1/\gamma}, \text{ for any } x > 0. \quad (1.3)$$

The parameter  $\gamma$  is the tail index and governs tail behavior, with higher values indicating heavier tails. For more details on these models, we can refer to [Bingham *et al.* (1987)], [Rolski *et al.* (1999)] and [Reiss and Thomas (2007)]. It has been shown that for large initial reserve  $u$ , the ruin probability  $\phi(u)$  can be approximated, under the assumption, by

$$\phi(u) := \left(\frac{c}{\lambda} - \mu_1\right)^{-1} \int_u^\infty (1 - F_1(x)) dx, \quad (1.4)$$

(see, e.g., [Asmussen (2000)]). This latter equation can be rewritten as

$$\phi(u) = \frac{\mu_2}{\omega - \mu_1}, \quad (1.5)$$

where  $\mu_2 := \mathbb{E}(Y) < \infty$ , with  $Y := (X - u)_+ = \max(X - u, 0)$  and  $\omega := c/\lambda$ . Then, the distribution function of the stop-loss variable  $Y$  denoted by  $F_2$  also satisfies the relationship (1.3) with the same index  $-1/\gamma < 0$ , i.e.

$$\lim_{t \rightarrow \infty} \frac{1 - F_2(tx)}{1 - F_2(t)} = x^{-1/\gamma} \text{ for any } x > 0. \quad (1.6)$$

Now let  $Q_i$ ,  $i = 1, 2$  be the generalized inverse functions (or quantile functions) related to df  $F_i$ ,  $i = 1, 2$  and defined as follows for all  $s \in (0, 1]$ :

$$Q_i(s) := \inf \{x > 0 : F_i(x) \geq s\}.$$

From Corollary 1.2.10 (p. 23) in [?], we have for any  $x > 0$

$$\lim_{s \downarrow 0} \frac{Q_i(1 - sx)}{Q_i(1 - s)} = x^{-\gamma}, \quad i = 1, 2. \quad (1.7)$$

By a change of variable, the expected values  $\mu_i = \int_0^\infty (1 - F_i(x))dx$ ,  $i = 1, 2$  can be rewritten in terms of quantile function  $Q_i$  as follows:  $\mu_i = \int_0^1 Q_i(s)ds$ ,  $i = 1, 2$ . Thus, the probability of ruin in infinite time can be rewritten as follows:

$$\phi(u) = \frac{\int_0^1 Q_2(s)ds}{\omega - \int_0^1 Q_1(s)ds}. \quad (1.8)$$

Notice that for fixed large  $u$ , we have

$$1 - F_1(x) \sim 1 - F_2(x) \quad \text{as } x \rightarrow \infty,$$

and therefore

$$Q_1(1-s) \sim Q_2(1-s) \quad \text{as } s \downarrow 0.$$

This paper is organized as follows. In section 2, we present some preliminaries on classical estimators of the probability of ruin  $\phi(u)$ . As these estimates suffer greatly from under-coverage or have a robustness problem, especially when losses are contaminated by large variations in the arrival of claims, we introduce, in subsection 3.1, a robust infinite-time estimator of the probability of ruin for heavy-tailed insured losses. Using extreme value methodology, we establish its asymptotic distribution in subsection 3.2. In subsection 4.1, we perform a simulation study to illustrate the behavior of our robust estimator compared to the classical estimator in terms of absolute bias and mean squared error. In subsection 4.2, we present a contamination study in which the robustness of the estimator is evaluated. All proofs are reported in section 5.

## 2 Estimating the ruin probability $\phi(u)$

First, we set a large initial reserve  $u$ . Let  $(X_1, \dots, X_n)$  and  $(Y_1, \dots, Y_n)$  be two independent samples of risks  $X$  and  $Y$  respectively. The non-parametric estimators of the distribution functions  $F_1$  and  $F_2$  are respectively defined as follows  $F_{1,n}(x) = n^{-1} \sum_{i=1}^n \mathbb{I}_{\{X_i \leq x\}}$  and  $F_{2,n}(y) = n^{-1} \sum_{i=1}^n \mathbb{I}_{\{Y_i \leq y\}}$ . Thus, their corresponding empirical quantile functions are expressed by  $Q_{i,n}(s) = \inf\{x; F_{i,n}(x) \geq s\}$ ,  $i = 1, 2$  where  $\mathbb{I}_S$  is the indicator function of the set  $S$ . Let's denote by  $X_{1,n} \leq \dots \leq X_{n,n}$  and  $Y_{1,n} \leq \dots \leq Y_{n,n}$  the order statistics associated respectively with the samples  $(X_1, \dots, X_n)$  and  $(Y_1, \dots, Y_n)$ . Therefore,  $Q_{1,n}(t) = X_{j,n}$  and  $Q_{2,n}(t) = Y_{j,n}$  for all  $t \in ((j-1)/n, j/n]$ , and for all  $j = 1, \dots, n$ .

To this end, a natural candidate for the empirical estimator of  $\phi(u)$  is obtained by replacing in (1.8) the real quantiles  $Q_1(\cdot)$  and  $Q_2(\cdot)$  by their respective sample quantiles  $Q_{1,n}(\cdot)$  and  $Q_{2,n}(\cdot)$ . We obtain the following "traditional" non-parametric estimator of the probability of ruin:

$$\bar{\phi}_n(u) = \frac{\bar{Y}}{\omega - \bar{X}}. \quad (2.9)$$

where  $\bar{X} := \frac{1}{n} \sum_{j=1}^n X_j$  and  $\bar{Y} := \frac{1}{n} \sum_{j=1}^n Y_j$  are the sample estimators the mean of  $\mu_1$  and  $\mu_2$  respectively.

Note that for  $\gamma \geq 1$ , the expected value of  $X$  (respectively  $Y$ ) does not exist. Consequently, we will focus exclusively on distributions whose tail indices lie in the unit interval  $0 < \gamma < 1$ .

Next, the random variable  $\sqrt{n}(\bar{\phi}_n(u) - \phi(u))$  can be rewritten as:

$$\frac{1}{(\omega - \bar{X})(\omega - \mu_1)} \sqrt{n}((\bar{Y} - \mu_1)(\omega - \mu_1) + (\bar{X} - \mu_1)\mu_2).$$

According to the law of large numbers (LLN), the random variable  $(\omega - \bar{X})$  converges in probability to  $(\omega - \mu_1)$ . Using asymptotic theory for L-statistics (e.g. [Shorack and Wellner (1986)]), and the underlying distributions with a sufficient number of finite moments, we obtain from [Jone and Zitikis(2007)] the following asymptotic normality result:

$$\sqrt{n} (\bar{\phi}_n(u) - \phi(u)) \xrightarrow{D} \mathcal{N}(0, \sigma_\phi^2), \text{ as } n \rightarrow \infty, \quad (2.10)$$

where

$$\sigma_\phi^2 = \frac{1}{(\omega - \mu_1)^4} \left( (\omega - \mu_1)^2 \sigma_2^2 + \mu_2^2 \sigma_1^2 + \mu_2(\omega - \mu_1) \sigma_{1,2} \right) < \infty,$$

with  $\sigma_1^2$  and  $\sigma_2^2$  represent respectively the variances of  $X$  and  $Y$  and  $\sigma_{1,2} = \int_0^1 \int_0^1 (\min(s, t) - st) dQ_1(1-s) dQ_2(1-t)$ .

Note that in the case where the tail index  $\gamma$  is in the lower half of the unit interval, i.e.  $0 < \gamma < \frac{1}{2}$ , the second-order moments for the two random variables  $X$  and  $Y$  are finite. Consequently, the asymptotic normality of  $\phi_n(u)$  in (2.10) holds. This result is not respected when the loss distribution is heavy-tailed with an index  $\gamma$  located in the upper half of the unit interval, i.e.  $\frac{1}{2} < \gamma < 1$ , since the asymptotic variance  $\sigma_\phi^2$  is infinite, which is due in this case to the infinitability of the second-order moments of the loss  $X$ . In this case,  $\phi(u)$  must be estimated using another approach that would guarantee asymptotic normality.

To remedy this situation, [Rassoul (2014)] used extreme value theory taking into account Hill's estimator [Hill (1975)] estimator of the tail index  $\gamma$  and introduced a semi-parametric estimator for the probability of ruin  $\phi(u)$  for heavy-tailed losses with infinite second-order moments.

The estimation of  $\gamma$  has been extensively studied in the literature, and  $\gamma$  is a positive-tail index. Hill's estimator is the most popular estimator of the positive-tail index  $\gamma$  in extreme value theory, and is defined as follows:

$$\hat{\gamma}_{1,n,k} := k^{-1} \sum_{i=1}^k j (\log X_{n-j+1,n} - \log X_{n-j,n}),$$

for an intermediate sequence  $k = k(n)$ , i.e., are sequences such that  $k \rightarrow \infty$  and  $k/n \rightarrow 0$ , as  $n \rightarrow \infty$ . On the other hand, the Hill estimator associated with the stop loss sample  $(Y_1, \dots, Y_n)$  is given as follows:

$$\hat{\gamma}_{2,n,\ell} := \ell^{-1} \sum_{i=1}^{\ell} j (\log Y_{n-j+1,n} - \log Y_{n-j,n}),$$

where  $\ell = \ell(n)$  is another intermediate sequences satisfying  $\ell \rightarrow \infty$  and  $\ell/n \rightarrow 0$ , as  $n \rightarrow \infty$ . In extreme value theory, Hill's estimator has been extensively studied, improved and even generalized to any real parameter  $\gamma$  (see e.g. [Dekkers et al (1989)], [Beirlant et al. (1999)], [Lo and Fall (2011)]). Its weak consistency was established under the condition of regular variation by [Mason (1982)] assuming only that the underlying distribution varies regularly at infinity. [Deheuvels et al (1988)] proved the strong consistency of Hill's estimator.

However, the asymptotic normality of Hill's estimator has been studied, under various conditions relating to the tail of the distribution, by many researchers, including [Csörgő et al. (1985)], [Beirlant and Teugels (1989)], [Dekkers et al (1989)].

Note that  $\mu_1$  and  $\mu_2$  can be rewritten respectively as follows :

$$\mu_1 = \int_0^{1-k/n} Q_1(s) ds + \int_{1-k/n}^1 Q_1(s) ds \quad (2.11)$$

and

$$\mu_2 = \int_0^{1-\ell/n} Q_2(s) ds + \int_{1-\ell/n}^1 Q_2(s) ds. \quad (2.12)$$

Let us define respectively the following estimators for  $Q_1(s)$  and  $Q_2(s)$ ,  $s \in [0, 1)$ :

$$\widehat{Q}_{1,n,k}(s) = \begin{cases} Q_{1,n}(s) & \text{for } 0 \leq s \leq 1 - k/n, \\ Q_{1,n,k}^W(s) & \text{for } 1 - k/n < s < 1, \end{cases}$$

and

$$\widehat{Q}_{2,n,\ell}(s) = \begin{cases} Q_{2,n}(s) & \text{for } 0 \leq s \leq 1 - \ell/n, \\ Q_{2,n,\ell}^W(s) & \text{for } 1 - \ell/n < s < 1, \end{cases}$$

where  $Q_{1,n}(s)$  and  $Q_{2,n}(s)$  are the above empirical quantile estimators,  $Q_{1,n,k}^W(s) = ((1-s)n/k)^{-\widehat{\gamma}_{1,n,k}^H} X_{n-k,n}$  and  $Q_{2,n,\ell}^W(s) = ((1-s)n/\ell)^{-\widehat{\gamma}_{2,n,\ell}^H} Y_{n-\ell,n}$  are respectively the Weissman's estimators ([?]) of high quantiles  $Q_1(s)$  and  $Q_2(s)$  for  $s \rightarrow 1$ .

Replacing in (2.11),  $Q_1(s)$  (resp. in (2.12),  $Q_2(s)$ ) by  $\widehat{Q}_{1,n,k}(s)$  (resp. by  $\widehat{Q}_{2,n,\ell}(s)$ ), we arrive respectively at the following alternative estimators for the means  $m_1$  and  $m_2$  when losses are heavy tailed with tail index  $\gamma$  in the upper haft of unit interval ( $1/2 < \gamma < 1$ ):

$$\widehat{\mu}_{1,n,k} = n^{-1} \sum_{i=1}^{n-k} X_{i,n} + \frac{k}{n} \frac{X_{n-k,n}}{(1 - \widehat{\gamma}_{1,n,k})},$$

and

$$\widehat{\mu}_{2,n,\ell} = n^{-1} \sum_{i=1}^{n-\ell} Y_{i,n} + \frac{\ell}{n} \frac{Y_{n-\ell,n}}{(1 - \widehat{\gamma}_{2,n,\ell})}.$$

These estimators of means were first studied by [Peng (2001)] and also generalized in [?], [Necir *et al.* (2010)], [?], [Deme *et al.* (2015)] and [Deme *et al.* (2021)] to assess financial and actuarial risk measures.

As in (2.9), substituting  $\widehat{\mu}_{1,n}$  and  $\widehat{\mu}_{2,n}$  with  $\mu_1$  and  $\mu_2$ , respectively, on the right-hand side of the Equation (1.5), [Rassoul (2014)] introduced the following alternative estimator for the ruin probability  $\phi(u)$ :

$$\widetilde{\phi}_{n,k,\ell}(u) := \frac{\widehat{\mu}_{2,n,\ell}}{\omega - \widehat{\mu}_{1,n,k}} \quad \text{for } 1/2 < \gamma < 1. \quad (2.13)$$

[Rassoul (2014)] established the asymptotic normality of the estimator  $\widetilde{\phi}_{n,k,\ell}(u)$  under certain restrictive assumptions. Finally, an asymptotic normal of  $\phi(u)$ , for  $0 < \gamma < 1$  takes the following form:

$$\widehat{\phi}_n(u) := \begin{cases} \overline{\phi}_n(u), & \text{for } 0 < \gamma \leq 1/2, \\ \widetilde{\phi}_{n,k,\ell}(u), & \text{for } 1/2 < \gamma < 1. \end{cases}$$

Note that the alternative ruin probability estimator  $\widetilde{\phi}_{n,k,\ell}(u)$  is associated to the the Hill estimators  $\widehat{\gamma}_{1,n,\ell}$  and  $\widehat{\gamma}_{2,n,\ell}$ .

It is well known that these Hill estimators are both pseudo-maximum likelihood estimators based on the exponential approximation of normalized log-spacings, i.e.,  $V_{1,j} := j(\log X_{n-j+1,n} - \log X_{n-j,n})$ , for  $j = 1, \dots, k$  and  $V_{2,j} := j(\log Y_{n-j+1,n} - \log Y_{n-j,n})$ , for  $j = 1, \dots, \ell$ , see, eg. [Beirlant *et al.* (2004)], [Beirlant *et al.* (1999)]. Clearly, these Hill estimators depend respectively on the choice of sample fractions  $k$ ,  $\ell$  and their influence functions are slowly increasing but not bounded. Consequently, these estimators are not very robust to large values of  $V_{bullet,j}$ , which makes the estimator of the probability of ruin  $\widehat{\phi}_{n,k,\ell}(u)$  sensitive. This constitutes a serious problem in terms of bias and root mean square error (RMSE). To overcome this problem, we introduce in the next section a robust estimator of the probability of ruin  $\phi(u)$  for heavy-tailed distributions whose index lies in the upper half of the unit interval, and establish its asymptotic properties.

### 3 Robust Estimator and main results

#### 3.1 Robust Estimator for the ruin probability $\phi(u)$

To solve the aforementioned problem of the classical Hill estimator, [Fabián and Stehlík (2009)] proposed a sensible distribution known as the t-Hill estimator (t-score or score moment estimate) for the tail index of any distribution that varies the tail regularly. In addition, [Jordanova and Pancheva (2012)] discovered the limiting distribution of the t-Hill estimator in the case where the rank  $S = k$ ,  $\ell$  of the higher-order statistic is  $o(n)$  and proved its asymptotic normality.

This estimator of the score moment has been studied in [Stehlík *et al.*(2010)] and [Stehlík *et al.*(2012)]. According to these authors, this estimator is more robust than the classic Hill estimator. Recently, several studies on t-Hill have been published, see [Beran *et al.* (2014)] and [Jordanova *et al.* (2016)]. In order to improve the quality of the averages  $\mu_1$  and  $\mu_2$  given respectively in (2.11) and (2.12), which allows us to improve the quality of the infinite-time ruin probability for a heavy-tailed distribution, instead of implementing the Hill estimator, we propose to estimate the tail index  $\gamma$  by the so-called t-score moment procedure, in order to obtain a robust result.

The formula of the t-Hill estimators of  $\gamma$  are given by:

$$\widehat{\gamma}_{1,n,k}^{tH} =: \left( \frac{1}{k} \sum_{i=1}^k \frac{X_{n-k,n}}{X_{n-i+1,n}} \right)^{-1} - 1 \quad (3.14)$$

and

$$\widehat{\gamma}_{2,n,\ell}^{tH} =: \left( \frac{1}{\ell} \sum_{i=1}^{\ell} \frac{Y_{n-\ell,n}}{Y_{n-i+1,n}} \right)^{-1} - 1. \quad (3.15)$$

For other robust estimators of the tail index  $\gamma$ , we may refer the reader to [ [Juárez SF and Schucany WR (2004)], [Kim and Lee (2008)], [Peng and Welsh (2001)], [Vandewalle *et al.*(2017)]]].

As already mentioned, the  $\widehat{\phi}_{n,k,\ell}(u)$  estimator given in (1.5) is not robust. To this end, we provide a solution using the t-Hill estimator of  $\gamma$  to derive a robust estimator of the infinite-time probability of ruin for heavy-tailed  $\phi(u)$  distributions. We follow the same method and steps as [Rassoul (2014)] to write this new estimator, but instead of the simple tail index estimators  $\widehat{\gamma}_{1,n,k}^{tH}$  and  $\widehat{\gamma}_{2,n,\ell}^{tH}$  defining respectively in



(3.14) and (3.15).and we introduce the following robust estimators of the ruin probability  $\phi(u)$ :

$$\widehat{\phi}_{n,k,\ell}^{tH}(u) = \frac{\widehat{\mu}_{2,n,\ell}^{tH}}{\omega - \widehat{\mu}_{1,n,k}^{tH}}, \text{ for } 1/2 < \gamma < 1, \quad (3.16)$$

where

$$\widehat{\mu}_{1,n,k}^{tH} := n^{-1} \sum_{i=1}^{n-k} X_{i,n} + \frac{k}{n} \frac{X_{n-k,n}}{(1 - \widehat{\gamma}_{1,n,k}^{tH})}$$

and

$$\widehat{\mu}_{2,n,\ell}^{tH} := n^{-1} \sum_{i=1}^{n-\ell} Y_{i,n} + \frac{\ell}{n} \frac{Y_{n-\ell,n}}{(1 - \widehat{\gamma}_{2,n,\ell}^{tH})}.$$

In the next subsection, we establish the asymptotic properties of our proposed estimator of the probability of ruin.

### 3.2 Asymptotic Results of the estimator $\widehat{\phi}_{n,k,\ell}^{tH}(u)$

As usual in the extreme value framework, to prove asymptotic normality results, we need a second-order condition on the function  $\mathbb{U}_i(x) = Q_i(1 - 1/x)$ ,  $x > 1$ ,  $i = 1, 2$ , such as the following:

**Condition** ( $\mathcal{R}_{\mathbb{U}_i}$ ). There exist a function  $A_i(x) \rightarrow 0$  as  $x \rightarrow \infty$  of constant sign for large values of  $x$  and a second-order parameter  $\rho_i < 0$  such that, for every  $x > 0$

$$\lim_{t \rightarrow \infty} \frac{\log \mathbb{U}_i(tx) - \log \mathbb{U}_i(t) - \gamma \log(x)}{A_i(t)} = \frac{x^{\rho_i} - 1}{\rho_i}, \quad i = 1, 2.$$

Note that condition ( $\mathcal{R}_{\mathbb{U}_i}$ ) implies that  $|A_i|$  is regularly varying with index  $\rho_i$  (see, e.g., [Geluk and de Haan (1987)], [de Haan and Ferreira (2006)]). It is satisfied for most of the classical distribution functions such as the Pareto, Burr, and Fréchet ones.

**Theorem 3.1** *Assume that  $F_i$  satisfies ( $\mathcal{R}_{\mathbb{U}_i}$ ) with  $\gamma \in (1/2, 1)$ . Then for any sequence of integer  $k = k(n)$  and  $\ell = \ell(n)$  satisfying  $k \rightarrow \infty$ ,  $k/n \rightarrow 0$ ,  $\sqrt{k}A_1(n/k) \rightarrow 0$ ,  $\ell \rightarrow \infty$ ,  $\ell/n \rightarrow 0$ ,  $\sqrt{\ell}A_2(n/\ell) \rightarrow 0$  and  $\ell/k \rightarrow \theta < \infty$  as  $n \rightarrow \infty$ , we have:*

$$\frac{\sqrt{n} \left( \widehat{\phi}_{n,k,\ell}^{tH}(u) - \phi(u) \right)}{(k/n)^{1/2} \mathbb{U}_1(n/k)} \stackrel{\mathcal{D}}{=} \kappa_1 \sum_{i=1}^3 \mathbb{W}_{n,i} + \theta^{(1/2-\gamma)} \kappa_2 \sum_{i=1}^3 \overline{\mathbb{W}}_{n,i} + o_{\mathbb{P}}(1)$$

where

$$\left\{ \begin{array}{l} \mathbb{W}_{n,1} = -\sqrt{\frac{n}{k}} \int_0^{1-\frac{k}{n}} \frac{\mathbb{B}_n(s)}{Q_1(1-\frac{k}{n})} dQ_1(s), \\ \mathbb{W}_{n,2} = -\frac{\gamma}{(1-\gamma)} \sqrt{\frac{n}{k}} \mathbb{B}_n\left(1-\frac{k}{n}\right), \\ \mathbb{W}_{n,3} = \frac{\gamma(\gamma+1)^2}{(1-\gamma)^2} \int_0^1 s^{\gamma-1} \mathbb{B}_n(s) ds, \end{array} \right. \quad \left\{ \begin{array}{l} \overline{\mathbb{W}}_{n,1} = -\sqrt{\frac{n}{\ell}} \int_0^{1-\frac{\ell}{n}} \frac{\mathbb{B}_n(s)}{Q_2(1-\frac{\ell}{n})} dQ_2(s), \\ \overline{\mathbb{W}}_{n,2} = -\frac{\gamma}{(1-\gamma)} \sqrt{\frac{n}{\ell}} \mathbb{B}_n\left(1-\frac{\ell}{n}\right), \\ \overline{\mathbb{W}}_{n,3} = \frac{\gamma(\gamma+1)^2}{(1-\gamma)^2} \int_0^1 s^{\gamma-1} \mathbb{B}_n(s) ds, \end{array} \right.$$

with  $\kappa_1 = \mu_2 / (\omega - \mu_1)^2$ ,  $\kappa_2 = (\omega - \mu_1)^{-1}$  and  $\mathbb{B}_n(s)$ ,  $0 \leq s \leq 1$  is a sequence of Brownian, Bridges.

Now, by computing the asymptotic variances of the different processes appearing in Theorem 3.1, we deduce the following corollary:

**Corollary 3.1** *Under the assumptions of Theorem 3.1, we have*

$$\frac{\sqrt{n} \left( \widehat{\phi}_{n,k,\ell}^{tH}(u) - \phi(u) \right)}{(k/n)^{1/2} \mathbb{U}_1(n/k)} \xrightarrow{\mathcal{D}} \mathcal{N}(0, \sigma^2(\gamma, \theta))$$

where

$$\begin{aligned} \sigma^2(\gamma, \theta) &= \frac{\gamma^2}{(1-\gamma)^2(2\gamma-1)} \left( \kappa_1^2 + \kappa_2^2 \theta^{(1-2\gamma)} + 2\kappa_1\kappa_2 \right) \\ &\quad + \frac{\gamma^2(\gamma+1)^2}{(2\gamma+1)(1-\gamma)^4} \left( \kappa_1 + \kappa_2 \theta^{(1/2-\gamma)} \right)^2. \end{aligned}$$

**Remark 3.1** *In the case where  $k \sim \ell$ , as  $n \rightarrow \infty$ , we have  $\theta = 1$  and the asymptotic variance  $\sigma^2(\gamma, \theta)$  is equal to  $4\gamma^5(\kappa_1 + \kappa_2)^2 / (1-\gamma)^4(4\gamma^2 - 1)$ .*

## 4 Simulation Study

### 4.1 Performance and comparative study

In this simulation study, we examine the performance of the new estimator  $\widehat{\phi}_{n,k,\ell}^{tH}(u)$  given in (3.16) with the classical estimator  $\widehat{\phi}_{n,k,\ell}(u)$  proposed by [Rassoul (2014)] and defined in (1.5). Thus, we generate  $N = 1000$  samples  $(X_1, \dots, X_n)$  with the sample size  $n = 1000, 1500, 2000$  from a Pareto distribution function defined as:  $F_1(x) = 1 - x^{-1/\gamma}$ ,  $x \geq 1$  with extreme value index  $\gamma \in \{2/3, 3/4\}$ . For a given initial large reserve  $u = 1.5$ , we derive for each sample its corresponding excess of loss  $(Y_1, \dots, Y_n)$ , where  $Y_j = \max(X_j - u, 0)$ .

The ruin probability estimators  $\widehat{\phi}_{n,k,\ell}^{tH}(u)$  and  $\widehat{\phi}_{n,k,\ell}(u)$  are computed with the parameter  $\omega = c/\lambda = 18$  (in order to ensure that  $\mu_2 < \omega - \mu_1$ ) and with respectively the tail index estimators  $\gamma_{i,n,S}^H$  and  $\gamma_{i,n,S}^{tH}$ ,  $(i, S) \in \{(1, k), (2, \ell)\}$ , for different sample fractional numbers of top order statistics  $k = 1, \dots, n-1$  and  $\ell = 1, \dots, m_n - 1$ , where  $m_n$  is the number of positive values of  $Y_j \neq 0$ ,  $j = 1, \dots, n$ . Employing the algorithm of [Reiss and Thomas (2007)], Page 137, the optimal values  $k^*$  and  $\ell^*$  of the number of top extremes of  $k$  and  $\ell$  to compute the ruin probability estimators are respectively values  $k^*$  and  $\ell^*$  defined as:

$$k^* = \arg \min_k \frac{1}{k} \sum_{j=1}^k j^\delta \left| \widehat{\gamma}_{1,n,j}^\bullet - \text{median} \left( \widehat{\gamma}_{1,n,1}^\bullet, \dots, \widehat{\gamma}_{1,n,k}^\bullet \right) \right|, \quad 1 \leq k \leq n-1, \quad (4.17)$$

and

$$\ell^* = \arg \min_\ell \frac{1}{\ell} \sum_{j=1}^\ell j^\delta \left| \widehat{\gamma}_{2,n,j}^\bullet - \text{median} \left( \widehat{\gamma}_{2,n,1}^\bullet, \dots, \widehat{\gamma}_{2,n,\ell}^\bullet \right) \right|, \quad 1 \leq \ell \leq m_n - 1, \quad (4.18)$$

where  $0 \leq \delta < 1/2$  and  $\widehat{\gamma}_{1,n,k}^\bullet$  (respectively  $\widehat{\gamma}_{2,n,\ell}^\bullet$ ) is either the Hill's or the t-Hill's estimator of the tail index  $\gamma$  computed with the sample  $(X_1, \dots, X_n)$ , respectively with the excess sample  $(Y_1, \dots, Y_n)$ . By the way, choosing  $\delta = 1/4$ , we compute the optimal values  $k^*$  and  $\ell^*$  as in (4.17) and (4.18) for each tail index estimator used in the computation of their associated ruin probability estimators.

• Next, we compare the performance of the above-mentioned ruin probability estimators by computing the absolute value of the mean as well as the mean square errors (MSE) based on the  $N = 500$  simulated samples, and defined as follows:

$$\text{ABias}(\phi_{n,k^*,\ell^*}^\bullet(u)) := \left| \frac{1}{N} \sum_{j=1}^N \frac{\phi_{n,k^*,\ell^*}^{\bullet,j}(u)}{\phi(u)} - 1 \right|$$

and

$$\text{MSE}(\phi_{n,k^*,\ell^*}^\bullet(u)) := \frac{1}{N} \sum_{j=1}^N \left( \frac{\phi_{n,k^*,\ell^*}^{\bullet,j}(u)}{\phi(u)} - 1 \right)^2,$$

where  $\phi(u)$  is the true value of the ruin probability and  $\phi_{n,k^*,\ell^*}^{\bullet,j}(u)$  is the  $j$ -th value ( $j = 1, \dots, N$ ) of any ruin probability estimator  $\phi_{n,k^*,\ell^*}^\bullet(u)$  of  $\phi(u)$ , evaluated at their optimal numbers of higher-order statistics.

The point estimates of the probability of ruin at their optimal  $k^*$  values as well as their ABias and MSE are summarized in the following Tables 4.1.

$n = 1000$							
$\gamma = 2/3, \phi(u) = 0.1$				$\gamma = 3/4, \phi(u) = 0.1576$			
$\widehat{\phi}_{k^*,\ell^*,n}(u)$	0.0893	$\widehat{\phi}_{k^*,\ell^*,n}^{tH}(u)$	0.0943	$\widehat{\phi}_{k^*,\ell^*,n}(u)$	0.1504	$\widehat{\phi}_{k^*,\ell^*,n}^{tH}(u)$	0.1638
ABais	0.1069	ABais	0.0648	ABais	0.0615	ABais	0.0585
MSE	0.0114	MSE	0.0042	MSE	0.0037	MSE	0.0034
$n = 1500$							
$\gamma = 2/3, \phi(u) = 0.1$				$\gamma = 3/4, \phi(u) = 0.1576$			
$\widehat{\phi}_{k^*,\ell^*,n}(u)$	0.0905	$\widehat{\phi}_{k^*,\ell^*,n}^{tH}(u)$	0.0953	$\widehat{\phi}_{k^*,\ell^*,n}(u)$	0.1638	$\widehat{\phi}_{k^*,\ell^*,n}^{tH}(u)$	0.1523
ABais	0.0946	ABais	0.0545	ABais	0.0526	ABais	0.0520
MSE	0.0089	MSE	0.0029	MSE	0.0027	MSE	0.0023
$n = 2000$							
$\gamma = 2/3, \phi(u) = 0.1$				$\gamma = 3/4, \phi(u) = 0.1576$			
$\widehat{\phi}_{k^*,\ell^*,n}(u)$	0.0925	$\widehat{\phi}_{k^*,\ell^*,n}^{tH}(u)$	0.0995	$\widehat{\phi}_{k^*,\ell^*,n}(u)$	0.1649	$\widehat{\phi}_{k^*,\ell^*,n}^{tH}(u)$	0.1546
ABais	0.0751	ABais	0.0354	ABais	0.0502	ABais	0.0358
MSE	0.0056	MSE	0.0013	MSE	0.0025	MSE	0.0012

Table 4.1: Estimation results of  $\widehat{\phi}_{k^*,\ell^*,n}(u)$  and  $\widehat{\phi}_{k^*,\ell^*,n}^{tH}(u)$  estimators of the ruin probability  $\phi(u) = 0.1$  for  $u = 1.5$  and  $\omega = c/\lambda = 18$ , computed with optimal numbers of top statistics  $k^*$  and  $\ell^*$ , based on  $N = 1000$  samples of size  $n = 1000; 1500; 2000$ , from the distribution  $F_1(x) = 1 - x^{-1/\gamma}$ ,  $\gamma = 2/3; 3/4$ .

Examination of the table leads to two conclusions, whatever the situation. Firstly, we note that the absolute bias of both probability of ruin estimators decrease to zero when the sample size  $n$  becomes large. Secondly, we find that the MSEs of the  $\widehat{\phi}_{n,k,\ell}^{tH}(u)$  estimator converge faster to zero as  $n$  increases, compared with the MSEs of  $\widehat{\phi}_{n,k,\ell}(u)$ . In this context, these numerical results show that the estimator  $\widehat{\phi}_{n,k,\ell}^{tH}(u)$  is the best.

## 4.2 Comparative robustness study

One way of increasing robustness is to create a contamination model, which is considered to replace some of the variables of the data  $X$  with outliers. Thus, to assess the robustness of our estimator, a simulation was performed

with contaminated data for each estimator. The main points here are to consider a  $\varepsilon$ -contamination model, which consists in considering a Pareto distribution  $F_1(x) = 1 - x^{-1/\gamma}$  polluted by variables extracted from the other Pareto distribution  $F_{1,a}(x) = 1 - (\frac{x}{a})^{-1/\gamma}$  and to use the following mixing distribution:

$$F_1^c(x) = (1 - \varepsilon) \times F_1(x) + \varepsilon \times F_{1,a}(x) = 1 - (1 - \varepsilon)x^{-1/\gamma} + \varepsilon \left(\frac{x}{a}\right)^{-1/\gamma},$$

where  $\varepsilon \in (0, 1)$  is the contamination rate. Now, for a given  $\varepsilon = 10\%$  and  $a = 3$ , we generate also  $N = 1000$  samples of size  $n = 1000, 1500, 2000$  from the contaminated Pareto distribution  $F_1^c(x)$ . This kind of  $\varepsilon$ -contaminated model is used in [Fabián and Stehlík (2009)], [Brahim and Kenioua (2016)] and [Bouali *et al* (2021)] to evaluate insured.

Next, as in subsection 4.1, we compare the new estimator  $\widehat{\phi}_{n,k,\ell}^{tH}(u)$  with the classical estimator  $\widehat{\phi}_{n,k,\ell}(u)$  of the ruin probability  $\phi(u)$ , by calculating the Absolute bias and the MSE. The results are shown in Table 4.2. It turned out that the effect of contamination gets immediately apparent.

n = 1000							
$\gamma = 2/3, \phi(u) = 0.1$				$\gamma = 3/4, \phi(u) = 0.1576008$			
$\widehat{\phi}_{k^*,\ell^*,n}(u)$	0.0886	$\widehat{\phi}_{k^*,\ell^*,n}^{tH}(u)$	0.0995	$\widehat{\phi}_{k^*,\ell^*,n}(u)$	0.1424	$\widehat{\phi}_{k^*,\ell^*,n}^{tH}(u)$	0.1597
ABais	0.1145	ABais	0.0492	ABais	0.0985	ABais	0.0542
MSE	0.0131	MSE	0.0024	MSE	0.0097	MSE	0.0029
n = 1500							
$\gamma = 2/3, \phi(u) = 0.1$				$\gamma = 3/4, \phi(u) = 0.1576008$			
$\widehat{\phi}_{k^*,\ell^*,n}(u)$	0.0905	$\widehat{\phi}_{k^*,\ell^*,n}^{tH}(u)$	0.1012	$\widehat{\phi}_{k^*,\ell^*,n}(u)$	0.1724	$\widehat{\phi}_{k^*,\ell^*,n}^{tH}(u)$	0.1568
ABais	0.0981	ABais	0.0418	ABais	0.0974	ABais	0.0476
MSE	0.0096	MSE	0.0017	MSE	0.00948	MSE	0.0022
n = 2000							
$\gamma = 2/3, \phi(u) = 0.1$				$\gamma = 3/4, \phi(u) = 0.1576008$			
$\widehat{\phi}_{k^*,\ell^*,n}(u)$	0.0901	$\widehat{\phi}_{k^*,\ell^*,n}^{tH}(u)$	0.1008	$\widehat{\phi}_{k^*,\ell^*,n}(u)$	0.1422	$\widehat{\phi}_{k^*,\ell^*,n}^{tH}(u)$	0.1567
ABais	0.0947	ABais	0.0305	ABais	0.0939	ABais	0.0348
MSE	0.0089	MSE	0.0012	MSE	0.0088	MSE	0.0013

Table 4.2: Estimation results of  $\widehat{\phi}_{k^*,\ell^*,n}(u)$  and  $\widehat{\phi}_{k^*,\ell^*,n}^{tH}(u)$  estimators of the ruin probability  $\phi(u) = ..$  for  $u = ..$ , computed with optimal numbers of top statistics  $k^*$  and  $\ell^*$ , based on  $N = 1000$  samples of size  $n = 1000; 1500; 2000$ , from the distribution  $F_1^c(x) = 1 - (1 - \varepsilon)x^{-1/\gamma} + \varepsilon \left(\frac{x}{a}\right)^{-1/\gamma}$ ,  $\gamma = 2 - 3, 3/4$ .

## 5 Proofs

Let  $E_1, \dots, E_n$  be independent and identically distributed random variables from the unit Pareto distribution  $G$ , defined as  $G(t) = 1 - t^{-1}$ ,  $t \geq 1$ . For each  $n \geq 1$ , let  $E_{1,n} \leq \dots \leq E_{n,n}$  be the order statistics pertaining to  $E_1, \dots, E_n$ . Clearly  $X_{j,n} \stackrel{d}{=} \mathbb{U}_1(E_{j,n})$  and  $Y_{j,n} \stackrel{d}{=} \mathbb{U}_2(E_{j,n})$ ,  $j = 1, \dots, n$ . In order to use the results from [Csörgő *et al.* (1986)], a probability space  $(\Omega, \mathbb{A}, \mathbb{P})$  is constructed carrying a sequence  $\xi_1, \xi_2, \dots$  of independent random variables uniformly distributed on  $(0, 1)$  and a sequence of Brownian bridges  $\mathbb{B}_n(s)$ ,  $0 \leq s \leq 1$ ,  $n = 1, 2, \dots$  such that for all  $0 \leq \nu < 1/2$  and  $\lambda > 0$

$$\sup_{\lambda/n \leq s \leq 1 - \lambda/n} \frac{|\beta_n(s) - \mathbb{B}_n(s)|}{(s(1-s))^{1/2-\nu}} = O_{\mathbb{P}}(n^{-\nu}), \quad (5.19)$$

where  $\beta_n$  is following the uniform quantile process

$$\beta_n(t) = \sqrt{n}(t - V_n(t))$$

with  $V_n$  denoting the empirical uniform quantile function defined to be  $V_n(t) = \xi_{j,n}$ ,  $\frac{j-1}{n} < t \leq \frac{j}{n}$ ,  $j = 1, \dots, n$  and  $V_n(0) = 0$ .

### Proof of Theorem 3.1

Recall that:

$$\widehat{\phi}_{n,k,\ell}^{tH}(u) - \phi(u) = \frac{\widehat{\mu}_{2,n,\ell}^{tH}}{\omega - \widehat{\mu}_{1,n,k}^{tH}} - \frac{\mu_2}{\omega - \mu_1},$$

where

$$\widehat{\mu}_{1,n,k}^{tH} := \int_0^{1-k/n} Q_{1,n}(s) ds + \frac{k}{n} \frac{X_{n-k,n}}{1 - \widehat{\gamma}_{1,n,k}^{tH}},$$

and

$$\widehat{\mu}_{2,n,\ell}^{tH} := \int_0^{1-\ell/n} Q_{2,n}(s) ds + \frac{\ell}{n} \frac{Y_{n-\ell,n}}{1 - \widehat{\gamma}_{2,n,\ell}^{tH}}.$$

Then, one may be rewrite the random variable  $\widehat{\phi}_{n,k,\ell}^{tH}(u) - \phi(u)$  as follows:

$$\widehat{\phi}_{n,k,\ell}^{tH}(u) - \phi(u) = \frac{\mu_2}{(\omega - \widehat{\mu}_{1,n,k}^{tH})(\omega - \mu_1)} \left( \widehat{\mu}_{1,n,k}^{tH} - \mu_1 \right) + \frac{1}{\omega - \widehat{\mu}_{1,n,k}^{tH}} \left( \widehat{\mu}_{2,n,\ell}^{tH} - \mu_2 \right).$$

Under the assumption (1.6), we have from from Theorem 2.4.1 in [de Haan and Ferreira (2006)],  $X_{n-k,n} = Q_1(1 - k/n)\{1 + o_{\mathbb{P}}(1)\}$ , as  $n \rightarrow \infty$ . Since the relation (1.6) is equivalent to  $Q_1(1-s) = s^{-\gamma} \ell_{Q_1}(s)$ ,  $s \in (0, 1)$ , where  $\ell_{Q_1}$  is a slowly varying function, more precisely  $\ell_{Q_1}(sx)/\ell_{Q_1}(s) \rightarrow 1$ , as  $s \rightarrow 0$ , then  $(k/n)Q_1(1 - k/n) = (k/n)^{1-\gamma} \ell_{Q_1}(k/n)$ . Thus, for a given  $\gamma \in (1/2, 1)$ , we have from Proposition 1.3.6 in [Bingham *et al.* (1987)],  $(k/n)^{1-\gamma} \ell_{Q_1}(k/n) \rightarrow 0$ , as  $n \rightarrow \infty$ . Therefore, using the weak consistency of the estimator  $\widehat{\gamma}_{1,n,k}^{tH}$  to  $\gamma$  (see, [Jordanova *et al.* (2016)]), we obtain  $(k/n)X_{n-k,n}/(1 - \widehat{\gamma}_{1,n,k}^{tH}) \xrightarrow{\mathbb{P}} 0$ , as  $n \rightarrow \infty$ .

Also, under assumption we have from [Peng (2001)],

$$\frac{\sqrt{n} \int_0^{1-k/n} (Q_{1,n}(s) - Q_1(s)) ds}{(k/n)^{1/2} Q_1(1 - k/n)} \stackrel{d}{=} - \frac{\int_0^{1-k/n} \mathbb{B}_n(s) dQ_1(s)}{(k/n)^{1/2} Q_1(1 - k/n)} + o_{\mathbb{P}}(1). \quad (5.20)$$

Since the right term in (5.20) is bounded in probability, it comes for all large values of  $n$ ,

$$\int_0^{1-k/n} Q_{1,n}(s) ds = \int_0^{1-k/n} Q_1(s) ds + o_{\mathbb{P}}(1).$$

Remarking that  $k/n \rightarrow 0$ , as  $n \rightarrow \infty$ , we have  $\int_0^{1-k/n} Q_1(s) ds = \int_0^1 Q_1(s) ds \{1 + o_{\mathbb{P}}(1)\}$ , as  $n \rightarrow \infty$ . Which leads to the convergence in probability of  $\widehat{\mu}_{1,n,k}^{tH}$  to the mean  $\mu_1$ . Next, let's denote by  $\kappa_1 = \mu_2/(\omega - \mu_1)^2$  and  $\kappa_2 = 1/(\omega - \mu_1)$ . Then, for all  $n$  large enough, the random variable  $\widehat{\phi}_{n,k,\ell}^{tH}(u) - \phi(u)$  can be also represented as follows

$$\widehat{\phi}_{n,k,\ell}^{tH}(u) - \phi(u) \stackrel{d}{=} \kappa_1 \left( \widehat{\mu}_{1,n,k}^{tH} - \mu_1 \right) + \kappa_2 \left( \widehat{\mu}_{2,n,\ell}^{tH} - \mu_2 \right).$$

Consequently,

$$\begin{aligned} \frac{\sqrt{n} \left( \widehat{\phi}_{n,k,\ell}^{tH}(u) - \phi(u) \right)}{(k/n)^{1/2} Q_1(1-k/n)} &\stackrel{d}{=} \kappa_1 \frac{n^{1/2} (\widehat{\mu}_{1,n,k}^{tH} - \mu_1)}{(k/n)^{1/2} Q_1(1-k/n)} \\ &\quad + \kappa_2 \sqrt{\ell/k} \left( \frac{Q_2(1-\ell/n)}{Q_1(1-k/n)} \right) \frac{n^{1/2} (\widehat{\mu}_{2,n,\ell}^{tH} - \mu_2)}{(\ell/n)^{1/2} Q_2(1-\ell/n)}, \\ &:= A_{1,n} + A_{2,n}. \end{aligned}$$

Now, let We first compute the  $A_{1,n}$  term. By substituting  $\widehat{\mu}_{1,n,k}^{tH}$  and  $\mu_1$  with their expressions, we have

$$\begin{aligned} A_{1,n} &= \kappa_1 \frac{\sqrt{n} \int_0^{1-k/n} (Q_{1,n}(s) - Q_1(s)) ds}{(k/n)^{1/2} Q_1(1-k/n)} \\ &\quad + \kappa_1 \frac{n^{1/2} \left( \frac{(k/n) X_{n-k,n}}{1 - \widehat{\gamma}_{1,n,k}^{tH}} - \int_{1-k/n}^1 Q_1(s) ds \right)}{(k/n)^{1/2} Q_1(1-k/n)} \\ &:= A_{1,n}^{(1)} + A_{1,n}^{(2)}. \end{aligned}$$

From the statement in (5.20), we have

$$A_{1,n}^{(1)} \stackrel{d}{=} -\kappa_1 \mathbb{W}_{n,1} + o_{\mathbb{P}}(1), \quad (5.21)$$

where

$$\mathbb{W}_{n,1} := \frac{\int_0^{1-k/n} \mathbb{B}_n(s) dQ_1(s)}{(k/n)^{1/2} Q_1(1-k/n)}.$$

Next, remarking that  $\mathbb{U}_1(n/k) = Q_1(1-k/n)$  and  $X_{n-k,n} \stackrel{d}{=} \mathbb{U}_1(E_{n-k,n})$ , we have

$$A_{1,n}^{(2)} \stackrel{d}{=} \sum_{i=1}^4 T_{n,i},$$

where

$$\begin{aligned} T_{n,1} &= \frac{\kappa_1}{1 - \widehat{\gamma}_{1,n,k}^{tH}} \sqrt{k} \left[ \frac{\mathbb{U}_1(E_{n-k,n})}{\mathbb{U}_1(n/k)} - \left( \frac{k}{n} E_{n-k,n} \right)^\gamma \right], \\ T_{n,2} &= \frac{\kappa_1}{1 - \widehat{\gamma}_{1,n,k}^{tH}} \sqrt{k} \left( \left( \frac{k}{n} E_{n-k,n} \right)^\gamma - 1 \right), \\ T_{n,3} &= \frac{\kappa_1}{(1-\gamma)(1 - \widehat{\gamma}_{1,n,k}^{tH})} \sqrt{k} (\widehat{\gamma}_{1,n,k}^{tH} - \gamma), \\ T_{n,4} &= \kappa_1 \sqrt{k} \left[ \frac{1}{1-\gamma} - \frac{\int_1^{+\infty} s^{-2} \mathbb{U}_1(ns/k) ds}{\mathbb{U}_1(n/k)} \right]. \end{aligned}$$

We study each term separately.

Term  $T_{n,1}$ . According to [de Haan and Ferreira (2006)], Theorem 2.3.9), for any  $\delta > 0$ , we have

$$\sqrt{k} \left( \frac{\mathbb{U}_1(E_{n-k,n})}{\mathbb{U}_1(n/k)} - \left( \frac{k}{n} E_{n-k,n} \right)^\gamma \right) = \sqrt{k} A_1 \left( \frac{n}{k} \right) \left\{ \left( \frac{k}{n} E_{n-k,n} \right)^\gamma \frac{\left( \frac{k}{n} E_{n-k,n} \right)^\rho - 1}{\rho} + o_{\mathbb{P}}(1) \left( \frac{k}{n} E_{n-k,n} \right)^{\gamma+\rho\pm\delta} \right\},$$

Thus, since  $kE_{n-k,n}/n \rightarrow 1$ ,  $\sqrt{k} A_1(n/k) \rightarrow 0$  and  $\widehat{\gamma}_{1,n,k}^{tH} \xrightarrow{\mathbb{P}} \gamma$ , as  $n \rightarrow \infty$  (see, [Jordanova et al (2016)]), it readily follows that

$$T_{n,1} = o_{\mathbb{P}}(1). \quad (5.22)$$

Term  $T_{n,2}$ . The equality  $E_{n-k,n} \stackrel{d}{=} (1 - \xi_{n-k,n})^{-1}$  yields:

$$\begin{aligned} \sqrt{k} \left[ \left( \frac{k}{n} E_{n-k,n} \right)^\gamma - 1 \right] &\stackrel{d}{=} \sqrt{k} \left( \left( \frac{n}{k} (1 - \xi_{n-k,n}) \right)^{-\gamma} - 1 \right) \\ &= -\gamma \sqrt{k} \left( \frac{n}{k} (1 - \xi_{n-k,n}) - 1 \right) (1 + o_{\mathbb{P}}(1)) \quad \text{by a Taylor expansion} \\ &= -\gamma \sqrt{\frac{n}{k}} \beta_n \left( 1 - \frac{k}{n} \right) (1 + o_{\mathbb{P}}(1)) \\ &= -\gamma \sqrt{\frac{n}{k}} \left( \mathbb{B}_n \left( 1 - \frac{k}{n} \right) + O_{\mathbb{P}}(n^{-\nu}) \left( \frac{k}{n} \right)^{1/2-\nu} \right) (1 + o_{\mathbb{P}}(1)), \end{aligned}$$

for  $0 \leq \nu < 1/2$ , by [Csörgő *et al.* (1986)]. Thus, using again the weak consistency of  $\widehat{\gamma}_{1,n,k}^{tH}$  to  $\gamma$ , it follows that:

$$T_{n,2} \stackrel{d}{=} \kappa_1 \mathbb{W}_{n,2}(1 + o_{\mathbb{P}}(1)), \quad (5.23)$$

where

$$\mathbb{W}_{n,2} = -\frac{\gamma}{(1-\gamma)} \sqrt{\frac{n}{k}} \mathbb{B}_n \left(1 - \frac{k}{n}\right).$$

Term  $T_{n,3}$ . From the Proposition 1 in [Brahim and Kenioua (2016)], page 877, we have

$$\sqrt{k} \left( \widehat{\gamma}_{1,n,k}^{tH} - \gamma \right) \stackrel{d}{=} \gamma(\gamma+1)^2 \int_0^1 s^{\gamma-1} \mathbb{B}_n(s) ds + o_{\mathbb{P}}(1).$$

And by using again the consistency in probability of  $\widehat{\gamma}_{1,n,k}^{tH}$  to  $\gamma$ , we get for all  $n$  large enough:

$$T_{n,3} = \kappa_1 \frac{\gamma(\gamma+1)^2}{(1-\gamma)^2} \int_0^1 s^{\gamma-1} \mathbb{B}_n(s) ds + o_{\mathbb{P}}(1) = \kappa_1 \mathbb{W}_{n,3} + o_{\mathbb{P}}(1). \quad (5.24)$$

Term  $T_{n,4}$ . A change of variables and an integration by parts yield

$$\begin{aligned} T_{n,4} &= \kappa_1 \sqrt{k} \left\{ \frac{1}{1-\gamma} - \int_1^\infty x^{-2} \frac{\mathbb{U}_1(nx/k)}{\mathbb{U}_1(n/k)} dx \right\} \\ &= -\kappa_1 \sqrt{k} \int_1^\infty x^{-2} \left( \frac{\mathbb{U}_1(nx/k)}{\mathbb{U}_1(n/k)} - x^\gamma \right) dx. \end{aligned}$$

Theorem 2.3.9 in [de Haan and Ferreira (2006)] entails that, for  $\gamma \in (1/2, 1)$ ,

$$\begin{aligned} T_{n,4} &= -\kappa_1 \sqrt{k} A_1 \left( \frac{n}{k} \right) \int_1^\infty x^{\gamma-2} \frac{x^\rho - 1}{\rho} dx (1 + o_{\mathbb{P}}(1)) \\ &= \kappa_1 \sqrt{k} A_1 \left( \frac{n}{k} \right) \frac{1}{(1-\gamma)(\gamma+\rho-1)} (1 + o_{\mathbb{P}}(1)). \\ &= o_{\mathbb{P}}(1), \quad \text{by } \sqrt{k} A_1(n/k) \rightarrow 0. \end{aligned} \quad (5.25)$$

Combining (5.21), (5.22), (5.23), (5.24) and (5.25), we get

$$A_{n,1} \stackrel{d}{=} \kappa_1 \left( \mathbb{W}_{n,1} + \mathbb{W}_{n,2} + \mathbb{W}_{n,3} \right) + o_{\mathbb{P}}(1). \quad (5.26)$$

Now let's compute the  $A_{2,n}$  term. We first have:  $Q_2(1 - \frac{\ell}{n}) \sim Q_1(1 - \frac{\ell}{n}) = Q_1(1 - \frac{\ell k}{k n}) \sim (\frac{\ell}{k})^{-\gamma} Q_1(1 - \frac{k}{n})$ , as  $n \rightarrow \infty$ . Since by assumption  $\ell/k \rightarrow \theta > 0$ , then  $Q_2(1 - \frac{\ell}{n}) \sim \theta^{-\gamma} Q_1(1 - \frac{k}{n})$ , as  $n \rightarrow \infty$ . This leads for large values of  $n$  to

$$A_{2,n} \stackrel{d}{=} \theta^{(1/2-\gamma)} \kappa_2 \frac{n^{1/2} (\widehat{\mu}_{2,n,\ell}^{tH} - \mu_2)}{(\ell/n)^{1/2} Q_2(1 - \ell/n)}.$$

Further, Substituting  $\widehat{\mu}_{2,n,\ell}^{tH}$  and  $\mu_2$  with their expressions and using similar arguments as those developed to show the expression  $A_{1,n}$  in (5.26) together with  $Y_{n-\ell,n} \stackrel{d}{=} \mathbb{U}_2(E_{n-\ell,n})$ , it comes:

$$A_{n,2} \stackrel{d}{=} \theta^{(1/2-\gamma)} \kappa_2 (\overline{\mathbb{W}}_{n,1} + \overline{\mathbb{W}}_{n,2} + \overline{\mathbb{W}}_{n,3}) + o_{\mathbb{P}}(1), \quad (5.27)$$

where

$$\begin{aligned} \overline{\mathbb{W}}_{n,1} &:= -\frac{\int_0^{1-\ell/n} \mathbb{B}_n(s) dQ_2(s)}{(\ell/n)^{1/2} Q_2(1 - \ell/n)}, \\ \overline{\mathbb{W}}_{n,2} &:= -\frac{\gamma}{1-\gamma} \sqrt{\frac{n}{\ell}} \mathbb{B}_n \left(1 - \frac{\ell}{n}\right), \\ \overline{\mathbb{W}}_{n,3} &:= \frac{\gamma(\gamma+1)^2}{(1-\gamma)^2} \int_0^1 s^{\gamma-1} \mathbb{B}_n(s) ds. \end{aligned}$$

Finally, for all  $n$  large enough, the Theorem 3.1 holds with

$$\frac{\sqrt{n} \left( \widehat{\phi}_{n,k,\ell}^{tH}(u) - \phi(u) \right)}{(k/n)^{1/2} Q_1(1 - k/n)} \stackrel{d}{=} \kappa_1 \sum_{i=1}^3 \mathbb{W}_{n,i} + \theta^{(1/2-\gamma)} \kappa_2 \sum_{i=1}^3 \overline{\mathbb{W}}_{n,i} + o_{\mathbb{P}}(1)$$

■

### Proof of Corollary 3.1

From Theorem 3.1, we need to compute the asymptotic variance  $\sigma^2(\gamma, \theta)$  of the limiting process. More precisely, have

$$\begin{aligned} \sigma^2(\gamma, \theta) = & \lim_{n \rightarrow \infty} \left( (\kappa_1)^2 \left\{ \mathbb{E}(\mathbb{W}_{n,1}^2) + \mathbb{E}(\mathbb{W}_{n,2}^2) + \mathbb{E}(\mathbb{W}_{n,3}^2) \right\} \right. \\ & + (\kappa_2)^2 \theta^{(1-2\gamma)} \left\{ \mathbb{E}(\overline{\mathbb{W}}_{n,1}^2) + \mathbb{E}(\overline{\mathbb{W}}_{n,2}^2) + \mathbb{E}(\overline{\mathbb{W}}_{n,3}^2) \right\} \\ & + 2(\kappa_1)^2 \left\{ \mathbb{E}(\mathbb{W}_{n,1}\mathbb{W}_{n,2}) + \mathbb{E}(\mathbb{W}_{n,1}\mathbb{W}_{n,3}) + \mathbb{E}(\mathbb{W}_{n,2}\mathbb{W}_{n,3}) \right\} \\ & + 2(\kappa_2)^2 \theta^{(1-2\gamma)} \left\{ \mathbb{E}(\overline{\mathbb{W}}_{n,1}\overline{\mathbb{W}}_{n,2}) + \mathbb{E}(\overline{\mathbb{W}}_{n,1}\overline{\mathbb{W}}_{n,3}) + \mathbb{E}(\overline{\mathbb{W}}_{n,2}\overline{\mathbb{W}}_{n,3}) \right\} \\ & + 2\kappa_1\kappa_2 \theta^{(1/2-\gamma)} \left\{ \mathbb{E}(\mathbb{W}_{n,1}\overline{\mathbb{W}}_{n,1}) + \mathbb{E}(\mathbb{W}_{n,1}\overline{\mathbb{W}}_{n,2}) + \mathbb{E}(\mathbb{W}_{n,1}\overline{\mathbb{W}}_{n,3}) \right\} \\ & + 2\kappa_1\kappa_2 \theta^{(1/2-\gamma)} \left\{ \mathbb{E}(\mathbb{W}_{n,2}\overline{\mathbb{W}}_{n,1}) + \mathbb{E}(\mathbb{W}_{n,2}\overline{\mathbb{W}}_{n,2}) + \mathbb{E}(\mathbb{W}_{n,2}\overline{\mathbb{W}}_{n,3}) \right\} \\ & \left. + 2\kappa_1\kappa_2 \theta^{(1/2-\gamma)} \left\{ \mathbb{E}(\mathbb{W}_{n,3}\overline{\mathbb{W}}_{n,1}) + \mathbb{E}(\mathbb{W}_{n,3}\overline{\mathbb{W}}_{n,2}) + \mathbb{E}(\mathbb{W}_{n,3}\overline{\mathbb{W}}_{n,3}) \right\} \right). \end{aligned}$$

Recall that  $Q_1(1-su)/Q_1(1-u) \rightarrow s^{-\gamma}$ ,  $Q_2(1-su)/Q_2(1-u) \rightarrow s^{-\gamma}$  and  $Q_1(1-u) \sim Q_2(1-u)$  as  $u \rightarrow 0$ ,  $u \rightarrow 0$  and  $0u \rightarrow 0$ . Following [Deme *et al.* (2015)] with some algebraic operations, we get, as  $n \rightarrow \infty$ :

$$\begin{aligned} \mathbb{E}(\mathbb{W}_{n,1}^2) &\rightarrow \frac{2\gamma}{2\gamma-1}, & \mathbb{E}(\mathbb{W}_{n,2}^2) &\rightarrow \frac{\gamma^2}{(1-\gamma)^2}, \\ \mathbb{E}(\mathbb{W}_{n,3}^2) &= \frac{\gamma^2(\gamma+1)^2}{(2\gamma+1)(\gamma-1)^4}, & \mathbb{E}(\mathbb{W}_{n,1}\mathbb{W}_{n,2}) &\rightarrow \frac{\gamma}{1-\gamma}, \\ \mathbb{E}(\mathbb{W}_{n,1}\mathbb{W}_{n,3}) &= 0 + o(1), & \mathbb{E}(\mathbb{W}_{n,2}\mathbb{W}_{n,3}) &= 0 + o(1), \\ \mathbb{E}(\overline{\mathbb{W}}_{n,1}^2) &\rightarrow \frac{2\gamma}{2\gamma-1}, & \mathbb{E}(\overline{\mathbb{W}}_{n,2}^2) &\rightarrow \frac{\gamma^2}{(1-\gamma)^2}, \\ \mathbb{E}(\overline{\mathbb{W}}_{n,3}^2) &= \frac{\gamma^2(\gamma+1)^2}{(2\gamma+1)(\gamma-1)^4}, & \mathbb{E}(\overline{\mathbb{W}}_{n,1}\overline{\mathbb{W}}_{n,2}) &\rightarrow \frac{\gamma}{1-\gamma}, \\ \mathbb{E}(\overline{\mathbb{W}}_{n,1}\overline{\mathbb{W}}_{n,3}) &= 0 + o(1), & \mathbb{E}(\overline{\mathbb{W}}_{n,2}\overline{\mathbb{W}}_{n,3}) &= 0 + o(1), \\ \mathbb{E}(\mathbb{W}_{n,1}\overline{\mathbb{W}}_{n,1}) &\rightarrow \frac{\gamma}{1-\gamma} \left( \frac{1}{2\gamma-1} - \theta^{(1-\gamma)} \right) \theta^{(\gamma-1/2)}, & \mathbb{E}(\mathbb{W}_{n,1}\overline{\mathbb{W}}_{n,2}) &\rightarrow \frac{\gamma}{1-\gamma} \theta^{1/2}, \\ \mathbb{E}(\mathbb{W}_{n,1}\overline{\mathbb{W}}_{n,3}) &= 0 + o(1), & \mathbb{E}(\mathbb{W}_{n,2}\overline{\mathbb{W}}_{n,1}) &\rightarrow \frac{\gamma(1-\gamma\theta^{(1-\gamma)})}{(1-\gamma)^2} \theta^{(\gamma-1/2)}, \\ \mathbb{E}(\mathbb{W}_{n,2}\overline{\mathbb{W}}_{n,2}) &\rightarrow \frac{\gamma^2}{(1-\gamma)^2} \theta^{1/2}, & \mathbb{E}(\mathbb{W}_{n,2}\overline{\mathbb{W}}_{n,3}) &= 0 + o(1), \\ \mathbb{E}(\mathbb{W}_{n,3}\overline{\mathbb{W}}_{n,1}) &= 0 + o(1), & \mathbb{E}(\mathbb{W}_{n,3}\overline{\mathbb{W}}_{n,2}) &= 0 + o(1), & \mathbb{E}(\mathbb{W}_{n,3}\overline{\mathbb{W}}_{n,3}) &= \frac{\gamma^2(1+\gamma)^2}{(1+2\gamma)(1-\gamma)^4}. \end{aligned}$$

## References

- [Asmussen (2000)] Asmussen, S. (2000). *Ruin Probabilities*, World Scientific, Singapore, New Jersey, London, Hong Kong.
- [Beirlant *et al* (2001)] Beirlant, J., Matthys, G., Dierckx, G. (2001). Heavy-tailed distributions and rating, *Astin Bull.*, **31**, 37–58.
- [Beran *et al* (2014)] Beran J., Schell D., Stehlik M. (2014). The harmonic moment tail index estimator: asymptotic distribution and robustness, *Ann. Inst. Statist. Math.*, **66**(1), 193–220.



- [Beirlant *et al.* (2004)] Beirlant, J., Goegebeur, Y., Teugels, J. and Segers, J. (2004). Statistics of extremes: theory and applications. *Wiley Series in Probability and Statistics. John Wiley and Sons Ltd.*
- [Beirlant *et al.* (1999)] Beirlant, J., Dierckx, G. Y., and Matthys G. (1999). Tail index estimation and an exponential regression model. *Extremes*, **2**, 177–200.
- [Beirlant and Teugels (1989)] Beirlant, J., Teugels, J. (1989). Asymptotic normality of Hill’s estimator. In: Extreme Value Theory (Oberwolfach, 1987), *Lecture Notes in Statist*, **51**, 148–155.
- [Bingham *et al.* (1987)] Bingham, N. H., Goldie, C.M., Teugels, J.L. (1987). Regular variation, Cambridge.
- [Bouali *et al.* (2021)] Bouali, D. L., Bouali1, Benatia1, F., Brahimi1, B., Chesneau, C. (2021). Robust Estimator of Conditional Tail Expectation of Pareto-Type Distribution, *J. Stat. Theory Pract.* **15**, 16 <https://doi.org/10.1007/s42519-020-00153-0>.
- [Brahim and Kenioua (2016)] Brahim, B. and Kenioua, Z. (2016). Robust estimator of distortion risk premiums for heavy-tailed losses, *Afrika Statistika.*, **11** (1), 869–882.
- [Brazauskas *et al.* (2008)] Brazauskas, V., Jones, B., Puri, M., Zitikis, R. (2008). Estimating conditional tail expectation with actuarial applications in view, *J. Statist. Plann. Inference*, **138**, 3590-3604.
- [Čížek *et al.* (2005)] Čížek, P., Härdle, W., Weron, R.: Statistical Tools for Finance and Insurance. Springer, Berlin-Heidelberg (2005).
- [Csörgő *et al.* (1985)] Csörgő, S., Deheuvels, P., Mason, D.M. (1985). Kernel estimates of the tail index of a distribution, *Ann. Statist.*, **13**, 1050-1077.
- [Csörgő *et al.* (1986)] Csörgő, M., Csörgő, S., Horváth, L., Mason, D.M. (1986). Weighted empirical and quantile processes, *Ann. Probab.*, **14**, 31–85.
- [de Haan and Ferreira (2006)] de Haan, L., Ferreira, A. (2006). *Extreme value theory: an introduction*, Springer.
- [Deheuvels *et al.* (1988)] Deheuvels, P., Haeusler, E., Mason, D. (1988). Almost sure convergence of the Hill estimator, *Math. Proc. Camb. Philos. Soc*, 104, 371–381.
- [Dekkers *et al.* (1989)] Dekkers, A., Einmahl, J., de Haan, L. (1989). A moment estimator for the index of an extreme-value distribution, *Ann. Statist*, 17, 1833–1855.
- [Deme *et al.* (2013a)] Deme, E., Girard, S., Guillou, A. (2013a). Reduced-bias estimator of the Proportional Hazard Premium for heavy-tailed distributions, *Insurance Math. Econom.*, **52**, 550-559.
- [Deme *et al.* (2013b)] Deme E.H., Gardes L. and Girard S. (2013b). On the estimation of the second order parameter for heavy-tailed distributions. *REVSTAT*, **113**, 277–299 .
- [Deme *et al.* (2015)] Deme, E.H., Girard, S., Guillou A., (2015) Reduced-biased estimators of the Conditional Tail Expectation for heavy-tailed distributions, *Mathematical statistics and limit theorems, Springer, 105-123*
- [Deme *et al.* (2021)] Deme E.H., Allaya M., Deme S, Dhaker A. and Dabye A. S. (2021). Estimation of risk measures from heavy-tailed distributions. *east journal of theoretical statistics*. volume **62**,1, p. 35
- [Fabián and Stehlík (2009)] Fabián Z. and Stehlík M. (2009). On robust and distribution sensitive Hill like method, IFAS research report, **43**.
- [Geluk and de Hann (1987)] Geluk, J. L., de Haan, L., (1987). Regular variation, extensions and Tauberian theorems: CWI tract 40, Center for Mathematics and Computer Science, P.O. Box 4079, 1009 AB Amsterdam, The Netherlands.
- [Hill (1975)] Hill, B.M. (1975). A simple approach to inference about the tail of a distribution, *Ann. Statist.*, **3**, 1136-1174.
- [Jordanova and Pancheva (2012)] Jordanova PK., Pancheva EI. (2012). Weak asymptotic results for t-Hill estimator, *Comptes rendus de l’académie bulgare des sciences*, **65**(12),1649–1656

- [Jordanova *et al* (2016)] Jordanova P., Fabián Z., Hermann P, Strelec L., Rivera A., Girard S., Torres S., Stehlík M. (2016). Weak properties and robustness of t-Hill estimators, *Extremes*, **19**(4),591–626.
- [Juárez SF and Schucany WR (2004)] Juárez SF., Schucany WR. (2004). Robust and efficient estimation for the generalized Pareto distribution. *Extremes*. **7**(3),237–251.
- [Kim and Lee (2008)] Kim M., Lee S. (2008). Estimation of a tail index based on minimum density power divergence, *J. Multivariate Anal.*, **99**(10), 2453–2471.
- [Jone and Zitikis(2007)] Jones, B. L. and Zitikis, R., (2007). Risk measures, distortion parameters and their empirical estimation. *Insurance Math. Econom.*, **41**(2), 279–297.
- [Lo and Fall (2011)] Lo, G.S, Fall, A.M. (2011). Another look at Second order condition in Extreme Value Theory, *Afr. Stat.*, **6**, 346–370.
- [Mason (1982)] Mason, D.(1982). Laws of large numbers for sums of extreme values, *Ann. Probab.*, **10**, 754–764.
- [Matthys *et al.* (2004)] Matthys, G., Delafosse, E., Guillou, A., Beirlant, J. (2004). Estimating catastrophic quantile levels for heavy-tailed distributions, *Insurance Math. Econom.*, **34**, 517-537.
- [Necir *et al.* (2010)] Necir, A., Rassoul, A., Zitikis, R. (2010). Estimating the conditional tail expectation in the case of heavy-tailed losses, *J. Probab. Stat.*, **ID 596839**, 17 pages.
- [Panjer and Willmot (1992)] Panjer, H., Willmot, G.E. (1992). *Insurance Risk Models*, Society of actuaries, Michigan.
- [Peng (2001)] Peng, L. (2001). Estimating the mean of a heavy tailed distribution. *Stat. Prob. Lett* **52**, 255–264.
- [Peng and Welsh (2001)] Peng L., Welsh AH. (2001), Robust estimation of the generalized Pareto distribution, *Extremes*, **4**(1),53–65.
- [Rassoul (2014)] Rassoul, A. (2014). Estimation of the Ruin Probability in Infinite Time for Heavy Right-Tailed Losses.
- [Rolski *et al* (1999)] Rolski, T., Schmidli, H., Schmid, V., Teugels, J.L. (1999). *Stochastic Processes for Insurance and Finance*, John Wiley and Sons, New York.
- [Reiss and Thomas (2007)] Reiss, R.D., Thomas, M., (2007). *Statistical Analysis of Extreme Values with Applications to Insurance, Finance, Hydrology and Other Fields*, 3rd ed. Birkhäuser.
- [Shorack and Wellner (1986)] Shorack, G. and Wellner J. (1986). *Empirical Processes with Applications to Statistics*. John Wiley Sons, New York.
- [Stehlík *et al*(2010)] Stehlík M., Potocký R., Waldl H., Fabián Z. (2010). On the favorable estimation for fitting heavy tailed data, *Comput. Statist.*, **25**(3),485–503.
- [Stehlík *et al*(2012)] Stehlík M., Fabián Z., Střelec L. (2012). Small sample robust testing for normality against Pareto tails, *Comm. Statist. Simulation Comput.*, **41**(7),1167–1194.
- [Vandewalle and Beirlant (2006)] Vandewalle, B., Beirlant, J. (2006). On univariate extreme value statistics and the estimation of reinsurance premiums. *Insurance, Mathematics and Economics*, **38**, 441-459.
- [Vandewalle *et al*(2017)] Vandewalle B., Beirlant J., Christmann A., Hubert M. (2007). A robust estimator for the tail index of Pareto-type distributions, *Comput. Statist. Data Anal.*, **51**(12), 6252–6268.
- [Weissman (1978)] Weissman, I.(1978). Estimation of parameters and large quantiles based on the k largest observations, *J. Amer. Statist. Assoc.*, **73**, 812–815.