



HAL
open science

The N-terminus of the cauliflower mosaic virus aphid transmission protein P2 is involved in transmission body formation and microtubule interaction

Christiane Then, Aurélie Bak, Alexandre Morisset, Beatriz Dáder, Marie Ducouso, Jean-Luc Macia, Martin Drucker

► To cite this version:

Christiane Then, Aurélie Bak, Alexandre Morisset, Beatriz Dáder, Marie Ducouso, et al.. The N-terminus of the cauliflower mosaic virus aphid transmission protein P2 is involved in transmission body formation and microtubule interaction. *Virus Research*, 2021, 297, pp.198356. 10.1016/j.virusres.2021.198356 . hal-04185405

HAL Id: hal-04185405

<https://hal.science/hal-04185405v1>

Submitted on 22 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

The N-terminus of the cauliflower mosaic virus aphid transmission protein P2 is involved in transmission body formation and microtubule interaction

5 Christiane Then^{1,2}, Aurélie Bak^{1,3}, Alexandre Morisset¹, Beatriz Dáder^{1,4}, Marie Ducousso¹, Jean-Luc Macia¹, Martin Drucker^{1,5,6}

¹ INRAE Centre Occitanie - Montpellier, France

² Present address: Julius-Kühn-Institut, Braunschweig, Germany

10 ³ Present address: KWS Gateway Research Center, St Louis, Missouri, USA

⁴ Present address: ETSIAAB, Universidad Politécnica de Madrid, Madrid, Spain

⁵ Present address: INRAE Centre Grand Est – Colmar, France

⁶ Corresponding author: martin.drucker@inrae.fr

Abstract

15 Cauliflower mosaic virus (CaMV) is transmitted by aphids using the non-circulative transmission
mode: when the insects feed on infected leaves, virus particles from infected cells attach rapidly to
their stylets and are transmitted to a new host when the aphids change plants. Mandatory for CaMV
transmission, the viral helper protein P2 mediates as a molecular linker binding of the virus particles
20 to the aphid stylets. P2 is available in infected plant cells in a viral inclusion that is specialized for
transmission and named the transmission body (TB). When puncturing an infected leaf cell, the
aphid triggers an ultra-rapid viral response, necessary for virus acquisition and called transmission
activation: The TB disrupts and P2 is redistributed onto cortical microtubules, together with virus
particles that are simultaneously set free from virus factories and join P2 on the microtubules to
form the so-called mixed networks (MNs). The MNs are the predominant structure from which
25 CaMV is acquired by aphids. However, the P2 domains involved in microtubule interaction are not
known. To identify P2 regions involved in its functions, we generated a set of P2 mutants by alanine
scanning and analyzed them in the viral context for their capacity to form a TB, to interact with
microtubules and to transmit CaMV. Our results show that contrary to the previously characterized
P2-P2 and P2-virion binding sites in its C-terminus, the microtubule binding site is contained in the
30 N-terminal half of P2. Further, this region is important for TB formation since some P2 mutant
proteins did not accumulate in TBs but were retained in the viral factories where P2 is translated.
Taken together, the N-terminus of P2 is not only involved in vector interaction as previously
reported, but also in interaction with microtubules and in formation of TBs.

Keywords

35 Plant virus, aphid vector, transmission, cell biology, microtubules, cauliflower mosaic virus.

Introduction

Cauliflower mosaic virus (CaMV) is like hundreds of other plant viruses transmitted by aphids (reviewed in ^{1,2}). The aphid vectors acquire CaMV particles (virions) when feeding on infected plants and transport them attached to the aphid mouthparts (the stylets) to a new host plant, where the virions are inoculated during subsequent feeding activity. In the stylets, the icosahedral virions bind to a receptor belonging to the stylin proteins and localized at the stylet tips ^{3,4}. However, virions alone can not bind to the vector stylets. For this, they need P2, the virus-encoded transmission helper protein of CaMV. P2 complexes with virions to form a transmission unit that binds to the stylets ^{5,6}. CaMV acquisition from infected tissue is a complicated process because P2 is not complexed with virions in infected cells. Rather, virions localize mainly in distinct cytoplasmic virus-induced inclusions, so-called virus factories (VF), whereas P2 is found in transmission bodies (TB), another type of cytoplasmic viral inclusions, where it colocalizes with viral protein P3 ⁷⁻⁹. A consequence of this set-up is that P2 and virions must form ad hoc transmission units when aphids probe infected plant cells ⁷. This can happen either in the infected cell or in the aphid stylets, where it allows for sequential acquisition of CaMV: first attachment of P2 on stylet receptors, then binding of virions on stylet-associated P2 ¹⁰. Both transmission pathways are facilitated by a viral-cellular reaction induced by the aphid feeding activity ^{11,12}. In their quest for nutrient-filled sap transported within the phloem vessels, aphids insert their stylets into cells located in the epidermal and mesophyll layers for a sampling of their contents. The stylet punctures wound but do not kill the visited cells and trigger instantaneous dissolution of the transmission bodies and the redistribution of P2 onto cortical microtubules as P2 networks. Simultaneously, virions are dispatched from the factories and associate with the P2 networks to form so-called mixed networks, composed of transmission units and free P2. Transmission units and free P2 are readily ingested by the aphids and attach to the stylets. Aphids may acquire more P2 and transmission units during subsequent punctures, and, after taking-off and alighting on another plant, inoculate them to start a new infection. The sequential acquisition and the vector-induced transmission mode, coined transmission activation, have important consequences on transmission biology (discussed in ^{13,14}). Remarkably, transmission activation can also be induced by abiotic stresses such as incubation with CO₂ or azide that cause formation of P2 networks ¹².

Formation of the P2-virion transmission unit is mandatory for aphid transmission of CaMV, whereas interaction with microtubules is only required in the cellular context, because firstly, aphids transmit CaMV after feeding on purified P2-virion preparations, devoid of microtubules ¹⁵ and secondly, inhibiting formation of P2 networks in infected cells by drugs coincides with inhibition of transmission ¹².

Little is known about the structure and functional domains of P2. Structure predictions and limited experimental data suggest that the 159 amino acids 18 kD P2 protein contains an N-terminal knob domain, composed of short β -sheets and α -helices (Fig. 1). The knob seems to be separated by a disordered stretch from the rod-like C-terminus, which contains two experimentally confirmed α -helices ¹⁵. The α -helices of three P2 molecules form parallel coiled-coils and promote auto-assembly of P2 into a trimer. The trimers can assemble to higher-order structures, so-called paracrystals, indicating that there is a second, yet unknown self-interaction domain in the P2 molecule ¹⁵. The α -helices are also involved in formation of the transmissible complex because they interact with the virion ⁵. For this, the P2 C-terminal α -helices bind to the N-terminal α -helices of P3, which is besides a component of TBs also a virus-associated protein ¹⁶ forming a mesh on the capsid surface, composed of viral capsid protein P4 ^{17,18}. The N-terminus of P2 is required for interaction with the vector, because substitutions of amino acid 6 of P2 modify or abolish transmission but do not compromise formation of the transmissible complex ¹⁹. P2 was reported to interact with microtubules in plants, in insect cells and *in vitro* ^{20,21}. The P2-microtubule interaction is necessary for formation of transmission bodies ²¹, and for efficient CaMV acquisition by aphids

85 ^{11,12}. However, the microtubule-binding domain of P2 is unknown. Likewise, besides the P3-
interaction domain, no other P2 regions interacting with cellular or viral factors are identified.

We here set out to better characterize the functional domains of P2. We were in particular
interested to identify the hitherto unknown microtubule and vector binding regions of P2. For this,
P2 mutants were created by alanine-scanning and tested for their capacity to form mixed networks
90 and to transmit the virus.

Material and methods

Aphids and plants

A non-viruliferous clonal *Myzus persicae* (Sulz) population was reared under controlled conditions
(22/18 °C day/night with a photoperiod of 14/10 h day/night) on eggplant. Turnip plants (*Brassica*
95 *rapa* cv. 'Just Right') were grown at 24/19 °C day/night with a photoperiod of 16/8 h day/night.

Cloning

For transient expression in plant protoplasts, the alanine substitutions Ala1 to Ala12 were inserted
with the QuikChange mutagenesis kit (Agilent Technologies) into the pCK-P2 plasmid, which
permits expression of wild type P2 under control of 35S promoter ¹¹.

100 For baculovirus expression in *Sf9* cells, the P2 sequence with a his-tag added to the N-terminus
[HP2, ¹⁵] was PCR-amplified using pCK-P2 as matrix. The primers used were Apa-Start-his (5'-
TCCCGGGCCCATGCACCATCACCATCACCATGGAGGTTCTAGCATTACGGGTCAACCGCA
TG-3') coding for an *ApaI* site, a his6-tag, a GGS linker and the N-terminus of P2 and P2-Sac2 (5'-
105 CATCCGCGGTTAGCCAATAATATTCTTTAATCC-3') coding the C-terminus of P2 and a *SacII*
site. The PCR product was digested with *ApaI* and *SacII* and inserted between the *ApaI* and *SacII*
restriction sites of the baculovirus transfer vector pOETIC_6xHis (Oxford Expression
Technologies). This yielded a plasmid coding for HP2 under control of the polh promoter. The
alanine substitutions were then introduced with the QuikChange mutagenesis kit.

For expression of P2-Ala in the viral context, the P2 sequence in the infectious CaMV-coding
110 plasmid pGreen-35S-BJI ²² was replaced by an *ApaI* site. The different P2-Ala sequences were
PCR-amplified from pOETIC-P2-Ala plasmids using primers with additional *ApaI* sites and
inserted into pGreen-35S-BJI.

All inserts were verified by sequencing.

115 *Baculoviruses and protein expression*

Recombinant baculoviruses coding P2-Ala and P2-Ala-GFP mutants were obtained by
recombination in *Sf9* cells using the flashBAC system (Oxford Expression Technologies) and
pOETIC_6xHis plasmids. Baculovirus stocks were produced as indicated by the manufacturer. For
protein production, *Sf9* cells cultivated at 28 °C in TNM-FH insect medium (Sigma) enriched with 5
120 % fetal calf serum, were infected at near confluence with the virus stocks and cultivated for another
48-72 h. Cells were then harvested by centrifugation, resuspended in SES buffer ²³, ultrasonicated to
extract the proteins, and stored at -20 °C or -80 °C until use.

Plant inoculation

125 Two weeks old turnip plants were rub-inoculated with 15-30 µg of pGreen plasmid coding for wild
type CaMV or P2-Ala1-6 mutants. Plants were observed for symptoms every 2-3 days up to 50 days
after inoculation (dpi). Subsequent inoculations were made by rub-inoculation with crude extracts
prepared from infected plants as described ¹¹. The mutant viruses were thus serially passaged 2 to 6
times. The P2 sequence in infected plants was verified by PCR sequencing and plants containing
130 viral sequences with additional mutations were not used for experiments.

Aphid transmission tests

135 For transmission experiments after membrane acquisition, aphids starved for 1 h were allowed to
acquire CaMV for 15 min across stretched Parafilm membranes from suspensions containing
purified CaMV particles²⁴, and extracts from P3- and P2-Ala mutant expressing *Sf9* cells in 15 %
sucrose/SES buffer as described⁷. Then at least 10 aphids per plant were placed onto 20 one-week
140 old turnip seedlings for at least 1 h inoculation period. Aphids were killed with insecticide as
described²⁵. After 3 weeks, the percentage of symptomatic plants was scored visually. For plant to
plant transmission assays, 2-3 detached medium sized infected leaves with well-established,
comparable symptoms were used. Starved aphids were placed on the leaves for 1 h acquisition time.
Then at least 10 aphids were placed per plant on turnip seedlings for 1 h inoculation. Aphids were
killed as described above and test plants observed for symptoms visually for up to 50 dpi. Plant
leaves and *Sf9* extracts were checked for P2 content by Western blot and material devoid of P2 was
excluded from analysis.

145

SDS-PAGE and Western blot

Sf9 extracts or total plant extracts, obtained by grinding frozen leaf discs in 50 mM HEPES pH 8.0,
were mixed with 4x Laemmli buffer, heated for 5 min at 105 °C and separated by discontinuous
6/13.5 % SDS-PAGE²⁶. Gels were stained with Coomassie Blue or proteins were transferred onto
150 nitrocellulose membranes with a semidry blotting apparatus. Membranes were stained with Ponceau
Red S and antigens revealed by Western blot using the NBT-BCIP color reaction for revelation as
described¹².

Immunofluorescence microscopy

155 Protoplasts were fixed with 1 % glutaraldehyde, immobilized on polylysine-coated slides,
permeabilized for 1 h with 0.1-0.3 % Triton-X100, bleached for 15 min with 2 mg/ml NaBH₄ and
blocked in BB buffer (3 % BSA, 0.1 % Triton-X100, 0.05 % NaN₃ in PBS). Cells were then
incubated with a 1:200 dilution of anti-rabbit P2, P4 or anti-rat P6, and with a 1:100 dilution of anti-
mouse tubulin antibody in BB for 1 h at 37 °C or overnight at 4 °C. After rinsing with PBS, the
160 slides were incubated with a 1:200 dilution of Alexa Fluor 488 or 594-conjugated goat anti-rabbit or
goat anti-rat and a 1:75 dilution of Alexa 488 or 594-conjugated goat anti-mouse secondary
antibodies (Thermo Fisher Scientific) in BB for 1 h at 37 °C or overnight at 4 °C. Slides were
mounted for observation in Fluoromount medium containing DAPI.

Observations were made with a Zeiss LSM700 confocal microscope operated in sequential or
165 single channel mode. Alexa Fluor 594 dye was excited with a 555 nm laser, and the mirror was set
to record emission from 555-620 nm. Alexa Fluor 488 fluorophores were excited with a 488 nm
laser, and emission was collected from 490-530 nm. For collecting fluorescence of DAPI-colored
nuclei, a 405 nm laser was used for excitation and emission was captured from 405-530 nm. Images
were captured at 0.29-0.34 μ m intervals using a 63x oil-immersion objective. Raw images were
170 processed using proprietary Zen (Zeiss) and open source Image J software (NIH) to adjust
brightness and contrast.

Antibodies

The following primary antibodies were used: rabbit anti-P2²⁷, rabbit anti-P3⁷ rabbit anti-P6⁹, rat
175 anti-P6 (directed against a peptide and prepared by Eurogentec), monoclonal mouse anti- α -tubulin
DM1A²⁸, and rabbit anti-P4 (Loewe). Alexa 488 and 594 (Thermo Fisher Scientific) and alkaline
phosphatase (Santa Cruz) conjugates of secondary antibodies were used for immunofluorescence
and Western blot analyses, respectively.

180 **Protoplasts**

Protoplasts were prepared from infected turnip leaves by enzymatic digestion as described¹². For stress treatment, protoplasts were exposed to 0.02 % azide for 40 min, or to gaseous CO₂ for 15 min as described¹². For transfection, protoplasts were prepared by enzymatic digestion and transfected with 30 µg plasmid using the PEG method as described²⁹. Protoplasts were cultivated in Gamborg medium at 25 °C.

PCR and sequencing of CaMV and baculoviruses

For extraction of CaMV DNA, leaves were frozen and ground in 50 mM HEPES pH 8.0, centrifuged for 10 min at 14,000 g, and 10-100x diluted supernatants used for PCR with primers pGREEN-P2-F (5'-AAT-AGG-AAA-CGG-TGC-TTC-ATC-CTC-T-3') and pGREEN-P2-R (5'-TTG-TTA-CAG-GGG-CAA-TCA-TTG-ATG-A-3') flanking P2. For extraction of baculovirus DNA, 100 µl of virus stocks were lysed with 1 volume 10 mM Tris-HCl pH 7.6, 10 mM EDTA, 0.25 % SDS, extracted with 1 volume chloroform and the aqueous phase precipitated with sodium acetate and ethanol. DNA pellets was resuspended in 30 µl of water and aliquots used for PCR with primers pOET-F (5'-GGA-GAT-AAT-TAA-AAT-GAT-AAC-C-3') and pOET-R (5'-CAA-CAA-CGC-ACA-GAA-TCT-AGC-GC-3') flanking P2. PCR products were sequenced using the same primers.

Statistics

200 Fisher's exact test with 2 nominal variables (infection status and CaMV mutant) was used to compare transmission rates. Tests were run using the R implementation on <https://biostatv.sentiweb.fr/>?module=tests/fisher>.

Results

We wanted to identify functional domains in the P2 sequence. For this, we used the alanine-scanning technique and replaced three wild-type (wt) amino acids by three alanine residues approximately every ten amino acids all along the P2 sequence (Fig. 1). As P2 mutations are often lethal in the viral context, thus precluding analysis^{20,30}, the alanine mutations were first introduced in pCK-P2, a plasmid for transient expression of P2. The plasmids were used to transfect protoplasts purified from turnip plants infected with CaMV B-JI ΔP2²¹. This recombinant virus is deleted of the entire P2 open reading frame and does not express P2, but is otherwise fully infectious. CaMV B-JI ΔP2 infected protoplasts form *de novo* regular TB when transfected with pCK-P2¹¹ and can therefore be used to study P2 functions. We used this system and immunofluorescence to determine whether the P2 alanine mutants could still form regular TBs and, as a functional test, associate with microtubules after CO₂ or azide stress, as does wt P2¹². Figure 2A,B shows that wt P2 formed one or several normal TBs in unstressed cells and decorated microtubules in stressed cells as reported^{11,12}. The alanine mutants showed different P2 distributions (Fig. 2C-H) that are summarized in Table 1: P2-Ala1 to Ala6 accumulated in diffuse or dense perinuclear inclusions, and stress treatment induced association, to varying degrees, of the mutant proteins with microtubules: P2-Ala6 decorated microtubules, P2-Ala1 was found close to microtubules, Ala2 and Ala5 were found close to microtubules or decorating them, and P2-Ala3 and Ala4 did not colocalize with microtubules. Alanine substitutions in the second half of the P2 sequence (P2-Ala7 to Ala12) did not induce an unusual phenotype: all P2 mutants formed normal-looking TB and stress treatment induced relocalization of mutant P2 from inclusions to microtubule networks.

225 Because the P2 mutants P2-Ala1 to Ala6 showed, contrary to P2-Ala7 to Ala12, an aberrant P2 location and, except P2-Ala6, aberrant MT interaction, we examined them more closely to better define P2 regions necessary for TB formation and microtubule interaction. For this, we introduced

230 P2-Ala1 to Ala6 in pGreen-35S-B-JI, an infectious CaMV-encoding plasmid, and rub-inoculated
turnip plants with the plasmids. Mosaic and leaf-curling symptoms, typical of CaMV infection,
appeared in plants inoculated with wt and all mutant viruses except those inoculated with CaMV-
P2-Ala4, which was not infectious, and was excluded from this part of the analysis (Table 2A).
Infection rates using plasmids were comparable for wt CaMV and P2-Ala1 mutant virus, and lower
for all other mutant viruses (Table 2A). Plants displayed symptoms 22 to 32 days after plasmid
235 inoculation with all viruses except CaMV-P2-Ala2, where symptoms appeared about 20 days later
(Table 2A). On subsequent serial passages using extracts from infected leaves as inoculum, the time
span from inoculation to appearance of symptoms (latency) shortened considerably for all CaMV
mutants. Latency periods as for wild type virus were observed after one serial passage for CaMV-
P2-Ala1 and Ala2, whereas CaMV-P2-Ala3, Ala5 and Ala6 took more than one passage to reach
latency periods comparable to wild type CaMV (Table 2B).

240 PCR analysis of the genome region containing P2 indicated that the P2 sequence was gradually
deleted from the genome of CaMV-P2-Ala2, Ala3, and Ala6, whereas the P2 sequence of the P2-
Ala1 and P2-Ala5 mutants was stably maintained (Table 2A and Fig. 3A). Partial and complete
deletions were observed within the same infected plants and accumulated late in infection, starting
58 to 67 days after inoculation with plasmids, which corresponds to 12 (Ala2), 33 (Ala3), and 36
245 (Ala6) days after appearance of symptoms. Deletions were also observed after serial passages using
leaf extracts that had been screened by PCR to contain the full length P2 sequence before being
used for inoculation. In this case, deletions appeared 23 to 41 days after inoculation, i.e. 11 to 19
days after appearance of symptoms (Table 2B). Several PCR products of different sizes were often
detected in one plant, suggesting that genomes with different deletions in P2 co-existed for a certain
250 time. There was a tendency that deletions appeared faster during later passages. Figure 1S shows for
CaMV-P2-Ala6 how P2 deletions appeared during infection and after mechanical serial passages.
Sequencing of PCR products indicated partial and complete deletions of the P2 sequence from the
genome. Partial deletions introduced often a premature stop codon as was the case for the P2-Ala2
(P2-Ala2-1) and P2-Ala3 (P2-Ala3-1 and -3) mutants shown in Fig. 3B. Interestingly, one of the
255 two *ApaI* sites flanking P2 (used for insertion of P2 mutants into the viral genome) was often
conserved (Fig. 3B). Besides deletions, also rearrangements or short insertions were observed
occasionally.

We tested by Western blot accumulation of P2 and of key viral proteins, as a proxy to estimate
the effect of the different P2-Ala mutations on P2 accumulation and on virus infection. For this we
260 chose the capsid- and TB-associated protein P3, capsid protein P4 and the VF matrix protein P6.
Concerning P2, only P2-Ala3 accumulated to wt P2 levels. P2 protein was not detected in plants
inoculated with CaMV-P2-Ala1, P2 accumulation was very low in CaMV-P2-Ala5 and low in
CaMV-P2-Ala2- and P2-Ala6-infected plants (Fig. 4). P3, P4 and P6 accumulated similarly in
mutant and wt-infected plants, indicating that – once symptoms appeared – the mutant P2 proteins
265 had no major effect on accumulation of these proteins and thus on infection. However, as indicated
above, mechanically passaging the different viruses using tissue extracts resulted reproducibly in
deletion of the P2-Ala2, Ala3 and Ala6 sequences from the genome, suggesting that the mutations
destabilized the genome (Table 2).

Then we looked at the ability of the mutant P2 proteins to form TBs in the context of an
270 authentic viral infection and, as a functional test, to interact with microtubules after stress treatment.
For this, protoplasts were prepared from infected plants and processed either directly or after
treatment with azide or CO₂ for immunofluorescence against P2, P4, P6 and α -tubulin (TUA) to
evaluate P2 location with regard to VFs, virions and microtubules. Figure 5A shows that wt P2
accumulated in one or several TBs that were clearly distinct from the often perinuclear VFs. Stress
275 treatment induced dissolution of TB and relocalization of wt P2 in the form of networks, whereas
VFs remained unchanged. The wt P2 networks colocalized with microtubules (Fig. 5B). Also a
portion of the virions formed, as judged by P4 labeling, networks under stress condition (Fig. 5C),

as reported ^{11,12}. No TBs or any other P2-containing structures were detected in cells infected with CaMV-P2-Ala1, in line with the results from Western blots, where no P2 protein was detected (Fig. 5A,B), and stress treatment did not induce P4 networks (Fig. 5C). P2-Ala2 accumulated in often perinuclear inclusions. The P2 inclusions labeled positive for P6, suggesting that P2 was at least partially retained in VFs in these cells (Fig. 5A). Occasionally, the inclusions stained also positive for TUA (Fig. 4B) as did wt TBs (¹² and data not shown), indicating that P2-Ala2 could interact with tubulin. Indeed, stress treatment induced P2-Ala2 networks on microtubules and also P4 networks (Fig. 5B,C). We observed also another phenotype in CaMV-P2-Ala2 infected cells, which appeared after several mechanical passages of the virus. In these plants, P2-Ala2 accumulated diffuse in the cytoplasm, but P2 interacted still with microtubules after stress treatment (Fig. 2S). Sequencing revealed that the P2 sequence contained an additional point mutation changing amino acid 10 from a tyrosine to a phenylalanine. P2-Ala3 accumulated in infected cells in one or a few inclusions of various size, some of which were apposed to the nucleus. P2/P6 double-labeling showed that some of the P2-Ala3 inclusions stained positive for P6, indicating that some P2-Ala3 was retained in VFs (Fig. 5A). The P2-Ala3 inclusions contained sometimes also TUA (Fig. 5B). Stress treatment did not induce formation of P2-Ala3 networks on microtubules; the P2-Ala3 inclusions seemed unchanged by the treatment (Fig. 5B), and also virions did not form networks after stress treatment (Fig. 5C). CaMV-P2-Ala5-infected cells displayed P2-Ala5 in often perinuclear inclusions that stained positive for P6 and were thus VFs (Fig. 5A). Stress treatment induced partial P2-Ala5 relocalization on microtubules (Fig. 5B), suggesting that TB formation, but not microtubule interaction was impaired by the mutation. Also a part of the virions localized on networks after stress treatment (Fig. 5C). CaMV-P2-Ala6 formed one or several P2-containing inclusions in infected plant cells. Many of them labeled also positive for P6, suggesting that most of P2-Ala6 was retained in VFs (Fig. 5A). Stress treatment induced P2-Ala6 and virion networks (Fig. 5B,C), indicative of microtubule interaction of P2-Ala6 and virions.

Next we tested aphid transmission of the mutant viruses in plant-to-plant transmission assays. Only CaMV-P2-Ala2, Ala5 and Ala6 were transmissible (Table 3). Transmission rates of the three mutants were significantly lower compared to wt CaMV (93±8 % for wt CaMV, 2±3 % for CaMV-P2-Ala2, 1±3 % for CaMV-P2-Ala5, 21±17 % for CaMV-P2-Ala6; $p < 0.001$, Fisher's exact test). Transmission rates of CaMV-P2-Ala6 dropped at later stages of the infection, whereas they remained stable for wt CaMV (Fig. 6). This seemed to correlate with the appearance of deletions of partial or total P2 sequence and concomitant lower P2 protein levels in infected plants (Fig. 2S). CaMV genomes with full length P2 sequence and incomplete P2 sequences coexisted often for some time in infected plants, and transmission was only recorded when at least some full length P2 was present. Whether the same was true for CaMV-P2-Ala2 and Ala5, was difficult to determine because of the low transmission rates. To test whether non-transmissibility of P2-Ala3 was due to the context of the viral infection in the plant or whether it was an intrinsic property of the mutant protein, aphid transmission assays after membrane acquisition of recombinant P2, P3 and purified virions were performed. At the same time we tested also transmission after membrane acquisition of P2-Ala1 that did not accumulate in infected plants, and of P2-Ala4 that was not infectious in the viral context, and where in both cases plant-to-plant transmission experiments were not possible. Table 4 shows that none of the mutant P2 proteins supported virus transmission.

320 Discussion

In this work we started a functional analysis of the CaMV aphid transmission protein P2. This protein has many exciting features but it is still hardly characterized functionally or structurally. This is mainly due to its intrinsic properties: its insolubility precludes structural characterization by crystallization, its toxicity impedes production in all current recombinant protein expression systems except the *Sf9*/baculovirus system, and many P2 mutants are either unstable or lethal in the viral context, again hindering its characterization. To circumvent this problem, we here implemented successfully a protoplast system using cells infected with a P2 deletion mutant of

CaMV and provided P2 *in trans* by transfection with P2 expression plasmids. These cells display all features of a genuine CaMV infection but do not express P2 and do not contain TBs. TBs will only
330 form after P2 is expressed from the plasmid. This approach avoided interference of P2 mutants with establishment of viral infection, as has been reported for other CaMV P2 mutants that are either non-infectious or revert to wild type^{30,31}. Using this approach, we tested twelve P2 alanine scanning mutants distributed along the P2 sequence for their capacity to form TBs and to interact with microtubules after stress treatment. We found that P2-Ala7 to Ala12, all mutated in the second half
335 of the protein, formed normal TBs. Stress induced P2 networks on microtubules for all these six mutants. This indicates that microtubule binding activity, required for P2 export from VFs towards newly forming TBs²¹ and for relocalization of P2 from TBs to microtubules during vector acquisition¹², is not mediated by the C-terminal half of P2.

For the P2 alanine substitutions localized in the N-terminus (P2-Ala1 to Ala6), we found variable microtubule interactions. P2-Ala3 and Ala4 did not colocalize with microtubules. P2-Ala1 was detected close to but not on microtubules, Ala2 and Ala5 decorated microtubules or were detected close to microtubules after stress treatment. Finally, P2-Ala6 decorated microtubules like wt P2. These results indicate that the entire P2 N-terminus is involved in microtubule binding, with the region comprising amino acids 35 to 50 contributing most to it, since alanine substitutions in this
345 region (P2-Ala3 and Ala4) abolished microtubule interaction completely. Interestingly, P2-Ala3 inclusions in infected plants contained some apparently soluble tubulin (Fig. 5B) as do wt TBs under certain conditions¹². This suggests that P2 interaction with microtubules is somewhat different from that with soluble tubulin. Additional analysis is necessary to define more precisely P2's microtubule- and tubulin-binding domain(s).

Basically similar results concerning P2 distribution were obtained when the P2 alanine mutants were introduced in the viral genome. A notable difference was that P2-Ala1 protein could not be detected in infected cells. Apparently, it was unstable in the viral context, as has been reported before for other P2 mutants^{9,32}. Further, the diffuse cytoplasmic distribution of P2-Ala2 and Ala3 was less often observed in infected plants than in transfected protoplasts. Rather, P2-Ala2 and Ala3
355 accumulated, like P2-Ala5 and Ala6, in VFs. The differences might be due to different translation sites of P2: P2 translated from the viral 35S RNA is produced, like all other viral proteins, in or close to VFs and in close contact with P6 and other viral proteins³³, whereas P2 from plasmid-derived mRNA is probably translated free in the cytoplasm. Whatever the reason for the discrepancies, the presence of the mutant P2s in VFs indicated that they were not exported correctly
360 from VFs to emerging TBs²¹. One explanation is that missing or aberrant P2-microtubule interaction prevented P2 export and TB formation. Support for this interpretation comes from the observation that P2 was also retained in VFs after incubation of CaMV-transfected protoplasts with the microtubule depolymerizing drug oryzalin²¹. Alternatively or additionally, a putative P2-P6 interaction region for which there is genetic and biochemical evidence^{34,35}, might localize to the N-terminus of P2 and be perturbed in the mutants. There were also slight differences in microtubule
365 interaction between transfected protoplasts and protoplasts derived from infected plants: P2 found close to microtubules but not decorating them (P2-Ala2 and Ala5) was less often observed in infected than in transfected cells. We assume again that the less strictly controlled expression of P2 in the transfected cells might account for the differences.

In aphid transmission assays using plants inoculated with the mutant viruses as virus source, CaMV-P2-Ala2 and Ala5 were barely and CaMV-P2-Ala6 moderately transmissible, compared to wt CaMV. The lower transmission rates might be explained partially by lower accumulation of the mutant proteins in infected tissue. However, the transmission rates correlated also nicely with microtubule interaction: P2 from barely transmissible CaMV-P2-Ala2 and Ala5 associated or
375 decorated with microtubules weakly, whereas P2 from highly transmissible wt CaMV and moderately transmissible CaMV-P2-Ala6 decorated microtubules strongly. P2 from CaMV-P2-Ala3 did not interact at all with microtubules and was not transmissible, although it accumulated to

similar levels like wt P2. CaMV-P2-Ala1 and Ala4 could not be tested in plant-to-plant transmission because the mutant virus did not accumulate P2 (CaMV-P2-Ala1) or was not infectious (CaMV-P2-Ala4). Therefore these two P2 mutants and the plant-to-plant non transmissible P2-Ala3 were produced as recombinant proteins and tested for transmission in membrane feeding assays where aphids were allowed acquisition assay periods on suspensions containing mutant P2 and virions. Again, no transmission was observed. Because these P2 mutants had in common that they did not or only weakly interact with microtubules in transfected protoplasts, the results indicate that the microtubule interaction domain might overlap with the yet undefined stylet-binding domain, which is localized in P2's N-terminus ¹⁹.

An interesting observation was that the sequences of P2-Ala2, Ala3 and Ala6 were unstable in the genome and were partially or completely deleted from the viral genome whereas wt P2, P2-Ala1 and Ala5 sequences were stable. Partial deletion of the P2 sequence often introduced premature stop codons. Besides deletions also rearrangements were observed. Taken together, this indicates that expression of P2-Ala2, Ala3 and Ala6 was noxious to the virus. Wild type P2 is regarded as a toxic protein whose expression is tightly regulated ^{30,31,36}. According to this hypothesis, the alanine substitutions in P2-Ala2, Ala3 and Ala6 increased P2 toxicity and their elimination by deletion restored viral fitness. Expression of P2-Ala4 might have been too toxic to allow CaMV replication, explaining why it was not infectious. Finally, P2-Ala5 had similar properties like wt P2, and P2-Ala1 could not exhibit its possible toxicity because it did not accumulate in infected cells. Therefore, the latter two mutants were maintained in the genome.

It remains enigmatic why P2 is maintained in the CaMV genome at all because this protein has, except transmission, no other function during infection of the plant and because P2 deletion mutants are as infectious as wild type CaMV. One should expect that selection pressure eliminates P2 from the genome just as GFP and other transgenic insertions into viral genomes are often unstable ^{37,38}. One explanation is that the replication initiation site and the splice acceptor site in the P2 sequence boost replication and counterbalance costs of P2 expression. Since the observed partial and complete deletions removed also the genetic elements in the P2 sequence, they do not seem to be required for CaMV infection when there is no P2 gene or accumulation.

Complete deletions conserved one of the two *ApaI* sites used for insertion of P2 sequences. We propose the repetitive *ApaI* sequences were used as a scaffold to excise the P2 sequence in several steps, probably during reverse transcription or synthesis of the +strand DNA. A similar observation was made for the C4-184 mutant where the deletion site eliminating a large portion of the P2 sequence is surrounded by a stretch of highly similar sequences ³⁹.

Taken together, we present here evidence for a modular organization of P2 that will be fascinating to explore further. Its C-terminus is dedicated to viral interactions, i.e. P2 self-interaction ¹⁵ and binding to P3 and virions ⁵. Its N-terminus controls interactions with the environment, i.e. vector-binding and microtubule interaction. Interestingly, functions housed by the modules overlap; domains for P2 self-interaction and P2-P3 interaction are contained in the same region of the C-terminus, and those for vector-binding and microtubule interaction map to the same stretch in the N-terminus of P2.

Acknowledgments

We thank Sophie Le Blaye for plant care. All authors declare that there is no conflict of interest.

420 **Funding**

Our work was financed by INRA SPE department (to M.D.), ANR grant 12-BSV7-005-01 (to M.D.) and Human Frontier Science Program grant RGP0013/2015 (to M.D.). B.D. was supported by the AgreenSkills fellowship program which received funding from the EU's Seventh Framework Program under grant agreement N° FP7-609398 (AgreenSkills+ contract). The funding sources had
425 no role in the study design, in the collection, analysis and interpretation of data; in the writing of the report, and in the decision to submit the article for publication.

Author contributions

Christiane Then: conceptualization, validation, investigation, writing – original draft, writing – review & editing, visualization, supervision. Aurélie Bak: conceptualization, investigation.
430 Alexandre Morisset: validation, investigation. Beatriz Dáder: investigation, validation, writing – review & editing. Marie Ducouso: validation, investigation. Jean-Luc Macia: validation, investigation. Martin Drucker: conceptualization, writing – original draft, writing – review & editing, visualization, formal analysis, supervision, project administration, funding acquisition.

References

1. Blanc, S., Drucker, M., and Uzest, M. (2014). Localizing viruses in their insect vectors. *Annu. Rev. Phytopathol.* 52, 403–425. 10.1146/annurev-phyto-102313-045920.
2. Dáder, B., Then, C., Berthelot, E., Ducouso, M., Ng, J.C.K., and Drucker, M. (2017). Insect transmission of plant viruses: multilayered interactions optimize viral propagation. *Insect Sci.* 10.1111/1744-7917.12470.
3. Uzest, M., Gargani, D., Drucker, M., Hébrard, E., Garzo, E., Candresse, T., Fereres, A., and Blanc, S. (2007). A protein key to plant virus transmission at the tip of the insect vector stylet. *Proc. Natl. Acad. Sci. U. S. A.* 104, 17959–17964. 10.1073/pnas.0706608104.
4. Webster, C.G., Pichon, E., van Munster, M., Monsion, B., Deshoux, M., Gargani, D., Calevro, F., Jimenez, J., Moreno, A., Krenz, B., et al. (2018). Identification of Plant Virus Receptor Candidates in the Stylets of Their Aphid Vectors. *J. Virol.* 92. 10.1128/JVI.00432-18.
5. Leh, V., Jacquot, E., Geldreich, A., Hermann, T., Leclerc, D., Cerutti, M., Yot, P., Keller, M., and Blanc, S. (1999). Aphid transmission of cauliflower mosaic virus requires the viral PIII protein. *EMBO J.* 18, 7077–7085. 10.1093/emboj/18.24.7077.
6. Woolston, C.J., Covey, S.N., Penswick, J.R., and Davies, J.W. (1983). Aphid transmission and a polypeptide are specified by a defined region of the cauliflower mosaic virus genome. *Gene* 23, 15–23. 10.1016/0378-1119(83)90212-3.
7. Drucker, M., Froissart, R., Hébrard, E., Uzest, M., Ravallec, M., Espérandieu, P., Mani, J.-C., Pugnière, M., Roquet, F., Fereres, A., et al. (2002). Intracellular distribution of viral gene products regulates a complex mechanism of cauliflower mosaic virus acquisition by its aphid vector. *Proc. Natl. Acad. Sci. U. S. A.* 99, 2422–2427. 10.1073/pnas.042587799.
8. Espinoza, A.M., Medina, V., Hull, R., and Markham, P.G. (1991). Cauliflower mosaic virus gene II product forms distinct inclusion bodies in infected plant cells. *Virology* 185, 337–344. 10.1016/0042-6822(91)90781-6.
9. Khelifa, M., Journou, S., Krishnan, K., Gargani, D., Espérandieu, P., Blanc, S., and Drucker, M. (2007). Electron-lucent inclusion bodies are structures specialized for aphid transmission of cauliflower mosaic virus. *J. Gen. Virol.* 88, 2872–2880. 10.1099/vir.0.83009-0.
10. Palacios, I., Drucker, M., Blanc, S., Leite, S., Moreno, A., and Fereres, A. (2002). Cauliflower mosaic virus is preferentially acquired from the phloem by its aphid vectors. *J. Gen. Virol.* 83, 3163–3171.
11. Bak, A., Gargani, D., Macia, J.-L., Malouvet, E., Vernerey, M.-S., Blanc, S., and Drucker, M. (2013). Virus factories of cauliflower mosaic virus are virion reservoirs that engage actively in vector transmission. *J. Virol.* 87, 12207–12215. 10.1128/JVI.01883-13.
12. Martinière, A., Bak, A., Macia, J.-L., Lautredou, N., Gargani, D., Doumayrou, J., Garzo, E., Moreno, A., Fereres, A., Blanc, S., et al. (2013). A virus responds instantly to the presence of the vector on the host and forms transmission morphs. *eLife* 2, e00183. 10.7554/eLife.00183.
13. Drucker, M., and Then, C. (2015). Transmission activation in non-circulative virus transmission: a general concept? *Curr. Opin. Virol.* 15, 63–68. 10.1016/j.coviro.2015.08.006.

14. Froissart, R., Michalakakis, Y., and Blanc, S. (2002). Helper component-transcomplementation in the vector transmission of plant viruses. *Phytopathology* 92, 576–579.
15. Hébrard, E., Drucker, M., Leclerc, D., Hohn, T., Uzest, M., Froissart, R., Strub, J.M., Sanglier, S., van Dorselaer, A., Padilla, A., et al. (2001). Biochemical characterization of the helper component of Cauliflower mosaic virus. *J. Virol.* 75, 8538–8546. 10.1128/JVI.75.18.8538-8546.2001.
16. Leclerc, D., Burri, L., Kajava, A.V., Mougeot, J.L., Hess, D., Lustig, A., Kleemann, G., and Hohn, T. (1998). The open reading frame III product of cauliflower mosaic virus forms a tetramer through a N-terminal coiled-coil. *J. Biol. Chem.* 273, 29015–29021.
17. Hoh, F., Uzest, M., Drucker, M., Plisson-Chastang, C., Bron, P., Blanc, S., and Dumas, C. (2010). Structural insights into the molecular mechanisms of cauliflower mosaic virus transmission by its insect vector. *J Virol* 84, 4706–4713. doi:10.1128/JVI.02662-09.
18. Plisson, C., Uzest, M., Drucker, M., Froissart, R., Dumas, C., Conway, J., Thomas, D., Blanc, S., and Bron, P. (2005). Structure of the mature P3-virus particle complex of Cauliflower mosaic virus revealed by cryo-electron microscopy. *J Mol Biol* 346, 267–277. 10.1016/j.jmb.2004.11.052.
19. Moreno, A., Hébrard, E., Uzest, M., Blanc, S., and Fereres, A. (2005). A single amino acid position in the helper component of cauliflower mosaic virus can change the spectrum of transmitting vector species. *J. Virol.* 79, 13587–13593. 10.1128/JVI.79.21.13587-13593.2005.
20. Blanc, S., Schmidt, I., Vantard, M., Scholthof, H.B., Kuhl, G., Esperandieu, P., Cerutti, M., and Louis, C. (1996). The aphid transmission factor of cauliflower mosaic virus forms a stable complex with microtubules in both insect and plant cells. *Proc. Natl. Acad. Sci. U. S. A.* 93, 15158–15163.
21. Martinière, A., Gargani, D., Uzest, M., Lautredou, N., Blanc, S., and Drucker, M. (2009). A role for plant microtubules in the formation of transmission-specific inclusion bodies of Cauliflower mosaic virus. *Plant J. Cell Mol. Biol.* 58, 135–146. 10.1111/j.1365-313X.2008.03768.x.
22. Khelifa, M., Massé, D., Blanc, S., and Drucker, M. (2010). Evaluation of the minimal replication time of Cauliflower mosaic virus in different hosts. *Virology* 396, 238–245. 10.1016/j.virol.2009.09.032.
23. Blanc, S., Cerutti, M., Usmany, M., Vlak, J.M., and Hull, R. (1993). Biological activity of cauliflower mosaic virus aphid transmission factor expressed in a heterologous system. *Virology* 192, 643–650. 10.1006/viro.1993.1080.
24. Hull, R., Shepherd, R.J., and Harvey, J.D. (1976). Cauliflower Mosaic Virus: an Improved Purification Procedure and Some Properties of the Virus Particles. *J. Gen. Virol.* 31, 93–100. 10.1099/0022-1317-31-1-93.
25. Martinière, A., Macia, J.-L., Bagnolini, G., Jridi, C., Bak, A., Blanc, S., and Drucker, M. (2011). VAPA, an innovative “virus-acquisition phenotyping assay” opens new horizons in research into the vector-transmission of plant viruses. *PLoS One* 6, e23241. doi:10.1371/journal.pone.0023241.
26. Laemmli, U.K. (1970). Cleavage of Structural Proteins during the Assembly of the Head of Bacteriophage T4. *Nature* 227, 680–685. 10.1038/227680a0.

27. Blanc, S., Schmidt, I., Kuhl, G., Esperandieu, P., Lebourier, G., Hull, R., Cerutti, M., and Louis, C. (1993). Paracrystalline structure of cauliflower mosaic virus aphid transmission factor produced both in plants and in a heterologous system and relationship with a solubilized active form. *Virology* 197, 283–292. 10.1006/viro.1993.1589.
28. Blose, S.H., Meltzer, D.I., and Feramisco, J.R. (1984). 10-nm filaments are induced to collapse in living cells microinjected with monoclonal and polyclonal antibodies against tubulin. *J. Cell Biol.* 98, 847–858. 10.1083/jcb.98.3.847.
29. Yoo, S.-D., Cho, Y.-H., and Sheen, J. (2007). Arabidopsis mesophyll protoplasts: a versatile cell system for transient gene expression analysis. *Nat. Protoc.* 2, 1565–1572. 10.1038/nprot.2007.199.
30. Froissart, R., Uzest, M., Ruiz-Ferrer, V., Drucker, M., Hébrard, E., Hohn, T., and Blanc, S. (2004). Splicing of Cauliflower mosaic virus 35S RNA serves to downregulate a toxic gene product. *J. Gen. Virol.* 85, 2719–2726. 10.1099/vir.0.80029-0.
31. Kiss-László, Z., Blanc, S., and Hohn, T. (1995). Splicing of cauliflower mosaic virus 35S RNA is essential for viral infectivity. *EMBO J.* 14, 3552–3562.
32. Nakayashiki, H., Tsuge, S., Kobayashi, K., Okuno, T., and Furusawa, I. (1993). Reasons for the low accumulation level of aphid transmission factor protein in infected leaves with an aphid-non-transmissible cauliflower mosaic virus isolate, CM1841. *J. Gen. Virol.* 74 (Pt 11), 2469–2472.
33. Haas, M., Bureau, M., Geldreich, A., Yot, P., and Keller, M. (2002). Cauliflower mosaic virus: still in the news. *Mol. Plant Pathol.* 3, 419–429. 10.1046/j.1364-3703.2002.00136.x.
34. Geldreich, A., Haas, G., Kubina, J., Bouton, C., Tanguy, M., Erhardt, M., Keller, M., Ryabova, L., and Dimitrova, M. (2017). Formation of large viroplasm and virulence of Cauliflower mosaic virus in turnip plants depend on the N-terminal EKI sequence of viral protein TAV. *PLoS One* 12, e0189062. 10.1371/journal.pone.0189062.
35. Qiu, S.G., Wintermantel, W.M., Sha, Y., and Schoelz, J.E. (1997). Light-dependent systemic infection of solanaceous species by cauliflower mosaic virus can be conditioned by a viral gene encoding an aphid transmission factor. *Virology* 227, 180–188.
36. Bouton, C., Geldreich, A., Ramel, L., Ryabova, L.A., Dimitrova, M., and Keller, M. (2015). Cauliflower mosaic virus Transcriptome Reveals a Complex Alternative Splicing Pattern. *PLoS One* 10, e0132665. 10.1371/journal.pone.0132665.
37. Chen, C.-C., Chen, T.-C., Raja, J.A.J., Chang, C.-A., Chen, L.-W., Lin, S.-S., and Yeh, S.-D. (2007). Effectiveness and stability of heterologous proteins expressed in plants by Turnip mosaic virus vector at five different insertion sites. *Virus Res.* 130, 210–227. 10.1016/j.virusres.2007.06.014.
38. Gadea, G., Bos, S., Krejbich-Trotot, P., Clain, E., Viranaicken, W., El-Kalamouni, C., Mavingui, P., and Desprès, P. (2016). A robust method for the rapid generation of recombinant Zika virus expressing the GFP reporter gene. *Virology* 497, 157–162. 10.1016/j.virol.2016.07.015.
39. Howarth, A.J., Gardner, R.C., Messing, J., and Shepherd, R.J. (1981). Nucleotide sequence of naturally occurring deletion mutants of cauliflower mosaic virus. *Virology* 112, 678–685. 10.1016/0042-6822(81)90313-5.
40. Kelley, L.A., Mezulis, S., Yates, C.M., Wass, M.N., and Sternberg, M.J.E. (2015). The

Phyre2 web portal for protein modeling, prediction and analysis. *Nat. Protoc.* *10*, 845–858.
10.1038/nprot.2015.053.





Figure 1: Structure prediction of P2. (A) Secondary structure prediction discerns between β -strands, α -helices and disordered regions. The confidence of the prediction is indicated for each amino acid. The blue boxes show the position of the alanine substitution. (B) Three-dimensional model of P2. The colors represent the order of the amino acids from the N-terminus (dark blue) to the C-terminus (red). The overall structures suggests an N-terminal knob, separated by an unstructured stretch from a rod-like C-terminus. All predictions were made with the PHYRE2 program⁴⁰.



445 **Figure 2:** Ectopic expression of P2 alanine (Ala) mutants in CaMV- Δ P2-infected protoplasts. CaMV- Δ P2-infected protoplasts were transfected with plasmids coding for wt P2 or the alanine
 450 mutants P2-Ala1 to Ala12. After overnight incubation the protoplasts were processed directly (top panel) or after CO₂ stress (bottom panel) for immunofluorescence against P2 (red) and α -tubulin (TUA, green). Nuclei are counterstained with DAPI (blue). (A) Wild type P2 accumulates in a typical transmission body and associates (B) with microtubules after CO₂ treatment. (C-H) P2 alanine mutants show a wt phenotype (C), accumulate in perinuclear irregular dense (E) or less dense (G) inclusions and are, upon stress treatment still able to decorate microtubules (D,F) or are found only close to, but not on microtubules (H). Shown are confocal maximum projections (top panels) or single channels of single optical sections (small images in the bottom panel) of the
 455 regions outlined in the projections. The arrowheads point to colocalizing P2 and α -tubulin label represented in the insets in panels B, D, F and H. The following P2-Ala mutants are shown: P2-Ala8 (C,D), Ala3 (E), Ala2 (F) and Ala1 (G,H). Scale bar 5 μ m.




Figure 3: Deletion of the P2 sequence from the CaMV genome. (A) PCR detection of deletions. Extracts from infected plants were used in PCR with primers hybridizing at the end of P1 and the beginning of P3 of the CaMV genome. Wild type virus containing the entire P2 sequence yielded a 789 bp amplicon, and a genome deleted of the entire P2 sequence a 312 bp product. P₀ corresponds to plants initially inoculated with plasmid DNA, and P₅ to the last (5th) passage tested. The samples presented between P₀ and P₅ are from intermediate passages. Wild type P2 as well as P2-Ala1 and Ala5 were stable throughout the number of passages tested, whereas P2-Ala2, Ala3 and Ala6 sequences were deleted from the genome. Note that P2-Ala2 and Ala6 genomes contained deletions comprising almost the entire P2 sequence, whereas P2-Ala3 maintained ~150 bases of the P2 open reading frame. (B) Examples of deletions and rearrangements of the region comprising P2. PCR amplicons obtained with the same primers as in (A) were gel-purified and sequenced. The first image shows the genome region including P2, the *Apal* restriction sites used for inserting the P2-Ala mutant sequences and the position of the alanine substitutions. Premature stop codons created by the deletions are shown in the deleted sequences. For comparison, the natural P2 deletion mutant C4-184 is shown at the bottom.






Figure 4: Expression of viral proteins. Total extracts of infected leaves prepared after at least one serial passage of the viruses were analyzed by Western blot using the indicated antisera. Wild type and mutant CaMV-infected leaves presenting similar symptoms were used for analysis. This corresponds for wild type and P2-Ala1 leaves harvested at 19 dpi, for Ala2 at 27 dpi, for Ala3 at 36 dpi, for Ala5 at 36 dpi, and for Ala6 at 40 dpi. P2-Ala1 and Ala2 extracts were analyzed on the same membrane and show therefore the same wt extract. P2-Ala3, Ala5 and Ala6 extracts were analyzed independently on separate membranes, each together with a wt extract. The black vertical lines in the P2-Ala2 and Ala5 blots indicate that wt and mutant extracts were charged on the same gel, but not on neighboring lanes. Poinceau Red stain of the Large RuBisCO subunit (Rub) served as loading control. Experiments were repeated 3 times with similar results.



485 **Figure 5:** Phenotype of P2 alanine mutants in mutant CaMV-infected plant cells. Protoplasts purified from infected plants were processed for immunofluorescence against (A) P2 (red) and P6 (green), (B) P2 (red) and TUA (green) and (C) P4 (red) and P6 (green) either directly (unstressed) or after stress treatment. Nuclei are presented in blue (DAPI stain) in some images to show perinuclear localization of viral inclusions. Shown are confocal maximum projections and in the insets single channels of single sections of the projection. Scale bar 10 μ m.



490 **Figure 6:** Transmission of P2-Ala mutants related to age of infection. Starved aphids were placed for 60 min on infected turnip leaves for virus acquisition and then transferred to healthy turnip seedlings for inoculation. Ten aphids per test plant and 20 test plants were used for each assay. Each point represents one transmission assay realized at the day after appearance of symptoms indicated in the x-axis.

Tables

495 **Table 1:** Phenotype of P2-Ala mutants. Wild type or mutant P2 were transiently expressed in protoplasts infected with CaMV B-JI Δ P2, and the distribution of P2 in untreated cells and in cells treated with CO₂ analyzed by immunofluorescence microscopy. TB, transmission body.

P2 mutant	Untreated cells	Cells treated with CO ₂	
	P2 localization	P2 decorates microtubules	P2 is found close to microtubules
wild type	In normal TB	+	-
Ala1	In perinuclear aggregates	-	+
Ala2	Diffuse in cytoplasm	+	+
Ala3	Diffuse in cytoplasm	-	-
Ala4	In perinuclear aggregates	-	-
Ala5	In perinuclear aggregates	+	+
Ala6	In perinuclear aggregates	+	-
Ala7	In TB	+	-
Ala8	In TB	+	-
Ala9	In TB	+	-
Ala10	In TB	+	-
Ala11	In TB	+	-
Ala12	In TB	+	-

500 **Table 2:** Infectivity and stability of CaMV P2-Ala mutants. Turnip seedlings in the two juvenile
 leaf stage were for initial inoculation rub-inoculated with plasmids encoding the CaMV genome
 harboring wild type or mutant P2 and observed daily for symptoms. For the subsequent serial
 passages, extracts prepared from infected leaves were mechanically inoculated to test plants. The
 505 extracts were verified by PCR for presence of the P2 sequence before inoculation. dpi, days after
 inoculation, das, days after appearance of symptoms, n.a. not applicable, n.t. not tested.

A: Plasmid inoculation

CaMV mutant	Plants infected/inoculated	Infected plants (%)	Appearance of symptoms (dpi)	P2 sequence stable	Start degradation of P2 sequence	
					dpi	das
wild type	9/12	75	22-25	Yes	-	-
P2-Ala1	11/16	69	28-29	Yes	-	-
P2-Ala2	3/32	9	43-46	No	58-62	12-19
P2-Ala3	3/15	20	24-27	No	60-65	33-41
P2-Ala4	0/21	0	n.a.	n.a.	n.a.	n.a.
P2-Ala5	4/20	20	22-28	Yes	-	-
P2-Ala6	2/15	13	26-32	No	67-71	35-45

B: Inoculation with leaf extracts

CaMV mutant	Appearance of symptoms (dpi)		P2 sequence stable	Start degradation of P2 sequence (dpi)			
				1 st passage		5 th passage	
	1 st passage	5 th passage	dpi	das	dpi	das	
wild type	12	n.t.	Yes	-	-	-	-
P2-Ala1	13	11	Yes	-	-	-	-
P2-Ala2	13	12	No	27	14	27	15
P2-Ala3	22	11	No	41	19	23	12
P2-Ala4	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.
P2-Ala5	21	11	Yes	-	-	-	-
P2-Ala6	17	12	No	34	17	23	11

515 **Table 3:** Transmission of CaMV P2-Ala mutants. Starved aphids were allowed to feed for (A) 60 min to acquire CaMV from infected turnip leaves or (B) 15 min to acquire purified CaMV particles, recombinant his-tagged P2 (HP2) and P3 by membrane feeding, and then transferred to turnip seedlings for inoculation. Ten aphids per test plant were used for plant-to-plant transmission assays and for transmission tests after membrane acquisition feeding. n.a., not applicable.

A: Plant-to-plant transmission

CaMV mutant	Infected plants/total plants	n experiments	% Transmission (min-max)
Wild type	373/403	9	93 (75-100)
P2-Ala1	0/220	6	0*
P2-Ala2	6/328	13	2 (0-8)
P2-Ala3	0/256	8	0
P2-Ala4	n.a.	n.a.	n.a.
P2-Ala5	3/496	8	1 (0-10)
P2-Ala6	83/400	10	21 (0-70)

*no P2 protein detected.

520 B: Transmission after membrane acquisition

P2 mutant	Infected plants/total plants	n experiments	% Transmission (min-max)
HP2	74/259	8	29 (0-95)
HP2-Ala1	0/160	5	0
HP2-Ala3	0/260	8	0
HP2-Ala4	0/200	6	0