



# Fundamental constants explain submillimeter liquid shear forces

Laurence Noirez

## ► To cite this version:

Laurence Noirez. Fundamental constants explain submillimeter liquid shear forces. Science Advances , 2023, 9 (34), 10.1126/sciadv.adj8336 . hal-04188976

**HAL Id: hal-04188976**

**<https://hal.science/hal-04188976>**

Submitted on 27 Aug 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Fundamental constants explain submillimeter liquid shear forces

Laurence Noirez

From fundamental physical constants to the identification of liquid shear elasticity, over the past decade, new ways have emerged to understand viscosity.

Copyright © 2023 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. Distributed under a Creative Commons Attribution NonCommercial License 4.0 (CC BY-NC).

An unexpected link between bio-friendly viscosity and universal constants shows that liquids provide a daily reminder of our deep cosmic origins (1). This fundamental link also highlights the cohesive nature of liquids, in agreement with the energy associated with the shear elastic forces recently identified at the mesoscopic scale (2), and therefore the possibility of collective effects. The research by Kostya Trachenko featured in this week's issue of *Science Advances* is an invitation to no longer look at the formless matter called liquids in the same way, as a new approach to understanding fluid flow is emerging (1, 2).

## VISCOSITY AS AN EVOLVING CONCEPT

Viscosity characterizes the ability to flow, and its significance has long been recognized. For example, knowing viscosity of physiological fluids is commonly used as an indicator to understand and treat diseases. But, let us examine how it has been defined and the assumptions that underlie it.

The modern concept of viscosity has neither always been evident nor intuitive. In the mid-19th century, French engineer, Henri Navier, introduced the idea of viscosity as a constant representing the resistance of a layer to a wall (3). Navier also noticed that at small scales, this resistance of a fluid to flow depends on the substrate. He noted, for example, that water flows three or four times more slowly near glass than near copper. On a larger scale, the agreement between experimental results and theory gave credence to the hypothesis that the liquid has zero velocity at the

substrate, leading to the no-slip boundary assumption. Therefore, the concept of macroscopic viscosity evolved, breaking free from experimental conditions. However, on a small scale, it is no longer valid (for example, for micro- and nanofluidics) and reminds us that the transfer mechanism of interfacial stresses is still largely unknown. The link between measured viscosity and the strength of the molecular interactions has not been easily demonstrated. This might explain the difficulty of theoreticians to develop a precise law for the interaction in liquids. The challenge is thus to find a starting point, a common characteristic, that is free from liquid molecular parameters that are highly variable in liquids and make a general theoretical framework difficult to develop.

Trachenko has succeeded in identifying a nearly constant value for viscosity of various liquids at the nexus of liquid and gas states. This viscosity is independent of the molecular interactions and structure (1). Regardless of a liquid's nature, this common characteristic—that the lowest viscosity is obtained by varying the temperature and pressure—is not a coincidence but an indication of the fundamental parameters that define the lower allowed viscosity. This enables Trachenko to define a fundamental kinematic viscosity that depends only on three fundamental physical constants:  $\hbar$ ,  $m_e$ , and  $m_p$ , where  $\hbar$  is Planck's constant,  $m_e$  is the electron mass, and  $m_p$  the proton mass (1, 4). Even separated by about 15 decades, the lower kinematic viscosity points to its fundamental origin in the more stable components of the universe.

Why does the relationship to the fundamental constants seem valid for the lowest

viscosity only? Recent experimental advances might provide an answer by demonstrating that when the boundary conditions are ignored conventional measurements only partially access liquid properties.


## WHAT SMALL-SCALE MEASUREMENTS TELL US

Recent measurements show that at the submillimeter scale, viscosity is not universal but depends on how a liquid interacts with a substrate (2, 5). This result echoes the Navier observation about water's behavior on glass and copper (3). More precisely, conventional viscosity or viscoelasticity measurements do not guarantee the necessary no-slip condition of the liquid at the wall.

When liquid/substrate wetting improves toward non-slip boundary conditions, the mesoscopic fluid response reveals a fundamentally different response (2, 5). The fluid resists before flowing. Liquids are thus endowed with "static" shear elasticity. This is important because the ability of liquids to support solid-like shear stress is an indicator of their cohesion energy. Liquid shear elasticity challenges the very definition of what a liquid should be. Shear elasticity is a collective property. It involves long-range interactions (Fig. 1). Considering the mesoscopic shear elasticity allows a new understanding of liquids and the discovery of new properties on basis of collective excitations mostly applied only in solids until recently.

## THEORY AND EXPERIMENT CONVERGE

The first experimental identification of liquid shear elasticity at the micron scale occurred in the late 1980s (6). The research was extended to the submillimeter scale



**Fig. 1. Liquid properties such as viscosity and shear elasticity echo the history of our distant stellar origin via their dependence on high-energy particles (1, 4).** The image shows thermal traces left in a furrow in water illustrating long-range interactive forces in liquids (2, 5, 8, 9). Credit: P. Baroni, M. Warburton, and L. Noirez.

about 15 years later with the identification of liquid shear elasticity (2).

Concurrent to, but independent of, the work to identify liquid shear elasticity, Trachenko was applying insights from the Maxwell-Frenkel viscoelastic theory to show that nothing prevents liquids from propagating a gapped shear wave (7, 8). Working with A. Zaccane, Trachenko has now developed an atomic-level mathematical theory of the (visco)elasticity of confined liquids clarifying that confinement increases liquid shear elasticity by suppressing long-wavelength shear waves (8). As happens in science, theoretical formalism and experimental measurements converged and established the existence of liquid shear elasticity.

### LIQUIDS NOT SO DIFFERENT FROM SOLIDS

Liquid shear elasticity is accessible at the submillimeter scale and under low perturbative mechanical conditions. Why? Moving away from near-equilibrium conditions (e.g., under flow or by increasing the probed length scale or the magnitude of the shear excitation) makes its measurement hardly accessible and restores conventional viscous behavior. Should we believe that the elasticity collapses over a stress threshold? Again, our perception is wrong because

liquids may not completely dissipate the mechanical energy. What becomes of it then? If energy is stored in a liquid, does it appear as a thermodynamic signature?

The first microthermographic images of liquids such as glycerol, polymer melts, or water revealed that liquids emit a modulated thermal signal when subjected to a mechanical shear stress wave (1, 5). Thermal waves propagate and split the liquid in hot and cold bands several tenths of a micron wide, varying synchronously with the applied stress wave. The mechanically induced temperatures are fundamentally linked to the energy of the intermolecular distances. It is a collective effect analogous to the so-called thermoelasticity of solids, i.e., requiring strong cohesive forces.

What makes these elastic correlations so solid that thermoelastic behavior is identifiable in the so-called viscous regime? This thermomechanical coupling demonstrates that liquid molecules are not dynamically free, as was often assumed at the beginning of theoretical approaches to liquids. Instead, these molecules bind as in solids, with this cohesive energy expressible through fundamental physical constants. The insight that liquids are more similar to solids than to gases (although liquids flow and solids do not) turns out to be central to understanding the properties of liquids (9). With the

link between liquid properties and fundamental constants established, we can start asking what would happen to liquids and liquid-based life if the constants are given different values (1, 4). To enable liquid-based life, these constants need to lie in a constrained bio-friendly window.

This fundamental research will enable new applications in disciplines such as controlled fluid transport, thermal exchangers, health, and biomedicine. Revising the definition of liquids is indispensable to extend our ability to control this state of matter. In practice, medicine is already taking advantage of this promise as it is now possible to also access the elasticity of blood (10), a "remembrance" of our distant past where only elementary particles existed (1, 4).

### REFERENCES

1. K. Trachenko, Constraints on fundamental physical constants from bio-friendly viscosity and diffusion. *Sci. Adv.* **9**, eadh9024 (2023).
2. L. Noirez, Probing submillimeter dynamics to access static shear elasticity, Encyclopedia of polymers and composites, in *Polymers and polymeric composites: A reference series*, S. Palsule, Ed (Springer, 2020).
3. H. Navier, *Mémoire sur les lois des mouvements des fluides* (Mémoire de l'Académie Royale des Sciences de l'Institut de France, 1846), vol. 9, pp. 389.
4. K. Trachenko, V. Brazhkin, Minimal quantum viscosity from fundamental physical constants. *Sci. Adv.* **6**, eaba3747 (2020).
5. E. Kume, L. Noirez, Thermal shear waves induced in mesoscopic liquids at low frequency mechanical deformation. *J. Non-Equilib. Thermodyn.* **47**, 155–163 (2022).
6. B. V. Derjaguin, U. B. Bazarov, K. T. Zandanova, O. R. Budaev, The complex shear modulus of polymeric and small-molecule liquids. *Polymer* **30**, 97–103 (1989).
7. K. Trachenko, V. V. Brazhkin, Collective modes and thermodynamics of the liquid state. *Rep. Prog. Phys.* **79**, 016502 (2016).
8. A. Zaccane, K. Trachenko, Explaining the low-frequency shear elasticity of confined liquids. *Proc. Natl. Acad. Sci. U.S.A.* **117**, 19653–19655 (2020).
9. M. Kayanattila, Z. Huanga, D. Gitarica, S.W. Epp, Rubber-like elasticity in laser-driven free surface flow of a Newtonian fluid. *Proc. Natl. Acad. Sci. U.S.A.* **120**, e2301956120 (2023).
10. U. Windberger, P. Baroni, L. Noirez, Capillary-size flow of human blood plasma: Revealing hidden elasticity and scale dependence. *J. Biomed. Mater. Res.* **110**, 298–303 (2022).

10.1126/sciadv.adj8336

## Fundamental constants explain submillimeter liquid shear forces

Laurence Noirez

*Sci. Adv.*, **9** (34), eadj8336.  
DOI: 10.1126/sciadv.adj8336

### View the article online

<https://www.science.org/doi/10.1126/sciadv.adj8336>

### Permissions

<https://www.science.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of service](#)