



HAL
open science

A dense subset of $M_n(R)$ containing diagonalizable matrices

Flavien Mabilat

► To cite this version:

| Flavien Mabilat. A dense subset of $M_n(R)$ containing diagonalizable matrices. 2023. hal-04190482

HAL Id: hal-04190482

<https://hal.science/hal-04190482>

Preprint submitted on 29 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A DENSE SUBSET OF $M_n(\mathbb{R})$ CONTAINING DIAGONALIZABLE MATRICES

FLAVIEN MABILAT^a

ABSTRACT. In this note, we consider matrices similar to X -form matrices, which are the matrices for which only the diagonal and the anti-diagonal elements can be different from zero. First, we give a characterization of these matrices using the minimal polynomial. Then, we prove that the set of matrices similar to X -form matrices over \mathbb{R} and \mathbb{C} are dense and we give a characterization of the interior of this set.

Keywords : X -form matrix, minimal polynomial, Jordan normal form, companion matrix.

Declarations of interest: none

“La mémoire est aussi menteuse que l’imagination, et bien plus dangereuse avec ses petits airs studieux.”

Françoise Sagan, *Derrière l’épaule*

1. INTRODUCTION

In this note, all fields considered are commutative. Let \mathbb{K} be an arbitrary commutative field. The set of all square matrices of size n over \mathbb{K} is denoted $M_n(\mathbb{K})$, 0_n is the zero matrix of $M_n(\mathbb{K})$. If $A \in M_n(\mathbb{K})$ we denote $\pi_A(X)$ the minimal polynomial of A (with the convention π_A monic polynomial) and $\chi_A(X) = \det(XI_n - A)$ the characteristic polynomial of A (with this definition $\chi_A(X)$ is a monic polynomial). We use the convention $\prod_{i=1}^0 a_i = 1$. An elementary Jordan matrix is a matrix composed of zeroes everywhere except for the diagonal, which is filled with a fixed element $\lambda \in \mathbb{K}$, and for the superdiagonal, which is composed of ones. The Frobenius companion matrix of the monic polynomial

$$P(X) = X^n + \sum_{i=0}^{n-1} a_i X^i \text{ is the square matrix defined as } C(P) = \begin{pmatrix} 0 & \cdots & 0 & -a_0 \\ 1 & \ddots & \vdots & -a_1 \\ & \ddots & 0 & \vdots \\ & & 1 & -a_{d-1} \end{pmatrix}.$$

A classical results states that the set of diagonalizable matrices is dense over \mathbb{C} but not over \mathbb{R} . Here, we want to find a subset of $M_n(\mathbb{K})$ containing all diagonalizable matrices which is dense in the cases $\mathbb{K} = \mathbb{R}, \mathbb{C}$ and we want to find the interior of his set. For this, we will study the objects introduced in the following definitions :

Definition 1.1. *i) A X -form matrix is a square matrix for which only the diagonal and the anti-diagonal elements can be different from zero.*

ii) An endomorphism u of a finite dimensional vector space E is X -formable if there is a basis of E with respect to which the matrix of u is a X -form matrix.

iii) A X -formable matrix is a matrix similar to a X -form matrix.

First, we give some of the easy properties verifying by that kind of matrices :

- Matrices of size 1 or 2 are X -form matrices;

- X -form matrices of odd size has an eigenvalue belonging to \mathbb{K} ;
- If A is a X -form matrix then A^t is a X -form matrix;
- If A is diagonalizable then A is X -formable;
- The set of X -form matrices of size n is a vector subspace of $M_n(\mathbb{K})$;
- If A and B are X -form matrices then AB is a X -form matrix;
- If A is an invertible X -form matrix then A^{-1} is a X -form matrix (this follows from the last two points and the equality $A^{-1} = \frac{-1}{a_0}(A^{r-1} + \sum_{i=1}^{r-1} a_i A^{i-1})$ with $\pi_A(X) = X^r + \sum_{i=0}^{r-1} a_i X^i$, $a_0 \neq 0$ since A is invertible).

We also have some other properties in the case of central-symmetric X -form matrices (see [6, 7]) and also in the case of block central-symmetric X -form matrices (see [8]).

A classical result states that a square matrix is diagonalizable if and only if its minimal polynomial is a product of distinct linear factors over \mathbb{K} . It is natural to find a similar characterization in the case of X -formable matrices. This seems to be not very difficult but we have been unable to locate such a result in the litterature. Hence, our first objective is to prove the following result :

Theorem 1.2. *Let \mathbb{K} be a commutative field and $A \in M_n(\mathbb{K})$. A is X -formable if and only if*

$$\pi_A(X) = \prod_{i=1}^r P_i(X) \prod_{i=1}^q (X - \lambda_i)^{n_i},$$

with $r, q \geq 0$, P_i irreducible monic polynomials of degree 2, $\lambda_i \in \mathbb{K}$, $1 \leq n_i \leq 2$, $P_i \neq P_j$ and $\lambda_i \neq \lambda_j$ for $i \neq j$.

To simplify the proof, we will use the following notation : a polynomial verifies the property (\mathcal{P}) if it has the same factorization as in the previous theorem.

Then, we will consider some topological properties of X -formable matrices over \mathbb{R} and \mathbb{C} related to our initial goal. The main results of this text gathered in the following theorem give an answer to this problem :

Theorem 1.3. $n \in \mathbb{N}^*$. *Let F be the set of X -formable matrices of size n over \mathbb{C} and G the set of X -formable matrices of size n over \mathbb{R} .*

i) F is dense in $M_n(\mathbb{C})$.

ii) The interior of F is $\{A \in M_n(\mathbb{C}), \chi_A \text{ verifies } (\mathcal{P})\} = \{A \in M_n(\mathbb{C}), \chi_A \text{ has only simple or double roots}\}$.

iii) G is dense in $M_n(\mathbb{R})$.

iv) The interior of G is $\{A \in M_n(\mathbb{R}), \chi_A \text{ verifies } (\mathcal{P})\}$.

2. X -FORMABLE MATRICES

In this subpart, \mathbb{K} is an arbitrary commutative field. We begin by an easy lemma.

Lemma 2.1. *A matrix A is X -formable if and only if it is similar to a block-diagonal matrix in which each block has a size less than 2.*

Proof. First, we consider the case $n = 2m$. Let A be a X -formable matrix of size n . Let u be an endomorphism of \mathbb{K}^n whose matrix in the canonical basis is A . It exists $\mathcal{B} = (e_1, \dots, e_{2m})$ a basis of \mathbb{K}^n such that the matrix of u in \mathcal{B} is a X -form matrix. The matrix of u in the basis $(e_1, e_{2m}, e_2, e_{2m-1}, \dots, e_m, e_{m+1})$ is a block-diagonal matrix in which each block has a size less than 2. Let A be a matrix similar to a block-diagonal matrix in which each block has a size less than 2 and u be an endomorphism of \mathbb{K}^n whose matrix in the canonical basis is A . It exists $\mathcal{B}' = (f_1, \dots, f_{2m})$ a basis of \mathbb{K}^n such that the matrix of u

in \mathcal{B}' is a block-diagonal matrix in which each block has a size less than 2. The matrix of u in the basis $(f_1, f_3, \dots, f_4, f_2)$ is a X -form matrix. The proof of the case $n = 2m + 1$ is similar. \square

Note that this result, combined with the normal matrices reduction theorem (see [3] Theorem 4, p 271), allows us to see that normal matrices on \mathbb{R} are X -formable.

2.1. Proof of theorem 1.2.

Proof. Let A be a X -formable matrix of size n . By the previous lemma, it exists a basis of \mathbb{K}^n such that the matrix of u in this basis is a block-diagonal matrix C in which each block has a size less than 2. Thus, π_A is the least common multiple of the minimal polynomials of the diagonal blocks of C . The minimal polynomials of matrix of size 1 or 2 are one of the following type :

- irreducible monic polynomial of degree 2;
- $(X - \lambda)^2$;
- $(X - \lambda)$;
- $(X - \lambda_1)(X - \lambda_2)$, $\lambda_1 \neq \lambda_2$.

Hence, they verify (\mathcal{P}) . Thus, π_A verifies (\mathcal{P}) .

Now, we want to prove the remaining implication of the theorem.

First, we consider a matrix A of size n such that π_A verifies (\mathcal{P}) and such that π_A has only one irreducible factor (over \mathbb{K}). We have three cases :

- $\pi_A = (X - \lambda)$ then A is diagonalizable, and so X -formable.
- $\pi_A = (X - \lambda)^2$. By the decomposition theorem of Jordan, A is similar to a block diagonal matrix

$$H = \begin{pmatrix} J_1 & & \\ & \ddots & \\ & & J_l \end{pmatrix} \text{ in which } J_i \text{ is an elementary Jordan matrix. Besides, each } J_i \text{ is a square}$$

matrix of size $n_i \in \{1, 2\}$. Indeed, $\pi_A = \pi_H = \text{lcm}(\pi_{J_i}, 1 \leq i \leq l)$ and $\pi_{J_i} = (X - \lambda)^{n_i}$. Hence, A is X -formable, by lemma 2.1.

- π_A is an irreducible monic polynomial of degree 2. By the decomposition theorem of Frobenius, A is similar to a block diagonal matrix

$$H = \begin{pmatrix} C(R_1) & & \\ & \ddots & \\ & & C(R_t) \end{pmatrix},$$

with R_i monic polynomials verifying R_i divides R_{i+1} . Since $\pi_A = \pi_H = \text{lcm}(R_i, 1 \leq i \leq t)$ and π_A irreducible, $R_i = \pi_A$. Thus, H is block-diagonal matrix in which each block has size 2 and A is X -formable, by lemma 2.1.

Now, we proceed by induction on the size n of the matrix. If $n = 1$, then the result is true. Suppose it exists $n \geq 1$ such that all matrices of size less than n whose minimal polynomial verifies (\mathcal{P}) are X -formable. Let A be a matrix of size $n + 1$ whose minimal polynomial verifies (\mathcal{P}) and u the endomorphism canonically associated to A . If π_A has only one irreducible factor, then the result is true by the previous discussion. Suppose π_A has several irreducible factors. It exists a monic polynomial P such that $\pi_A = P \frac{\pi_A}{P}$, P has degree greater or equal to 1, P and $\frac{\pi_A}{P}$ are relatively prime polynomials. P and $\frac{\pi_A}{P}$ verify (\mathcal{P}) . Consider $F = \text{Ker}(P(u))$ and $G = \text{Ker}(\frac{\pi_A}{P}(u))$. Hence, by the kernel lemma¹, $\mathbb{K}^{n+1} = F \oplus G$

¹This translation of the French name "lemme des noyaux" seems to be the most used name for this result.

and F, G are invariant subspaces of u .

By induction assumption, $u|_G$ (restriction of u to G) and $u|_F$ are X -formable. Hence, u is X -formable and theorem 1.2 is proved. \square

2.2. Some additional elements about theorem 1.2.

In the case of an algebraically closed field we have the following result :

Corollary 2.2. *Let \mathbb{K} be an algebraically closed field and $A \in M_n(\mathbb{K})$. The following assertions are equivalent :*

i) A is X -formable.

ii) $\pi_A(X) = \prod_{i=1}^q (X - \lambda_i)^{n_i}$ with $q \geq 1$, $\lambda_i \in \mathbb{K}$, $1 \leq n_i \leq 2$ and $\lambda_i \neq \lambda_j$ for $i \neq j$.

iii) All the Jordan blocks appearing in the Jordan normal form of A have their size equal to 1 or 2.

If A is a X -formable matrix then there is not a unique X -form matrix similar to A . For example, $A = \begin{pmatrix} 1 & 1 \\ 0 & 2 \end{pmatrix}$ is a X -form matrix but A is also similar to $\begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$ which is also a X -form matrix.

We now give the two following examples :

- We consider $\mathbb{K} = \mathbb{R}$ and $B = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$. $\pi_B(X) = (X - 1)^3$. Hence, by the theorem 1.2, B is not X -formable.

- We consider $\mathbb{K} = \mathbb{R}$ and $C = \begin{pmatrix} -22 & 47 & -19 & 18 \\ 1 & 3 & -3 & -5 \\ 14 & -23 & 7 & -16 \\ -15 & 27 & -9 & 17 \end{pmatrix}$. $\pi_C(X) = (X - 2)(X - 3)(X^2 + 2)$.

Hence, by the theorem 1.2, C is X -formable. For instance,

$$C = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 0 & 2 & 1 & 2 \\ 0 & 3 & 0 & 1 \\ 1 & 0 & 1 & 1 \end{pmatrix} \begin{pmatrix} 4 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & -3 & -1 & 0 \\ -2 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} -1 & 1 & 0 & 2 \\ 1 & -2 & 1 & -1 \\ 4 & -7 & 2 & -4 \\ -3 & 6 & -2 & 3 \end{pmatrix}.$$

We conclude this part by giving some elements about the product of X -formable matrices. We have the following result :

Theorem 2.3 (Botha, [1] Theorem 2.1). *Let \mathbb{K} be any field such that the characteristic of \mathbb{K} is different from 2 and such that $\mathbb{K} \neq \mathbb{F}_3$ (the field with 3 elements). Then every matrix over \mathbb{K} is a product of two diagonalizable matrices.*

Since diagonalizable matrices are X -formable, this result covers a lot of cases. Here, we consider the case of \mathbb{F}_3 . By the decomposition theorem of Frobenius, it is sufficient to consider companion matrices. We have the following equality :

$$\begin{pmatrix} 0 & \cdots & 0 & a_0 \\ 1 & \ddots & \vdots & a_1 \\ & \ddots & 0 & \vdots \\ & & 1 & a_{d-1} \end{pmatrix} = \begin{pmatrix} & & & 1 \\ & \cdots & & \\ 1 & & & \end{pmatrix} \begin{pmatrix} & & & 1 & a_{d-1} \\ & \ddots & & \vdots & \\ 1 & & & a_1 & \\ 0 & \cdots & 0 & a_0 & \end{pmatrix}.$$

The first matrix on the right is diagonalizable since its minimal polynomial is $(X-1)(X+1)$ and $1 \neq -1$ in \mathbb{F}_3 . If $a_0 = \pm 1$ then the second matrix on the right is X -formable since its minimal polynomial divides $(X-1)^2(X+1)^2$ (theorem 1.2). If $a_0 = 0$ then the second matrix on the right is X -formable since its minimal polynomial divides $X^2(X-1)^2(X+1)^2$ (theorem 1.2). Hence, every matrix over \mathbb{F}_3 is a product of two X -formable matrices.

3. SOME TOPOLOGICAL ASPECTS OF THE SET OF X -FORMABLE MATRICES

The aim of this section is to prove theorem 1.3. Here, we suppose $\mathbb{K} = \mathbb{R}, \mathbb{C}$. The set of polynomials of degree less than n over \mathbb{K} is denoted $\mathbb{K}_n[X]$. We use the following norm, if $A = (a_{i,j})_{1 \leq i, j \leq n} \in M_n(\mathbb{K})$ we denote $\|A\|_\infty = \max(|a_{i,j}|, 1 \leq i, j \leq n)$ (we recall that all norms are equivalent on $M_n(\mathbb{K})$). We start by some preliminary results.

3.1. Preliminary lemmas. In this subsection, we give some results concerning polynomials over \mathbb{K} . If $P(X) = \sum_{k=0}^n a_k X^k$, we denote $\|P\|_\infty = \max(|a_k|, 0 \leq k \leq n)$ and $D(z, r) = \{w \in \mathbb{C}, |z - w| < r\}$. We begin by the following well-known result :

Theorem 3.1 (Continuity of the roots of a polynomial, [4]). *Let P be a polynomial, z_i its distinct roots with $i = 1, \dots, p$, m_i the multiplicity of the root z_i ($m_1 + \dots + m_p = \deg(P)$). Then for any $\epsilon > 0$ such that $D(z_i, \epsilon) \cap D(z_j, \epsilon) = \emptyset$ for any $i \neq j$ there exists $\eta > 0$ such that any polynomial Q whose coefficients differ from those of P only by less than η has exactly m_i roots (distinct or not) in $D(z_i, \epsilon)$.*

Now we can prove the following result :

Proposition 3.2. $n \in \mathbb{N}^*$. *The set \hat{F} of polynomials over \mathbb{C} of degree n whose roots are simple or double is an open subset of $\mathbb{C}_n[X]$.*

Proof. Let $P \in \mathbb{C}_n[X]$ with $\deg(P) = n$. We assume that P has only simple and double roots. We denote z_i its distinct roots with $i = 1, \dots, p$ and m_i the multiplicity of the root z_i . We have $m_i \leq 2$ and $m_1 + \dots + m_p = n$.

Since \mathbb{C} is a separated normed space, it exists $\epsilon > 0$ such that $D(z_i, \epsilon) \cap D(z_j, \epsilon) = \emptyset$ for any $i \neq j$. Hence, by the previous theorem it exists $\eta > 0$ such that any polynomial Q verifying $\|P - Q\|_\infty < \eta$ has exactly m_i roots (distinct or not) in $D(z_i, \epsilon)$. So, in each $D(z_i, \epsilon)$, Q has exactly one root of multiplicity less than two or two roots of multiplicity one. Besides, Q has n roots (with multiplicity) in $\bigcup_{i=1}^p D(z_i, \epsilon)$ and $\deg(Q) \geq n$.

Hence, a polynomial T belonging to $\mathbb{C}_n[X]$ and verifying $\|P - T\|_\infty < \eta$ has degree n and all its roots are simple or double. Thus, \hat{F} is an open subset of $\mathbb{C}_n[X]$. □

This result is no longer true if we replace \mathbb{C} by \mathbb{R} . For instance, we can consider the polynomial sequence $P_n = X^2 + \frac{1}{n}X + \frac{1}{n}$. For all $n \in \mathbb{N}^*$, P_n is irreducible, since its discriminant is $\frac{1}{n}(\frac{1}{n} - 4) < 0$. However, this sequence converges to X^2 . Hence, the complementary of the set of polynomials over \mathbb{R} of degree 2 whose roots are simple or double is not closed. So, this set is not an open set. However, we can still have a similar result :

Proposition 3.3. $n \in \mathbb{N}^*$. *The set \hat{G} of polynomials over \mathbb{R} of degree n which verify (\mathcal{P}) is an open subset of $\mathbb{R}_n[X]$.*

Proof. Let $P \in \mathbb{R}_n[X]$ with $\deg(P) = n$. We assume that P verifies (\mathcal{P}) . We denote z_i its distinct roots in \mathbb{C} with $i = 1, \dots, p$ and m_i the multiplicity of the root z_i . Since P verify (\mathcal{P}) , we have $m_i \leq 2$ and $m_1 + \dots + m_p = n$. Besides, if $z_i \in \mathbb{C} - \mathbb{R}$ then $m_i = 1$ and \bar{z}_i is also a root of P of multiplicity 1.

Since \mathbb{C} is a separated normed space, it exists $\epsilon > 0$ such that $D(z_i, \epsilon) \cap D(z_j, \epsilon) = \emptyset$ for any $i \neq j$. Hence, by the previous theorem it exists $\eta > 0$ such that any polynomial Q in $\mathbb{R}[X] \subset \mathbb{C}[X]$ verifying $\|P - Q\|_\infty < \eta$ has exactly m_i roots (distinct or not) in $D(z_i, \epsilon)$. In particular, Q has n roots in \mathbb{C} (with multiplicity) belonging to $\bigcup_{i=1}^p D(z_i, \epsilon)$ and $\deg(Q) \geq n$. We have several possible cases for the roots of Q belonging to $D(z_i, \epsilon)$:

- $D(z_i, \epsilon)$ contains exactly one real root of multiplicity less than two or two real roots of multiplicity one.
- $D(z_i, \epsilon)$ contains exactly one root λ of multiplicity one belonging to $\mathbb{C} - \mathbb{R}$. Then, $D(\bar{z}_i, \epsilon)$ contains exactly one root $\bar{\lambda}$ of multiplicity one. Thus, the polynomial $(X - \lambda)(X - \bar{\lambda})$ is an irreducible real factor of Q and this factor appears only one time in Q .
- $D(z_i, \epsilon)$ contains exactly two roots of multiplicity one, λ and μ , belonging to $\mathbb{C} - \mathbb{R}$. In this case, $z_i \in \mathbb{R}$ (since the non-real roots of P has multiplicity one). Besides, $\bar{\lambda}$ and $\bar{\mu}$ are roots of Q of multiplicity one and they also belong to $D(z_i, \epsilon)$ (since $z_i \in \mathbb{R}$). Hence, $\mu = \bar{\lambda}$. Thus, the polynomial $(X - \lambda)(X - \bar{\lambda})$ is an irreducible real factors of Q and this factor appears only one time in Q .
- $D(z_i, \epsilon)$ contains exactly one real root of multiplicity one x and one root of multiplicity one λ belonging to $\mathbb{C} - \mathbb{R}$. In this case, $z_i \in \mathbb{R}$ (since the non-real roots of P has multiplicity one). Besides, $\bar{\lambda}$ is a root of Q of multiplicity one which belongs to $D(z_i, \epsilon)$ (since $z_i \in \mathbb{R}$). Hence, this case is not possible.

Hence, a polynomial T belonging to $\mathbb{R}_n[X]$ and verifying $\|P - T\|_\infty < \eta$ has degree n and verifies (\mathcal{P}) . Thus, \hat{G} is an open subset of $\mathbb{R}_n[X]$. □

Remark 3.4. In the first case considered in the proof, $z_i \in \mathbb{R}$. Indeed, suppose z_i is a non-real root of P . $D(z_i, \epsilon)$ contains only one root of multiplicity one (since the non-real roots of P has multiplicity one). \bar{z}_i is a root of P of multiplicity one. Hence, $D(\bar{z}_i, \epsilon)$ contains exactly one root of Q . We set x this root. x is necessarily real since otherwise $D(z_i, \epsilon)$ would contain a non-real root which would be \bar{x} . Besides, $|\bar{z}_i - x| = |z_i - x| \leq \epsilon$. Thus, $x \in D(z_i, \epsilon) \cap D(\bar{z}_i, \epsilon)$. This is not possible since $D(z_i, \epsilon) \cap D(\bar{z}_i, \epsilon) = \emptyset$. Thus, $z_i \in \mathbb{R}$.

3.2. Proof of theorem 1.3.

Proof. We consider the following continued map
$$\varphi : \begin{array}{ccc} M_n(\mathbb{K}) & \longrightarrow & \mathbb{K}_n[X] \\ A & \longmapsto & \chi_A \end{array} .$$

i) The set of diagonalizable matrices $D_n(\mathbb{C})$ is included in F . Moreover, $D_n(\mathbb{C})$ is dense in $M_n(\mathbb{C})$. Hence, F is dense in $M_n(\mathbb{C})$.

ii) \hat{F} is an open subset of $\mathbb{C}_n[X]$ (proposition 3.2). Hence,

$$\varphi^{-1}(\hat{F}) = \{A \in M_n(\mathbb{C}), \chi_A \text{ verifies } (\mathcal{P})\} = \{A \in M_n(\mathbb{C}), \chi_A \text{ has only simple or double roots}\}$$

is an open subset of $M_n(\mathbb{C})$. Besides, $\varphi^{-1}(\hat{F})$ is included in F (by the theorem 1.2 and the theorem of Cayley-Hamilton). Hence, $\varphi^{-1}(\hat{F})$ is included in the interior of F .

Let $A \in F$ such that χ_A doesn't verify (\mathcal{P}) . By corollary 2.2, all the Jordan blocks appearing in the

Jordan normal form of A have their size equal to 1 or 2. Since, χ_A doesn't verify (P) , it exists λ such that $(X - \lambda)^3$ divides χ_A . Thus, one of the following occurs :

- a) the Jordan normal form of A contains two blocks $J_\lambda = \begin{pmatrix} \lambda & 1 \\ 0 & \lambda \end{pmatrix}$;
- b) the Jordan normal form of A contains three blocks (λ) ;
- c) the Jordan normal form of A contains one block $J_\lambda = \begin{pmatrix} \lambda & 1 \\ 0 & \lambda \end{pmatrix}$ and one block (λ) .

We consider each case separately :

- If a) occurs. It exists $P \in GL_n(\mathbb{C})$ such that $A = P \begin{pmatrix} J_\lambda & & \\ & J_\lambda & \\ & & B \end{pmatrix} P^{-1}$. For all $n \in \mathbb{N}^*$, we

$$\text{define } A_n = P \begin{pmatrix} \lambda & 1 & \frac{1}{n} & 0 \\ 0 & \lambda & 0 & \frac{1}{n} \\ 0 & 0 & \lambda & 1 \\ 0 & 0 & 0 & \lambda \\ & & & & B \end{pmatrix} P^{-1}. (A_n) \text{ converges to } A. \text{ However, for all } n \in \mathbb{N}^*, (X - \lambda)^3$$

divides π_{A_n} . Hence, $A_n \notin F$ (theorem 1.2). Thus, A doesn't belong to the interior of F .

- If b) occurs. It exists $P \in GL_n(\mathbb{C})$ such that $A = P \begin{pmatrix} \lambda & & & \\ & \lambda & & \\ & & \lambda & \\ & & & B \end{pmatrix} P^{-1}$. For all $n \in \mathbb{N}^*$, we

$$\text{define } A_n = P \begin{pmatrix} \lambda & \frac{1}{n} & 0 \\ 0 & \lambda & \frac{1}{n} \\ 0 & 0 & \lambda \\ & & & B \end{pmatrix} P^{-1}. (A_n) \text{ converges to } A. \text{ However, for all } n \in \mathbb{N}^*, (X - \lambda)^3$$

divides π_{A_n} . Hence, $A_n \notin F$ (theorem 1.2). Thus, A doesn't belong to the interior of F .

- If c) occurs. It exists $P \in GL_n(\mathbb{C})$ such that $A = P \begin{pmatrix} J_\lambda & & \\ & \lambda & \\ & & B \end{pmatrix} P^{-1}$. For all $n \in \mathbb{N}^*$, we define

$$A_n = P \begin{pmatrix} \lambda & 1 & 0 \\ 0 & \lambda & \frac{1}{n} \\ 0 & 0 & \lambda \\ & & & B \end{pmatrix} P^{-1}. (A_n) \text{ converges to } A. \text{ However, for all } n \in \mathbb{N}^*, (X - \lambda)^3 \text{ divides}$$

π_{A_n} . Hence, $A_n \notin F$ (theorem 1.2). Thus, A doesn't belong to the interior of F .

Hence, the interior of F is equal to $\varphi^{-1}(\hat{F})$.

iii) Let $A \in M_n(\mathbb{R})$. If A is triangularizable then it exists a sequence (A_n) of diagonalizable matrices such that (A_n) converges to A .

Suppose now A is not triangularizable. Thus, $\pi_A = \prod_{i=1}^r P_i^{n_i} \prod_{i=1}^l (X - \lambda_i)^{m_i}$ with $r \geq 1$, $l \geq 0$, $\lambda_i \in \mathbb{R}$, $\lambda_i \neq \lambda_j$ for $i \neq j$, $n_i, m_i \geq 1$ and P_i irreducible monic polynomials of degree 2, $P_i \neq P_j$ for $i \neq j$. Therefore, A is similar to a block-diagonal matrix in which each block has its minimal polynomial equals to $P_i^{n_i}$ or $(X - \lambda_i)^{m_i}$ (by the kernel lemma). Hence, it is sufficient to consider square matrices whose minimal polynomial is the power of an irreducible monic polynomial of degree 2 (since the case of triangularizable matrices has already been considered).

Let $B \in M_n(\mathbb{R})$ such that $\pi_B = P^m$ with P an irreducible monic polynomial of degree 2 and $m \leq \frac{n}{2}$. It exists $\lambda \in \mathbb{C} - \mathbb{R}$ such that $P(X) = (X - \lambda)(X - \bar{\lambda})$. By the decomposition theorem of Frobenius, B is similar to a block diagonal matrix

$$H = \begin{pmatrix} C(R_1) & & \\ & \ddots & \\ & & C(R_t) \end{pmatrix},$$

with R_i real polynomials verifying R_i divides R_{i+1} . Since $\pi_B = \pi_H = \text{lcm}(R_i, 1 \leq i \leq t)$ and $\pi_B = P^m$ with P irreducible, R_i is a power of P . Thus, it exists $1 \leq r_i \leq m$ such that $R_i = P^{r_i} = (X - \lambda)^{r_i}(X - \bar{\lambda})^{r_i}$. We set $R_{i,n} = \prod_{j=1}^{r_i} (X - \lambda + \frac{j}{n^2})(X - \bar{\lambda} + \frac{j}{n^2})$. $(R_{i,n})$ converges to R_i . Hence, $C(R_{i,n})$ converges to $C(R_i)$. Besides, $R_{i,n}$ verifies (\mathcal{P}) for all $n \in \mathbb{N}^*$ and $\pi_{C(R_{i,n})} = R_{i,n}$. Thus, $C(R_{i,n})$ is X -formable (theorem 1.2).

So, it exists a sequence of X -formable matrices which converges to B .

Hence, G is dense in $M_n(\mathbb{R})$.

iv) \hat{G} is an open subset of $\mathbb{R}_n[X]$ (proposition 3.3). Hence, $\varphi^{-1}(\hat{G}) = \{A \in M_n(\mathbb{R}), \chi_A \text{ verifies } (\mathcal{P})\}$ is an open subset of $M_n(\mathbb{R})$. Besides, $\varphi^{-1}(\hat{G})$ is included in G (by the theorem 1.2 and the theorem of Cayley-Hamilton). Hence, $\varphi^{-1}(\hat{G})$ is included in the interior of G .

Let $A \in G$ such that χ_A doesn't verify (\mathcal{P}) . A is similar to a block-diagonal matrix in which each block has a size less than 2 (lemme 2.1). Besides, each triangularizable block has a Jordan normal form. Hence, A is similar to a block-diagonal matrix C in which each block is a Jordan block of size 1 or 2 or a square matrix of size 2 whose characteristic polynomial is an irreducible monic polynomial of degree 2. Since, χ_A doesn't verify (\mathcal{P}) , it exists λ such that $(X - \lambda)^3$ divides χ_A or it exists an irreducible monic polynomial P of degree 2 such that P^2 divides χ_A . Thus, one of the following occurs :

- a) C contains two blocks $J_\lambda = \begin{pmatrix} \lambda & 1 \\ 0 & \lambda \end{pmatrix}$;
- b) C contains three blocks (λ) ;
- c) C contains one block $J_\lambda = \begin{pmatrix} \lambda & 1 \\ 0 & \lambda \end{pmatrix}$ and one block (λ) ;
- d) C contains two blocks $T = \begin{pmatrix} x & y \\ z & t \end{pmatrix}$ and $O = \begin{pmatrix} u & v \\ w & h \end{pmatrix}$ which have the same irreducible characteristic polynomial $S(X) = X^2 + \alpha X + \beta$. In particular, $y, z, v, w \neq 0$.

We can study the first three cases in the same way as that used previously.

If d) occurs. It exists $P \in GL_n(\mathbb{R})$ such that $A = P \begin{pmatrix} T & & \\ & O & \\ & & B \end{pmatrix} P^{-1}$. For all $n \in \mathbb{N}^*$, we define $A_n = P \begin{pmatrix} x & y & 0 & \frac{1}{n} \\ z & t & 0 & 0 \\ & u & v & \\ & w & h & \\ & & & B \end{pmatrix} P^{-1}$. The minimal polynomial of $Z_n = \begin{pmatrix} x & y & 0 & \frac{1}{n} \\ z & t & 0 & 0 \\ 0 & 0 & u & v \\ 0 & 0 & w & h \end{pmatrix}$ is equal to the minimal

polynomial of A_n . Besides, S divides the minimal polynomial of Z_n . $S(Z_n) = \begin{pmatrix} 0 & 0 & \frac{w}{n} & \frac{x+h+\alpha}{n} \\ 0 & 0 & 0 & \frac{z}{n} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \neq 0_4$

(since $w \neq 0$) and $S^2(Z_n) = 0_4$. Hence, $\pi_{Z_n} \neq S$ and π_{Z_n} divides S^2 . Thus, $\pi_{Z_n} = S^2$, since S is irreducible. Hence, $\pi_{A_n} = S^2$. So, π_{A_n} doesn't verify (\mathcal{P}) and $A_n \notin G$ (theorem 1.2). However, (A_n) converges to A . Thus, A doesn't belong to the interior of G .

Hence, the interior of G is equal to $\varphi^{-1}(\hat{G})$. □

We have therefore, as announced, a dense subset of $M_n(\mathbb{R})$ and $M_n(\mathbb{C})$ and a complete characterization of the interior and of this set. We conclude by noticing a curious similarity between the diagonalizable matrices and the X -formable matrices. For this, we will use the following notation : a polynomial verifies the property (\mathcal{Q}) if it is a product of distinct linear factors over \mathbb{K} . Let $A \in M_n(\mathbb{K})$ with $\mathbb{K} = \mathbb{R}, \mathbb{C}$, we have :

- A is diagonalizable if and only if π_A verifies (\mathcal{Q}) ; A belongs to the interior of the set of diagonalizable matrices if and only if χ_A verifies (\mathcal{Q}) ;
- A is X -formable if and only if π_A verifies (\mathcal{P}) ; A belongs to the interior of the set of X -formable matrices if and only if χ_A verifies (\mathcal{P}) .

4. SOME OPEN PROBLEMS

We collect here some open problems related to X -formable matrices. The first problem is related to theorem 1.3. Indeed, we know some topological properties of the set of X -formable matrices. Therefore, it is natural to want to look for other.

Problem 4.1. *Find other topological properties of the set of X -formable matrices over \mathbb{R} and \mathbb{C} .*

We have a certain number of results concerning the maximum dimension of a vector subspace all of whose elements verify a given property (such as Gerstenhaber's theorem, see [2]). This naturally leads to the following problem :

Problem 4.2. *What is the maximal dimension of a vector subspace of $M_n(\mathbb{K})$ containing only X -formable elements ?*

In the case of diagonalizable matrices we have the following result :

Theorem 4.3 (Klarès's criterion, [5] p 125). *Let $B \in M_n(\mathbb{K})$ with \mathbb{K} an algebraically closed field. We set $Ad_B : M \in M_n(\mathbb{K}) \mapsto BM - MB$. B is diagonalizable if and only if $Ker(Ad_B) = Ker(Ad_B^2)$.*

This leads to the formulation of the problem below :

Problem 4.4. *Can we find a "similar" criterion for X -formable matrices ?*

REFERENCES

- [1] J. D. Botha, *Products of diagonalizable matrices*, Linear algebra and its applications, Vol. 273 1-3, (1998), pp 65-82.
- [2] M. Gerstenhaber, *On nilalgebras and linear varieties of nilpotent matrices (I)*, Amer. J. Math., Vol. 80, (1958), pp 614-622.
- [3] X. Gourdon, *Les maths en tête : Algèbre*, 2^{ème} édition, Ellipses, 2009.
- [4] D. Leborgne, *Calcul différentiel complexe*, Presses universitaires de France, 1996.
- [5] R. Mansuy, R. Mneimné, *Algèbre linéaire: Réduction des endomorphismes*, De Boeck Supérieur, 2016.
- [6] A. M. Nazari, E. Afshari, B.A. Omid, *Properties of Central Symmetric X -Form Matrices*, Iranian Journal of Mathematical Sciences and Informatics, Vol. 6, No. 2 (2011), pp 9-20.

- [7] A. Sadeghi, M.S. Solary, *The exponential functions of central-symmetric X-form matrices*, Journal of Mathematical Modeling, Volume 4, Issue 1 (2016), pp 19-34.
- [8] A. Sadeghi, *On the Function of Block Anti Diagonal Matrices and Its Applications*, International Journal of Mathematical Modelling & Computations, Vol. 06, No. 02 (2016), pp 105- 117.

^a, LABORATOIRE DE MATHÉMATIQUES DE REIMS, UMR9008 CNRS ET UNIVERSITÉ DE REIMS CHAMPAGNE-ARDENNE,
U.F.R. SCIENCES EXACTES ET NATURELLES MOULIN DE LA HOUSSE - BP 1039 51687 REIMS CEDEX 2, FRANCE
Email address: flavien.mabilat@univ-reims.fr