

Viscous theory for three-layer Rayleigh-Taylor instability.

Antoine Simon, Jorge-César Brändle de Motta, Christophe
Dumouchel & Marie-Charlotte Renout

CNRS UMR 6614 - CORIA
University of Rouen Normandy



HILIMBA: 2 labs (CORIA/LOMC), 2 PhD students, 320 k€

ICMF, Kobe, Japan, 2-7 April 2023

1 - Rayleigh-Taylor instability (RTI)

Hydrodynamic instability

- ▶ at a **fluid-fluid** interface
- ▶ when **accelerated** in the direction of the density gradient

$$\frac{(2) : \rho_2 > \rho_1}{(1) : \rho_1} \quad \uparrow \vec{\nabla} \rho \quad \uparrow \vec{a}$$

$$\frac{(2) : \rho_2 > \rho_1}{(1) : \rho_1} \quad \uparrow \vec{\nabla} \rho \quad \downarrow \vec{g}$$

Digitations of **various** shapes

- ▶ (1) \equiv oil / (2) \equiv water $\rightarrow \rho_2 - \rho_1 \sim 10^2 \text{ kg/m}^3$



$t = 0,04\text{s}$



$t = 0,10\text{s}$



$t = 0,16\text{s}$



$t = 0,24\text{s}$

- ▶ (1) \equiv air / (2) \equiv water $\rightarrow \rho_2 - \rho_1 \sim 10^3 \text{ kg/m}^3$




$t = 0,04\text{s}$



$t = 0,09\text{s}$



$t = 0,16\text{s}$



$t = 0,21\text{s}$

1 - 2-layer fluid system \rightarrow 3-layer fluid system

Reference: S Chandrasekhar, *Hydrodynamic and Hydromagnetic Stability*, 1961, Chapter X.

Fields of observation: geophysics, astrophysics and nuclear safety
 \rightarrow **multi-layer** fluid systems

Case of study: a **3-layer** fluid system immersed in a gravity field

$$\begin{array}{c} (3) : \rho_3 \\ \hline (2) : \rho_2 \\ \hline (1) : \rho_1 \end{array} \quad \downarrow \rho g$$

\rightarrow 2 unstable interfaces if $\rho_{i+1} > \rho_i$ for $i \in \{1; 2\}$

Focus: RTI at the **lower** interface

$$\begin{array}{c} (3) : \rho_3 < \rho_2 \\ \hline (2) : \rho_2 > \rho_1 \\ \hline (1) : \rho_1 \end{array} \quad \begin{array}{c} \downarrow \vec{\nabla} \rho \\ \uparrow \vec{\nabla} \rho \end{array} \downarrow \rho g$$

1 - Review of the literature


Theoretical studies

1. KO Mikaelian Phys. Rev. A 1983
→ inviscid model, surface tension ignored, analytical solutions.
2. S Parhi & G Nath Int. J. Engng Sci. 1991
→ viscous model, solving method, stability criterion.
3. KO Mikaelian Phys. Rev. E 1996
→ viscous model, solving method.

Experimental work

1. JW Jacobs & SB Dalziel J. Fluid Mech. 2005
→ miscible fluids, method of barrier removal, turbulent initial flow.
2. R Adkins, EM Shelton, MC Renoult, P Carles, C Rosenblatt Phys. Rev. Fluids 2017
→ non-miscible fluids, method of magnetic levitation, sinusoidal initial interface deformation.

1 - Motivation



Set of images versus time for one experimental run from Adkins et al.

Growth rate measurements at the lower interface


- ▶ assuming exponential growth
- ▶ **lower** values than the ones predicted by the inviscid Mikaelian theory.

Sources of discrepancy

1. Fluids **viscosity**
2. **Sides** effects
3. Interfaces **coupling**

→ **viscous** theory and **comparison** to experiments and numerical simulations.

2 - Formulation



Assumptions:

- ▶ **Incompressible** fluids.
- ▶ **Newtonian** fluids.
- ▶ **Non miscible** fluids.
- ▶ Extreme layers of **infinite** depth.
- ▶ **Isothermal** conditions.

Condition for RTI occurrence:

$$\rho_3 > \rho_2 \text{ and/or } \rho_2 > \rho_1.$$


2 - Equations

Volume equations in each phase $i \in \{1; 2; 3\}$:

$$\vec{\nabla} \cdot \vec{u}_i = 0 \quad (\text{mass balance})$$

$$\rho_i \left(\vec{u}_{i,t} + (\vec{u}_i \cdot \vec{\nabla}) \vec{u}_i \right) = -\vec{\nabla} p_i + \mu_i \vec{\Delta} \vec{u}_i + \rho_i \vec{g} \quad (\text{momentum balance})$$

Jump conditions at each interface $ii + 1$
with $i \in \{1; 2\}$:



$$\vec{u}_i \cdot \vec{n}_i = \vec{v}_{ii+1} \cdot \vec{n}_i = \vec{u}_{i+1} \cdot \vec{n}_i \quad (\text{mass balance})$$

$$(\rho_{i+1} - \rho_i + \gamma_{ii+1} \kappa_{ii+1}) \vec{n}_i = (\bar{\tau}_{i+1} - \bar{\tau}_i) \cdot \vec{n}_i \quad (\text{momentum balance})$$

$$\vec{u}_i \times \vec{n}_i = \vec{u}_{i+1} \times \vec{n}_i \quad (\text{no slip})$$

with $\vec{n}_i = (-z_{ii+1,x}, -z_{ii+1,y}, 1)^T$, $\kappa_{ii+1} = \frac{z_{ii+1,x,x}}{(1+z_{ii+1,x}^2)^{\frac{3}{2}}} + \frac{z_{ii+1,y,y}}{(1+z_{ii+1,y}^2)^{\frac{3}{2}}}$ and

$$\bar{\tau}_i = \mu_i (\vec{\nabla} \vec{u}_i + (\vec{\nabla} \vec{u}_i)^T)$$

2 - Basic state

Definition:

- ▶ Fluids at **rest**: $\vec{u}_i^* = \vec{0}$
- ▶ Interface 12 **plane**: $z_{12}^* = 0$
- ▶ Interface 23 **plane**: $z_{23}^* = h$

Solution for the pressure:

$$p_1^*(z) = -\rho_1 g z + p_c$$

$$p_2^*(z) = -\rho_2 g z + p_c$$

$$p_3^*(z) = -\rho_3 g z + gh(\rho_3 - \rho_2) + p_c$$

with $p_c = p_1^*(z = 0)$ a constant.

2 - Perturbed state

Definition:

$$\begin{aligned} \bar{u}_i(x, y, z, t) &= \bar{u}_i^* + \bar{u}'_i(x, y, z, t) = \bar{u}'_i \text{ with } u'_i \ll 1 \\ \rho_i(x, y, z, t) &= \rho_i^*(z) + \rho'_i(x, y, z, t) \text{ with } \rho'_i \ll \rho_i^* \\ z_{ii+1}(x, y, t) &= z_{ii+1}^* + z'_{ii+1}(x, y, t) \text{ with } z'_{ii+1} \ll z_{ii+1}^* \end{aligned}$$

Volume equations in each phase $i \in \{1; 2; 3\}$:

Mass balance: $u_{ij,j} = 0$

Linearized momentum balance: $\rho_i u_{ij,t} = -\rho'_{i,j} + \mu_i \Delta u_{ij} \quad j \in \{x; y; z\}$

Linearized jump conditions at $z = (i - 1)h$, $i \in \{1; 2\}$:

$$\begin{aligned} u_{iz} &= z'_{ii+1,t} = u_{i+1z} \\ u_{ix} &= u_{i+1x} \\ u_{iy} &= u_{i+1y} \\ \rho'_{i+1} - \rho'_i - (\rho_{i+1} - \rho_i) g z'_{ii+1} - \gamma_{ii+1} (z'_{ii+1,x,x} + z'_{ii+1,y,y}) &= \\ &= 2(\mu_{i+1} u_{i+1z,z} - \mu_i u_{iz,z}) \\ \mu_{i+1} (u_{i+1z,x} + u_{i+1x,z}) &= \mu_i (u_{iz,x} + u_{ix,z}) \\ \mu_{i+1} (u_{i+1z,y} + u_{i+1y,z}) &= \mu_i (u_{iz,y} + u_{iy,z}) \end{aligned}$$

2 - Dispersion relation

- ▶ Solutions searched under the form: $e^{(ik_x x + ik_y y + \omega t)}$ with $k^2 = k_x^2 + k_y^2$ the wavenumber of the perturbation and ω the time coefficient (complex number).
- ▶ Volume equations $\rightarrow u_{iz}$:

$$u_{1z} = (C_1 e^{kz} + D_1 e^{q_1 z}) e^{(ik_x x + ik_y y + \omega t)}$$

$$u_{2z} = (A_2 e^{-kz} + B_2 e^{-q_2 z} + C_2 e^{kz} + D_2 e^{q_2 z}) e^{(ik_x x + ik_y y + \omega t)}$$

$$u_{3z} = (A_3 e^{-kz} + B_3 e^{-q_3 z}) e^{(ik_x x + ik_y y + \omega t)}$$

with $C_1, D_1, A_2, B_2, C_2, D_2, A_3$ and B_3 **8 unknown** coefficients and $q_i = \sqrt{k^2 + \omega \rho_i / \mu_i}$ the **modified** wavenumber in phase i .

- ▶ 4 equations at each interface \rightarrow **8** equations \rightarrow **closed** problem

Dispersion relation: solution of $M \cdot (C_1, D_1, A_2, B_2, C_2, D_2, A_3, B_3)^T = 0$

where M is a 8×8 matrix dependant of 9 dimensionless numbers (Π theorem: 12 physical parameters - 3 fundamental units).

2 - Solutions

Non-trivial solutions are solutions of

$$|M| = 0$$

There is one **analytical** solution, stable $\forall k$:

$$\Omega_1 = \frac{\omega_1}{\sqrt{gk}} = \sqrt{\frac{g}{k^3} \frac{\rho_i}{\mu_i}} = \boxed{-Re_2^{-1}}$$

And 3 or 4 other solutions, depending on k , determined **numerically**.

Limit case: disregarding the viscosity and the surface tension
→ Mikaelian relation


$$a(At_{12}, At_{23}, H) \Omega^4 + b(At_{12}, At_{23}, H) \Omega^2 + c(At_{12}, At_{23}, H) = 0$$

with $At_{ij+1} = (\rho_{i+1} - \rho_i) / (\rho_{i+1} + \rho_i)$ the Atwood numbers and $H = kh$ the dimensionless thickness → 4 **analytical** solutions $\forall k$.

3 - Operating conditions

Let us consider the 2 cases studied in Adkins et al.

- ▶ (1) \equiv Oil / (2) \equiv Water / (3) \equiv Oil
- ▶ (1) \equiv Oil / (2) \equiv Water / (3) \equiv Air




Variable quantities: h and $k \rightarrow$ various H .

3 - 2 limit cases for comparison

- 2 separate **2-layer** fluid systems (considering the fluid viscosities)

(3)/(2)/(1) is compared to (2)/(1) and (3)/(2)

- ▶ (2)/(1) is gravitationally **unstable**.
- ▶ (3)/(2) is gravitationally **stable**.




- 3-layer **inviscid** model of Mikaelian

3 - Solutions for 2 separate 2-layer fluid systems

g^{\dagger} -unstable

vs

g^{\dagger} -stable




k_c : cutoff wavenumber due to surface tension.

$\Re(w) > 0 \rightarrow$ unstable solution; $\Im(w) \neq 0 \rightarrow$ oscillations.

3 - All solutions on the same graph!

$k < k_c$: 1 unstable + 2 stables


$k > k_c$: 4 stables



3 - Solutions for the 3-layer fluid system, $h=1$ mm


3 totally **stable** solutions: **1 analytically**, **2 numerically** determined

Other solutions: 2 **stable** for $k > k_c$, 1 **unstable** for $k < k_c$.



3 - Effect of h on the unstable solution


ω^* : maximum growth rate for the **viscous 2-layer** fluid model



Decreasing h (i.e. increasing interface coupling) decreases ω .

3 - Comparison to experimental values

ω^* : maximum growth rate for the **inviscid 3-layer** fluid model



The viscous model improves the prediction as h is reduced.


4 - Conclusion & Ongoing work

Conclusion

- ▶ Development of a theory for 3-layer RTI that considers the viscosity of the fluids and all the solutions of the dispersion relation.
- ▶ Improvement of the theoretical prediction.

Ongoing work

- ▶ Comparison of interface deformations with numerical simulations.



OPENFOAM simulations. (1) \equiv Huile/(2) \equiv Eau/(3) \equiv Mercure.
Set of images versus time.

- ▶ Inclusion of the magnetic field for a viscous model for liquid metal batteries (\vec{g} -stable stratification).

Future work \rightarrow New experiments at Case Western Reserve University?

Thank you for your attention.

Any questions?