

Supplementary Material

Logarithmic Mathematical Morphology: theory and applications

Guillaume Noyel



-
- *G. Noyel was with the University Jean Monnet, Hubert Curien Laboratory, Saint-Etienne, France.
E-mail: guillaume.noyel@univ-st-etienne.fr*

Supplementary Material

Logarithmic Mathematical Morphology: theory and applications

APPENDIX

PROOFS OF PROPOSITIONS 1 (P. 4) AND 2 (P. 4)

In this section, the proofs of propositions 1 and 2 are given. They were previously demonstrated in [18].

Proof of proposition 1: δ_b^Δ is a dilation (def. 1) because as the addition Δ preserves the order \leq [16], we have $\forall f, g \in \overline{\mathcal{F}}_M, \forall x \in D$,

$$\begin{aligned} \delta_b^\Delta(f \vee g)(x) &= \vee_{h \in D} \{[(f \vee g)(x - h)] \Delta b(h)\} \\ &= \vee_{h \in D} \{[f(x - h) \Delta b(h)] \vee [g(x - h) \Delta b(h)]\} \\ &= [\vee_{h \in D} \{f(x - h) \Delta b(h)\}] \vee [\vee_{h \in D} \{g(x - h) \Delta b(h)\}] \\ &= \delta_b^\Delta(f)(x) \vee \delta_b^\Delta(g)(x). \end{aligned}$$

For the least element O of $\overline{\mathcal{F}}_M$, we have $\forall x \in D$:

$$\begin{aligned} \delta_b^\Delta(O)(x) &= \delta_b^\Delta(f_{-\infty})(x) = \vee_{h \in D} \{-\infty(x - h) \Delta b(h)\} \\ &= \vee_{h \in D} \{-\infty(1 - b(h)/M) + b(h)\} = -\infty \\ &= O(x). \end{aligned}$$

Similarly, ε_b^Δ is an erosion because we have $\forall f, g \in \overline{\mathcal{F}}_M, \varepsilon_b^\Delta(f \wedge g) = \varepsilon_b^\Delta(f) \wedge \varepsilon_b^\Delta(g)$ and $\varepsilon_b^\Delta(I) = \varepsilon_b^\Delta(f_M) = M = I$. Finally, the pair $(\varepsilon_b^\Delta, \delta_b^\Delta)$ is an adjunction because, $\forall f, g \in \overline{\mathcal{F}}_M$, we have:

$$\begin{aligned} \delta_b^\Delta(f) \leq g &\Leftrightarrow \forall x \in D, \vee_{h \in D} \{f(x - h) \Delta b(h)\} \leq g(x) \\ &\Leftrightarrow \forall x, h \in D, f(x - h) \Delta b(h) \leq g(x) \\ &\Leftrightarrow \forall y, h \in D, f(y) \leq g(y + h) \Delta b(h) \\ &\Leftrightarrow \forall y \in D, f(y) \leq \wedge_{h \in D} \{g(y + h) \Delta b(h)\} \\ &\Leftrightarrow f \leq \varepsilon_b^\Delta(g). \end{aligned}$$

□

Proof of proposition 2: The logarithmic-erosion ε_b^Δ and dilation δ_b^Δ are dual by their *negative function* because for $b \in \overline{\mathcal{F}}_M$ we have, $\forall f, g \in \overline{\mathcal{F}}_M, \forall x \in D$,

$$\begin{aligned} (\delta_b^\Delta(f^*))^*(x) &= \Delta(\vee_{h \in D} \{\Delta f(x - h) \Delta b(h)\}) \\ &= \wedge_{h \in D} \{f(x - h) \Delta b(h)\} \\ &= \wedge_{h \in D} \{f(x + h) \Delta \bar{b}(h)\} \\ &= \varepsilon_b^\Delta(f)(x). \end{aligned}$$

Similarly, we have $(\varepsilon_b^\Delta(f^*))^* = \delta_b^\Delta(f)$. □

PROOF OF PROPOSITION 3 (P. 5)

In this section, the proof of proposition 3 is given. It was previously demonstrated in [18]. The dilation δ_b and the erosion ε_b are mappings of the lattice $\overline{\mathbb{R}}^D$, whereas the logarithmic-dilation δ_b^Δ and erosion ε_b^Δ are mappings of the lattice $\overline{\mathcal{F}}_M$. In order to link these operations, a bijective mapping (i.e. an isomorphism) is needed between these two lattices. Such an isomorphism $\xi : \overline{\mathcal{F}}_M \rightarrow \overline{\mathbb{R}}^D$ and its inverse $\xi^{-1} : \overline{\mathbb{R}}^D \rightarrow \overline{\mathcal{F}}_M$ were both defined in [44] by $\xi(f) = -M \ln(1 - f/M)$ and $\xi^{-1}(f) = M(1 - \exp(-f/M))$.

We also need to demonstrate the following lemma.

Lemma 1. *Let $f, g \in \overline{\mathcal{F}}_M$ be two functions. The isomorphism ξ transforms the LIP-sum Δ and the LIP-difference \triangle between functions into the usual sum $+$ and difference $-$, respectively. We have*

$$\xi(f \Delta g) = \xi(f) + \xi(g), \tag{A.1}$$

$$\xi(f \triangle g) = \xi(f) - \xi(g). \tag{A.2}$$

Proof of lemma 1: Let $f, g \in \overline{\mathcal{F}}_M$ be two functions. There is:

$$\begin{aligned}
\xi(f \triangle g) &= -M \ln(1 - (f \triangle g)/M) \\
&= -M \ln((1 - f/M)(1 - g/M)) \\
&= -M \ln(1 - f/M) - M \ln(1 - g/M) \\
&= \xi(f) + \xi(g), \\
\xi(\triangle g) &= -M \ln(1 + g/(M - g)) = -M \ln(M/(M - g)) \\
&= M \ln(1 - g/M) = -\xi(g), \\
\xi(f \triangle g) &= \xi(f \triangle (\triangle g)) = \xi(f) + \xi(\triangle g) = \xi(f) - \xi(g).
\end{aligned}$$

□

We can now establish the proof of proposition 3.

Proof of proposition 3: Let f be a function and b a structuring function of $\overline{\mathcal{F}}_M$. Let $\hat{f} : D \rightarrow \overline{\mathbb{R}}$ be a function equal to $\hat{f} = -\ln(1 - f/M)$. As increasing bijections, the isomorphisms ξ and ξ^{-1} distribute over infima and suprema. The dilation δ_b^{\triangle} can therefore be expressed for all $x \in D$ by:

$$\begin{aligned}
\delta_b^{\triangle}(f)(x) &= \xi^{-1} \circ \xi[\vee_{h \in D} \{f(x-h) \triangle b(h)\}] \\
&= \xi^{-1} [\vee_{h \in D} \{\xi[f(x-h) \triangle b(h)]\}] \\
&= \xi^{-1} [\vee_{h \in D} \{\xi[f(x-h)] + \xi[b(h)]\}] \\
&= \xi^{-1} [\delta_{\xi(b)}[\xi(f)]]
\end{aligned} \tag{A.3}$$

From (A.3), we have:

$$\begin{aligned}
\delta_b^{\triangle}(f)(x) &= M \left[1 - e^{\frac{-1}{M} \vee_{h \in D} \{-M \ln[1 - \frac{f(x-h)}{M}] - M \ln[1 - \frac{b(h)}{M}]\}} \right] \\
&= M \left[1 - e^{\frac{-1}{M} \vee_{h \in D} \{M(-\ln[1 - \frac{f(x-h)}{M}] - \ln[1 - \frac{b(h)}{M}])\}} \right] \\
&= M \left[1 - e^{\frac{-M}{M} \vee_{h \in D} \{-\ln[1 - \frac{f(x-h)}{M}] - \ln[1 - \frac{b(h)}{M}]\}} \right] \\
&= M \left[1 - e^{-\vee_{h \in D} \{f(x-h) + b(h)\}} \right] \\
&= M \left[1 - e^{-\delta_b^{\triangle}(f)(x)} \right].
\end{aligned}$$

Similarly, we have

$$\varepsilon_b^{\triangle}(f) = \xi^{-1} [\varepsilon_{\xi(b)}[\xi(f)]] = M[1 - e^{-\varepsilon_b^{\triangle}(f)}].$$

This ends the proof. □

PROOF OF PROPOSITION 4 (P. 6)

Proof of proposition 4: From (15), we have $\forall x \in D$:

$$\begin{aligned}
c_{1_b}(f)(x) &= \vee \{f(x+h) \triangle b(h), h \in D_b\} \\
&= \vee \{f(x-h) \triangle b(-h), -h \in D_b\} \\
&= \vee \{f(x-h) \triangle (\triangle \bar{b}(h)), h \in D_{\bar{b}}\} \\
&= \delta_{\triangle \bar{b}}^{\triangle}(f)(x).
\end{aligned}$$

By comparing (8) with (16), we have for all $x \in D$:

$$c_{2_b}(f)(x) = \varepsilon_b^{\triangle}(f)(x).$$

From (14), we deduce that $Asp_b^{\triangle}(f) = \delta_{\triangle \bar{b}}^{\triangle}(f) \triangle \varepsilon_b^{\triangle}(f)$. □

MAP OF ASPLUND DISTANCES: LINK BETWEEN DOUBLE PROBING AND LOGARITHMIC-OPENING AND CLOSING

Let us study the location of the supremum or infimum of the translated probes $b \in \overline{\mathcal{F}}_M$ such that they are in contact with the function f from below or from above, respectively. In a similar way to that used by Heijmans et al. in [1], the mapping $\tau_{h,v}$ of horizontal and vertical translations on $\overline{\mathcal{F}}_M$ is defined by $(\tau_{h,v}(f))(x) = f(x-h) \triangle v$, where $h \in D$ and $v \in]-\infty, M[$. For the lower probes, the supremum of the translated probes b which are in contact with the function f from below is the supremum of the translated probes $\tau_{h,v}(b)$ which are less or equal to the function f . It is equal to $\Gamma_b^{\triangle}(f) = \sup \{\tau_{h,v}(b) \mid h \in D, v \in]-\infty, M[, \tau_{h,v}(b) \leq f\}$. As demonstrated in [2], $\Gamma_b^{\triangle}(f)$ is an opening.




Fig. 1. (a) An image f is analysed by a probe b from above and below. (b) The $mlub$ $c_{1_b}(f)$, the $mglb$ $c_{2_b}(f)$ and the map of Asplund distances $Asp_b^\Delta(f)$ between the image and the probe. (c) The supremum of the probe b such that the probe is in contact with the function f from below is the logarithmic-opening γ_b^Δ of the function f . (d) The infimum of the probe b such that the probe is in contact with the function f from above is the logarithmic-closing $\varphi_{\Delta\bar{b}}^\Delta(f)$ of the function f by the structuring function $\Delta\bar{b}$.

Similarly, for the upper probes, the infimum of the translated probes b which are in contact with the function f from above is equal to $\Phi_b^\Delta(f) = \inf \{ \tau_{h,v}(b) \mid h \in D, v \in]-\infty, M[, \tau_{h,v}(b) \geq f \}$. As demonstrated in [2], $\Phi_b^\Delta(f)$ is a closing.

Proposition A.1. *The supremum of the translated probes b which are in contact with the function f from below, $\Gamma_b^\Delta(f)$, is equal to the logarithmic-opening of f , $\gamma_b^\Delta(f)$:*

$$\Gamma_b^\Delta(f)(x) = \gamma_b^\Delta(f)(x). \quad (\text{A.4})$$

The infimum of the translated probes b which are in contact with the function f from above, $\Phi_b^\Delta(f)$, is equal to the logarithmic-closing of f , $\varphi_{\Delta\bar{b}}^\Delta(f)$:

$$\Phi_b^\Delta(f)(x) = \varphi_{\Delta\bar{b}}^\Delta(f)(x). \quad (\text{A.5})$$

Fig. 1c and 1d illustrate the supremum or infimum of the translated probes b which are in contact with the function f from below or from above, respectively. In Fig. 1c, one can notice that the opening $\gamma_b^\Delta(f)$ of f is very close to the image f . In Fig. 1d, the closing $\varphi_{\Delta\bar{b}}^\Delta(f)$ of f is similar to the image f with the exception of both valleys of f , where the structuring function f is very dissimilar to f .

Proof of proposition A.1:

Let f and b be a function and a probe of $\overline{\mathcal{F}}_M$, respectively. For $h \in D$ and $v \in]-\infty, M[$, the mapping $\tau_{h,v}$ of horizontal and vertical translations on $\overline{\mathcal{F}}_M$ is defined by $(\tau_{h,v}(f))(x) = f(x-h) \Delta v$. The supremum of the translated probes b which are in contact with the function f from below is equal to:

$$\begin{aligned} \Gamma_b^\Delta(f)(x) &= \sup \{ \tau_{h,v}(b)(x) \mid h \in D, v \in]-\infty, M[, \tau_{h,v}(b) \leq f \} \\ &= \sup \{ b(x-h) \Delta v \mid h \in D, v \in]-\infty, M[, x' \in D, b(x'-h) \Delta v \leq f(x') \} \\ &= \vee_{h \in D} \{ b(x-h) \Delta v(h) \mid v(h) = \sup_{x' \in D} \{ v, b(x'-h) \Delta v \leq f(x') \} \} \\ &= \vee_{h \in D} \{ b(x-h) \Delta v(h) \mid v(h) = \sup_{x' \in D} \{ v, b(x') \Delta v \leq f(h+x') \} \} \\ &= \vee_{h \in D} \{ b(x-h) \Delta c_{2_b} f(h) \} \quad (\text{from (16)}) \\ &= \vee_{h \in D} \{ b(x-h) \Delta \varepsilon_b^\Delta(f)(h) \} \quad (\text{from Prop. 4}) \\ &= \vee_{h \in D} \{ \varepsilon_b^\Delta(f)(h) \Delta b(x-h) \} \\ &= \vee_{h \in D} \{ \varepsilon_b^\Delta(f)(x-h) \Delta b(h) \} \\ &= \delta_b^\Delta \varepsilon_b^\Delta(f)(x) \quad (\text{from (7)}) \\ &= \gamma_b^\Delta(f)(x). \quad (\text{from (9)}) \end{aligned}$$

Such a relation is an opening one.

The infimum of the translated probes b which are in contact with the function f from above is equal to:

$$\begin{aligned}
\Phi_b^{\Delta}(f)(x) &= \inf \{ \tau_{h,v}(b)(x) \mid h \in D, v \in]-\infty, M[, \tau_{h,v}(b) \geq f \} \\
&= \inf \{ b(x-h) \Delta v \mid h \in D, v \in]-\infty, M[, x' \in D, b(x'-h) \Delta v \geq f(x') \} \\
&= \wedge_{h \in D_b} \{ b(x-h) \Delta v(h) \mid v(h) = \inf_{x' \in D_b} \{ v, b(x'-h) \Delta v \geq f(x') \} \} \\
&= \wedge_{h \in D_b} \{ b(x-h) \Delta v(h) \mid v(h) = \inf_{x' \in D_b} \{ v, b(h) \Delta v \geq f(x'+h) \} \} \\
&= \wedge_{h \in D_b} \{ b(x-h) \Delta c_{1_b} f(h) \} \quad (\text{from (15)}) \\
&= \wedge_{h \in D_b} \{ b(x-h) \Delta \delta_{\Delta \bar{b}}^{\Delta}(f)(h) \} \quad (\text{from Prop. 4}) \\
&= \wedge_{h \in D_b} \{ \delta_{\Delta \bar{b}}^{\Delta}(f)(h) \Delta b(x-h) \} \\
&= \wedge_{h \in D_b} \{ \delta_{\Delta \bar{b}}^{\Delta}(f)(x-h) \Delta b(h) \} \\
&= \wedge_{-h \in D_b} \{ \delta_{\Delta \bar{b}}^{\Delta}(f)(x+h) \Delta b(-h) \} \\
&= \wedge_{h \in D_b} \{ \delta_{\Delta \bar{b}}^{\Delta}(f)(x+h) \Delta (\Delta \bar{b}(h)) \} \\
&= \varepsilon_{\Delta \bar{b}}^{\Delta} \delta_{\Delta \bar{b}}^{\Delta}(f)(x) \quad (\text{from (8)}) \\
&= \varphi_{\Delta \bar{b}}^{\Delta}(f)(x) \quad (\text{from (10)}).
\end{aligned}$$

Such a relation is a closing one. □

PROOFS OF EQUATIONS (20) AND (21) (P. 7)

Proof of equation 20: Let B_0 a flat structuring element with the same support D_{b_0} as the symmetric and constant structuring element b_0 . $b_0 : D \rightarrow \mathbb{R}$ is defined for all $x \in D_{b_0}$, where $D_{b_0} \subset D$, by $b_0(x) = b_0$ and $b_0(-x) = b_0(x)$. For all $x \in D$, we have:

$$\begin{aligned}
Asp_{b_0}^{\Delta}(f)(x) &= \delta_{\Delta \bar{b}_0}^{\Delta}(f)(x) \Delta \varepsilon_{b_0}^{\Delta}(f)(x) \\
&= \vee_{h \in D_{\bar{b}_0}} \{ f(x-h) \Delta (\Delta \bar{b}_0) \} \Delta \wedge_{h \in D_{b_0}} \{ f(x+h) \Delta b_0 \} \\
&= \left(\vee_{h \in D_{b_0}} \{ f(x-h) \} \Delta b_0 \right) \Delta \left(\wedge_{h \in D_{b_0}} \{ f(x+h) \} \Delta b_0 \right) \\
&= \vee_{h \in D_{b_0}} \{ f(x-h) \} \Delta \wedge_{h \in D_{b_0}} \{ f(x+h) \} \\
&= \delta_{B_0}(f)(x) \Delta \varepsilon_{B_0}(f)(x) \\
&= \varrho_{B_0}^{LIP}(f)(x).
\end{aligned}$$

□

Proof of equation 21: Let $Y \subset D$ be a constant (i.e. flat) zone of a function f and let $X = Y \ominus D_b$ be the eroded flat zone by the domain D_b of the structuring function b . \ominus represents the binary erosion [3], [4]. We have, for all $x \in X$:

$$\begin{aligned}
Asp_b^{\Delta}(f)(x) &= c_{1_b} f(x) \Delta c_{2_b} f(x) \\
&= \vee_{h \in D_b} \{ f(x+h) \Delta b(h) \} \Delta \wedge_{h \in D_b} \{ f(x+h) \Delta b(h) \} \\
&= (f(x) \Delta \vee_{h \in D_b} \{ b(h) \}) \Delta (f(x) \Delta \wedge_{h \in D_b} \{ b(h) \}) \\
&= \Delta \vee_{h \in D_b} \{ b(h) \} \Delta \wedge_{h \in D_b} \{ b(h) \} \\
&= \wedge_{h \in D_b} \{ b(h) \} \Delta \vee_{h \in D_b} \{ b(h) \} \\
&= b_{sup} \Delta b_{inf}.
\end{aligned}$$

b_{sup} and b_{inf} are the supremum and the infimum, respectively, of the structuring element b . In the eroded flat zone X , the Asplund metric is therefore equal to the constant $b_{sup} \Delta b_{inf}$. □

PROOFS OF PROPERTIES 1 (P. 9) AND 2 (P. 9)

Proof of property 1: Let $f \in \overline{\mathcal{F}}_M$ be a function and $b^l, b^r \in \overline{\mathcal{F}}_M$ be structuring functions. From (8), we have $\forall x \in D$, $\wedge_{h \in D_b} \{ f(x+h) \Delta b^l(h) \} = \varepsilon_{b^l}^{\Delta}(f)(x)$ and from (19), $c_{2_b}(f) = \varepsilon_b^{\Delta}(f)$. From (25), we deduce that:

$$\begin{aligned}
E(b^l, f)(x) &= \wedge_{h \in D_b} \{ f(x+h) \Delta b^l(h) \} \Delta c_{2_b}(f)(x) \\
&= \varepsilon_{b^l}^{\Delta}(f)(x) \Delta \varepsilon_b^{\Delta}(f)(x).
\end{aligned}$$

In a similar way, we have $E(b^r, f)(x) = \varepsilon_{b^r}^{\Delta}(f)(x) \Delta \varepsilon_b^{\Delta}(f)(x)$. □

Proof of property 2: From (8), we have $\forall c \in]-\infty, M[$ and $\forall x \in D$:

$$\begin{aligned}
\varepsilon_b^\Delta(f \Delta c)(x) &= \wedge_{h \in D_b} \{(f \Delta c)(x+h) \Delta b(h)\} \\
&= \wedge_{h \in D_b} \{f(x+h) \Delta c \Delta b(h)\} \\
&= \wedge_{h \in D_b} \{f(x+h) \Delta b(h)\} \Delta c \\
&= \varepsilon_b^\Delta(f)(x) \Delta c.
\end{aligned} \tag{A.6}$$

From (28), one deduces that:

$$\begin{aligned}
E(b^l, f \Delta c) &= \varepsilon_{b^l}^\Delta(f \Delta c) \Delta \varepsilon_b^\Delta(f \Delta c) \\
&= (\varepsilon_{b^l}^\Delta(f) \Delta c) \Delta (\varepsilon_b^\Delta(f) \Delta c) \\
&= \varepsilon_{b^l}^\Delta(f) \Delta \varepsilon_b^\Delta(f) \Delta c \Delta c \\
&= \varepsilon_{b^l}^\Delta(f) \Delta \varepsilon_b^\Delta(f) \\
&= E(b^l, f).
\end{aligned}$$

In a similar way, we obtained $E(b^r, f \Delta c) = E(b^r, f)$. As a result, from (27), we have $E(b, f \Delta c) = E(b, f)$. \square

PROOF OF PROPERTY 3 (P. 10)

Proof of property 3: Let $f \in \overline{\mathcal{F}}_M$ be a function and $b, b_r \in \overline{\mathcal{F}}_M$ be structuring functions. Similarly as in (A.6), we have $\forall c \in]-\infty, M[$ and $\forall x \in D$:

$$\begin{aligned}
\delta_b^\Delta(f \Delta c)(x) &= \vee_{h \in D_b} \{(f \Delta c)(x-h) \Delta b(h)\} \\
&= \vee_{h \in D_b} \{f(x-h) \Delta c \Delta b(h)\} \\
&= \vee_{h \in D_b} \{f(x-h) \Delta b(h)\} \Delta c \\
&= \delta_b^\Delta(f)(x) \Delta c.
\end{aligned} \tag{A.7}$$

From (9), (A.6) and (A.7), we have $\forall c \in]-\infty, M[$:

$$\begin{aligned}
\gamma_b^\Delta(f \Delta c) &= \delta_b^\Delta \left[\varepsilon_b^\Delta(f \Delta c) \right] \\
&= \delta_b^\Delta \left[\varepsilon_b^\Delta(f) \Delta c \right] \\
&= \delta_b^\Delta \left[\varepsilon_b^\Delta(f) \right] \Delta c \\
&= \gamma_b^\Delta(f) \Delta c.
\end{aligned} \tag{A.8}$$

From (30), one deduces $\forall c \in]-\infty, M[$ that:

$$\begin{aligned}
G_b^\Delta(f \Delta c) &= \gamma_b^\Delta(f \Delta c) \Delta \gamma_{b_r}^\Delta(f \Delta c) \\
&= \gamma_b^\Delta(f) \Delta c \Delta \left[\gamma_{b_r}^\Delta(f) \Delta c \right] \\
&= \gamma_b^\Delta(f) \Delta c \Delta \gamma_{b_r}^\Delta(f) \Delta c \\
&= \gamma_b^\Delta(f) \Delta \gamma_{b_r}^\Delta(f) \\
&= G_b^\Delta(f).
\end{aligned}$$

\square

PROOF OF PROPERTY 4 (P. 11)

Proof of property 4: Let $f \in \overline{\mathcal{F}}_M$ be a function and $b, b_r \in \overline{\mathcal{F}}_M$ be structuring functions. From (32) and (A.8), we have $\forall c \in]-\infty, M[$:

$$\begin{aligned}
R_b^\Delta(f \Delta c) &= [f \Delta c] \Delta \gamma_b^\Delta(f \Delta c) \\
&= f \Delta c \Delta \left[\gamma_b^\Delta(f) \Delta c \right] \\
&= f \Delta c \Delta \gamma_b^\Delta(f) \Delta c \\
&= f \Delta \gamma_b^\Delta(f) \\
&= R_b^\Delta(f).
\end{aligned}$$

\square

PROOF OF PROPOSITION 5 (P. 11)

Proof of proposition 5: First of all, let us remind the definition of the LIP-additive Asplund metric with tolerance which was introduced in [5].

Definition A.1 (LIP-additive Asplund metric with tolerance). *Let $(1 - p)$ be a percentage of points of D to be discarded and D' the set of these discarded points. The LIP-additive Asplund metric with tolerance between two functions f and $g \in \mathcal{F}_M$ is defined by:*

$$d_{asp,p}^{\Delta}(f, g) = c_{1,p} \Delta c_{2,p}. \quad (\text{A.9})$$

The constants $c_{1,p}$ and $c_{2,p}$ are equal to:

$$c_{1,p} = \inf\{c, \forall x \in D, \gamma_{(f|_{D \setminus D'}, g|_{D \setminus D'})}^{\Delta}(x) \leq c\},$$

$$c_{2,p} = \sup\{c, \forall x \in D, c \leq \gamma_{(f|_{D \setminus D'}, g|_{D \setminus D'})}^{\Delta}(x)\}.$$

A percentage $(1 - p)/2$ of the points $x \in D$ with the greatest, respectively lowest contrast values $\gamma_{(f,g)}^{\Delta}(x) = f(x) \Delta g(x)$ are discarded.

Remark A.1. In practice, for a function $f : D \rightarrow]\infty, M[$ defined on a discrete grid D , e.g. $D \subset \mathbb{Z}^n$, the number of points to be suppressed are selected as follows. Let $\#D$ be the cardinal of D . The number of points n_{suppr} to be suppressed is equal to $n_{suppr} = \text{round}[(1 - p)\#D]$, where *round* is the rounding operator of any real number to its nearest integer. For the constant $c_{1,p}$ and $c_{2,p}$, the number of points to be suppressed are respectively equal to:

$$n_1 = \text{round}(n_{suppr}/2), \quad (\text{A.10})$$

$$n_2 = n_{suppr} - n_1, \quad (\text{A.11})$$

Let us now recall the rank filter definitions. Let $\zeta_{b,k} : \mathbb{R}^D \rightarrow \mathbb{R}^D$ be the erosion-rank filter of rank k by the structuring function b . Let $\vartheta_{b,k}$ be the dilation-rank filter of rank k by the structuring function b . Both operations are defined by:

$$\zeta_{b,k}(f)(x) = \bigwedge^k \{f(x+h) - b(h), h \in D_b\}, \quad (\text{A.12})$$

$$\vartheta_{b,k}(f)(x) = \bigvee^k \{f(x-h) + b(h), h \in D_b\}. \quad (\text{A.13})$$

The LIP-erosion-rank filter $\zeta_{b,k}^{\Delta} : \overline{\mathcal{F}}_M^D \rightarrow \overline{\mathcal{F}}_M^D$ and the LIP-dilation-rank filter $\vartheta_{b,k}^{\Delta}$ are defined as follows:

$$\zeta_{b,k}^{\Delta}(f)(x) = \bigwedge^k \{f(x+h) \Delta b(h), h \in D_b\}, \quad (\text{A.14})$$

$$\vartheta_{b,k}^{\Delta}(f)(x) = \bigvee^k \{f(x-h) \Delta b(h), h \in D_b\}. \quad (\text{A.15})$$

The map of LIP-additive Asplund distances with tolerance (to noise extrema) is defined as follows.

Definition A.2 (Map of LIP-additive Asplund distances with tolerance [5]). *Let $f \in \mathcal{F}_M$ be a grey-level image and $b : D_b \rightarrow]-\infty, M[$ a probe. Let $(1 - p)$ be a percentage of points of D_b to be discarded. The map of Asplund distances $Asp_b^{\Delta} : \mathcal{F}_M \rightarrow \mathcal{F}_M$ with a tolerance (to extrema) p is defined by:*

$$Asp_b^{\Delta}(f)(x) = d_{asp,p}^{\Delta}(f|_{D_b(x)}, b). \quad (\text{A.16})$$

For each point $x \in D$, the distance $d_{asp}^{\Delta}(f|_{D_b(x)}, b)$ is computed in the neighbourhood $D_b(x)$ centred in x and the template b is acting like a structuring function. $f|_{D_b(x)}$ is the restriction of f to $D_b(x)$. $Asp_b^{\Delta}(f) : D \rightarrow \mathcal{F}_M$ is the map of Asplund distances between the image f and the probe b .

In order to establish the link with MM, we will express the map of Asplund distances with neighbourhood operations. From Eq. A.9, for each $x \in D$, the map expression becomes

$$d_{asp,p}^{\Delta}(f|_{D_b(x)}, b) = c_{1,p}(f)(x) \Delta c_{2,p}(f)(x),$$

where

$$c_{1,p}(f)(x) = \inf\{c, \forall h \in D_b, \gamma_{(f|_{D_b(x) \setminus D'_b(x)}, b|_{D_b \setminus D'_b(x)})}^{\Delta}(h) \leq c\}$$

$$c_{2,p}(f)(x) = \sup\{c, \forall h \in D_b, c \leq \gamma_{(f|_{D_b(x) \setminus D'_b(x)}, b|_{D_b \setminus D'_b(x)})}^{\Delta}(h)\}.$$

This leads to the following definition.

Definition A.3 (LIP-additive maps of the least upper and of the greatest lower bounds with tolerance). *Let $f \in \overline{\mathcal{F}}_M$ be a function and $b : D_b \rightarrow]\infty, M[$ a probe. For any $x \in D$, let $(1 - p)$ be a percentage of points of the neighbourhood $D_b(x)$ to be*

discarded and $D'_b(x)$ the set of these discarded points. The map of the least upper bounds (mlub) $c_{1b,p} : \overline{\mathcal{F}}_M \rightarrow \overline{\mathcal{F}}_M$ and the map of the greatest lower bounds (mglb) $c_{2b,p} : \overline{\mathcal{F}}_M \rightarrow \overline{\mathcal{F}}_M$, of f by b are defined, for any $x \in D$, by:

$$c_{1b,p}(f)(x) = \inf \left\{ c, \forall h \in D_b, \quad \gamma_{(f|_{D_b(x) \setminus D'_b(x)}, b|_{D_b \setminus D'_b(x)})}^\Delta(h) \leq c \right\}, \quad (\text{A.17})$$

$$c_{2b,p}(f)(x) = \sup \left\{ c, \forall h \in D_b, \quad c \leq \gamma_{(f|_{D_b(x) \setminus D'_b(x)}, b|_{D_b \setminus D'_b(x)})}^\Delta(h) \right\}. \quad (\text{A.18})$$

The number of points to be suppressed n_1 and n_2 for the mlub $c_{1b,p}$ and for the mglb $c_{2b,p}$ are equal to $n_1 = \text{round}(n_{\text{suppr}}/2)$, and $n_2 = n_{\text{suppr}} - n_1$, respectively, where $n_{\text{suppr}} = \text{round}[(1-p)\#D_b]$ and D_b is the cardinal of $\#D_b$.

From (A.17), one deduces, for any $x \in D$, that:

$$\begin{aligned} c_{1b,p}(f)(x) &= \bigvee \left\{ \gamma_{(f|_{D_b(x) \setminus D'_b(x)}, b|_{D_b \setminus D'_b(x)})}^\Delta(h), h \in D_b \right\} \\ &= \bigvee^{n_1} \left\{ \gamma_{(f|_{D_b(x)}, b)}^\Delta(h), h \in D_b \right\} \\ &= \bigvee^{n_1} \{ f(x+h) \triangle b(h), h \in D_b \} \\ &= \bigvee^{n_1} \{ f(x-h) \triangle b(-h), -h \in D_b \} \\ &= \bigvee^{n_1} \{ f(x-h) \triangle (\triangle \bar{b}(h)), h \in D_{\bar{b}} \} \\ &= \vartheta_{\triangle \bar{b}, n_1}^\Delta(f)(x) \quad (\text{from (A.13)}). \end{aligned}$$

Similarly, from (A.18), one deduces, for any $x \in D$, that:

$$\begin{aligned} c_{2b,p}(f)(x) &= \bigwedge \left\{ \gamma_{(f|_{D_b(x) \setminus D'_b(x)}, b|_{D_b \setminus D'_b(x)})}^\Delta(h), h \in D_b \right\} \\ &= \bigwedge^{n_2} \{ f(x+h) \triangle b(h), h \in D_b \} \\ &= \zeta_{b, n_2}^\Delta(f)(x) \quad (\text{from (A.12)}). \end{aligned}$$

As a consequence, the following proposition holds.

Proposition A.2 (i.e. proposition 5 (p. 8)). *Let $f \in \overline{\mathcal{F}}_M$ be a function defined on a discrete grid, e.g. $D \subset \mathbb{Z}^n$. Let $b \in \overline{\mathcal{F}}_M$ be a structuring function, where for all $x \in D_b$, $D_b \subset D$, $b(x) > -\infty$. Let $(1-p)$ be a percentage of points of D_b to be discarded. The map of LIP-additive Asplund distances with a tolerance p between the function f and the structuring function b is equal to:*

$$\text{Asp}_{b,p}^\Delta(f) = \vartheta_{\triangle \bar{b}, n_1}^\Delta(f) \triangle \zeta_{b, n_2}^\Delta(f). \quad (\text{A.19})$$

The number of points to be suppressed n_1 and n_2 for the mlub $\vartheta_{\triangle \bar{b}, n_1}^\Delta$ and for the mglb ζ_{b, n_2}^Δ are equal to $n_1 = \text{round}(n_{\text{suppr}}/2)$, and $n_2 = n_{\text{suppr}} - n_1$, respectively, where $n_{\text{suppr}} = \text{round}[(1-p)\#D_b]$ and $\#D_b$ is the cardinal of D_b . For the mlub and the mglb of f , $c_{1b,p}(f)$ and $c_{2b,p}(f)$, we have:

$$c_{1b,p}(f) = \vartheta_{\triangle \bar{b}, n_1}^\Delta(f), \quad (\text{A.20})$$

$$c_{2b,p}(f) = \zeta_{b, n_2}^\Delta(f). \quad (\text{A.21})$$

□

PROOF OF PROPERTY 5 (P. 16)

Proof of property 5:

Let $f \in \overline{\mathcal{F}}_M$ be a function and $b, b_\theta, b_\theta^l, b_\theta^c, b_\theta^r \in \overline{\mathcal{F}}_M$ be structuring functions.

From (A.14), we have $\forall c \in]-\infty, M[$ and $\forall x \in D$:

$$\begin{aligned} \zeta_b^\Delta(f \triangle c)(x) &= \bigwedge_{h \in D_b}^k \{ (f \triangle c)(x+h) \triangle b(h) \} \\ &= \bigwedge_{h \in D_b}^k \{ f(x+h) \triangle c \triangle b(h) \} \\ &= \bigwedge_{h \in D_b}^k \{ f(x+h) \triangle b(h) \} \triangle c \\ &= \zeta_b^\Delta(f)(x) \triangle c. \end{aligned} \quad (\text{A.22})$$

From (A.6), (A.22) and knowing that the LIP-addition of a constant $\triangle c$ preserves the order \leq , one deduces that:

$$\begin{aligned}
\dot{c}_{b_\theta, k}(f \triangle c) &= \bigwedge \left\{ \varepsilon_{b_\theta}^{\triangle}(f \triangle c), \bigwedge [\zeta_{b_\theta, k}^{\triangle}(f \triangle c), \zeta_{b_\theta, k}^{\triangle}(f \triangle c)] \right\} \\
&= \bigwedge \left\{ \varepsilon_{b_\theta}^{\triangle}(f) \triangle c, \bigwedge [\zeta_{b_\theta, k}^{\triangle}(f) \triangle c, \zeta_{b_\theta, k}^{\triangle}(f) \triangle c] \right\} \\
&= \bigwedge \left\{ \varepsilon_{b_\theta}^{\triangle}(f) \triangle c, \bigwedge [\zeta_{b_\theta, k}^{\triangle}(f), \zeta_{b_\theta, k}^{\triangle}(f)] \triangle c \right\} \\
&= \bigwedge \left\{ \varepsilon_{b_\theta}^{\triangle}(f), \bigwedge [\zeta_{b_\theta, k}^{\triangle}(f), \zeta_{b_\theta, k}^{\triangle}(f)] \right\} \triangle c.
\end{aligned} \tag{A.23}$$

From (34), (A.22) and (A.23), we have:

$$\begin{aligned}
E^k(b_\theta^l, f \triangle c) &= \zeta_{b_\theta, k}^{\triangle}(f \triangle c) \triangle \dot{c}_{b_\theta, k}(f \triangle c) \\
&= \zeta_{b_\theta, k}^{\triangle}(f) \triangle c \triangle [\dot{c}_{b_\theta, k}(f) \triangle c] \\
&= \zeta_{b_\theta, k}^{\triangle}(f) \triangle \dot{c}_{b_\theta, k}(f) \\
&= E^k(b_\theta^l, f).
\end{aligned} \tag{A.24}$$

Similarly, from (35), we have:

$$E^k(b_\theta^r, f \triangle c) = E^k(b_\theta^r, f). \tag{A.25}$$

From (36), (A.24) and (A.25), we have:

$$\begin{aligned}
E^k(b_\theta, f \triangle c) &= \bigvee \{E^k(b_\theta^l, f \triangle c), E^k(b_\theta^r, f \triangle c)\} \\
&= \bigvee \{E^k(b_\theta^l, f), E^k(b_\theta^r, f)\}. \\
&= E^k(b_\theta, f)
\end{aligned} \tag{A.26}$$

As a consequence from (37), we have:

$$\begin{aligned}
E^k(b, f \triangle c) &= \bigwedge \{E^k(b_\theta, f \triangle c) \mid \theta \in \Theta\} \\
&= \bigwedge \{E^k(b_\theta, f) \mid \theta \in \Theta\} \\
&= E^k(b, f).
\end{aligned} \tag{A.27}$$

Finally, from (38), we have :

$$\begin{aligned}
e_b^k(f \triangle c) &= \bigwedge \{E^k(b_i, f \triangle c) \mid i \in \llbracket 1 \dots I \rrbracket\} \\
&= \bigwedge \{E^k(b_i, f) \mid i \in \llbracket 1 \dots I \rrbracket\} \\
&= e_b^k(f).
\end{aligned} \tag{A.28}$$

□

REFERENCES

- [1] H. Heijmans and C. Ronse, "The algebraic basis of mathematical morphology I. Dilations and erosions," *Comput. Vision Graphics and Image Process.*, vol. 50, no. 3, pp. 245 – 295, Jun. 1990.
- [2] C. Ronse and H. Heijmans, "The algebraic basis of mathematical morphology: II. Openings and closings," *CVGIP: Image Understanding*, vol. 54, no. 1, pp. 74 – 97, 1991.
- [3] J. Serra and N. Cressie, *Image Analysis and Mathematical Morphology*. Orlando, FL, USA: Academic, 1982, vol. 1.
- [4] J. Serra, *Image analysis and mathematical morphology: Theoretical advances*. San Diego, CA, USA: Academic, 1988, vol. 2.
- [5] G. Noyel and M. Jourlin, "Functional asplund metrics for pattern matching, robust to variable lighting conditions," *Image Analysis & Stereology*, vol. 39, no. 2, pp. 53–71, 2020.