



HAL
open science

Laser-light and interactions with particles (LIP), 2022

Fabrice R.A. Onofri, Gérard Gouesbet, Maciej Kolwas, Daniel Jakubczyk

► **To cite this version:**

Fabrice R.A. Onofri, Gérard Gouesbet, Maciej Kolwas, Daniel Jakubczyk. Laser-light and interactions with particles (LIP), 2022. Journal of Quantitative Spectroscopy and Radiative Transfer, 2023, 309, pp.108710. 10.1016/j.jqsrt.2023.108710 . hal-04193911

HAL Id: hal-04193911

<https://hal.science/hal-04193911>

Submitted on 1 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1
2
3 **Editorial**
4 **Laser-light and interactions with particles (LIP), 2022**

5 Fabrice R.A. Onofri*
6 Aix-Marseille Université, CNRS, IUSTI UMR 7343, 13 453, Marseille, France

7
8 Gérard Gouesbet
9 CORIA-UMR 6614, Normandie Université, CNRS, Université et INSA de Rouen, Campus
10 Universitaire du Madrillet, Saint-Etienne-du-Rouvray, 76800, France

11
12 Maciej Kolwas
13 Institute of Physics, Al. Lotnikow 32/46, 02-668 Warsaw, Poland

14
15 Daniel Jakubczyk
16 Institute of Physics, Al. Lotnikow 32/46, 02-668 Warsaw, Poland

17
18 *Corresponding author : fabrice.onofri@univ-amu.fr

19
20
21 **Keywords:**

22 Lorenz-Mie theory and shaped beams, Bessel functions, optical forces, plasmon-polariton
23 excitation and photonic molecules, optical wireless communication, optical particle
24 characterization methods

25
26
27 This “Laser-light and interactions with particles (LIP), 2022”, or “LIP2022”, topical issue brings
28 together papers selected after the organization, but not exclusively, of the eponymous conference
29 which took place from August 21st to 26th, 2023, in Warsaw, under the chairmanship of Professors D.
30 Jakubczyk and M. Kolwas [1]. The biennial LIP conference series is a direct descendant of the former
31 "Optical Particle Sizing (OPS), theory and practice" conference, launched and chaired by G. Gouesbet
32 and G. Gréhan in Rouen in 1987. OPS was renamed as "Optical Particle Characterization Conference
33 (OPC)" after 1990. The OPS and OPC conferences were mainly devoted to optical measurements,
34 using laser sources, of the size of individual particles and, subsequently, to the characterization of their
35 composition or temperature. In 2012, to take account of the most recent developments in the field, it
36 was decided to rename the conference series to “LIP” and broaden its topics to laser beams
37 description, including acoustical and quantum beams; mechanical effects of light for particle
38 manipulation, optical tweezer, force measurement; near-field, far-field and time-resolved scattering
39 computations and laboratory experiments; plasmonic and other resonances; scattering properties of
40 complex shaped particles and aggregates: morphology; multiple scattering in random, dense or
41 turbulent. Since the beginning applications are also welcomed in the following domains: two- and
42 multiphase flow characterization; aerosol and atmospheric sciences: nucleation, transport,
43 characterization; plasma and soft matter physics: colloids, suspensions, aggregation; biomedical
44 optical engineering: cell manipulation and properties, microrheology; remote sensing and free space
45 communication and optical particle characterization [2-5].

46
47 The present topical issue contains 16 papers that are briefly introduced in what follows. They are
48 organized in seven categories or topics: computation of complex functions related to light scattering
49 problems, description of particular structured beams in free space, scattering of nanoemitters and
50 photonic molecules, optical forces, shaped beams and their propagation in the atmosphere, evaluation
51 of asymptotic light scattering models and optical particle characterization methods.

52
53 We start with two papers by Ambrosio et al. [6, 7], dealing with the computation of complex
54 functions related to light scattering problems. Since the beginning of electromagnetic theory, the

55 numerical calculation of solutions has been a key issue. The case of the Lorenz-Mie theory (LMT),
56 whose formalism was established more than a century old [8, 9], is well known. The applicability of
57 this theory, which gives the general expressions for the scattering of a plane wave by spherical,
58 homogeneous and isotropic scatterers, became only real from the end of the 1970s, beginning of the
59 1980s. This, thanks to the spreading of personal computers, but also its partial rewriting, in terms of
60 the ratio of complex functions and their derivatives as well the development of stable numerical
61 algorithms using specific upward and downward recurrences [10-12]. This is precisely what it is
62 proposed by Ambrosio et al [6], for specific expressions involving products between (Ricatti-)Bessel
63 functions, and their derivatives, that are of interest to compute photophoresis effects on lossy
64 concentric spherical particles. This approach is extended by the same authors, in a second article [7],
65 for expressions of complex functions appearing in the calculation of the photophoretic asymmetry
66 factors of infinite cylinders. The latter, illuminated by plane waves or light-sheets, being, either
67 isolated or positioned near conducting corners or planar boundaries.

68
69 The next two papers [13, 14] deal with the description of structured beams in the frame of the
70 Generalized Lorenz-Mie theory (GLMT) [15]. Generally speaking, GLMT gives general expressions
71 for the scattering of shaped beams by spherical homogeneous or concentric particles, spheroids...,
72 with a separation variable method and the introduction of beam shape coefficients (BSCs) [15]. In the
73 first paper, Ambrosio [13] introduces an analytical approach to overcome numerical limitations in the
74 description of frozen waves, i.e. a specific class of non-diffracting and stationary localized wave field
75 whose intensity envelope remains static [16]. The two major improvements concern the determination
76 of the Fourier coefficients of compositions of generalized Legendre functions and the exact solution to
77 integrals involving Bessel functions for an on-axis configuration. The second paper deals with the
78 description of Ince-Gaussian beams (IGBs, named for Edward L. Ince's polynomials) [17] in the
79 frame of the GLMT[15]. IGBs beams are exact solutions of the paraxial wave equation, like Hermite-
80 Gaussian, Laguerre-Gaussian or hypergeometric-Gaussian beams for instance. The Hermite-Gaussian
81 and Laguerre-Gaussian modes are a special case of the IG modes that are natural solutions to the
82 paraxial wave equation (PWE) in elliptic cylindrical coordinates. In the present work, Votto et al.
83 succeed to derive with the finite series method the BSCs of IGBs beams from the BSCs Laguerre-
84 Gaussian beams, opening perspectives for a better description of frozen waves in light scattering
85 problems notably.

86
87 The beam shape also matters when dealing with long distance optical wireless communication and
88 LIDAR remote sensing, due to the diffraction of the beam, the presence of aerosol and raindrops [18],
89 as well as the turbulence of the atmosphere [19]. Dong et al. [18] consider the energy attenuation
90 characteristics of high-order Bessel vortex beams [20] under rainfall atmospheric conditions. An
91 optical vortex is a structured light wave which has an isolated point singularity. A polarization vortex
92 (vector beam) has spatially variant polarizations with a polarization singularity, whereas a phase
93 vortex has a spiral phase front with a phase singularity or screw dislocation [21]. The fact is that
94 optical vortices has recently gained increasing interest in optical particle manipulation, laser
95 machining, optical microscopy and optical communications. In the present paper, the authors show,
96 through numerical simulation using GLMT framework, that the beam extinction caused by raindrop is
97 significantly reduced when using Bessel vortex beam instead of classical Gaussian beams. The authors
98 also show that increasing the topological charge and axial cone angle of Bessel vortex laser beam
99 increases their penetrating ability in rainfall environment, like the use of near-infrared beams rather
100 than visible or mid-infrared ones. Then, Gökçe et al [19] investigate the propagation of multimode
101 beams whose wavefront is distorted by the turbulence of the atmosphere itself, limiting the quality and
102 range of optical communications [22]. The model developed [19] accounts particularly for the
103 structure constants of atmosphere, the number of modes of the laser beam, the bi-static distance, the
104 focal length and radius of the coupling lenses at the optical fiber outputs. Numerical simulations reveal
105 that increasing the number of modes enhances notably the coupling efficiency performance when
106 compared with Gaussian beams. The same hold for the source size, so has to the size and the focal
107 length of the coupling lenses, and the wavelength in a certain extent.

109 The next two papers deal with the scattering of nanoemitters [23] and photonic molecules [24]. The
110 work by G. Burlak G. and Medina-Ángel [23] is part of the studies devoted to the linear and nonlinear
111 plasmonic properties of nanohybrids and nanocomposites made of metallic nanoparticles and
112 nanoemitters emitters (e.g. [25, 26]). The model developed herein reveals that the inclusion in these
113 structures of dispersed single-walled carbon nanotubes considerably changes the properties of the
114 plasmon-polariton fields in such a hybrid system. This makes possible to control the properties of local
115 optical fields that allows creating of miniature low threshold coherent sources with tunable
116 polarization. Numerical simulations show also that the strength of coupling of nanoemitter with
117 plasmon-polaritons is maximal when the field interconnections appear in the entire carbon nanotubes
118 array. We then move to a work by Y. E. Geints that is focused on the optical properties of miniature
119 resonant microcavities whose efficiency is sufficient to provide whispering-gallery modes. These
120 “photonic molecules (PM)” [27], composed of “atoms” (e.g., cylinders or spheres) are studied when
121 excited simultaneously via two light channels (tapered fibers). The current numerical study, restricted
122 to mirror-symmetric PM, reports that for a seven-atom silicon microcylinder cyclic-PM a tenfold
123 intensity amplification/suppression of several supermodes from the Stokes and anti-Stokes bands of
124 PM spectrum can be reached [26].

125
126 In many situations laser diagnostics can be considered as non-intrusive. This is no longer the case
127 when the laser beams ionize or heat up matter or, as the following two papers [28] [29] show, induce
128 mechanical forces through radiation pressure. Gouesbet et al. [28] study forces exerted on
129 quadrupoles, in terms of BSCs and Mie coefficients, within the framework of GLMT. These forces are
130 categorized in mixing forces and recoil forces, gradient and non-gradient forces. The authors also
131 propose to introduce a third category, in which the so-called scattering forces are actually decomposed
132 in scattering forces and non-standard forces [28]. The second paper on optical forces is from the same
133 authors, it is the last paper of a series aiming to the partition of optical forces exerted on arbitrary sized
134 particles in the framework of GLMT [29].

135
136 Although there are various electromagnetic approaches for calculating the scattering properties of
137 fractal aggregates [30-32], they are still limited in terms of maximum dimension of the aggregates or
138 the size and absorptivity of the monomers. The computational efforts they require with respect to
139 analytical models like the Rayleigh-Debye-Gans approximation for fractal aggregates (RDG-FA)[33-
140 36] is also a critical aspect for inverse methods requiring the creation of large database. RDG-FA, that
141 is intrinsically based on Rayleigh theory of the scattering by a dipole, also neglects electromagnetic
142 coupling within the aggregate. But it provides simple formula for the interpretation of the behavior of
143 the optical structure factor (OSF) with key parameters such as the number of monomers, the fractal
144 dimension or the radius of gyration [35, 37], that are at the basis of numerous reasoning in various
145 research fields. In the present paper [38], Argentin et al employ the discrete dipole approximation and
146 phasor approach [39] to study the role of the coupling on the OSF for the specific case of soot
147 particles. The later have an intermediate fractal dimension (1.6 to 1.8) and high monomer absorptivity
148 (carbonaceous material) that partially violate the basic assumptions of this theory. However, via a
149 numerical study, the authors succeed to propose correction functions allowing to slightly extend the
150 range of validity of RDG-FA for soot particles. These functions are thought to be useful for related
151 laser diagnostics. The Geometrical Optics (GO) is another asymptotic approach whose hypothesis
152 regarding the scatterer size is totally opposite. It basically assumes that the scattered is much larger
153 than the wavelength and that other optical properties (like the Fresnel coefficients for instance) change
154 slowly regarding the wavelength scale. The fact is that, during the last decades, the interest in GO and
155 physical optics approximations (e.g. [40-47]), and their hybridization (e.g. [47, 48]), has considerably
156 raised. This is certainly due to the fact that electromagnetic models stalled in the race towards particles
157 with large size parameters (i.e. a few hundred). In the work by Zhang et al [49], the authors evaluate
158 the characteristic size of a particle that is the most relevant to the accuracy of the GOA in the case of
159 faceted dielectric particle generated from platonic solids. This model, introduced more than one
160 decade ago, is not based on a conventional ray tracing method with differential geometry
161 considerations, e.g. [44, 50-52]. As a hybrid model, it uses conventional GO to calculate the particle
162 near-field and physical optics to calculate the corresponding far-field. It comes out from numerical
163 comparisons with a rigorous invariant imbedding T-matrix method [53], that accuracy of this hybrid

164 model is determined by the individual facet size and the maximum dimension of the particle. The
165 curvature of the particle surface, that acts as a divergent factor [50, 54] in the calculation of the
166 intensity of the scattered rays or the focusing of the indent rays, is not considered in this work.

167
168 This topical issue ends with three papers on optical particle characterization methods. The first paper,
169 by Shopa et al [55], deals with experiments on the analysis of the migration of nanoparticle towards
170 the surface of a slowly evaporating colloidal droplet trapped in an electrodynamic trap. For this
171 purpose, the authors track the change in the laser induced fluorescence of the doped nanoparticles as
172 well as in the fine resonance structures of the scattering patterns and whispering gallery modes of the
173 droplet. Moreover, despite some successes, the authors stress on a possible direct interaction of the
174 pumping beam with the nanoparticles self-organization. The second paper, by Brunel et al [56],
175 proposes a model accounting for spherical aberrations introduced by the imaging lens of
176 interferometric particle imaging (IPI) systems. Note that IPI systems were initially designed to
177 characterize the size and velocity of flowing droplet or bubble [57-59]. The correction model
178 introduced here decomposes the aberrating phase term introduced by the imaging system over a basis
179 of Gaussian functions. The predictions and corrections achieved with this model are in excellent
180 agreement with experimental data for on-axis or off-axis rough particles, or for neighbored rough
181 particles whose interferometric images overlap, producing complex speckle-like images. These
182 developments are powerful not only for the correction of IPI images, but also all 3D reconstructions
183 based on multi-view setups and back projections from different 2D images. In the last paper, by Peng
184 et al [60], the authors investigate the robustness of the analysis of Global rainbow signals [61, 62] for
185 signal-to-noise-ratios in the range 60 down to 5dB. The inversion method used is a conventional
186 parametric analysis assuming the particle distribution shape (normal, log-normal, and bimodal
187 normal)[63] but it is coupled with a more innovative empirical mode decomposition (EMD) for
188 denoising. The numerical study, as well as preliminary experimental results, shows that the EMD
189 method helps a lot in the ideal case of an additive white noise.

190
191 To conclude this editorial, as guest editors and on behalf of the committees of LIP conferences, we
192 want to thank the JQSRT editorial board, for the quality of their work and their unwavering support.
193 We also thank all authors of selected papers presented herein and the anonymous reviewers who have
194 helped us, collectively, to improve the readability and impact of all selected papers. We also take this
195 opportunity to announce that the 14th edition of LIP conference will be organized by the School of
196 Physics of Xidian University, in Xi'an, China, on September 19th—22nd, 2024, see at [http://lip-](http://lip-conference.org/)
197 [conference.org/](http://lip-conference.org/), under the supervision of the conference conveners, L. Guo, Y. Han, F. Onofri and
198 G. Gouesbet.

199
200



201
202 **Fabrice Onofri** is senior researcher (directeur de recherche) at the French National Center for
203 Scientific Research (CNRS). He is mainly working on the optical and electromagnetic characterization
204 of particle and particle systems. Over the years, his research has found applications in various fields
205 ranging from aerosols and multiphase flows to soft matter and plasma physics, in the frame of national
206 and international academic as well as industrial granted projects.

207
208



209
210
211
212
213
214
215
216
217
218
219
220
221

Gérard Gouesbet is a Professor Emeritus of the National Institute of Applied Sciences of Rouen. He has worked in particular on the modeling of turbulent and multiphase flows, on the nonlinear dynamics of systems, on combustion and plasmas, on optical particle characterization and light scattering. He is the driving force behind the development of generalized Lorenz-Mie theories, and the winner of the 2020 van de Hulst award. He published nearly 400 papers in peer-reviewed journals and book chapters. He is also involved in philosophical studies, e.g. his book in philosophy of sciences entitled "Hidden world in quantum physics, Dover publications" and his recently achieved tetralogy on the philosophy of violence (written in French)



222
223
224
225
226
227
228
229
230

Maciej Kolwas is professor in the Institute of Physics of the Polish Academy of Science. His scientific activity started in field of quantum optics studying two photon phenomena - Hanle effect, photon bunching, nonlinear phenomena inside laser resonators. Then his interest concentrated on study of thermodynamics of micro and nano droplets of pure liquids and colloidal suspensions with the laser light scattering. He held several administrative positions at the Institute of Physics- head of group, head of division, rector of International Post Graduate Studies. He was active member of physical societies acting on the end as President of Polish Physical Society and President European Physical Society.

231
232



233
234
235
236
237

Daniel Jakubczyk is an associate professor and head of the Group of Optical Characterisation of Micro- and Nanoobjects at the Institute of Physics, Polish Academy of Sciences. His research activities are essentially experimental. They have included the formation of light-induced sodium nanodroplets, near-field scanning optical microscopy, thermodynamics of evaporation of micro- and nanodroplets, surface thermodynamics of evaporating microdroplets of suspensions/nanofluids,

238 application of machine learning to optical recognition of nanofluids. He received his Ph.D. and
239 habilitation degree at the Institute of Physics PAS.

240

241 References

- 242 1. D. Jakubczyk, M. Kolwas, F. Onofri, and G. Gouesbet, "Laser-light and Interactions with
243 Particles (LIP2022) - Optical Particle Characterization follow-up !, August 21-26th, 2022, Warsaw,
244 Poland," <http://lip-conference.org/>.
- 245 2. G. Gouesbet, and F. R. A. Onofri, "Special issue: Laser-light and interactions with particles
246 (LIP), 2020," *J. Quant. Spectrosc. Radiat. Transf.* **274**, 107857 (2021).
- 247 3. G. Gouesbet, and F. Onofri, "Laser-light and Interactions with Particles (LIP), 2016," *J. Quant.*
248 *Spectrosc. Radiat. Transf.*
- 249 4. G. Gouesbet, P. Yang, and F. Onofri, "Afterword. Laser-light and interactions with particles
250 (LIP), 2018," *J. Quant. Spectrosc. Radiat. Transf.* **225**, 45-49 (2019).
- 251 5. G. Gouesbet, and M. Brunel, "Lasers and interactions with particles, 2012: Optical particle
252 characterization follow-up," *J. Quant. Spectrosc. Radiat. Transf.* **126**, 1-6 (2013).
- 253 6. L. A. Ambrosio, J. Wang, and G. Gouesbet, "On a class of definite integrals with products of
254 (Ricatti-)Bessel functions and their derivatives," *J. Quant. Spectrosc. Radiat. Transf.* **299**, 108512
255 (2023).
- 256 7. L. A. Ambrosio, G. Gouesbet, and J. Wang, "On analytical solutions to classes of definite
257 integrals with products of Bessel functions of the first kind and their derivatives," *J. Quant. Spectrosc.*
258 *Radiat. Transf.* **293**, 108387 (2022).
- 259 8. L. Lorenz, "Lysbevaegelsen i og uden for en hal plane lysbølger belyst kulge," *Vidensk. Selk.*
260 *Skr.* **6**, 1-62 (1898).
- 261 9. G. Mie, "Beiträge zur optik trüber medien, speziell kolloidaler metallösungen," *Ann. Phys.*
262 **330**, 377-445 (1908).
- 263 10. W. J. Wiscombe, "Improved Mie scattering algorithms," *Appl. Opt.* **19**, 1505-1509 (1980).
- 264 11. C. F. Bohren, and D. R. Huffman, *Absorption and scattering of light by small particles* (Wiley
265 & Sons, 1998).
- 266 12. S. C. Barber, and P. W. Hill, *Light Scattering by particles : computational methods* (World
267 Scientific, 1990).
- 268 13. L. A. Ambrosio, "Analytical description of on-axis zero-order continuous frozen waves in the
269 generalized Lorenz-Mie theory," *J. Quant. Spectrosc. Radiat. Transf.* **296**, 108442 (2023).
- 270 14. L. F. Votto, A. Chafiq, G. Gouesbet, L. A. Ambrosio, and A. Belafhal, "Ince-Gaussian beams
271 in the generalized Lorenz-Mie theory through finite series Laguerre-Gaussian beam shape
272 coefficients," *J. Quant. Spectrosc. Radiat. Transf.* **302**, 108565 (2023).
- 273 15. G. Gouesbet, and G. Gréhan, *Generalized Lorenz-Mie Theories* (Springer, 2011).
- 274 16. M. Zamboni-Rached, "Stationary optical wave fields with arbitrary longitudinal shape by
275 superposing equal frequency Bessel beams: Frozen Waves," *Opt. Express* **12**, 4001-4006 (2004).
- 276 17. M. A. Bandres, and J. C. Gutiérrez-Vega, "Ince-Gaussian beams," *Opt. Lett.* **29**, 144-146
277 (2004).
- 278 18. K. Dong, W. Jiang, M. Cheng, C. Shi, Y. Cao, and L. Guo, "Attenuation of Bessel vortex
279 beam transmission in the rain environment," *J. Quant. Spectrosc. Radiat. Transf.* **304**, 108620 (2023).
- 280 19. M. C. Gökçe, Y. Baykal, and Y. Ata, "Coupling efficiency of multimode beam to fiber in
281 atmospheric turbulence," *J. Quant. Spectrosc. Radiat. Transf.* **303**, 108590 (2023).
- 282 20. F. G. Mitri, "Electromagnetic Wave Scattering of a High-Order Bessel Vortex Beam by a
283 Dielectric Sphere," *IEEE Transactions on Antennas and Propagation* **59**, 4375-4379 (2011).
- 284 21. J. Wang, "Advances in communications using optical vortices," *Photon. Res.* **4**, B14-B28
285 (2016).
- 286 22. M. Uysal, and H. Nouri, "Optical wireless communications — An emerging technology," in
287 *2014 16th International Conference on Transparent Optical Networks (ICTON)(2014)*, pp. 1-7.
- 288 23. G. Burlak, and G. Medina-Ángel, "Extended dynamics and lasing of nanoemitters enhanced
289 by dispersing single-walled carbon nanotubes," *J. Quant. Spectrosc. Radiat. Transf.* **296**, 108463
290 (2023).
- 291 24. Y. E. Geints, "Phase-controlled supermodes in symmetric photonic molecules," *J. Quant.*
292 *Spectrosc. Radiat. Transf.* **302**, 108524 (2023).

- 293 25. N. Zhang, C. Han, X. Fu, and Y.-J. Xu, "Function-Oriented Engineering of Metal-Based
294 Nanohybrids for Photoredox Catalysis: Exerting Plasmonic Effect and Beyond," *Chem* **4**, 1832-1861
295 (2018).
- 296 26. M. R. Singh, D. G. Schindel, and A. Hatef, "Dipole-dipole interaction in a quantum dot and
297 metallic nanorod hybrid system," *Appl. Phys. Lett.* **99**, 181106 (2011).
- 298 27. M. Bayer, T. Gutbrod, J. P. Reithmaier, A. Forchel, T. L. Reinecke, P. A. Knipp, A. A.
299 Dremin, and V. D. Kulakovskii, "Optical Modes in Photonic Molecules," *Phys. Rev. Lett.* **81**, 2582-
300 2585 (1998).
- 301 28. G. Gouesbet, V. S. De Angelis, and L. A. Ambrosio, "Optical forces and optical force
302 categorizations exerted on quadrupoles in the framework of generalized Lorenz–Mie theory," *J. Quant.*
303 *Spectrosc. Radiat. Transf.* **298**, 108487 (2023).
- 304 29. G. Gouesbet, V. S. De Angelis, and L. A. Ambrosio, "Optical forces and optical force
305 partitions exerted on arbitrary sized spherical particles in the framework of generalized Lorenz–Mie
306 theory," *J. Quant. Spectrosc. Radiat. Transf.* **307**, 108661 (2023).
- 307 30. D. W. Mackowski, and M. I. Mishchenko, "Calculation of the T matrix and the scattering
308 matrix for ensembles of spheres," *J. Opt. Soc. Am. A* **13**, 2266-2278 (1996).
- 309 31. M. A. Yurkin, and A. G. Hoekstra, "The discrete-dipole-approximation code ADDA:
310 Capabilities and known limitations," *J. Quant. Spectrosc. Radiat. Transf.* **112**, 2234-2247 (2011).
- 311 32. B. T. Draine, and P. J. Flatau, "Discrete Dipole Approximation Scattering (DDSCAT) code,"
312 <http://code.google.com/p/ddscat/>.
- 313 33. Köylü Ü.Ö., Faeth G.M., Farias T.L., and Carvalho M.G., "Fractal and Projected Structure
314 Properties of Soot Aggregates," *Combustion and flame* **100**, 621-633 (1995).
- 315 34. T. L. Farias, Ü. Ö. Köylü, and M. G. Carvalho, "Range of validity of the Rayleigh–Debye–
316 Gans theory for optics of fractal aggregates," *Appl. Opt.* **35**, 6560-6567 (1996).
- 317 35. C. M. Sorensen, "Light Scattering by Fractal Aggregates: A Review," *Aerosol Science and*
318 *Technology* **35**, 648-687 (2001).
- 319 36. R. A. Dobbins, and C. M. Megaridis, "Absorption and scattering of light by polydisperse
320 aggregates," *Appl. Opt.* **30**, 4747-4754 (1991).
- 321 37. J. Mroczka, M. Wozniak, and F. R. A. Onofri, "Algorithms and methods for the analysis of the
322 optical structure factor of fractal aggregates,," *Metrology and Measurement Systems* **XIX**, 459-470
323 (2012).
- 324 38. C. Argentin, M. J. Berg, M. Mazur, R. Ceolato, and J. Yon, "Electromagnetic coupling and
325 determination of the structure factor of fractal aggregates," *J. Quant. Spectrosc. Radiat. Transf.* **296**,
326 108451 (2023).
- 327 39. M. J. Berg, C. M. Sorensen, and A. Chakrabarti, "Reflection symmetry of a sphere's internal
328 field and its consequences on scattering: a microphysical approach," *J. Opt. Soc. Am. A* **25**, 98-107
329 (2008).
- 330 40. E. A. Hovenac, "Calculation of far-field scattering from nonspherical particles using a
331 geometrical optics approach," *Appl. Opt.* **30**, 4739-4746 (1991).
- 332 41. J. A. Lock, "Ray scattering by an arbitrarily oriented spheroid. II. Transmission and cross-
333 polarization effects," *Appl. Opt.* **35**, 515-531 (1996).
- 334 42. J. A. Lock, "Ray scattering by an arbitrarily oriented spheroid. I. Diffraction and specular
335 reflection," *Appl. Opt.* **35**, 500-514 (1996).
- 336 43. C. L. Adler, J. A. Lock, I. P. Rafferty, and W. Hickok, "Twin-rainbow metrology. I.
337 Measurement of the thickness of a thin liquid film draining under gravity," *Appl. Opt.* **42**, 6584-6594
338 (2003).
- 339 44. O. N. Stavroudis, *The mathematics of Geometrical and Physical Optics* (Wiler-VCH Verlag &
340 Co. KGaA, 2006).
- 341 45. F. R. A. Onofri, S. Radev, M. Sentis, and S. Barbosa, "Physical-optics approximation of near-
342 critical-angle scattering by spheroidal bubbles," *Opt. Lett.* **37**, 4780-4782 (2012).
- 343 46. H. Yu, F. Xu, and C. Tropea, "Spheroidal droplet measurements based on generalized rainbow
344 patterns," *J. Quant. Spectrosc. Radiat. Transf.* **126**, 105-112 (2013).
- 345 47. M. P. L. Sentis, F. R. A. Onofri, L. Méès, and S. Radev, "Scattering of light by large bubbles:
346 Coupling of geometrical and physical optics approximations," *J. Quant. Spectrosc. Radiat. Transf.*
347 **170**, 8-18 (2016).

- 348 48. L. Bi, P. Yang, G. W. Kattawar, Y. Hu, and B. A. Baum, "Scattering and absorption of light
349 by ice particles: Solution by a new physical-geometric optics hybrid method," *J. Quant. Spectrosc.*
350 *Radiat. Transf.* **112**, 1492-1508 (2011).
- 351 49. Y. Zhang, J. Ding, P. Yang, and G. Videen, "Evaluating the accuracy of single-scattering
352 computations by the geometric optics approximation using Platonic solids," *J. Quant. Spectrosc.*
353 *Radiat. Transf.* **303**, 108591 (2023).
- 354 50. H. C. van de Hulst, *Light Scattering by Small Particles* (Dover Publications, 1957).
- 355 51. H. Yu, F. Xu, and C. Tropea, "Optical caustics associated with the primary rainbow of oblate
356 droplets: simulation and application in non-sphericity measurement," *Opt. Express* **21**, 25761-25771
357 (2013).
- 358 52. Q. Duan, F. R. A. Onofri, X. e. Han, and K. F. Ren, "Generalized rainbow patterns of oblate
359 drops simulated by a ray model in three dimensions," *Opt. Lett.* **46**, 4585-4588 (2021).
- 360 53. M. I. Mishchenko, "Comprehensive thematic T-matrix reference database: a 2017–2019
361 update," *J. Quant. Spectrosc. Radiat. Transf.* **242**, 106692 (2020).
- 362 54. K. F. Ren, F. Onofri, C. Rozé, and T. Girasole, "Vectorial complex ray model and application
363 to two-dimensional scattering of plane wave by a spheroidal particle," *Opt. Lett.* **36**, 370-372 (2011).
- 364 55. Y. Shopa, M. Kolwas, I. Kamińska, G. Derkachov, K. Nyandey, T. Jakubczyk, T.
365 Wojciechowski, A. Derkachova, and D. Jakubczyk, "Luminescent nanoparticles in a shrinking
366 spherical cavity – probing the evaporating microdroplets of colloidal suspension – optical lattices and
367 structural transitions," *J. Quant. Spectrosc. Radiat. Transf.* **296**, 108439 (2023).
- 368 56. M. Brunel, A. Abad, B. Delestre, Q. Frodello, R. Brunel, S. Coëtmellec, and M. Talbi,
369 "Analysis and numerical correction of aberration in interferometric particle imaging," *Journal of*
370 *Quantitative Spectroscopy and Radiative Transfer* **302**, 108579 (2023).
- 371 57. Kawaguchi T., Akasaka Y., and Maeda M., "Size measurements of droplets and bubbles by
372 advanced interferometric laser imaging technique," *Meas. Sci. Technol.* **13**, 308-316 (2002).
- 373 58. A. R. Glover, S. M. Skippon, and R. D. Boyle, "Interferometric laser imaging for droplet
374 sizing: a method for droplet-size measurement in sparse spray systems," *Appl. Opt.* **34**, 8409-8421
375 (1995).
- 376 59. H. Shen, S. Coëtmellec, G. Gréhan, and M. Brunel, "Interferometric laser imaging for droplet
377 sizing revisited: elaboration of transfer matrix models for the description of complete systems," *App.*
378 *Opt.* **51**, 5357-5368 (2012).
- 379 60. W. Peng, C. Li, T. Li, L. Huang, N. Li, and C. Weng, "Anti-noise and denoising performance
380 of global rainbow processing," *J. Quant. Spectrosc. Radiat. Transf.* **304**, 108619 (2023).
- 381 61. J. P. A. J. van Beeck, D. Giannoulis, L. Zimmer, and M. L. Riethmuller, "Global rainbow
382 thermometry for droplet-temperature measurement," *Opt. Lett.* **24**, 1696-1698 (1999).
- 383 62. X. Wang, Y. Wu, D. Xu, B. Wen, Q. Lv, and X. Wu, "Synthetic aperture rainbow
384 refractometry," *Opt. Lett.* **47**, 5272-5275 (2022).
- 385 63. M. Ouattara, F. Lamadie, M. P. L. Sentis, and F. R. A. Onofri, "Droplet sizing and mixture
386 fraction measurement in liquid flows with rainbow-angle diffractometry," *Appl. Opt.* **56**, 8109-8120
387 (2017).
- 388