



HAL
open science

Concentrations of bisphenols and phthalate esters in the muscle of Mediterranean striped dolphins (*Stenella coeruleoalba*)

Morgana Vighi, Asunción Borrell, Wissam Sahyoun, Sopheak Net, Alex Aguilar, Baghdad Ouddane, Odei Garcia-Garin

► To cite this version:

Morgana Vighi, Asunción Borrell, Wissam Sahyoun, Sopheak Net, Alex Aguilar, et al.. Concentrations of bisphenols and phthalate esters in the muscle of Mediterranean striped dolphins (*Stenella coeruleoalba*). *Chemosphere*, 2023, 339, pp.139686. 10.1016/j.chemosphere.2023.139686. hal-04195906

HAL Id: hal-04195906

<https://hal.science/hal-04195906v1>

Submitted on 4 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Journal Pre-proof

Concentrations of bisphenols and phthalate esters in the muscle of Mediterranean striped dolphins (*Stenella coeruleoalba*)

Morgana Vighi, Asunción Borrell, Wissam Sahyoun, Sopheak Net, Alex Aguilar, Baghdad Ouddane, Odei Garcia-Garin



PII: S0045-6535(23)01953-7

DOI: <https://doi.org/10.1016/j.chemosphere.2023.139686>

Reference: CHEM 139686

To appear in: *ECSN*

Received Date: 15 March 2023

Revised Date: 20 July 2023

Accepted Date: 29 July 2023

Please cite this article as: Vighi, M., Borrell, Asunció., Sahyoun, W., Net, S., Aguilar, A., Ouddane, B., Garcia-Garin, O., Concentrations of bisphenols and phthalate esters in the muscle of Mediterranean striped dolphins (*Stenella coeruleoalba*), *Chemosphere* (2023), doi: <https://doi.org/10.1016/j.chemosphere.2023.139686>.


This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 Published by Elsevier Ltd.

CRedit author statement

Morgana Vighi: Conceptualization, Investigation, Writing - Original Draft, Visualization, Supervision; **Asunción Borrell:** Conceptualization, Methodology, Resources, Writing - Review & Editing, Supervision; **Wissam Sahyoun:** Methodology, Formal analysis; **Sopheak Net:** Methodology, Resources, Writing - Review & Editing, Supervision; **Alex Aguilar:** Conceptualization, Resources, Writing - Review & Editing, Supervision; **Baghdad Ouddane:** Methodology, Resources; **Odei Garcia-Garin:** Conceptualization, Methodology, Formal analysis, Investigation, Data curation, Writing - Review & Editing, Visualization, Project administration, Funding acquisition.

Journal Pre-proof



Journal Pre-proof

1 **Concentrations of bisphenols and phthalate esters in the muscle of Mediterranean striped**
2 **dolphins (*Stenella coeruleoalba*)**

3

4 Morgana Vighi^{1,2*}, Asunción Borrell^{1,2}, Wissam Sahyoun³, Sopheak Net³, Alex Aguilar^{1,2}, Baghdad
5 Ouddane³, Odei Garcia-Garin^{1,2}

6 ¹Department of Evolutionary Biology, Ecology and Environmental Sciences, Faculty of Biology.
7 University of Barcelona, 08028, Barcelona, Spain

8 ²Biodiversity Research Institute (IRBio). University of Barcelona, 08028, Barcelona, Spain

9 ³Université de Lille, Faculté des Sciences et Technologies, Laboratoire LASIRE (UMR 8516 CNRS),
10 Cité Scientifique, 59655 Villeneuve d'Ascq Cedex, France

11 * Corresponding author: morgana.vighi@gmail.com. C/ Sant Baltasar, 18, 3, 08014, Barcelona, Spain

12

13 **Highlights**

14 .

- 15 1. Muscle samples were obtained from 30 stranded Mediterranean striped dolphins.
16 2. Six bisphenols and 6 phthalate esters were detected in the samples.
17 3. BPZ was detected in all the samples, at the highest concentrations.
18 4. Concentrations did not relate with biological traits but varied with time.
19 5. Results confirm dolphin exposure to plasticizers in an already anthropized basin.

20

21 **Abstract**

22 Bisphenols (BPs) and phthalate esters (PAEs) are important compounds for the plastics industry, also
23 called “everywhere chemicals” due to their ubiquity in daily use products. Both chemical groups are
24 well-known environmental contaminants, whose presence has been reported in all environmental

25 compartments, and whose effects, mainly associated to endocrine disruption, are detrimental to living
26 organisms. Cetaceans, due to their long life-span, low reproduction rate and high position in the trophic
27 web, are especially vulnerable to the effects of contaminants. However, little is known about BP and
28 PAE concentrations in cetacean tissues, their potential relation to individual biological variables, or
29 their trends over time. Here, the concentration of 10 BPs and 13 PAEs was assessed in the muscle of
30 30 striped dolphins (*Stenella coeruleoalba*) stranded along the Spanish Catalan coast (NW
31 Mediterranean) between 1990 and 2018. Six BP and 6 PAE compounds were detected, of which only
32 4,4'-(cyclohexane-1,1-diyl)diphenol (BPZ) was detected in all the samples, at the highest
33 concentration (mean 16.06 $\mu\text{g g}^{-1}$ lipid weight). Sex or reproductive condition were largely uninfluential
34 on concentrations: only dimethylphthalate (DMP) concentrations were significantly higher in
35 immature individuals than in adults, and the overall PAE concentrations were significantly higher in
36 males than in females. Temporal variations were only detected in bis(4-hydroxyphenyl)ethane (BPE),
37 diethylphthalate (DEP) and dimethylphthalate (DMP), whose concentrations were lower, and 9,9-
38 Bis(4-hydroxyphenyl)fluorene (BPFL), which were higher, respectively, in samples taken between
39 2014-2018, probably reflecting shifts in the production and use of these chemicals. These results
40 provide the first assessment of concentrations of several BP and PAE compounds in the muscle of an
41 odontocete cetacean.

42

43 **Keywords**

44 Pollutants; Plastic additives; Plasticizers; Contamination; Marine Mammals; Cetaceans

45 1. Introduction

46 Bisphenols (BPs) and phthalic acid esters (PAEs) have been called “everywhere chemicals”
47 due to their widespread use in the manufacture and processing of plastic and resin products and their
48 ubiquity in human body, wildlife and the environment (Net et al, 2015; Warner and Flaws, 2018;
49 Sangeetha et al, 2021). Indeed, they are important chemical components for the plastics industry, where
50 BPs (and more specifically Bisphenol A (2,2-bis(4-hydroxyphenyl)propane, henceforth BPA), along
51 with its analogues) are used in the manufacture of hard, transparent plastics such as water bottles, as
52 well as in the production of polycarbonate, polyacrylate, flame retardants, solvents, and other products
53 used for coating, such as epoxy resins, and are present in a large variety of everyday products (Staples
54 et al, 1998; Rochester et al, 2015; Mustieles et al, 2020; Torres-García et al, 2022). On the other hand,
55 PAEs are used to manufacture soft, flexible and transparent plastics, such as plastic polyvinyl chloride
56 (PVC) products, and a very broad range of products including building materials, toys, personal care
57 products (PCP), food packaging, cosmetics, medical devices (Graham, 1973), and many other daily
58 use products, where their content can be up to 10-60 % by weight (Net et al, 2015).

59 Although plastic materials are chemically inert, the chemical additives used to improve their
60 qualities may be released to the environment during the plastic weathering processes caused by solar
61 radiation, sea corrosion, and chemical and biological degradation (Koelmans et al, 2014; Net et al,
62 2015). Moreover, BPs and PAEs may also enter the environment during the production process and
63 ultimately be transported at sea through river discharge or atmospheric deposition (Xie et al., 2007).
64 The constant raise of plastic production, and consequent plastic pollution in the worldwide oceans
65 (Jambeck et al, 2015; Geyer et al, 2017; Lambert et al, 2020) is thus expected to increase the
66 concentration of plastic additives such as BPs and PAEs in the marine environment, as a result of their
67 direct deposition, and/or their leaching from the large amounts of marine plastic litter that reach the
68 oceans every day.

69 Indeed, both BPs and PAEs are well-known environmental pollutants, which have been
70 detected in every natural compartment, including air, soil, sediments, fresh and marine waters (Xie et
71 al, 2007; Zeng et al., 2009; Ben Sghaier et al, 2017; Paluselli et al, 2018; Moustafa et al, 2022; Torres-
72 García et al, 2022; Xing et al, 2022), as well as in the tissues of several animal species (Kang et al,
73 2007; Koelmans, 2014; Nehring et al, 2017; Routti et al., 2021). Both *in vivo* and *in vitro* studies have
74 proven their toxic effects to living organisms. *In vitro* studies with laboratory animals and
75 epidemiological studies in humans showed that BPA, several of its analogues, and most PAEs, cause
76 severe endocrine effects (Basak et al, 2020), as well as potential neurological and carcinogenic effects
77 (e.g., Benjamin et al, 2017; Li et al, 2018; Pelch et al, 2019; Han et al, 2021). Although most BPs and
78 PAEs are subject to efficient metabolic degradation and elimination (Kang et al, 2006; Net et al, 2015),
79 the increased levels of environmental and dietary exposure to which living organisms are subject may
80 maintain their tissue concentrations at chronic or sub-chronic levels, and trigger detrimental effects
81 (Silva et al, 2018; Warner and Flaws, 2018).

82 Cetaceans are particularly vulnerable to the effects of lipophilic contaminants. Being mammals
83 with a long life-span, low reproduction rate, and high position in the trophic web, they are subject to
84 both the bioaccumulation and biomagnification of pollutants (Loganathan, 2012). Despite the proven
85 toxic effects of BPs and PAEs to mammals, and their potentially high concentrations in the marine
86 environment, their concentrations in the tissues of marine mammals have been determined only by a
87 handful of studies so far. Most of the published research focuses on the assessment of PAEs and their
88 metabolites in the tissues of two main species: the bottlenose dolphin (*Tursiops truncatus*, *i.e.*, Dziobak
89 et al, 2021, 2022; Hart et al, 2018, 2020) and the fin whale (*Balaenoptera physalus*, *i.e.*, Bains et al,
90 2017; Fossi et al, 2012, 2014, 2016; Garcia-Garin et al, 2022); while other marine mammal species
91 have been scarcely studied (e.g., Rian et al, 2020; Routti et al, 2021), and the concentrations of BPs in
92 these organisms have been barely investigated (*i.e.*, Montoro-Martínez et al, 2021; Page-Karjian et al,
93 2020). Still little is known regarding the concentrations of these contaminants, and especially of BPs

94 and PAEs other than the more commonly studied BPA and di-(2-ethylhexyl) phthalate (DEHP), in
95 marine mammals, their potential relation with biological variables, or their trends over time.


96 With the aim of contributing to fill such information gap, here we analysed 10 BPs and 13 PAEs
97 in the muscle of 30 striped dolphins (*Stenella coeruleoalba*) stranded along the Mediterranean coast
98 of Spain between 1990 and 2018. Our results are discussed taking into account the potential influence
99 of biological variables, such as sex and maturation stage of the individuals, and of the year of death,
100 which might reflect the increasing levels of plastic pollution in the Mediterranean Sea and the
101 regulation of some of the analysed compounds, like BPA and DEHP. Although the Mediterranean
102 striped dolphin is currently not in a concerning conservation status, cetaceans are often subject to a
103 variety of human impacts that, combined, may compromise the healthy survival of their populations
104 (Hart et al, 2020, Dziobaak et al, 2022). The pathways and levels of exposure to these increasingly
105 widespread pollutants of potentially vulnerable marine fauna such as cetaceans deserve in-depth
106 investigation.

107 **2. Materials and methods**

108 *2.1. Study area and sampling*

109 Muscle samples were collected from 30 striped dolphins found stranded along the Spanish
110 Catalan coast, in the NW Mediterranean Sea (Fig.1), during three separate periods between 1990 and
111 2018 (*i.e.*, 1990, 2007-08, and 2014-18), and preserved in the biological tissue bank of the University
112 of Barcelona (BMA Tissue Bank). Since muscle represents approximately 50% of dolphins weight
113 and its average lipid content is 6%, this tissue was considered representative enough to assess the levels
114 of BPs and PAEs in the dolphins. The cause of death for most dolphins was related with the
115 morbillivirus epizootics occurred in the Mediterranean Sea during 1990 and 2006-07 periods. As age
116 information was only available from 7 out of the 30 dolphins, individuals were classified according to

117 their sexual maturation stage, defined according to their body length. Individuals whose body was
118 shorter than 1.3 m were classified as immature, and individuals whose body was longer than 1.8 m
119 were classified as adult (Table 1). Overall, sampled individuals included 24 adults (12 males and 12
120 females), and 6 immature individuals (2 males and 4 females, Table 1). Muscle samples, stored at -20
121 $^{\circ}\text{C}$, were subsampled using stainless steel material to obtain individual subsamples of 25-30 g, which
122 were placed in glass bottles and preserved frozen until analysis. To avoid contamination, no plastic
123 material was used during this procedure.



124
125 Figure 1: Map of the Western Mediterranean Sea highlighting (in pink) the area of the Spanish Catalan coast, where the
126 striped dolphins sampled in this study were found stranded.

127

128 Table 1: Sex, body length (in m), maturation stage and age (when determined) of the sampled striped dolphins, along with
 129 the lipid content and overall phthalate and bisphenol compounds concentrations found in their muscle (expressed in $\mu\text{g g}^{-1}$
 130 lipid weight (lw)).

Sample	Year	Sex	Length	Maturation stage	Age	Lipid content (%)	Σ PAE	Σ BP
2007_1	2007	M	2.1	MATURE	NA	1.24	72.499	1.277
2007_3	2007	F	1.8	MATURE	NA	1.21	9.564	17.251
2007_4	2007	F	1.95	MATURE	NA	0.69	30.430	32.939
2007_6	2007	M	1.85	MATURE	NA	1.10	23.231	17.249
2008_1	2008	F	2.1	MATURE	NA	15.46	3.568	0.369
2008_6	2008	F	1.89	MATURE	NA	1.08	46.250	26.104
2008_7	2008	M	1.85	MATURE	NA	1.04	7.923	42.594
9033	1990	M	1.93	MATURE	16	1.87	26.645	2.714
9034	1990	F	1.96	MATURE	14	1.64	3.542	8.971
9040	1990	M	1.81	MATURE	7	1.77	14.713	13.499
9042	1990	M	1.87	MATURE	10	1.67	4.463	50.728
9043	1990	F	1.8	MATURE	9	1.85	1.547	4.665
9046	1990	F	1.82	MATURE	30	1.58	5.192	9.069
9054	1990	F	2	MATURE	23	1.77	5.020	17.188
9067	1990	M	1.91	MATURE	NA	1.86	36.006	28.954
N_1_18	2018	F	2	MATURE	NA	1.04	2.089	1.905
N_168_17	2017	M	1.93	MATURE	NA	1.30	18.476	5.756
N_21_18	2018	M	1.82	MATURE	NA	1.40	15.992	0.924
N_286_15	2015	F	1	IMMATURE	NA	2.88	13.259	10.305

N_291_14	2014	F	1.04	IMMATURE	NA	16.78	1.832	0.154
N_352_14	2014	M	1.3	IMMATURE	NA	1.39	0.006	15.661
N_355_14	2014	F	1.07	IMMATURE	NA	3.10	9.681	33.913
N_370_14	2014	M	2.04	MATURE	NA	1.42	4.522	2.240
N_407_15	2015	M	2.1	MATURE	NA	1.22	6.141	20.854
N_45_17	2017	F	1.23	IMMATURE	NA	3.02	80.258	13.801
N_488_17	2017	M	1.98	MATURE	NA	1.38	16.826	24.986
N_49_15	2015	F	1.9	MATURE	NA	2.06	11.118	5.772
N_497_17	2017	F	1.9	MATURE	NA	8.77	1.806	3.242
N_523_17	2017	M	1.03	IMMATURE	NA	2.70	8.303	96.396
N_77_17	2017	F	1.81	MATURE	NA	1.41	1.372	4.375

131

132 *2.2. Sample preparation*

133 After 48 h lyophilization, a subsample of 4 g (dry weight, dw), was taken from the central
134 portion (to avoid possible contamination) of each muscle sample and homogenised. Each subsample
135 was then transferred in 50 mL centrifuge tubes and diluted in a mixture of 10 mL acetonitrile, 10 mL
136 Milli-Q water, 100 µL internal standard (d₄-DIBP), and extraction salts (composed by 1 g NaCl, 4 g
137 MgSO₄, 1 g C₆H₅Na₃O₇·2H₂O and 0.5 g C₆H₆Na₂O₇·1.5H₂O). After stirring in a vortex mixer with
138 ceramic homogenizers and centrifugating for 5 minutes at 5000 rpm, 8 mL of each extract were
139 transferred to centrifuge tubes containing the dispersive solid phase extraction clean-up salts (400 mg
140 PSA, 400 mg GCB, 1200 mg MgSO₄ and 400 mg C18EC). The mixture was then vortexed and
141 centrifuged again at 5000 rpm for 5 minutes. Five (5) mL of the supernatant were transferred to a vial,
142 and the extraction was repeated a second time using 10 mL ethyl acetate. The two resulting extracts
143 were then combined and reduced under a gentle nitrogen stream to 200 µL and 50 µL for PAEs and

144 BPs, respectively. Prior to PAE GC-MS/MS analyses, extracts were stored frozen in glass tubes. For
145 BPs, each extract was derivatized with 50 μ L of BSTFA containing 1% TMCS and heated at 65 °C
146 during 2 h. After derivatization, samples were kept at room temperature during 15 min prior to GC-
147 MS analysis (Sghaier et al, 2017).

148 Due to the lipophilic nature of PAEs and BPs, it is generally recommended to report their
149 concentrations on a lipidic weight (lw) basis to prevent any bias related to the individual, specific, or
150 tissue-related differences in lipid concentrations (Krahn et al, 2003). Thus, following Garcia-Garin et
151 al (2020), 1 g dw of each sample was used to determine the lw gravimetrically by a 15 min sonication
152 in a 15 mL solution of hexane:acetone (1:1). Two extractions were performed and the resulting extracts
153 were mixed. After the solvent evaporation under nitrogen stream, the remaining lipids were dried at
154 90 °C until they reached a constant weight.

155 2.3. Instrumental analysis, standards and reagents

156 Ten BP and thirteen PAE compounds were analysed, namely: BPA, 4,4'-(1-
157 Phenylethylidene)bisphenol (BPAP), 1,1-Bis(4-hydroxyphenyl)-1,1-diphenylmethane (BPBP), 2,2-
158 Bis(4-hydroxy-3-methylphenyl)propane (BPC), bis(4-hydroxyphenyl)ethane (BPE), (bis(4-
159 hydroxyphenyl)methane (BPF), 9,9-Bis(4-hydroxyphenyl)fluorene (BPFL), 4,4'-(1,3-
160 Phenylenediisopropylidene)bisphenol (BPM), 4,4'-(1,4-Phenylenediisopropylidene)bisphenol (BPP),
161 4,4'-(cyclohexane-1,1-diyl)diphenol (BPZ), dimethylphthalate (DMP), diethylphthalate (DEP), di-
162 butylphthalate (DBP), bis(2-methoxyethyl) phthalate (DMEP), dipentylphthalate (DPP), bis(2-
163 ethoxyethyl) phthalate (DEEP), benzyl butyl phthalate (BBP), phthalic acid dicyclohexyl ester
164 (DCHP), bis(4-methyl-2-pentyl) phthalate (BMPP), di-*n*-hexyl phthalate (DHP), bis(2-*n*-butoxyethyl)
165 phthalate (DBEP), bis(2-ethylhexyl) phthalate (DEHP) and di-*n*-octyl phthalate (DNOP). Analytical
166 standards of BPs and PAEs were purchased from TCI Europe (Zwijndrecht, Belgium) and Restek
167 (Lisses, France), respectively. Deuterated BPA (BPA d16) and labelled phthalic acid diisobutyl ester

168 (d₄-DIBP) were obtained from Sigma-Aldrich (St. Louis, MO, USA); acetonitrile and ethyl acetate
169 (for HPLC isocratic grade) for trace analysis were provided by VWR; the extraction salts were
170 provided by Merck Acetone (SupraSolv®) and Agilent Technologies; and the BSTFA + TMCS (99:1)
171 solution was provided by Supelco.

172 An Agilent 7890 GC coupled to an Agilent 7000C Triple Quadrupole GC/MS system (Agilent
173 Technologies, France) were used for analyses. An Agilent DB-17ms, 30 m × 0.25 mm, 0.25 μm
174 column, with a constant flow of 1.2 mL helium (99.999% purity) min⁻¹, was used to produce analyte
175 separation. The solvent vent injection mode was used through an ultra-inert inlet liner, with a glass
176 wool frit from Agilent, to inject samples in a multimode inlet. The injector operated with an injection
177 volume of 3 μL and a temperature cycle of: 0.8 min at 50 °C (solvent evaporation), raise to 300 °C at
178 720 °C min⁻¹; 5 min hold; cooling down to 280 °C; 10 min hold. The quenching and collision gas were
179 helium (99.999% purity; flow rate of 2.25 mL min⁻¹) and nitrogen (99.999% purity; flow rate of 1.5
180 mL min⁻¹), respectively. The temperature cycle of the oven included: 2 min at 40 °C, a first raise to
181 220 °C at 30 °C min⁻¹, a second raise to 260 °C at 5 °C min⁻¹ and a last raise to 280 °C at 20 °C min⁻¹,
182 7 min hold; for a total run time of 24 min. Electron impact ionisation (EI) at 70 eV ionising energy,
183 and the MRM mode were set to operate the triple quadrupole mass spectrometer, at temperatures of
184 280 °C for the transfer line, 230 °C for the ion source and 150 °C for quadrupoles 1 and 2.

185 2.4. Quality assurance

186 To prevent sample contamination, all procedures were performed in clean laboratory
187 conditions: samples and standards preparation, extraction, clean-up and the pre-concentration step
188 were carried out in a laminar flow cabinet. Glassware was washed with detergent (Decon), rinsed with
189 ultra-pure water (Milli-Q water) then washed with acid solution (HCl) and again rinsed with Milli-Q
190 water follow by acetone and was dried at 120 °C prior to use. Analytical blanks were included every
191 10 samples to minimize the error of quantification. Targeted compounds in blanks should be lower

192 than limit of quantification (LOQ) value: if they were higher than LOQ in the samples, the
193 concentration detected in the blank was subtracted from the corresponding batch of samples (Net et al,
194 2015). BPs and PAEs were lower than LOQ in the procedural blanks. Limit of detection (LOD) and
195 LOQ for each compound were estimated as 3-times and 10-times the baseline of blank chromatograms,
196 respectively, and were confirmed by experimental analysis. LOQs for BPs and PAEs were in the range
197 of 0.73 – 4.36 and 0.012 – 0.062 ng g⁻¹ dw, respectively. The recovery rate of each BP and PAE was
198 calculated by spiking the targeted compounds into a real matrix of interest. The mean BP recovery was
199 estimated at 99.7% with a rate of 108, 102, 110, 106, 94 and 97% for BPA, BPE, BPF, BPFL, BPP,
200 and BPZ, respectively. The mean PAE recovery was estimated at 92% with a rate of 104, 81, 86, 98,
201 76 and 106% for DMP, DEP, DBP, DEHP, DCHP and DNOP, respectively. Samples were not replicated
202 but the analytical procedure was repeated to check its replicability: Standard deviations obtained were
203 <5% when the concentration of the compound was higher than 0.1 ng g⁻¹ and ~12% when the
204 concentration of the compound was close to the LOQs. Calibration curves were obtained in the range
205 of 0.1 – 5000 µg l⁻¹ with the correlation coefficients of ≥ 0.994 .

206 The University of Lille, LASIRE laboratory (UMR CNRS 8516) provided the infrastructure
207 and support to perform all above analyses.

208 2.5. Statistical analysis

209 The distribution of each BP and PAE concentrations were tested for normality and
210 heteroscedasticity through a Shapiro Wilk and Levene tests, respectively. As concentrations were not
211 normally distributed, data were analysed using non-parametric tests, considering both each BP and
212 PAE compound individually, as well as the sum of all BP and of all PAE compounds. To exclude
213 potential biases, results obtained from mature individuals (n=24) were used to assess the relative
214 influence of sex and year of death, split in three groups (*i.e.*, 1990, 2007-2008 and 2014-2018), through
215 a Kruskal Wallis and a Spearman's rho correlation test, respectively. Results obtained from the 2014-

216 2018 samples (n=15) were used to assess the potential effect of the maturation stage through a Kruskal
 217 Wallis test. The level of significance was set at $p < 0.05$. Analyses were performed in R (R Core Team,
 218 2021).

219 3. Results


220 The overall concentration of BPs and PAEs detected in each sample are shown in Table 1.
 221 Minimum, maximum and mean concentrations of each BP and PAE, as well as their frequency of
 222 occurrence, are summarized in Table 2. Out of the 10 BPs and 13 PAEs analysed, 6 BPs and 6 PAEs
 223 were detected, respectively. The concentrations of the majority of selected BPs and PAEs were below
 224 the LOQ, meaning that these compounds were not detected in all the sampled dolphins. Indeed, BPZ
 225 was the only compound detected in all samples, at higher concentrations than any other compound
 226 (Table 2). DEP, DBP and DEHP were the only other compounds whose mean concentrations were
 227 higher than $1 \mu\text{g g}^{-1} \text{lw}$, and which were also detected in 80%, 73% and 63% of samples, respectively
 228 (Table 2, Fig. 2).

229 Table 2: Mean concentrations and ranges of individual and overall BP and PAE compound concentrations expressed on a
 230 lipid weight (lw) basis, and frequency of occurrence (FO) in the muscle of stranded striped dolphins. Values below LOQ
 231 are expressed as 0.

	Mean lw ($\mu\text{g g}^{-1}$)	Range ($\mu\text{g g}^{-1} \text{lw}$)	FO (%)
BPA	0.29	0-4.93	10
BPE	0.67	0-6.17	90
BPF	0.01	0-0.07	67
BPFL	0.05	0-0.44	47
BPP	0.03	0-0.94	7
BPZ	16.06	0.12-94.9	100
ΣBP	17.13	0.15-96.4	

DBP	7.24	0-26.59	73
DCHP	0.19	0-3.03	13
DEHP	6.85	0-48.9	63
DEP	1.62	0-8.04	80
DMP	0.08	0-1.38	38
DNOP	0.09	0-2.1	7
Σ PAE	16.08	0.01-80.29	

232



233

234 Figure 2: Relative contribution of the detected BP and PAE compounds split by sex, maturation stage, and sampling period.


235 Results of the comparison of BP and PAE concentrations between sexes, mature and immature
 236 individuals, and sampling periods are shown in table 3. No significant difference was detected between
 237 sexes for individual BPs or PAEs. However, the overall concentration of PAEs was significantly higher
 238 ($p = 0.024$, Kruskal-Wallis test) in samples from males than in those from females. No significant
 239 difference was detected between the concentrations of BPs or PAEs in mature and immature dolphins,
 240 with the exception of DMP, whose concentrations were significantly higher ($p = 0.023$, Kruskal-Wallis
 241 test) in immature individuals than in adults. Finally, a significant positive correlation between the

242 concentration of BPFL and the period of sampling ($\rho = 0.46, p = 0.024$), and significant negative
 243 correlations between the concentrations of BPE, DEP, and DMP and the period of sampling were
 244 detected ($\rho = -0.49, p = 0.016, \rho = -0.494, p = 0.014; \rho = -0.708, p < 0.001$, respectively) (Spearman's
 245 rho test, Table 3, Fig. 3).

246 Table 3. Results of the statistic tests performed to compare the concentrations of BPs and PAEs between males and females,
 247 mature and immature individuals, and the three sampling periods. The number of samples considered for each test is
 248 indicated in brackets. K-W refers to Kruskal-Wallis test. Significant results ($p < 0.05$) are marked with an asterisk (*).

	Sex (24 samples)		Maturation Stage (15 samples)		Sampling period (24 samples)		
	K-W chi-squared	<i>p</i> -value	K-W chi-squared	<i>p</i> -value	Spearmans' S	Spearmans' rho	<i>p</i> -value
BPA	2.087	0.1486	0.15873	0.6903	1910.3	0.169422	0.4287
BPE	0.27047	0.603	3.5619	0.05912	3422.3	-0.487936	0.01557*
BPF	0.65906	0.4169	3.0247	0.08201	3022	-0.3139107	0.1352
BPFL	0.15862	0.6904	1.2763	0.2586	1245.2	0.4585894	0.0242*
BPP	1	0.3173	1.5	0.2207	2336.8	-0.01599233	0.9409
BPZ	0.65333	0.4189	2	0.1573	3056	-0.328701	0.1168
ΣBP	0.40333	0.5254	2	0.1573	3007.2	-0.3074647	0.1439
DBP	1.0967	0.295	0.057613	0.8103	2395.2	-0.04140386	0.8477
DCHP	0.4943	0.482	1.5	0.2207	2671.3	-0.1614263	0.4511
DEHP	2.3774	0.1231	0.12963	0.7188	2376.3	-0.03319363	0.8776
DEP	3.4433	0.06351	0.031818	0.8584	3435.8	-0.4938182	0.01419*
DMP	0.14256	0.7057	5.1734	0.02294*	3928.4	-0.7079887	0.0001086*
DNOP	2.087	0.1486	0.66667	0.4142	2292.3	0.003369187	0.9875
ΣPAE	5.07	0.02434*	0	1	2489	-0.08217524	0.7027

249



251

252 Figure 3: Boxplots showing the significant correlations between BPs (BPFL and BPE, top graphs) and PAEs (DEP and
 253 DMP, bottom graphs) concentrations detected in samples from adult individuals and the period of sampling. Boxes show
 254 the first and the third quartiles, horizontal lines represent the median, and whiskers cover extreme values within the 1.5
 255 IQR value. Outliers are marked as dots.

256 4. Discussion

257 In this study, we assess the occurrence and concentration of BPs and PAEs in striped dolphins
 258 that stranded in the Western Mediterranean Sea across the 30-year period between 1990 and 2018, and
 259 provide the first record, to our knowledge, of the presence of a number of these compounds in the
 260 muscular tissue of cetaceans. We discuss our results in light of the scarce literature available on the
 261 concentration of these contaminants in the marine environment, and specifically on marine mammals.
 262 In the published literature, summarised in Table 4, these pollutants are assessed in cetaceans or marine
 263 mammals through a variety of tissues, including muscle, liver, blubber, skin, plasma and urine, which

264 only allows a speculative discussion of the different values observed by other studies. Taking the
265 limitations implied by the relatively low number of samples analysed into account, we also consider
266 the potential influence of factors liable of influencing the detected concentrations, such as sex,
267 maturation stage, and year of death of individuals. As information regarding the cause of death was
268 not available for all individuals, and most of the sampled dolphins were killed by the morbillivirus
269 epizootic, results could not be related to any potential toxic or detrimental effect to the health status of
270 dolphins.

271 Table 4. Concentrations of BP and PAE compounds assessed in the tissues of marine mammal species. All concentrations
272 are expressed in ng g^{-1} , with the exception of those in urine samples, which are expressed in ng mL^{-1} . The matrix in which
273 the compounds analysed, along with the number of samples and the specification on the basis on which concentrations are
274 calculated (dw: dry weight, ww: wet weight, lw: lipid weight) are indicated in the first column; the geographic area of each
275 study, along with the corresponding reference are indicated in the last column.

Species	#samples, matrix	BBP	DBP	DCHP	DEHP	MEHP	DEP	MEP	DMP	DNOP	BPA	BPF	BPS	Area, reference
Fin whale (<i>Balaenoptera physalus</i>)	5, Blubber (ww/lw) 30/10, skin + blubber biopsies (ww) 3, skin (dw) 6, Blubber (ww) 31, Muscle (dw)					58/177 55; 40								Italy ^{a,b} Mediterranean Sea/Sea of Cortez ^c Mediterranean Sea ^d Svalbard ^e Iceland ^f
Blue whale (<i>Balaenoptera musculus</i>)	7, Blubber (ww)	<10		<10	20		<37			<10				Svalbard ^e
Bowhead whale (<i>Balaena mysticetus</i>)	5, Blubber (ww)	<10		<10	n.a		<37			<10				Svalbard ^e
Minke whale (<i>Balaenoptera acutorostrata</i>)	2, Liver (ww)	29.7	10.7		86.2		15.1		2.5	6.7				Greenland ^g
Striped dolphin (<i>Stenella coeruleoalba</i>)	2, Skin (dw) 5, Muscle 30, Muscle (lw)				21460	1720								Mediterranean Sea ^d Canaries ^h This study
<i>Stenella spp.</i>	5, Blubber (dw)						70				<10			USA ⁱ
Risso's dolphin (<i>Grampus griseus</i>)	1, Skin (dw) 2, Muscle				1130	464		42			163	89	37	Mediterranean Sea ^d Canaries ^h
Bottlenose dolphin (<i>Tursiops truncatus</i>)	1, Skin (dw) 2, Muscle 46, Blubber (dw) 17/51/50, Urine (ng mL ⁻¹)		413		26068	1770		0			201	110	10	Mediterranean Sea ^d Canaries ^h U.S. ¹ U.S., ^j U.S. ^{k,1} U.S. ^m
Pilot whale (<i>Globicephala melas</i>)	3, Liver (ww)	28	15.1		103.6		23.2		2.50	5.6				Faroe ^f
Short-finned pilot whale (<i>Globicephala macrorhynchus</i>)	1, Muscle		969		335		0				732	81	12	Canaries ^h
Harbour porpoise (<i>Phocoena phocoena</i>)	100, Liver (ww)							5.99						Norway ⁿ

Pygmy sperm whale (<i>Kogia breviceps</i>)	1, Muscle	664		102	0		181	47	213	Canaries ^h
<i>Kogia spp</i>	22, Blubber (dw)				170		1650			U.S. ⁱ
Fraser's dolphin (<i>Lagenodelphis hosei</i>)	1, Muscle	552		330	98		72	67	28	Canaries ^h
White-beaked dolphin (<i>Lagenorhynchus albirostris</i>)	1, Blubber (dw)				1380 0		397400			U.S. ⁱ
Melon-headed whale (<i>Peponocephala electra</i>)	3, Blubber (dw)				500		<10			U.S. ⁱ
Polar bear (<i>Ursus maritimus</i>)	12, Fat tissue (ww)	<10	<10	n.a.	<37.5		<10			Svalbard ^e
	13, Plasma (ww)				<0.8	<0.8				Svalbard ^e
	2, Liver (ww)	31	12.8	142.6	22.7	51.4	22			Greenland ^f
Ringed seal (<i>Phoca hispida</i>)	4, Liver (ww)	128.6	7.3	117.2	21	3.7	7.6			Greenland ^f

276 ^aFossi et al (2012);

277 ^bFossi et al (2014);

278 ^cFossi et al (2016);

279 ^dBaini et al (2017) (also analysed MBzP, MBP, BBzP, DIOIP);

280 ^eRoutti et al (2021) (also assessed DMPP, DHxP, and DnNP, DnBP, DiBP, DiNP and DiDP in blubber of cetaceans and adipose tissue of polar bear; and

281 MMP, MEOHP, MiBP, MnBP, MBzB, MEHHP in the blood plasma of polar bear);

282 ^fGarcia-Garin et al (2022),

283 ^gVorkamp et al (2004),

284 ^hMontoto Martinez et al (2021),

285 ⁱPage-Karjian et al (2020) (they also analysed muscle samples from *Mesoplodon spp*, *G. griseus* and *Z. cavirostris* for BPA and DEP, which were not

286 detected);

287 ^jHart et al (2018) (also analysed MEOHP and MBP);

288 ^kHart et al, 2020 (also assessed MBP);

289 ^lDziobak et al (2021) (also assessed MBP, MBzP, MiBP, MEOHP, MEHHP, MMP),

290 ^mDziobak et al (2022);

291 ⁿRian et al (2020) (also assessed mIBP, mBP, PA, mMP, mHxP, mEHHP, mNP, mOP, mEOHP, mHpP, mBzP, mDP).

292

293 4.1 BP concentrations

294 Since its commercialization in the 1950s, BPA has been the most widely used BP during decades
295 (Staples et al, 1998). However, after several studies alerted about its adverse effects on reproduction,
296 development, and neural networks, as well as on the cardiovascular, metabolic, and immune systems
297 of living organisms, its use was discontinued for most applications in the European Union (EC,
298 2011/8/EU), as well as in Canada, United States, Japan, Brazil, China and Malaysia (Akhbarizadeh et
299 al, 2020). Industry soon after began to develop alternative compounds to replace it, and in 2016, 16
300 BP analogues were industrially applied, of which BPF, BPS and BPAF were the most widely used
301 (Chen et al, 2016). Due to their increased production and use, several of these compounds were soon
302 detected in foodstuff, household dust, and PCPs, and high concentrations were detected in both the
303 environment and human tissues (Pelch et al, 2019; Park et al, 2020).

304 In this study, we detected 6 of the 10 BPs analysed, which, in decreasing order of frequency, were:
305 BPZ, BPE, BPF, BPFL, BPA, BPP, and, in decreasing order of concentration: BPZ, BPE, BPA, BPFL,
306 BPP, BPF. Diet is the main intake route for marine mammals, which can ingest BPs either indirectly
307 through contaminated prey (*i.e.*, fish or molluscs) and water, or directly from accidentally ingested
308 pieces of plastics, microplastics, or other materials that may release BPs during their degradation
309 (Nehring et al, 2017). In smaller proportions, marine mammals may also be exposed to these
310 contaminants through the inhalation of polluted air (Nehring et al, 2017).

311 According to our results, BPZ was the most common and most concentrated compound (mean 16.06,
312 range 0.12-94.9 $\mu\text{g g}^{-1}$ lw), followed by BPE (mean 0.67, range 0-6.17 $\mu\text{g g}^{-1}$ lw), which was detected
313 in 90% of the samples, at slightly higher concentrations than the other four BP compounds detected.
314 Unfortunately, very little is known about the environmental concentration of both compounds. BPZ is
315 a component widely used in European countries for the manufacture of epoxy resins and polycarbonate
316 plastics, and has been detected in PCP (0.71-0.74 ng g^{-1}), foodstuff (0.03-0.08 $\mu\text{g g}^{-1}$), soft drinks (0.09
317 $\mu\text{g L}^{-1}$) and human biomonitoring samples (0.06 $\mu\text{g L}^{-1}$), as well as in indoor dust and sludge (at 0.001-

318 0.5, and up to 1.8 ng g⁻¹, respectively, Chen et al, 2016), which may explain its higher concentration
319 in all the analysed samples. On the other hand, BPE is used for polycarbonate and cyanate and epoxy
320 resins, and recent studies have shown that it is commonly detected in source and drinking water (0.83
321 ng L⁻¹, Zheng et al, 2019), as well as in municipal sewage sludge (0.980 µg g⁻¹, Sauer et al, 2021).
322 Moreover, according to Danzl et al (2009), BPE in seawater is slightly more resistant to biodegradation
323 than other compounds such as BPA, BPF or BPP, which may justify its presence in 90% of the samples
324 analysed.

325 The other four BPs were not detected in all the samples, as their lowest concentrations were below the
326 LOQ. Mean and maximum concentrations were below 1 µg g⁻¹ lw for all these compounds, except for
327 BPA, which was only detected in three individuals, but whose maximum concentration was one order
328 of magnitude higher than BPP, BPFL and two orders of magnitude higher than BPF. Indeed, although
329 in Europe the use of BPA has been banned from several products, this compound is still widely
330 produced and has been found in surface waters (0.25 µg L⁻¹, Zheng et al, 2019), sewage sludge (0.12–
331 6.50 µg g⁻¹, Sauer et al, 2021), bivalves, holothurians and fish from the Balearic Island, in the Western
332 Mediterranean Sea (0.29; 0.45 and 0.32 µg g⁻¹ ww, respectively, Rios-Fuster et al, 2022), samples of
333 herring gull feathers, Baltic grey seal fur, and human hair (0.1, 0.07 and 0.34 µg g⁻¹, respectively,
334 Nehring et al, 2017), as well as in the few cetacean species analysed (Page-Karjian et al, 2020,
335 Montoto-Martinez et al, 2021), which also included 5 striped dolphins (Table 4). Ozhan and Kocaman
336 (2019) detected BPA levels in the highly anthropized waters of the Black Sea, Bosphorus, and Sea of
337 Marmara in a range of 8.85–14.76 µg L⁻¹. However, BPA is rapidly degraded in seawater (Staples,
338 1998), where it can last only up to 30 days (Kang et al, 2007) and, similarly to other BPs, its excretion
339 rate in mammals is high (Kang et al, 2006), which possibly explains its low presence in the dolphins
340 analysed (*i.e.*, 3 out of 30).

341 BPFL was detected in almost half of the samples, at lower concentrations than all the previously
342 mentioned compounds. This compound is a novel BPA substitute used in the plastics industry as an
343 organic (fluorinate) synthesis intermediate (Zhang et al, 2021), and extensively applied to synthesize
344 polyesters, epoxy resins, polycarbonates, and in consumer products that are labelled BPA-free (Özkan-
345 Kotiloğlu et al, 2022). Aside from its detection in animal feeds (0.24 ng g⁻¹ dw, Wang et al, 2021) and
346 in environmental samples of water bodies and sediments (0.056-0.069 ng L⁻¹, Jin and Zhu, 2016), the
347 information regarding this compound is very limited. However, its increasing use may be the reason
348 of its presence in almost half of the sampled dolphins.

349 Finally, mean BPP and BPF concentrations were similarly low, but while BPP was detected only in 2
350 samples (7%), BPF was detected in almost 70% of them. Indeed, BPF is widely applied for the
351 manufacture of lacquers, varnishes, liners, adhesives plastics, and water pipes, as well as in dental
352 sealants, oral prosthetic devices, tissue substitutes and coatings for food packaging (Chen et al, 2016).
353 Both compounds have been detected in foodstuff (*e.g.*, 0.24 µg g⁻¹ BPP, and 1.13 µg g⁻¹ BPF,
354 respectively, in samples from mustard and ginger, Liao and Kannan, 2013), household dust (up to 2 ng
355 g⁻¹ BPP, and up to 1 µg g⁻¹ BPF, respectively, Chen et al, 2016), and PCPs (0.79-0.99 ng g⁻¹ BPP, and
356 0.820.99 µg g⁻¹ BPF, respectively, Chen et al, 2016), and BPF has also been detected in sewage sludge
357 (0-3.1 µg g⁻¹, Sauer et al, 2021), surface water (0.2 µg L⁻¹, Chen et al, 2016, 0.002 µg L⁻¹, Zheng et al,
358 2019; 0-0.075 µg L⁻¹, Sauer et al, 2021), marine fish (0.005 µg g⁻¹ ww, 0.001 µg g⁻¹ dw, Akhbarizadeh
359 et al, 2020), in bivalves, holothurians and fish in the Cabrera archipelago, not far from our sampling
360 area (0.29, 0.1 and 0.1 µg g⁻¹ ww, respectively, Rios-Fuster et al, 2022) and in the cetacean species
361 analysed by Montoto-Martinez et al (2021, Table 4). According to Danzl et al (2009), BPF in seawater
362 is more biodegradable than BPP, which may explain its lower concentrations, while its higher
363 frequency of detection may be justified by its wider use and applications. On the other hand, BPP is
364 one of the few BP analogues with a log K_{ow} value higher than 5 (Chen et al, 2016), which indicates a

365 tendency to accumulate in tissues, and thus be a liable factor contributing to the relatively higher
366 concentration found in the two samples where it was detected.

367 The scarcity of literature regarding environmental concentrations of BP compounds, especially in the
368 Mediterranean waters, as well as the almost totally lacking data on their concentration in marine
369 wildlife, and particularly cetaceans, prevents any comparison or further speculation on the
370 environmental factors determining the concentrations detected. Furthermore, the variety of tissues
371 analysed within the available studies on marine mammals, only allows a qualitative comparison
372 between the concentrations detected in the different cetacean species or geographical regions.
373 However, even at low concentrations, the presence of several BP compounds in often a large proportion
374 of the individuals analysed indicates that Mediterranean dolphins are somehow exposed to these
375 contaminants. Some of the BPs detected in the muscle of striped dolphins were also detected in the
376 Mediterranean waters, and bivalve, holothurians, and fish (Errico et al, 2017; Ozhan and Kocaman,
377 2019; Rios-Fuster et al, 2022) from the NW Mediterranean Sea. Interestingly, the concentrations found
378 by these authors were generally higher than those found here. The scattered nature of these data and
379 the different measurement methods applied prevent from assessing the potential occurrence of
380 bioconcentration or biomagnification of these pollutants. However, despite tissue levels of pollutants
381 may vary depending on their chemical properties and the species/individual metabolism, the presence
382 of BPs in the dolphin tissues can only be attributed to the pollution of the Mediterranean seawater,
383 either from plastic waste, industrial and domestic effluents, landfill wastes, and/or atmospheric
384 deposition.

385 4.2 PAE concentrations

386 DEHP was the first PAE to be synthesized, in the 1930s, for its application as a plasticizer for the
387 newly developed hard PVC (Graham, 1973). Since then, the industry of PAEs grew and diversified
388 along with their applications, which range from home furnishings to transportation, construction,

389 apparel, food, PCP, and the medical industry (Graham, 1973). However, research demonstrated that
390 also PAEs can leach from plastics to the environment and that the exposure to these chemicals can
391 produce toxic effects (Warner and Flaws, 2018). Several PAE compounds, such as DEHP, DBP and
392 DIBP, have been included as hazardous for the environment in the European REACH regulation to
393 protect human health from chemical risks (EU 1907/2006), and their use has been banned in the EU
394 from a number of applications (Net et al, 2015). They have also been included, along with a few others,
395 in the list of priority hazardous pollutants in China and the USA (CPSC, 2018), and are already subject
396 to some restrictions in the USA. Although the use of PAE has declined, they still make up over 55%
397 of world plasticizers, which are consumed at a growing rate of 3.5% per year (IHSMARKIT, 2020) and
398 are thus expected to end up in large quantities in the marine environment.

399 The analysis of PAEs is challenging, because the sample manipulation and the use of organic solvent
400 may lead to possible contamination of the samples and/or of the laboratory environment (Panio et al,
401 2020). In this study, we detected 6 of the 13 PAEs analysed, which in decreasing order of frequency,
402 were: DEP, DBP, DEHP, DMP, DCHP, DNOP, and, in decreasing order of concentration: DBP, DEHP,
403 DEP, DCHP, DNOP, DMP. Marine mammals are exposed to PAEs mainly through the ingestion of
404 materials that may release these compounds once they are made bioavailable by digestive surfactants
405 (Teuten et al, 2007), or through contaminated food, water and air. However, as PAEs are rapidly
406 degraded in the environment (Xie et al, 2007) and metabolised by living organisms (Mackintosh et al,
407 2004), and they do not appear to biomagnify in aquatic food webs (Mackintosh et al, 2004; Routti et
408 al, 2021), exposure through prey ingestion is expected to be limited (Hart et al, 2020).

409 In the NW Mediterranean Sea, PAE concentrations have been found to range from 100 to 527 ng L⁻¹
410 in seawater, from 12 to 610 ng g⁻¹ dw in sediment, and from 0.9 to 47 µg g⁻¹ dw in zooplankton
411 (Schmidt et al, 2021). Consistently with Garcia-Garin et al (2022) and Montoto-Martínez et al (2022),
412 the only two studies assessing PAEs in cetacean muscle, our results showed a high variability of the

413 overall individual PAE concentrations, and identified DEHP, DEP and DBP as the most common and
414 most concentrated compounds. Indeed, these compounds are among the most frequently detected PAEs
415 in surface waters (Net et al, 2015; Hart et al, 2020).

416 DEHP has been detected in the Mediterranean waters of the Gulf of Marseille at concentrations ranging
417 from 15.8 to 923.8 ng L⁻¹ (Paluselli et al, 2018), and was the more abundant PAE compound observed
418 in the surface sediments across the Gulf of Lion (Alkan et al, 2021). It was detected in the gonads of
419 sea urchins from Sardinia (17 ng g⁻¹, Raguso et al, 2022), in bivalves, holothurians and fish from the
420 Balearic Islands (mean 2.58, 1.48 and 0.88 µg g⁻¹ ww, respectively, Rios-Fuster et al, 2022) and in
421 biopsies taken from cetaceans (including two striped dolphins) in the Pelagos Sanctuary (Baini et al,
422 2016), where high concentrations of this compound and its metabolites were also detected in marine
423 neuston and plankton samples (Fossi et al, 2012; Baini et al, 2016). DEHP was also detected in the
424 cetacean samples analysed by Vorkamp et al (2004), Routti et al (2021), and Montoto-Martinez et al
425 (2021) in the North Atlantic Ocean (Table 4). As this compound is the most widely used to increase
426 flexibility of plastics (especially PVC), representing in 2015 over 35% of the global plasticizer market
427 (Routti et al, 2021), it is also the most studied, along with its metabolites. Given that DEHP has low
428 solubility in water and preferentially partitions to suspended particles and sediments, it is prone to
429 bioaccumulate in lipid-rich tissues and to biomagnify along marine food webs, the main exposure for
430 marine mammals happens through the food (Dziobak et al, 2021; Routti et al, 2021). These combined
431 factors may explain why this compound was detected at the highest concentrations in our samples.

432 On the other hand, studies on low molecular weight PAEs, such as DBP and DEP, in the tissues of
433 aquatic organisms are scarcer. Indeed, these compounds have been reported to accumulate at low level
434 in aquatic species (Net et al, 2015), and their half-life is short both in the marine environment (*i.e.*,
435 over 90% degradation within a week, Staples et al, 1997) and in fish tissues (Net et al, 2015). DEP was
436 detected in the waters of Marseille Bay at concentrations ranging from 3.3 to 50 ng L⁻¹ (Paluselli et al,

2018), in a small proportion of sediment samples from the Gulf of Lion (Alkan et al, 2021), in bivalves, holothurians and fish from the Balearic Islands (0.54, 0.49 and 0.17 $\mu\text{g g}^{-1}$ ww, respectively, Rios-Fuster et al, 2022), in 40% of the samples of cetacean muscle analysed by Montoto-Martínez et al (2021), as well as in the tissues of the dolphins analysed by Page-Karijan et al (2020) and in the muscle of the fin whales analysed by Garcia-Garin et al (2022). DBP was detected in the gonads of sea urchins from Sardinia (10 ng g^{-1} , Raguso et al, 2022), in bivalves, holothurians and fish from the Balearic Islands (0.78, 1.24, 0.72 $\mu\text{g g}^{-1}$ ww, respectively, Rios-Fuster et al, 2022), and in the samples of cetacean muscle analysed by Montoto-Martínez et al (2021) and Garcia-Garin et al (2022) (Table 4). The concentrations found in these studies were of comparable order of magnitude as those we detected, despite not all the cited studies analysed the same tissues, which prevents a rigorous comparison of concentrations, and the frequency of detection varied among studies and species, probably indicating that, while the metabolism of this compound is similar among cetacean species, its level of exposure varies between geographical areas. Its metabolite, MEP, was also detected in the samples of dolphin urine analysed by Hart et al (2018, 2020) and in the liver of harbour porpoises (Rian et al, 2020; Table 4).

DMP is also a low molecular weight PAE, which is used, along with DBP and DEP, as a component of industrial solvents, adhesives, waxes, inks, insecticides, PCPs and pharmaceutical products (Net et al, 2015). This compound was detected in the waters of the Marseille Bay at concentrations ranging between 0.8 and 11.9 ng L^{-1} (Paluselli et al, 2018) and in the sediments of the Gulf of Lion (Alkan et al, 2021). We detected this compound in half of our samples, at very low concentrations, similar to those detected in the liver of ringed seals and polar bears by Vorkamp et al (2004), and in the muscle of fin whales by Garcia-Garin et al (2022), who observed it in a higher proportion of samples (Table 4). Rian et al (2021) also detected the presence of its metabolite, mMP, in the liver of 69% of the 100 harbour porpoises they analysed. The apparent lower prevalence of this compound in our and other Mediterranean samples, as compared to those from other areas, is likely related to different rates of

462 DMP deposition in the Mediterranean Sea than in the Atlantic Ocean, which could be the result of
463 lower rates of production and use in the countries bordering the Mediterranean Sea.

464 Paluselli et al (2018) highlighted that the origin and aging of plastic materials may affect the
465 concentration of PAEs in both the seawater and the guts of marine organisms. Moreover, while
466 compounds like DEHP are denser than seawater and thus subject to sinking, phthalates can also leach
467 from positively buoyant polymers such as polyethylene (PE), polypropylene (PP), and polystyrene
468 (PS) (Paluselli et al, 2019). Thus, the highly variable concentrations detected, and the variability found
469 among studies may not only derive from different environmental deposition rates among compounds,
470 but also from different feeding habits of the cetacean species or of the individuals analysed (*i.e.*, pelagic
471 vs benthic), along with the intrinsic differences related to the assessment of these pollutants in different
472 tissues.

473 Finally, DCHP and DNOP were only detected in 4 and 2 dolphins analysed, respectively, at lower
474 concentrations than most other compounds. To our knowledge, the only study reporting these
475 compounds in a cetacean tissue is that from Garcia-Garin et al (2022), which detected DCHP in one
476 of the fin whale muscle samples analysed, while this compound was below the limit of quantifications
477 in the marine mammal samples analysed by Routti et al (2020) (Table 4). This result might indicate
478 that the exposure to these contaminants in the marine environment is relatively low, but the scarce
479 literature available prevents from relating the levels found to the local uses and production rates.

480 4.3 Influence of biological traits

481 *Sex*

482 None of the single BP or PAE compounds analysed was detected at significantly different
483 concentrations in samples from adult males and females. This result is consistent with the few
484 publications analysing these compounds in mammals of different sex, including marine mammals and

485 clinical studies in humans. Published research is consistent in stating that these compounds do not
486 bioaccumulate in tissues; on the contrary, they are largely excreted through the urine, faeces, bile, hair
487 or fur (Kang et al, 2006; Chen et al, 2016; Nehring et al, 2017). Dziobak et al (2021) and Rian et al
488 (2020) also showed that the concentrations of phthalate metabolites in the urine and liver of odontocete
489 species do not differ between sexes. However, the metabolism of each compound may vary according
490 to its chemical characteristics, as well as to individual variables and habits like the diet (Akhbarizadeh
491 et al, 2020). In humans, diet, lifestyle, and other individual habits, were proven to be the main factors
492 determining different metabolite concentrations in males and females (*e.g.*, Hartmann et al, 2015;
493 Huang et al, 2015; Silva et al, 2004). Moreover, the metabolism of female mammals often differs from
494 that of males as a consequence of their hormone cycle, pregnancy, and lactation, which can affect the
495 concentration of contaminants in their tissues (Aguilar et al, 1999). Indeed, laboratory studies show
496 that female rats metabolize BPA faster than males, and different BPA concentrations have been found
497 in the human serum, as a consequence of differences in the androgen-related metabolism (Kang et al,
498 2007). These factors may also explain why, although the concentration of none of the compounds was
499 significantly different between sexes, the summation of all PAE compound concentrations was overall
500 higher in samples from males than in those from female dolphins. However, given the limited number
501 of samples used to perform this analysis, this result is to be interpreted with caution.

502 *Maturation stage*

503 In contrast with certain organic compounds such as PCBs and chlorinated pesticides that show age-
504 related accumulation in cetaceans (*e.g.*, Tanabe et al, 1987), neither BPs nor PAEs tend to
505 bioaccumulate in animal tissues. Indeed, most of the few studies that assessed potential variations of
506 these compounds with the size or age of marine mammals did not detect any specific trend (*e.g.*,
507 Garcia-Garin et al, 2022, Montoto-Martinez et al, 2021). Our results showed no significant difference
508 between sexually mature and immature dolphins, with the exception of DMP, whose concentrations

509 were higher in samples from immature individuals. Indeed, out of the 15 samples analysed from the
510 2014-2018 period, this compound was only detected in three immature females, while its concentration
511 was below the LOQ in all the remaining samples from this period. Although the statistical significance
512 of the test is likely weak due to the limited number of samples, other authors have shown that adult
513 mammals tend to have a more efficient metabolism of some contaminants than immature specimens.
514 Indeed, Rian et al (2020) found that the concentration of a common metabolite of PAE, the phthalic
515 acid (PA), decreased with body length in dolphins, and Hart et al (2021) found that mean
516 concentrations of the overall PAE analysed and of MEHP were higher in calves than in adult dolphins,
517 possibly reflecting a more efficient biotransformation process of these compounds in adult individuals.
518 Maternal transfer during pregnancy and through lactation may also increase pollutant concentrations
519 in newborn or calf mammals (*e.g.*, Borrell et al 1995; Addison and Brodie, 1977; Borrell and Aguilar
520 2005; Fouin et al, 2012; Barbosa et al, 2018); however, this phenomenon has only been reported for
521 human milk and the more lipophilic phthalate diesters and metabolites, such as DEHP/MEHP, while
522 information on this regard for marine mammals is absent. As both BPs and PAEs are rapidly
523 metabolized and do not bioaccumulate in tissues, we believe unlikely that this process is relevant to
524 explain our results. Rather, the exposure to chemicals between adult and immature dolphins may be
525 influenced by dissimilarities in their diet, physiology and/or activity levels, as suggested by Hart et al
526 (2018) and Rian et al (2020).

527 4.3 Influence of year of death

528 The increased demand of plastic, and plastic-related materials, has led the production of plastic
529 additives such as BP and PAEs to raise exponentially during the past decades (Net et al, 2015). Indeed,
530 PAEs production reached 8 million tons in 2011, and in 2020 PAEs made up 55% of the world
531 consumption of plasticizers (IHSMARKIT, 2021). Given such a wider production, a higher amount of
532 these compounds is expected to reach the seas, either directly from spills during the production process,

533 or in the form of leachates dispersed from the large quantities of plastic or plastic-related items that
534 increasingly pollute the seas (Jambeck et al., 2015). However, the high liability of these compounds
535 limits their environmental persistence, and, indeed, according to several authors, atmospheric PAE
536 concentration in the North Atlantic did not change substantially from the 1970s (Giam et al, 1978;
537 Atlas and Giam, 1981; Bohlin-Nizzetto et al, 2018). Consistently, Garcia-Garin et al (2022) did not
538 detect any significant temporal trend between 1986 and 2015 in the concentration of PAE compounds
539 in samples of fin whale muscle.

540 Although our temporal analyses were based only on 24 samples and thus the observed trends must be
541 interpreted with caution, in line with the above findings, the concentrations of most compounds did
542 not vary significantly during the three periods (1990, 2007-2008 and 2014-2018), with the exception
543 of two BP and two PAE compounds. Interestingly, the concentration of three (BPE, DEP, DMP)
544 compounds showed a significant decreasing temporal trend, while concentrations of BPFL showed a
545 significant increase in the most recent samples. Indeed, the concentrations of BPFL were higher than
546 the LOQ only in one sample from the 90s, and in two samples from the 2007-2008 period, while they
547 were above the limit in most of samples from the 2014-2018 period. This result might be explained by
548 the relative novelty of this compound, which only began to be produced and used massively as a
549 replacement of BPA in the so-called “BPA-free” products (Özkan-Kotiloğlu et al, 2022), once the use
550 of BPA as additive was banned in numerous products in several countries and in the EU in the early
551 2010s.

552 BPE, in turn, was detected in 90% of samples, independently of the period, but its concentrations
553 showed a slightly decreasing trend, with the lowest concentrations occurring in the most recent samples
554 and the peak of concentrations detected in two samples from the 2007-2008 period. The slightly higher
555 persistence of this compound (Danzl et al, 2009) likely justifies its detection in most samples, however,
556 as little is known on its trends of production and use, we may only assume that this compound,

557 commonly used in cyanate resins (Zheng et al, 2019), has been recently replaced by other analogues
558 and is reaching the Mediterranean waters in reducing quantities.

559 Similarly, DMP was detected, although at very low concentrations, in all samples from the 1990s, at
560 slightly higher concentrations in only three samples from the 2007-2008 period, and again at very low
561 concentrations in only three samples from immature individuals from the 2014-2018 period. Finally,
562 DEP was detected in a consistent proportion of the individuals analysed from the three periods, but its
563 concentrations were substantially and significantly lower in the samples from the 2014-2018 period
564 than in those from the two earlier periods. Both compounds have been included in the lists of priority
565 pollutants in USA, EU and China (Net et al, 2015), which may imply that the rates at which they are
566 produced and used are every time more restricted. Indeed, while the market of high molecular weight
567 PAEs has kept generally steady in Europe, that of low molecular weight PAEs such as DMP and DEP
568 has been decreasing since the beginning of the 2000s: in 2017, low molecular weight PAEs still
569 represented 40% of the global plasticizers market, but only 11% of the European one (IHSMARKIT,
570 2021). The limited number of samples hampers a proper interpretation of results also in this case.
571 However, given the low persistence of these compounds in the marine environment, in contrast to the
572 behaviour of persistent organic pollutants that exhibit very slow decline in concentrations in cetacean
573 tissues (Loganathan et al, 1990), the lower frequency and concentrations observed in the most recent
574 samples might reflect a recent shift in the commercial production of these PAEs.

575 Indeed, our sampling period spans almost three decades, during which there have been significant
576 changes in the production and use of these chemicals. However, the observed temporal variability may
577 also be the result of the degradation of certain compounds (biodegradation or photodegradation), while
578 the non-detection of certain other compounds could be related to the presence of possible unknown
579 metabolites.

580 4.4 Implications for the Mediterranean striped dolphin population

581 The Mediterranean striped dolphin is currently classified by the IUCN red list under the “least concern”
582 category (IUCN, 2019). However, this population has experienced a series of mortality events, during
583 1990-1992, and 2006-2007, related with morbillivirus epizootics (Aguilar and Raga, 1993; Domingo
584 et al, 1990; 1995; Keck et al, 2010), and its individuals are known to carry in their tissues extremely
585 high levels of pollutants (Aguilar and Borrell, 1994; Aguilar, 2000). While, in principle, there should
586 be no direct correlation between contaminant burden and morbillivirus, it might be possible that
587 individuals suffering from heavy contamination have a weaker immune system and thus are more
588 susceptible to contract a virus. Unfortunately, this possibility could not be addressed here; however,
589 the concentrations we detected were relatively low and most probably did not have any health effect
590 on the dolphins. Despite the tissue concentration of organochlorine pollutants and mercury in striped
591 dolphins have shown decreasing trends (Aguilar and Borrell, 2005; Borrell et al, 2014), and the
592 concentration of plastics ingested by Mediterranean striped dolphins is generally low (Novillo et al,
593 2020), the main threat for this species in the Mediterranean is still associated with marine pollution
594 (IUCN, 2019). Indeed, the extreme anthropization of the basin causes these animals to be constantly
595 and increasingly exposed to a mixture of pollutants and subject to a variety of stress factors (Marsili
596 et al, 2018; Aznar-Alemany et al, 2021). Although the detected concentrations do not appear to be
597 concerningly high, and neither BPs nor PAEs seem to bioaccumulate in the tissues of dolphins, the
598 long-term exposure to these chemicals, combined with other multiple stressors, may potentially
599 produce adverse effects to long living animals such as cetaceans (Hart et al, 2020, Dziobaak et al,
600 2022).

601 **Conclusions**

602 Our results provide the first assessment of a number of BP analogues and PAE compounds in the
603 muscle of an odontocete cetacean, and indicate that Mediterranean dolphins are exposed to these
604 pollutants. While no significant relation was detected between the concentration of these compounds

605 and the dolphins' biological traits (sex and sexual maturation), some temporal trends emerged. The
606 limited number of samples available (n=30) allows only a partial interpretation of results, however, the
607 observed trends might reflect a shift in the commercial production and use of some BP and PAE
608 compounds, and the detected values contribute to enlarge the limited knowledge base on their levels
609 in the marine fauna. Given the increasing rate of production and use of these compounds, and the
610 scarce information regarding their presence in the marine environment, it is of utter importance to
611 continue investigating their levels in sensitive marine fauna like cetaceans, along with their sources,
612 intake routes, metabolism, and potential long-term effects.

613 **Acknowledgements**

614 Authors are grateful to the FSE “El FSE invierte en tu futuro” and the Fundació Montcelimar –
615 Universitat de Barcelona for supporting O. Garcia-Garin with a Ph.D. FPU scholarship (FPU17/00073)
616 and a mobility grant.

617 **References**

- 618 Addison, R.F., Brodie, P.F. 1977. Organochlorine Residues in Maternal Blubber, Milk, and Pup
619 Blubber from Grey Seals (*Halichoerus grypus*) from Sable Island, Nova Scotia. Journal of the
620 Fisheries Research Board of Canada 34,7, 937–941. Doi:10.1139/f77-146.
- 621 Aguilar, A. 2000. Population biology, conservation threats and status of Mediterranean striped
622 dolphins (*Stenella coeruleoalba*). Journal of Cetacean Research and Management 2,1,17-26.
- 623 Aguilar, A., and Borrell, A. 1994. Abnormally high polychlorinated biphenyl levels in striped dolphins
624 (*Stenella coeruleoalba*) affected by the 1990–1992 Mediterranean epizootic. Science of the Total
625 Environment 154,2-3, 237-247.

- 626 Aguilar, A., Borrell, A., Pastor, T. 1999. Biological factors affecting variability of persistent pollutant
627 levels in cetaceans. *Journal of Cetacean Research and Management* 83-116.
- 628 Aguilar, A., Borrell, A., 2005. DDT and PCB reduction in the western Mediterranean from 1987 to
629 2002, as shown by levels in striped dolphins (*Stenella coeruleoalba*). *Marine Environmental*
630 *Research* 59, 4, 391-404. Doi:10.1016/j.marenvres.2004.06.004
- 631 Aguilar, A., Raga, J.A. 1993. The striped dolphin epizootic in the Mediterranean Sea. *Ambio* 22, 8,
632 524-528.
- 633 Akhbarizadeh, R., Moore, F., Monteiro, C., Fernandes, J.O., Cunha, S.C. 2020. Occurrence, trophic
634 transfer, and health risk assessment of bisphenol analogues in seafood from the Persian Gulf.
635 *Marine Pollution Bulletin* 154, 111036, <https://doi.org/10.1016/j.marpolbul.2020.111036>.
- 636 Alkan, N., Alkan, A., Castro-Jiménez, J., Royer, F., Papillon, L., Ourgaud, M., Sempéré, R. 2021.
637 Environmental occurrence of phthalate and organophosphate esters in sediments across the Gulf
638 of Lion (NW Mediterranean Sea). *Science of The Total Environment* 760, 143412,
639 <https://doi.org/10.1016/j.scitotenv.2020.143412>.
- 640 Atlas, E., Giam, C.S. 1981. Global Transport of Organic Pollutants: Ambient Concentrations in the
641 Remote Marine Atmosphere. *Science* 211, 163–165.
- 642 Aznar-Aleman, O., Sala, B., Jobst, K.J., Reiner, E.J., Borrell, A., Aguilar, A., Eljarrat, E. 2021.
643 Temporal trends of halogenated and organophosphate contaminants in striped dolphins from the
644 Mediterranean Sea. *Science of The Total Environment* 753, 142205.
- 645 Bani, M., Martellini, T., Cincinelli, A., Campani, T., Minutoli, R., Panti, C., Finoia, M.G., Fossi, M.C.
646 2017. First detection of seven phthalate esters (PAEs) as plastic tracers in superficial

- 647 neustonic/planktonic samples and cetacean blubber. *Analytical Methods* 9,9, 1512–1520.
648 <https://doi.org/10.1039/c6ay02674e>
- 649 Barbosa, A.P.M., Méndez-Fernandez, P., Dias, P.S., Santos, M.C.O., Taniguchi, S., Montone, R.C.
650 2018. Transplacental transfer of persistent organic pollutants in La Plata dolphins (*Pontoporia*
651 *blainvillei*; Cetartiodactyla, Pontoporiidae). *Science of The Total Environment* 1,631-632,239-
652 245. Doi: 10.1016/j.scitotenv.2018.02.325.
- 653 Basak, S., Das, M.K., Duttaroy A.K. 2020. Plastics derived endocrine-disrupting compounds and their
654 effects on early development. *Birth Defects Research* 112, 1308-1325, 10.1002/bdr2.1741.
- 655 Ben Sghaier, R., Net, S., Ghorbel-Abid, I, Bessadok, S., Le Coz, M., Ben Hassan-Chehimi, D.,
656 Trabelsi-Ayadi, M., Tackx, M., Ouddane, B. 2017. Simultaneous detection of 13 Endocrine
657 Disrupting Chemicals in Water by a combined of SPE-BSTFA Derivatization and GC-MS in
658 transboundary rivers (France-Belgium). *Water, Air, & Soil Pollution* 228, 2. DOI
659 10.1007/s11270-016-3195-2.
- 660 Benjamin, S., Masai, E., Kamimura, N., Takahashi, K., Anderson, R.C., Faisal, P.A. 2017. Phthalates
661 impact human health: epidemiological evidences and plausible mechanism of action. *Journal of*
662 *Hazardous Materials* 340, 360–383. <https://doi.org/10.1016/j.jhazmat.2017.06.036>.
- 663 Bohlin-Nizzetto, P., Aas, W., Nikiforov, V. 2018. Monitoring of environmental contaminants in air
664 and precipitation. Annual report 2014. [https://nilu.brage.unit.no/nilu-](https://nilu.brage.unit.no/nilu-xmlui/handle/11250/2611257)
665 [xmlui/handle/11250/2611257](https://nilu.brage.unit.no/nilu-xmlui/handle/11250/2611257)
- 666 Borrell, A., Bloch, D., Desportes, G. 1995. Age trends and reproductive transfer of organochlorine
667 compounds in long-finned pilot whales from the Faroe Islands. *Environmental Pollution*, 88, 3,
668 283-292.

- 669 Borrell, A., Aguilar, A. 2005 Mother-calf transfer of organochlorine compounds in the common
670 dolphin (*Delphinus delphis*). Bulletin of environmental contamination and toxicology, 75, 1, 149-
671 156.
- 672 Borrell, A., Aguilar, A., Tornero, V., Drago, M. 2014. Concentrations of mercury in tissues of striped
673 dolphins suggest decline of pollution in Mediterranean open waters, Chemosphere, 107, 319-323,
674 <https://doi.org/10.1016/j.chemosphere.2013.12.076>.
- 675 Consumer Product Safety Commission (CPSC) 2018. Prohibition of Children’s Toys and Child Care
676 Articles Containing Specified Phthalates; 16 CFR Part 1307; Consumer Product Safety
677 Commission (CPSC): Bethesda, MA, USA, 2018.
- 678 Domingo M., Ferrer L., Pumarola M., Marco A. 1990. Morbillivirus in dolphins. Nature 348, 21.
- 679 Domingo, M., Vilafranca, M., Visa, J., Prats, N., Trudgett, A., Visser, I. 1995. Evidence for chronic
680 morbillivirus infection in the Mediterranean striped dolphin (*Stenella coeruleoalba*). Veterinary
681 Microbiology 44, 2-4, 229-239. [https://doi.org/10.1016/0378-1135\(95\)00016-4](https://doi.org/10.1016/0378-1135(95)00016-4).
- 682 Dziobak, M.K., Wells, R.S., Pisarski, E.C., Wirth, E.F., Hart, L.B. 2021. Demographic Assessment of
683 Mono(2-ethylhexyl) Phthalate (MEHP) and Monoethyl Phthalate (MEP) Concentrations in
684 Common Bottlenose Dolphins (*Tursiops truncatus*) From Sarasota Bay, FL, USA. GeoHealth 5,
685 1–13. <https://doi.org/10.1029/2020GH000348>
- 686 Dziobak, M.K.; Wells, R.S.; Pisarski, E.C.; Wirth, E.F.; Hart, L.B. A Correlational Analysis of
687 Phthalate Exposure and Thyroid Hormone Levels in Common Bottlenose Dolphins (*Tursiops*
688 *truncatus*) from Sarasota Bay, Florida (2010–2019). Animals 12, 824.
689 <https://doi.org/10.3390/ani12070824>.

- 690 Errico, S., Nicolucci, C., Migliaccio, M., Micale, V., Mita, D.G., Diano, N. 2017. Analysis and
691 occurrence of some phenol endocrine disruptors in two marine sites of the northern coast of Sicily
692 (Italy). *Marine Pollution Bulletin* 120, 1–2, 68–74.
693 <https://doi.org/10.1016/j.marpolbul.2017.04.061>.
- 694 European Commission. Commission directive 2011/8/EU of 28 January 2011 amending directive
695 2002/72/EC as regards the restriction of use of bisphenol A in plastic infant feeding bottles.
696 Official Journal of the European Union 10-05-2013.
- 697 Fossi, M.C., Coppola, D., Bainsi, M., Giannetti, M., Guerranti, C., Marsili, L., Panti, C., de Sabata, E.,
698 Clò, S. 2014. Large filter feeding marine organisms as indicators of microplastic in the pelagic
699 environment: The case studies of the Mediterranean basking shark (*Cetorhinus maximus*) and fin
700 whale (*Balaenoptera physalus*). *Marine Environmental Research* 100, 17–24.
701 <https://doi.org/10.1016/j.marenvres.2014.02.002>
- 702 Fossi, M.C., Marsili, L., Bainsi, M., Giannetti, M., Coppola, D., Guerranti, C., Caliani, I., Minutoli, R.,
703 Lauriano, G., Finoia, M.G., Rubegni, F., Panigada, S., Bérubé, M., Urban Ramirez, J., Panti, C.
704 2016. Fin whales and microplastics: The Mediterranean Sea and the Sea of Cortez scenarios.
705 *Environmental Pollution* 209, 68–78. <https://doi.org/10.1016/j.envpol.2015.11.022>
- 706 Fossi, M. C., Panti, C., Guerranti, C., Coppola, D., Giannetti, M., Marsili, L., Minutoli, R. 2012. Are
707 baleen whales exposed to the threat of microplastics? A case study of the Mediterranean fin whale
708 (*Balaenoptera physalus*). *Marine Pollution Bulletin* 64, 11, 2374–2379.
709 <https://doi.org/10.1016/j.marpolbul.2012.08.013>
- 710 Frouin, H., Lebeuf, M., Hammill, M., Fournier, M. 2012. Transfer of PBDEs and chlorinated POPs
711 from mother to pup during lactation in harp seals “*Phoca groenlandica*”. *Science of the Total*
712 *Environment* 417–418, 98–107. Doi:10.1016/j.scitotenv.2011.11.084

- 713 Garcia-Garin, O., Sala, B., Aguilar, A., Vighi, M., Víkingsson, G.A., Chosson, V., Eljarrat, E, Borrell,
714 A. 2020. Organophosphate contaminants in North Atlantic fin whales. *Science of the Total*
715 *Environment* 721, 137768. <https://doi.org/10.1016/j.scitotenv.2020.137768>.
- 716 Garcia-Garin, O., Sahyoun, W., Net, S., Vighi, M., Aguilar, A., Ouddane, B., Víkingsson, G.A.,
717 Chosson, V., Borrell, A., 2022. Intrapopulation and temporal differences of phthalate
718 concentrations in North Atlantic fin whales (*Balaenoptera physalus*) *Chemosphere* 300.
719 134453 <https://doi.org/10.1016/j.chemosphere.2022.134453>.
- 720 Geyer, R., Jambeck, J. R., Law, K.L. 2017. Production, use, and fate of all plastics ever made. *Science*
721 *Advances* 3, 7, 25–29. <https://doi.org/10.1126/sciadv.1700782>
- 722 Giam, C.S., Chan, H.S., Neff, G.S., Atlas, E.L. 1978. Phthalate ester plasticizers: A new class of marine
723 pollutant. *Science* 199,27, 419–420.
- 724 Graham, P.R. 1973. Phthalate ester plasticizers—why and how they are used. *Environmental Health*
725 *Perspectives* 3, 3–12.
- 726 Han, Y., Fei, Y., Wang, M., Xue, Y., Chen, H., Liu, Y. 2021. Study on the joint toxicity of BPZ, BPS,
727 BPC and BPF to Zebrafish. *Molecules* 26, 4180. <https://doi.org/10.3390/molecules26144180>
- 728 Hart, L. B., Beckingham, B., Wells, R. S., Alten Flagg, M., Wischusen, K., Moors, A., Kucklick, J.,
729 Pisarski, E., Wirth, E. 2018. Urinary Phthalate Metabolites in Common Bottlenose Dolphins
730 (*Tursiops truncatus*) From Sarasota Bay, FL, USA. *GeoHealth* 2, 10, 313–326.
731 <https://doi.org/10.1029/2018gh000146>
- 732 Hart, L. B., Dziobak, M.K., Pisarski, E. C., Wirth, E.F., Wells, R.S. 2020. Sentinels of synthetics – a
733 comparison of phthalate exposure between common bottlenose dolphins (*Tursiops truncatus*) and
734 human reference populations. *PloS ONE* 15, 1–16. <https://doi.org/10.1371/journal.pone.0240506>

- 735 Hartmann, C., Uhl, M., Weiss, S., Koch, H.M., Scharf, S., König, J. 2015. Human biomonitoring of
736 phthalate exposure in Austrian children and adults and cumulative risk assessment. *International*
737 *Journal of Hygiene and Environmental Health* 218, 5, 489–499.
738 <https://doi.org/10.1016/j.ijheh.2015.04.002>
- 739 Huang, P.C., Tsai, C.H., Liang, W.Y., Li, S.S., Pan, W.H., Chiang, H.C. 2015. Age and gender
740 differences in urinary levels of eleven phthalate metabolites in general Taiwanese population after
741 a DEHP episode. *PloS ONE* 10, 7, 1–17. <https://doi.org/10.1371/journal.pone.0133782>
- 742 IHSMarkit, 2021. Chemical Economics handbook – Plasticizers. Technical report. HIS Markit,
743 London, UK.
- 744 IUCN. 2019. The IUCN Red List of Threatened Species. Version 2019-1. Available at:
745 www.iucnredlist.org. (Accessed: 31 August 2022).
- 746 Jambeck, J.R., Geyer, R., Wilcox, C., Siegler, T.R., Perryman, M., Andrady, A., Law, K.L. 2015.
747 Plastic waste inputs from land into the ocean. *Science* 347, 6223, 768–771.
748 <https://doi.org/10.1126/science.1260352>
- 749 Jin, H., Zhu, L. 2016. Occurrence and partitioning of bisphenol analogues in water and sediment from
750 Liaohe River Basin and Taihu Lake, China, *Water Research* 103, 343-351,
751 <https://doi.org/10.1016/j.watres.2016.07.059>.
- 752 Kang, J.H., Katamaya, Y., Kondo, F. 2006. Biodegradation or metabolism of bisphenol A: From
753 microorganisms to mammals. *Toxicology*, 217, 81-90.
- 754 Kang, J.H., Aasi, D., Katayama, Y. 2007. Bisphenol A in the aquatic environment and its endocrine-
755 disruptive effects on aquatic organisms. *Critical Reviews in Toxicology* 37, 607-625. DOI:
756 10.1080/10408440701493103.

- 757 Keck, N., Kwiatek, O., Dhermain, F., Dupraz, F., Boulet, H., Danes, C., Laprie, C., Perrin, A., Godenir,
758 J., Micout, L. 2010. Resurgence of Morbillivirus infection in Mediterranean dolphins off the
759 French coast. *The Veterinary Record* 166, 654.
- 760 Koelmans, A.A., Besseling, E., Foekema, E.M. 2014. Leaching of plastic additives to marine
761 organisms. *Environmental pollution* 187, 49-54. <http://dx.doi.org/10.1016/j.envpol.2013.12.013>
- 762 Krahn, M.M., Ylitalo, G.M., Stein, J.E., Aguilar, A., Borrell, A. 2003. Organochlorine contaminants
763 in cetaceans: how to facilitate interpretation and avoid errors when comparing datasets. *Journal*
764 *of Cetacean Research and Management* 5, 2, 103–113.
- 765 Lambert, C., Authier, M., Dorémus, G., Laran, S., Panigada, S., Spitz, J., Van Canneyt, O., Ridoux,
766 V. 2020. Setting the scene for Mediterranean litterscape management: the first basin-scale
767 quantification and mapping of floating marine debris. *Environmental Pollution*, 263, 114430.
768 [10.1016/j.envpol.2020.114430](https://doi.org/10.1016/j.envpol.2020.114430)
- 769 Lee, S., Liao, C., Song, G.J., Ra, K., Kannan, K., Moon, H.B. 2015. Emission of bisphenol analogues
770 including bisphenol A and bisphenol F from wastewater treatment plants in Korea. *Chemosphere*
771 119, 1000-1006.
- 772 Li, Y., Perea, L., Coons, L.A., Burns, K.A., Ramsey, J.T., Pelch, K.E., Houtman, R., van Beuningen,
773 R., Teng, C.T., Korach, K.S. 2018. Differential in vitro biological action, coregulator interactions,
774 and molecular dynamic analysis of Bisphenol A, BPA, BPAF and BPS Ligand-ER α complexes.
775 <https://doi.org/10.1289/EHP2505>
- 776 Liao, C., Kannan, K. 2013. Concentrations and profiles of bisphenol A and other bisphenol analogues
777 in foodstuffs from the United States and their implications for human exposure. *Journal of*
778 *Agricultural and Food Chemistry* 15, 61(19): 4655-62. Doi: 10.1021/jf400445n.

- 779 Liao, C., Liu, F., Moon, H.B., Yamashita, N., Yun, S., Kannan, K. 2012. Bisphenol analogues in
780 sediments from industrialized areas in the United States, Japan, and Korea: spatial and temporal
781 distributions. *Environmental Science and Technology* 46, 21, 11558-11565.
- 782 Loganathan, B.G. 2012. Global contamination trends of persistent organic chemicals: An overview.
783 In. *Global Contamination Trends of Persistent Organic Chemicals*. (Eds. B.G. Loganathan, PKS.
784 Lam) CRC Press, Boca Raton, FL. Pp. 3-31.
- 785 Loganathan, B.G., Tanabe, S., Tanaka, H., Miyazaki, N., Amano, M., Tatsukawa, R. 1990.
786 Comparison of persistent residues in striped dolphin *Stenella coeruleoalba* from western North
787 Pacific in 1978-1986. *Marine Pollution Bulletin* 21, 435-439.
- 788 Mackintosh, C.E., Maldonado, J., Hongwu, J., Hoover, N., Chong, A., Ikonomou, M.G., Gobas,
789 F.A.P.C. 2004. Distribution of Phthalate Esters in a Marine Aquatic Food Web: Comparison to
790 Polychlorinated Biphenyls. *Environmental Science and Technology* 38, 7, 2011–2020.
791 <https://doi.org/10.1021/es034745r>
- 792 Marqueño, A., Pérez-Albadejo, E., Denslow, N.D., Bowden, J.A., Porte, C. 2021. Untargeted
793 lipidomics reveals the toxicity of bisphenol A bis(3-chloro-2-hydroxypropyl) ether and
794 bisphenols A and F in zebrafish liver cells. *Ecotoxicology and Environmental Safety* 219, 112311.
795 <https://doi.org/10.1016/j.ecoenv.2021.112311>
- 796 Marsili, L., Jiménez, B., Borrell, A. 2018. Persistent organic pollutants in cetaceans living in a hotspot
797 area: the Mediterranean Sea. In *Marine Mammal Ecotoxicology* (pp. 185-212). Academic Press.
- 798 Montoto-Martínez, T., De la Fuente, J., Puig-Lozano, R., Marques, N., Arbelo, M., Hernández-Brito,
799 J.J., Fernández, A., Gelado-Caballero, M.D. 2021. Microplastics, bisphenols, phthalates and

- 800 pesticides in odontocete species in the Macaronesian Region (Eastern North Atlantic). *Marine*
801 *Pollution Bulletin* 173. <https://doi.org/10.1016/j.marpolbul.2021.113105>
- 802 Moustafa A., Hamzeh M., Baroudi M., Ouddane B., Net S. 2022. 55 xenobiotic organic compounds in
803 Tripoli Landfill-Lebanon Leachate and their Fluxes to the Abou Ali River and Mediterranean
804 Sea. *Environmental Monitoring and Assessment* 194, 12. 10.1007/s10661-022-10522-w.
- 805 Mustieles, V., D’Cruz, S.C., Couderq, S., Rodríguez-Carrillo, A., Fini, J.B., Hofer, T., Steffensen, I.L.,
806 Dirven, H., Barouki, R., Olea, N., Fernández, M.F., David, A. 2020. Bisphenol A and its
807 analogues: a comprehensive review to identify and prioritize effect biomarkers for human
808 biomonitoring. *Environment International* 144, 105811. 10.1016/j.envint.2020.105811
- 809 Nehring, I., Staniszewska, M., Falkowska, L. 2017. Human Hair, Baltic Grey Seal (*Halichoerus*
810 *grypus*) Fur and Herring Gull (*Larus argentatus*) Feathers as Accumulators of Bisphenol A and
811 Alkylphenols. *Archives of Environmental Contaminant Toxicology* 72, 522-561.
812 10.1007/s00244-017-0402-0
- 813 Net, S., Sempéré, R., Delmont, A., Paluselli, A., Ouddane, B. 2015. Occurrence, Fate, Behavior and
814 Ecotoxicological State of Phthalates in Different Environmental Matrices (Critical Review).
815 *Environmental Science & Technology* 49, 4019-40. <https://doi.org/10.1021/es505233b35>.
- 816 Novillo, O., Raga, J.T., Tomas, J. 2020. Evaluating the presence of microplastics in striped dolphins
817 (*Stenella coeruleoalba*) stranded in the Western Mediterranean Sea. *Marine Pollution Bulletin*
818 160, 2020, 111557. <https://doi.org/10.1016/j.marpolbul.2020.111557>.
- 819 Owczarek K, Kudłak B, Simeonov V, Mazerska Z, Namieśnik J. 2018. Binary Mixtures of Selected
820 Bisphenols in the Environment: Their Toxicity in Relationship to Individual Constituents.
821 *Molecules* 23, 12, 3226. Doi: 10.3390/molecules23123226.

- 822 Özkan-Kotiloğlu, S., Arslan, P., Akca, G., Günel, A.C. 2022. Are BPA-free plastics safe for aquatic
823 life? – Fluorene-9-bisphenol induced thyroid-disrupting effects and histopathological alterations
824 in adult zebrafish (*Danio rerio*), *Comparative Biochemistry and Physiology Part C: Toxicology*
825 & *Pharmacology* 260, 109419, <https://doi.org/10.1016/j.cbpc.2022.109419>.
- 826 Ozhan, K., Kocaman, E. 2019. Temporal and Spatial Distributions of Bisphenol A in Marine and
827 Freshwaters in Turkey. *Archives of Environmental Contamination and Toxicology* 76, 246–254.
828 <https://doi.org/10.1007/s00244-018-00594-6>
- 829 Page-Karjian, A., Lo, C. F., Ritchie, B., Harms, C. A., Rotstein, D. S., Han, S., Hassan, S.M., Lehner,
830 A.F., Buchweitz, J.P., Thayer, V.G., Sullivan, J.M., Christiansen, E.F., Perrault, J.R. 2020.
831 Anthropogenic Contaminants and Histopathological Findings in Stranded Cetaceans in the
832 Southeastern United States, 2012–2018. *Frontiers in Marine Science* 7.
833 <https://doi.org/10.3389/fmars.2020.00630>
- 834 Paluselli, A., Fauvelle, V., Schmidt, N., Galgani, F., Net S., Sempéré, R. 2018. Distribution of
835 phthalates in Marseille Bay (NW Mediterranean Sea). *Science of the Total Environment* 621,
836 578-587.
- 837 Paluselli, A., Fauvelle, V., Galgani, F., Sempéré, R. 2019. Phthalate release from plastic fragments
838 and degradation in seawater. *Environmental Science & Technology* 53 (1), 166-175.
839 <https://doi.org/10.1021/acs.est.8b05083>
- 840 Panio, A., Fabbri Corsarini, S., Bruno, A., Lasagni, M., Labra, M., Saliu, F. 2020. Determination of
841 phthalates in fish fillets by liquid chromatography tandem mass spectrometry (LC-MS/MS): A
842 comparison of direct immersion solid phase microextraction (SPME) versus ultrasonic assisted
843 solvent extraction (UASE), *Chemosphere*, 255, 127034,
844 <https://doi.org/10.1016/j.chemosphere.2020.127034>.

- 845 Park, J.C., Lee, M.-C., Yoon, D.-S. Han, J., Kim, M., Hwang, U.-K., Jung, J.-H., Lee, J.-S. 2018.
846 Effects of bisphenol A and its analogs bisphenol F and S on life parameters, antioxidant system,
847 and response of defensome in the marine rotifer *Brachionus koreanus*. *Aquatic Toxicology*, 199,
848 21-29. <https://doi.org/10.1016/j.aquatox.2018.03.024>
- 849 Park, C., Song, H., Choi, J., Sim, S., Kojima, H., Park, J., Iida, M., Lee, Y. 2020. The mixture effects
850 of bisphenol derivatives on estrogen receptor and androgen receptor. *Environmental Pollution*,
851 260, 114036. <https://doi.org/10.1016/j.envpol.2020.114036>
- 852 Pelch, K.E., Li, Y., Perera, L., Thayer, K.A., Korach, K.S. 2019. Characterization of Estrogenic and
853 Androgenic Activities for Bisphenol A-like Chemicals (BPs): In Vitro Estrogen and Androgen
854 Receptors Transcriptional Activation, Gene Regulation, and Binding Profiles. *Toxicological*
855 *Science* 172, 1, 23–37. doi: 10.1093/toxsci/kfz173.
- 856 R Core Team. 2021. R: A language and environment for statistical computing. R Foundation for
857 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- 858 Raguso, C., Grech, D., Becchi, A., Ubaldi, P.G., Lasagni, M., Guala, I., Saliu, F. 2022. Detection of
859 microplastics and phthalic acid esters in sea urchins from Sardinia (Western Mediterranean Sea),
860 *Marine Pollution Bulletin*, 185, Part A, 114328,
861 <https://doi.org/10.1016/j.marpolbul.2022.114328>.
- 862 Rian, M.B., Vike-Jonas, K., Gonzalez, S.V., Ciesielski, T.M., Venkatraman, V., Lindstrøm, U., ...
863 Asimakopoulos, A.G. 2020. Phthalate metabolites in harbor porpoises (*Phocoena phocoena*) from
864 Norwegian coastal waters. *Environment International* 137, 105525.
865 <https://doi.org/10.1016/j.envint.2020.105525>

- 866 Rios-Fuster, B., Alomar, C., Paniagua González, G., Garcinuño Martínez, R.M., Soliz Rojas, D.L.,
867 Fernández Hernando, P., Deudero, S. 2022. Assessing microplastic ingestion and occurrence of
868 bisphenols and phthalates in bivalves, fish and holothurians from a Mediterranean marine
869 protected area. *Environmental Research* 214, 3, 114034.
870 <https://doi.org/10.1016/j.envres.2022.114034>.
- 871 Routti, H., Harju, M., Lühmann, K., Aars, J., Ask, A., Goksøyr, A., Kovacs, K.M., Lydersen, C. 2021.
872 Concentrations and endocrine disruptive potential of phthalates in marine mammals from the
873 Norwegian Arctic. *Environment International* 152. <https://doi.org/10.1016/j.envint.2021.106458>
- 874 Sangeetha, S., Vimalkumar, K., Loganathan, B.G. 2021. Environmental contamination and human
875 exposure to select endocrine-disrupting chemicals: A review. *Sustainable Chemistry* 2, 343-380.
- 876 Schmidt, N., Castro-Jiménez, J., Oursel, B., Sempéré, R. 2021. Phthalates and organophosphate esters
877 in surface water, sediments and zooplankton of the NW Mediterranean Sea: Exploring links with
878 microplastic abundance and accumulation in the marine food web. *Environmental Pollution* 1,
879 272, 115970. [10.1016/j.envpol.2020.115970](https://doi.org/10.1016/j.envpol.2020.115970).
- 880 Silva, M. J., Barr, D.B., Reidy, J.A., Malek, N.A., Hodge, C.C., Caudill, S. P., Brock, J.W., Needham,
881 L.L., Calafat, A.M. 2004. Urinary levels of seven phthalate metabolites in the U.S. population
882 from the National Health and Nutrition Examination Survey (NHANES) 1999-2000.
883 *Environmental Health Perspectives* 112, 3, 331–338. <https://doi.org/10.1289/ehp.6723>
- 884 Silva, D.C.V.R., Araújo, C.V.M., França, F.M., Neto, M.B., Paiva, T.C.B., Silva, F.T., Pompêo,
885 M.L.M., 2018. Bisphenol risk in fish exposed to a contamination gradient: triggering of spatial
886 avoidance. *Aquatic Toxicology* 197, 1–6. <https://doi.org/10.1016/j.aquatox.2018.01.016>

- 887 Staples, C.A., Dorn, P.B., Klecks, G.M., O'Block, S.T., Harris, L.R. 1998. A review of the
888 environmental fate, effects, and exposures of bisphenol A. *Chemosphere* 36, 10, 2149-2173.
889 [10.1016/s0045-6535\(97\)10133-3](https://doi.org/10.1016/s0045-6535(97)10133-3)
- 890 Staples, C.A., Peterson, D.R., Parkerton, T.F., Adams, W.J. 1997. Environmental fate of phthalate
891 esters: a literature review. *Chemosphere* 35, 4, 667-749.
- 892 Sghaier, R. Ben, Net, S., Ghorbel-Abid, I., Bessadok, S., Le Coz, M., Hassan-Chehimi, D. B., Trabelsi-
893 Ayadi, M., Tackx, M., Ouddane, B. 2017. Simultaneous Detection of 13 Endocrine Disrupting
894 Chemicals in Water by a Combination of SPE-BSTFA Derivatization and GC-MS in
895 Transboundary Rivers (France-Belgium). *Water, Air, and Soil Pollution* 228, 1.
896 <https://doi.org/10.1007/s11270-016-3195-2>
- 897 Tanabe, S., Loganathan, B.G., Subramaniam, A.N., Tatsukawa, R. 1987. Organochlorine residue in
898 shot-finned pilot whale: Possible use as tracer of biological parameters. *Marine Pollution Bulletin*
899 18, 561-563.
- 900 Teuten, E.L., Rowland, S.J., Galloway, T.S., Thompson, R.C. 2007. Potential for plastics to transport
901 hydrophobic contaminants. *Environmental Science and Technology* 41, 7759–7764.
- 902 Torres-García, J. L., Ahuactzin-Pérez, M., Fernández, F.J., Cortés-Espinosa, D.V. 2022. Bisphenol A
903 in the environment and recent advances in biodegradation by fungi. *Chemosphere* 303, 134940.
904 <https://doi.org/10.1016/j.chemosphere.2022.134940>.
- 905 Vorkamp, K., Dam, M., Riget, F., Fauser, P., Bossi, R., Hansen, A.B. 2004. Screening of “new”
906 contaminants in the marine environment of Greenland and the Faroe Islands. National
907 Environmental Research Institute, Denmark - 97 pp. - NERI Technical Report No 525. Available
908 at: <http://technical-reports.dmu.dk>

- 909 Wang, R., Tan, T., Liang, H., Huang, Y., Dong, S., Wang, P., Su, X. 2021. Occurrence and distribution
910 of bisphenol compounds in different categories of animal feeds used in China, *Emerging*
911 *Contaminants* 7, 179-186, <https://doi.org/10.1016/j.emcon.2021.08.001>.
- 912 Warner, G.R., Flaws, J.A. 2018. Bisphenol A and Phthalates: How Environmental Chemicals Are
913 Reshaping Toxicology. *Toxicological Sciences* 166, 2, 246-249. [10.1093/toxsci/kfy232](https://doi.org/10.1093/toxsci/kfy232).
- 914 Xie, Z., Ebinghaus, R., Temme, C., Lohmann, R., Cara, A., Ruck, W. 2007. Occurrence and air-sea
915 exchange of phthalates in the Arctic. *Environmental Science and Technology* 41, 13, 4555–4560.
916 <https://doi.org/10.1021/es0630240>
- 917 Xing, J., Zhang, S., Zhang, M., Hou, J. 2022. A critical review of presence, removal and potential
918 impacts of endocrine disruptors bisphenol A. *Comparative Biochemistry and Physiology Part C:*
919 *Toxicology & Pharmacology* 254, 109275. <https://doi.org/10.1016/j.cbpc.2022.109275>.
- 920 Yu, X., Xue, J., Yao, H., Wu, Q., Venkatesan, A.K., Halden, R.U., Kannan, K. 2015. Occurrence and
921 estrogenic potency of eight bisphenol analogs in sewage sludge from the U.S. EPA targeted
922 national sewage sludge survey. *Journal of Hazardous Materials* 299, 733-739.
- 923 Zeng, F., Cui, K.Y., Xie, Z.Y., Wu, L.N., Luo, D.L., Chen, L.X., Lin, Y.J., Liu, M., Sun, G.X. 2009.
924 Distribution of phthalate esters in urban soils of subtropical city, Guangzhou, China. *Journal of*
925 *Hazardous Materials* 164, 2, 1171–1178. <https://doi.org/10.1016/j.jhazmat.2008.09.029>

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Odei Garcia-Garin reports financial support was provided by European Social Fund.

Journal Pre-proof