



HAL
open science

Contribution of nuclear applications to better understand bioaccumulation of contaminants in aquaculture species

Marc Metian, Simon Pouil, François Oberhänsli, Paco Bustamante, Peter Swarzenski

► **To cite this version:**

Marc Metian, Simon Pouil, François Oberhänsli, Paco Bustamante, Peter Swarzenski. Contribution of nuclear applications to better understand bioaccumulation of contaminants in aquaculture species. 28th Annual Meeting SETAC (Society of Environmental Toxicology and Chemistry) Europe), May 2017, Rome, Italy. hal-04198152

HAL Id: hal-04198152

<https://hal.science/hal-04198152>

Submitted on 7 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Contribution of nuclear applications to better understand bioaccumulation of contaminants in aquaculture species

Marc Metian¹, Simon Pouil^{1,2}, François Oberhänsli¹, Paco Bustamante², Peter Swarzenski¹

¹International Atomic Energy Agency-Environment Laboratories (IAEA-EL), Principality of Monaco

²Littoral Environnement et Sociétés (LIENSs), University of La Rochelle, La Rochelle, France

E-mail contact: M.Metian@iaea.org ; mametian@gmail.com

1. Introduction

Wild- or farmed-marine fish are an important source of high-quality animal protein and long-chain polyunsaturated fatty acids, but are also a potential source of contaminants such as toxic trace elements that could be introduced to seafood, especially as one moves up the trophic food chain. Aquaculture of marine fish is regarded as the future of aquaculture, and thus, the safety of these farmed fish is paramount particularly as 50 % of fish consumed by humans are now farmed¹. Therefore, a better understanding of bioaccumulation kinetics and processes of such contaminants with current aquafarming practices is essential. Such an effort will yield important new information on the role the fish food or key environmental parameters on contamination of fish that could potentially impact the health of the cultured species and/or the human consumer. This is commonly performed in representative wild fish species.

In traditional aquaculture practices, caged fish can be fed different types of food such as natural prey, including, 1) forage fish, crustaceans, polychaetes or by-products of seafood processing, 2) artificial pellets produced by dedicated manufacturers with various formulations developed to improve the aquaculture performances. Common to all these feeding practices is that both natural prey and ingredients used in artificial pellet formulations are likely to contain contaminant residues. Thus, the forage fish and fish meal may be rich in organic mercury, and squid viscera meal (used as additives to stimulate growth) is a potentially high cadmium source^{2,3}.

Key environmental parameters have been shown to affect trace element bioaccumulation in aquatic organisms⁴. Among these factors, water salinity and temperature are key parameters in aquaculture as they both exert a strong influence on production performances (by affecting farmed fish physiology). Nevertheless, there is still limited information available about the influence of these factors on trace element trophic transfer in farmed fish.

In order to better understand the contamination risk of farmed species, a series of nuclear-based methods can be applied such as for example, high efficiency gamma-spectrometry. This is a very powerful and very sensitive approach that allows identifying the susceptibility of farm-raised fish to be contaminated by trace elements and radionuclides in particular through the measurement of the dietary assimilation efficiency (AE).

In this context, the aim of this presentation is to provide some examples of radiotracer studies such as the influence of food items and environmental conditions (salinity and temperature of the water) on a farmed fish, the turbot *Scophthalmus maximus* (reaching 72 000 tonnes in 2014) using radiotracer techniques

2. Materials and Methods

2.1. Origin and acclimation of the fish

Juvenile turbot was purchased from a fish farm (France Turbot, France) and shipped to the International Atomic Energy Agency premises in the Principality of Monaco. Fish were acclimated to laboratory conditions for several weeks. During the acclimation period, the fish were fed a daily ration of 2% of their biomass.

2.2. Radiolabelling of the different foods

For food radiolabelling, radiotracers of high specific activity were used. Live food were exposed to contaminants through dissolved phase for at least 4 days and pellets were dipped in radiolabelled seawater for 1h. Then, radioactive pellets were dried and stored in dry conditions. The use of gamma-emitting isotope allowed accurate measurements using environmentally realistic concentrations.

2.3. Assimilation efficiency

The experiments were performed in 20-L or 70-L aquaria (open-circuit, same conditions that during acclimation). Two weeks before the start of the experiments, juvenile turbot was randomly transferred from

the acclimation tank to the aquaria used for experiments and slits cut into the fins were used to facilitate individual recognition. Two hours after the unique radiolabelled feeding, all the turbot from the different experiments were counted using a high-resolution gamma-spectrometer system composed of 4 Germanium detectors to determine the ingested radioactivity. Then, turbot were counted after 24h and 48h of depuration. The assimilation efficiency (AE) was calculated as the percentage of radioactivity remaining in fish after 48h of depuration. Differences between AE were tested using non-parametric tests (Wilcoxon Mann-Whitney & Kruskal-Wallis). The level of significance for statistical analyses was always set at $\alpha = 0.05$.

3. Results and discussion

The quality and type of food is already known to directly influence trace element content in fish⁵. Here, we found that, using juvenile turbot, ¹⁰⁹Cd assimilation efficiency (AE) was significantly different depending the type of ingested food (Fig. 1A). These results can be explained by differences in Cd bioavailability in the foods as has been demonstrated in another species of fish, the grunt *Terapon jarbua*⁶.

Concerning the influence of salinity and temperature, our results shown that temperature had no significant influence on ^{110m}Ag AE in turbot (Fig. 1B). Nevertheless, a significant effect of salinity was observed in the ¹³⁷Cs (Fig. 1C). This finding can be related to the osmoregulation processes in marine fish that control the diffusive ion invasion and excreted the excess ions to maintain their osmolarity⁷. Based on these physiological mechanisms, the lower observed ¹³⁷Cs AE in turbot maintained in the highest salinity condition can be explained by (1) a limited entrance of dietary ¹³⁷Cs in the gastrointestinal tract, and/or (2) a higher excretion rate of ¹³⁷Cs.




Figure 1: Assimilation efficiencies, measured in turbot (n=5-8), of (A) ¹⁰⁹Cd in turbot fed with different foods, (B) ^{110m}Ag under two water temperature conditions and (C) ¹³⁷Cs under three water salinities.

Conclusions

In summary, this study has shown varying effects of food, water salinity and temperature in the AE of trace elements and radionuclides in farmed fish. Such findings set the stage for further investigations on the potential use of nuclear techniques in aquaculture and food safety.

References

- [1] Food and Agriculture Organisation (FAO). 2016. The state of world fisheries and aquaculture (SOFIA). FAO, Roma, Italy.
- [2] Dang F, Wang W-X. 2009. Assessment of tissue-specific accumulation and effects of cadmium in a marine fish fed contaminated commercially produced diet. *Aquat Toxicol Amst Neth* 95:248–255.
- [3] Wang W-X, Onsanit S, Dang F. 2012. Dietary bioavailability of cadmium, inorganic mercury, and zinc to a marine fish: Effects of food composition and type. *Aquaculture* 356–357:98–104.
- [4] Luoma SN, Rainbow PS. 2005. Why is metal bioaccumulation so variable? *Biodynamics as a unifying concept. Environ Sci Technol* 39:1921–1931.
- [5] Pouil S, Warnau M, Oberhänsli F, Teyssié J-L, Bustamante P, Metian M. 2016. Influence of food on the assimilation of essential elements (Co, Mn, and Zn) by turbot *Scophthalmus maximus*. *Mar Ecol Prog Ser* 550:207–218.
- [6] Zhang L, Wang W-X (2006) Significance of subcellular metal distribution in prey in influencing the trophic transfer of metals in a marine fish. *Limnol Oceanogr* 51:2008–2017.
- [7] Lorin-Nebel C, Boulo V, Bodinier C, Charmantier G (2006) The Na⁺/K⁺/2Cl⁻ cotransporter in the sea bass *Dicentrarchus labrax* during ontogeny: involvement in osmoregulation. *J Exp Biol* 209:4908–4922.

Acknowledgement - The IAEA is grateful for the support provided to its Environment Laboratories by the Government of the Principality of Monaco.