



HAL
open science

Influence of food increased Hg bioaccumulation in oyster when feeding on protozoans

Marc Metian, Simon Pouil, Christine Dupuis, Paco Bustamante, Jean-Louis Teyssié, Michel Warnau

► **To cite this version:**

Marc Metian, Simon Pouil, Christine Dupuis, Paco Bustamante, Jean-Louis Teyssié, et al.. Influence of food increased Hg bioaccumulation in oyster when feeding on protozoans. 9th International Conference on Marine Pollution and Ecotoxicology, Jun 2019, Hong-Kong, China. hal-04198174

HAL Id: hal-04198174

<https://hal.science/hal-04198174>

Submitted on 6 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

INFLUENCE OF FOOD INCREASED HG BIOACCUMULATION IN OYSTER WHEN FEEDING ON PROTOZOANS

Marc Metian¹, Simon Pouil¹⁻², Christine Dupuis², Paco Bustamante², Jean-Louis Teysié¹, Michel Warnau¹

¹International Atomic Energy Agency, Environment Laboratories, Monaco

²Littoral Environnement et Sociétés (LIENSs), CNRS-Université de La Rochelle, France

m.metian@iaea.org


Context

- Food is an important route of mercury uptake in marine organisms, including in bivalves
- Knowledge on mercury accumulation from their food is mainly based on experiments using phytoplankton as food
- Oysters are also feeding on a variety of organisms, including ciliates, in addition to phytoplankton


What is the influence of diet composition in the bioaccumulation of mercury in oysters ?
Does food types influence both inorganic mercury and methylmercury ?

Methods

- Pacific cupped oyster *Crassostrea gigas*
- Food: protist (*Uronema marina*) vs. phytoplankton (*Thalassiosira pseudonana*)
- Radiotracers: ²⁰³Hg and CH₃²⁰³Hg



Results and discussion



Oysters fed on ciliates assimilated $96 \pm 1\%$ and $31 \pm 2\%$ of the ingested CH₃Hg and iHg respectively while these elements were similarly assimilated in the oysters fed on phytoplankton ($78 \pm 3\%$ and $86 \pm 4\%$ for CH₃Hg and iHg, respectively).

Mercury assimilation in oyster is thus diet dependent (significant differences in AE, $p < 0.05$) and likely resulting from variations in Hg bioavailability in the two feed items tested and a gut passage time-dependent of the ingested matrix.