

Exponential stable manifold for the synchronized state of the abstract mean field system

W Oukil

▶ To cite this version:

W Oukil. Exponential stable manifold for the synchronized state of the abstract mean field system: Exponential stability for the mean field system. 2024. hal-04198672v3

HAL Id: hal-04198672 https://hal.science/hal-04198672v3

Preprint submitted on 8 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Exponential stable manifold for the synchronized state of the abstract mean field system

W. Oukil

Faculty of Mathematics. University of Science and Technology Houari Boumediene. BP 32 EL ALIA 16111 Bab Ezzouar, Algiers, Algeria.

March 8, 2024

Abstract

We study the exponential stability for the mean field systems in its synchronized state. More precisely, we study the linearized system and we obtain a exponential stable invariant manifold.

Keywords: Mean field theory, differential equation, exponential stability, periodic system.

AMS subject classifications: 34D05, 37B65, 34C15. 35L04.

1 Introduction and Main result

We study in this article the exponential stability on the mean field systems as the Winfree mode in [2, 3, 4, 5, 6, 7, 10, 13] in its synchronized state. In 1967 Winfree [13] proposed a mean field model describing the synchronization of a population of organisms or *oscillators* that interact simultaneously [1].

Our main result consists of two parts: "The linear part," where we will study the stability of a class of perturbed linear systems by decomposing the fundamental matrix. "The non-linear part" where we study the exponential stability of the coupled mean-field systems. In other words, we shows the existence of a stable limit cycle. The results are within the framework of perturbation theory. However, we clarify that the results in the non-linear case are not straightforward, even in the case of no perturbation.

1.1 Main Results: Linear part

Consider the following notation: Let be $(n,m) \in \mathbb{N}^* \times \mathbb{N}^*$ and

$$A(t) := \{a_{i,j}(t) \in \mathbb{R}, \ 1 \le i \le n, \ 1 \le j \le m, \ t \in I \subseteq \mathbb{R}\},\$$

a $n \times m$ matrix-valued function, we denote the sup-norm of A(t) as

$$||A|| = \max\{\sup_{t \in I} |a_{ij}(t)|, 1 \le i \le n, \ 1 \le j \le m\},\$$

and we denote by I_N the square identity matrix of order N.

1.1.1 Stability Assumption (H_{stab})

Let $\zeta(t)$ be a square matrix-valued function of order N that depends on time $t \in \mathbb{R}$ satisfying $||\zeta|| < +\infty$. We consider the following perturbed linear system:

$$\dot{Y}(t) = [b(t)I_N + \mathcal{A}(t) + \zeta(t)]Y(t), \quad t \ge t', \tag{1}$$

where $t' \in \mathbb{R}$ and $\mathcal{A}(t) = \{a_{i,j}(t)\}_{1 \leq i,j \leq N}$ is a continuous square matrixvalued function of order N with rank 1, and $b : \mathbb{R} \to \mathbb{R}$ a continuous scalar function. We consider the following assumption about b(t) and \mathcal{A} that we call the stability assumption

$$(H_{stab}) \quad \begin{cases} a_{i,j}(t) = a_j(t) \text{ for all } 1 \le i, j \le N, \\ b : \mathbb{R} \to \mathbb{R}, a_j : \mathbb{R} \to \mathbb{R}, j = 1, \dots, N, \\ \text{are a continuous 1-periodic functions,} \\ \int_0^1 b(s) + \sum_{j=1}^N a_j(s) ds = 0, \quad \text{and} \quad -\alpha := \int_0^1 b(s) ds < 0. \end{cases}$$

Define the *fundamental matrix* of a linear system in the following sense

Definition 1. Let $t \mapsto A(t)$ be a continuous square matrix-valued function of order N. The *fundamental matrix* of the linear system

$$\dot{y} = A(t)y, \quad t \in \mathbb{R},$$

is the matrix-valued function R(s;t) with $s,t \in \mathbb{R}$ that satisfies

- $\forall t_0 \in \mathbb{R}, \ R(t_0; t_0) = I_N,$
- $\forall t \in \mathbb{R}, \ \frac{d}{dt}R(t;t_0) = A(t)R(t;t_0).$

To gain further insights into the behavior of solutions of the linear system (1), we will introduce a class of matrices ζ referred to as "normalizing matrices," defined as follows

Definition 2. We say that the matrix ζ is a *normalizing* matrix if the system (1) has a solution $V(t) = (v_1(t), \ldots, v_N(t))$ such that

$$\inf_{t\in\mathbb{R}}||V(t)||>0,\quad\text{and}\quad\sup_{t\in\mathbb{R}}||V(t)||<+\infty.$$

We call V(t) a normalizing solution of (1) associated with the matrix ζ .

1.1.2 Linear Result

In the following two linear results, we will consider two cases: ζ being a normalizing matrix or arbitrary.

Main Result (I^{*l*}). Consider the system (1) with fundamental matrix R(s; t). Suppose that *b* and *A* satisfy the stability assumption (H_{stab}). Let $\beta \in (0, \alpha)$, then there exist K > 0 and $D_* > 0$ such that for any normalizing matrix ζ satisfying $||\zeta|| < D_*$ and for any $t \in \mathbb{R}$, there exists a linear form $\mathcal{L}_t : \mathbb{R}^N \to \mathbb{R}$ such that for any $Y \in \mathbb{R}^N$ and for any $s \geq t$, we have

• $\mathcal{L}_s(R(s;t)V(t)) = 1$ and $\mathcal{L}_t(Y) < K||Y||,$

•
$$\mathcal{L}_t(Y) = \mathcal{L}_s(R(s;t)Y),$$

• $||R(s;t)[Y - \mathcal{L}_t(Y)V(t)]|| < K||Y||\exp(-\beta(s-t)).$

where V(t) is a normalizing solution of (1) associated with the matrix ζ . Moreover, the operator R(s;t) admits the decomposition

$$R(s;t)Y = \mathcal{L}_t(Y)V(s) + R(s;t)[Y - \mathcal{L}_t(Y)V(t)]$$

In other words, the submanifold $\mathcal{W}_{stab} := \{Z \in \mathbb{R}^N, \mathcal{L}_t(Z) = 0\}$ of dimension N-1 is exponentially stable.

Main Result (II^{*l*}). [General Case] Consider the system (1) with fundamental matrix R(s;t). Suppose that *b* and *A* satisfy the stability assumption (H_{stab}) . Let $\beta \in (0, \alpha)$, then there exist K > 0 and $D_* > 0$ such that for any matrix ζ satisfying $||\zeta|| < D_*$ and for any $t \in \mathbb{R}$, there exists a linear form $\psi_t : \mathbb{R}^N \to \mathbb{R}$ such that for any $Y \in \mathbb{R}^N$ and for any $s \ge t$, we have

• $||R(s;t)[Y - \psi_t(Y)\mathbb{1}|| < K||Y|| \exp(-\beta(s-t)).$

where $\mathbb{1} := (1, \ldots, 1) \in \mathbb{R}^N$. Moreover, the fundamental matrix R(s; t) admits the decomposition

$$R(s;t)Y = \psi_t(Y)R(s;t)\mathbb{1} + R(s;t)[Y - \psi_t(Y)\mathbb{1}].$$

In other words, the submanifold $W_{stab} := \{Z \in \mathbb{R}^N, \psi_t(Z) = 0\}$ of dimension N-1 is exponentially stable. Furthermore, if there exists a solution that does not exponentially decay to zero, then

•
$$\psi_t(Y) = \psi_s(R(s;t)Y).$$

1.2 Main Results: Non-linear Part

The class of mean-field coupled systems we will study is given by the following two systems: the *non-perturbed periodic system*

$$\dot{x}_i = F(X, x_i), \quad i = 1, .., N, \quad t \ge t_0,$$
 (NP)

and the *perturbed system*

$$\dot{x}_i = F(X, x_i) + H_i(X), \quad i = 1, ..., N, \quad t \ge t_0,$$
 (P)

where $N \geq 2$ and $X = (x_1, \ldots, x_N)$. $F : \mathbb{R}^N \times \mathbb{R} \to \mathbb{R}$ and $H = (H_1, \ldots, H_N) : \mathbb{R}^N \to \mathbb{R}^N$ are C^1 functions.

1.2.1 Notations and Definitions

For $q, p \in \mathbb{N}^*$, let $g := (g_1, \ldots, g_p) : \mathbb{R}^q \to \mathbb{R}^p$ be a function. Consider the following semi-norm on the vector space of continuous functions from \mathbb{R}^q to \mathbb{R}^p :

$$||g||_B = \sup_{y \in B} \max_{1 \le i \le p} |g_i(y)|,$$

where $B = \{y = (y_1, \ldots, y_q) \in \mathbb{R}^q : \max |y_i - y_j| \leq 1\}$. This semi-norm is a norm on the vector space of continuous functions from B to \mathbb{R}^p . We denote $d^i g, i = 1, 2, \ldots$, as the *i*th derivative of g. We define

$$||dg||_B = \max_{\substack{1 \le i \le p \\ 1 \le j \le q}} ||\partial_j g_i||_B, \quad ||d^2g||_B = \max_{\substack{1 \le i \le p \\ 1 \le j, k \le q}} ||\partial_k \partial_j g_i||_B.$$

In the following, we will consider functions defined on \mathbb{R}^q initially, with q = N, and then restrict them to the set B.

Exponential stability for the mean field system

Let
$$g: \mathbb{R}^N \times \mathbb{R} \to \mathbb{R}, y = (y_1, \dots, y_N) \in \mathbb{R}^N$$
, and $z \in \mathbb{R}$. We define

$$\partial_i g(y,z) = \begin{cases} \frac{\partial}{\partial z} g(y,z), & i = N+1\\ \frac{\partial}{\partial y_i} g(y,z), & i := 1, ..., N. \end{cases}$$

For $q, p \in \mathbb{N}^*$, a function $g : \mathbb{R}^q \to \mathbb{R}^p$ is called 1-*periodic* according to the following definition:

Definition 3. [1-periodicity] Let $g : \mathbb{R}^q \to \mathbb{R}^p$ be a function, and denote $\mathbb{1} := (1, \ldots, 1) \in \mathbb{R}^q$. The function g is called 1-*periodic* if

$$g(y+1) = g(y), \quad \forall y \in \mathbb{R}^q.$$

It is to note that this definition does not imply that the function G is periodic with respect to each variable individually, which allows us to study a broader class of mean-field systems.

Denote Φ^t the flow of the system (P) (including (NP)). We now define the concept of a positively Φ^t -invariant set, as follows:

Definition 4. Suppose that the flow Φ^t of the system (P) exists for all $t \ge t_0$. We say that an open set $C \subset \mathbb{R}^N$ is *positively invariant* by the flow Φ^t , or C is positively Φ^t -invariant, if $\Phi^t(C) \subset C$ for all $t \ge t_0$.

1.2.2 Synchronization Hypotheses (H) and (H_*)

We will consider the following hypotheses (H) and (H_*) about the field of the system (P) (including (NP)), which are sufficient for obtaining synchronization and stability.

$$(H) \quad \begin{cases} F \text{ is of class } C^2, & \text{and} & \max\{||F||_B, ||dF||_B, ||d^2F||_B\} < +\infty, \\ F \text{ is } \mathbb{1}\text{-periodic and} & \min_{s \in [0,1]} F(s\mathbb{1}, s) > 0, \\ (H_*) \quad \int_0^1 \frac{\partial_{N+1} F(s\mathbb{1}, s)}{F(s\mathbb{1}, s)} ds < 0. \end{cases}$$

The hypothesis (H_*) is referred to as the "synchronization hypothesis" under the condition that (H) is satisfied. The hypothesis (H) implies that the function F(s1, s) is uniformly Lipschitz. When $X = x_01$ with $x_0 \in \mathbb{R}$ is chosen as the initial condition for the solution $X(t) = (x_1(t), \ldots, x_N(t))$ of the system (NP), then each oscillator $x_i(t)$ is a solution to the scalar differential equation

$$\dot{x} = F(x\mathbb{1}, x), \quad t \ge t_0.$$

By the Cauchy-Lipschitz theorem, there exists a unique solution, so $|x_i(t) - x_j(t)| = 0 < +\infty$ for all $t \ge 0$ and $1 \le i, j \le N$. Thus, the distance between the oscillators $\{x_i(t)\}_i$ remains bounded for all times $t \ge t_0$, which is a property of synchronization. The following result is a generalization of this property to a neighborhood of $x_0 \mathbb{1}$. The hypothesis (H_*) arises from the fact that when $H \approx 0$ and $x_i \approx x_j(\approx x)$, the system (P) satisfies

$$\frac{d}{dt}x_i \approx F(x\mathbb{1}, x)$$
, and $\frac{d}{dt}(x_i - x_j) \approx \partial_{N+1}F(x\mathbb{1}, x)(x_i - x_j)$.

More precisely, in [9, 11] we have the following Theorems I and II. The method used to obtain synchronization is based on the principle of comparison of differential equations. Assuming that the distance between the oscillators of a solution remains finite at an arbitrary time provides information about the system. Then, the principle of comparison of differential equations is used, implying that this arbitrary time is actually infinity. The first theorem demonstrates the existence of a solution and the state of synchronization. The second, shows periodic locking state .

Theorem (I). Consider the system (P). Suppose that F satisfies hypotheses (H) and (H_*) . Then, there exists $D_* > 0$ such that for all $D \in (0, D_*]$, there exists r > 0 and an open set C_r of the form

$$C_r := \Big\{ X = (x_i)_{i=1}^N \in \mathbb{R}^N : \exists \nu \in \mathbb{R}, \quad \max_i |x_i - \nu| < \Delta_r(\nu) \Big\},\$$

where $\Delta_r : \mathbb{R} \to (0, D]$ is a C^1 and 1-periodic function, such that for any H of class C^1 satisfying $||H||_B < r$, the following holds:

- 1. Existence of Solution: The flow Φ^t of the system (P) exists for all $X \in C_r$ and $t \ge t_0$.
- 2. Synchronization: The open set C_r is positively invariant under Φ^t . Furthermore, for all $X \in C_r$,

$$\min_{1 \le i \le N} \inf_{t \ge t_0} \frac{d}{dt} \Phi_i^t(X) > 0 \quad and \quad |\Phi_i^t(X) - \Phi_j^t(X)| < 2D,$$

for all $1 \le i, j \le N$ and $t \ge t_0$.

Theorem (II). Consider the system (P). Suppose that F satisfies hypotheses (H) and (H_{*}). Then, there exists $D_* > 0$ such that for all $D \in (0, D_*]$, there

exists r > 0 such that for any function H of class C^1 and $\mathbb{1}$ -periodic function satisfying $||H||_B < r$, there exists an open set C_r (defined as in Theorem I) and an initial condition $X_* \in C_r$ such that

$$\Phi_{i}^{t}(X_{*}) = \rho t + \Psi_{i,X_{*}}(t), \quad \forall i = 1, ..., N, \ \forall t \ge t_{0},$$

where $\rho > 0$ and $\Psi = (\Psi_1, \ldots, \Psi_N)$ with $\Psi_i : \mathbb{R} \to \mathbb{R}$ are C^1 and $\frac{1}{\rho}$ -periodic functions.

1.2.3 Main Results

The following result I shows the stability of the system (P).

Main Result (I). Consider the system (P). Suppose that F satisfies hypotheses (H) and (H_{*}), then there exists $D_* > 0$ such that for all $D \in (0, D_*]$, there exists r > 0, and there exists an open set C_r (as defined in Theorem (I)), such that for any function H of class C^1 satisfying $\max\{||H||_B, ||dH||_B\} < r$, we have

 $\exists M > 0, \ \forall X \in C_r, \exists \delta > 0, \forall Y \in C_r, ||X - Y|| < \delta:$ $||\Phi^t(X) - \Phi^t(Y)|| < M||X - Y||, \ \forall t \ge t_0,$

Furthermore, for all $X \in C$, there exists a submanifold $W_X \subset C_r$ of dimension N-1 at X such that

$$\begin{aligned} \exists \beta > 0, \ \exists K > 0, \ \forall X \in C_r, \ \forall Y \in \mathcal{W}_X : \\ ||\Phi^t(X) - \Phi^t(Y)|| < K \exp(-\beta(t - t_0))||X - Y||, \quad \forall t \ge t_0. \end{aligned}$$

A consequence of the Theorem (II) and the Main result I results is the following result.

Main Result (II). Consider the system (P). Suppose that F satisfies hypotheses (H) and (H_{*}), then there exists $D_* > 0$ such that for all $D \in (0, D_*]$, there exists r > 0, and there exists an open set C_r (as defined in Theorem (I)) such that for any function H of class C^1 and 1-periodic satisfying $\max\{||H||_B, ||dH||_B\} < r$, there exists a submanifold $W_{stab} \subset C_r$ of dimension N - 1 such that

$$\exists \beta > 0, \ \exists K > 0, \ \forall X \in \mathcal{W}_{stab} :$$
$$||\Phi^t(X) - \rho t - \Psi(t)|| < K \exp(-\beta(t - t_0)), \ \forall t \ge t_0,$$

where $\rho > 0$ and $\Psi = (\Psi_1, \ldots, \Psi_N)$ with $\Psi_i : \mathbb{R} \to \mathbb{R}$ are C^1 functions and $\frac{1}{\rho}$ -periodic.

Remark.

- 1. The last II result shows that the periodic orbit, stated in Theorem II, is a stable limit cycle.
- 2. In general case, when the field is an trigonometric polynomial, the mathematical existence of the limits (frequencies)

$$\lim_{t \to +\infty} \frac{x_i(t)}{t}, \quad i := 1, \dots, n,$$

is proved in [8]. That addresses the mathematical existence of equation (4) posed in [12]

3. Theorem I and the Main result I can be generalized for functions H(t, X) that depend on time t.

2 Proof of Linear Results I^l and II^l : Stability of Perturbed Linear Systems

To study the stability of coupled systems, as given by equation (P) (in particular (NP)), we will investigate the stability of a class of perturbed linear systems. More precisely, we will prove the two linear results I^l and II^l from the linear part. This will allow us, in Section 3, to prove the two results I and II. These perturbed linear systems satisfy an assumption called the "stability assumption", which is sufficient to decompose the phase space into a central manifold and another exponentially stable manifold.

For simplification, we define $c_b = \max_{t \in [0,1]} |b(t)|$ and $c_a = \max_{t \in [0,1]} \sum_{j=1}^N |a_j(t)|$. To further simplify, we denote $e(t,s) := \exp(\int_s^t b(x) dx)$ and $P(t,s) = \exp(\int_s^t b(x) + \sum_{j=1}^N a_j(x) dx)$, so that $\max_{s,t \in \mathbb{R}} P(t,s) \leq \exp(c_b + c_a)$. In the following, we denote by $\langle Y, Z \rangle$ the scalar product of Y and $Z \in \mathbb{R}^N$, and by Y^T the transpose of the vector $Y \in \mathbb{R}^N$. Recall that we denoted $\mathbb{1} = (1, \ldots, 1)^T \in \mathbb{R}^N$. Let's define

$$\begin{cases} A_*(s) = (a_1(s), \dots, a_N(s)), \\ \zeta_i(s) = (\zeta_{i,1}(s), \dots, \zeta_{i,N}(s)), \quad i = 1, \dots, N. \end{cases}$$
(2)

2.1 Proof Tools

To prove the linear results I^l and II^l , we consider in this section only the stability assumption (H_{stab}) . On the other hand, this allows us to deduce result II^l . Consider the following non-homogeneous system defined for all $t \ge t'$:

$$\begin{cases} \dot{Z}^{*}(t) = b(t)Z^{*}(t) + \zeta(t)[z_{N+1}(t)\mathbb{1} + Z^{*}(t) + e(t,t')Y], \\ \dot{z}_{N+1}(t) = [b(t) + \langle A_{*}(t), \mathbb{1} \rangle]z_{N+1}(t) \\ + \langle A_{*}(t), Z^{*}(t) + e(t,t')Y \rangle, \end{cases}$$
(3)

where $Z^*(t) = (z_1(t), \ldots, z_N(t))^T$, and A_* is defined by equation (2). The goal of introducing the above system is that the part $E(t', t) = R(t; t')[Y - \psi_{t'}(Y)\mathbb{1}]$ satisfies the following decomposition:

$$E(t',t)Y = z_{N+1}(t)\mathbb{1} + Z^*(t) + e(t,t')Y,$$

where $Z(t) := (Z^*(t)^T, z_{N+1}(t))^T$ with $z_{N+1}(t)$ and $Z^*(t) = (z_1(t), \ldots, z_N(t))^T$ is a solution of the coupled non-homogeneous system (3). The idea is to show that $||Z(t)|| < K \exp(-\beta(t-t'))$ with $\alpha > \beta > 0$. Here, it is worth noting that the initial condition of Z(t) satisfies $z_{N+1}(t')\mathbb{1} + Z^*(t') = -\psi_{t'}(Y)\mathbb{1}$. The problem to solve is to find this suitable initial condition. To do this, we impose that $Z^*(t') = \psi_{t'}(Y)(\mathbb{1}^T, 0)^T$. First, we will see in this section under what assumptions we will have exponential decay of Z(t) to zero. This will allow us to find a specific initial condition $\psi_{t'}(Y)(\mathbb{1}^T, 0)^T$, which in turn will lead to the linear form $\mathcal{L}_{t'}$ in Section 2.2. We will need three lemmas, with the third lemma 7 being the main one. The following lemma enables us to prove lemma 6, which in turn allows us to prove the main lemma 7.

Lemma 5. Let $b : \mathbb{R} \to \mathbb{R}$ be a periodic function such that

$$-\alpha := \int_0^1 b(s)ds < 0.$$

Let $\alpha > \beta > 0$, L > 0, and D > 0. Consider the following equation:

$$\frac{d}{dt}\Delta(t) = [b(t) + \beta]\Delta(t) + DL$$

then there exists $D_0 > 0$ such that for all $D < D_0$, the above equation has a solution $\Delta(t)$ that is 1-periodic and strictly positive, with $\max_{t \in [0,1]} \Delta(t) < 1$. The solution $\Delta(t)$ is given by

$$\Delta(t) = DL \frac{\int_t^{t+1} \exp(\int_s^{t+1} b(x) + \beta dx) ds}{1 - \exp(\beta - \alpha)}$$

Lemma 6. Consider the system (3) with b(t) and \mathcal{A} satisfying the stability assumption (H_{stab}) . Let $\beta \in (0, \alpha)$, then there exist L > 0 and $D_* > 0$ such that for any matrix ζ that is continuous and has a norm $||\zeta|| < D_*$, for any $Y \in \mathbb{R}^N$, and for any solution $Z(t) = (Z^*(t)^T, z_{N+1}(t))^T$ of system (3) with an initial condition $Z(t') = Z \in \mathbb{R}^N$, we have:

$$\forall T > t' : z_{N+1}(T) = 0$$

$$\implies ||Z(t)|| < L \exp(-\beta(t-t'))[||Z|| + ||Y||], \ \forall t \in [t',T].$$

Proof. Let $Z(t) = (Z^*(t)^T, z_{N+1}(t))^T$, where $Z^*(t) = (z_1(t), \ldots, z_N(t))^T$ is a solution to the linear system (3) with the initial condition $Z(t') = Z \in \mathbb{R}^N$. Suppose that $z_{N+1}(T) = 0$. By integrating (3), we obtain

$$z_{N+1}(t') = -\int_{t'}^{T} \langle A_*(s), Z^*(s) + e(s, t')Y \rangle P(t', s)ds.$$
(4)

Let $\beta < \alpha$. Let C = ||Z|| + ||Y||, so for all M > 1, we have ||Z|| < MC, and there exists $\epsilon > 0$ such that $||Z^*(t)|| < MC \exp(-\beta(t-t'))$ for every $t \in (t', t' + \epsilon)$. Define,

$$T_* = \sup \Big\{ t > t' : \quad ||Z^*(s)|| < MC \exp(-\beta(s - t')), \quad \forall s \in (t', t) \Big\},$$

Then, we have

$$||Z^*(t)|| < MC \exp(-\beta(t-t')) \quad \forall t \in [t', T_*[.$$
(5)

The strategy is to find a particular constant M such that $T_* \geq T$. By contradiction, assume that $T_* < T$. By integrating (3) and using (4), we obtain for all $t \in [t', T_*]$

$$|z_{N+1}(t)| = \left| P(t,t') \left[z_{N+1}(t') + \int_{t'}^{t} \langle A_{*}(s), Z^{*}(s) + e(s,t')Y \rangle P(t',s)ds \right] \right|$$

$$= P(t,t') \left| -\int_{t}^{T} \langle A_{*}(s), Z^{*}(s) + e(s,t')Y \rangle P(t',s)ds \right|$$

$$< \exp(c_{b} + c_{a})c_{a}C \int_{t}^{T} M \exp(-(\beta s - t')) + \exp(-\alpha(s - t') + c_{b})ds$$

$$< \exp(2c_{b} + c_{a})c_{a}C \frac{2M}{\beta} \exp(-\beta(t - t')).$$
(6)

Let ζ be a continuous matrix with $||\zeta|| = D < D_*$, then from equation (3), we have the following two inequalities for $z_i(t)$ and $-z_i(t)$

$$\frac{d}{dt}z_i(t) < b(t)z_i(t) + DC[\exp(2c_b + c_a)c_a\frac{2M}{\beta} + M + \exp(c_b))]\exp(-\beta(t - t')) < b(t)z_i(t) + DMC[\exp(2c_b + c_a)\frac{2c_a}{\beta} + 1 + \exp(c_b))]\exp(-\beta(t - t')).$$
(7)

$$\frac{d}{dt} - z_i(t) < b(t)(-z_i(t)) + DC[\exp(2c_b + c_a)c_a\frac{2M}{\beta} + M + \exp(c_b))]\exp(-\beta(t - t')) < b(t)(-z_i(t)) + DMC[\exp(2c_b + c_a)\frac{2c_a}{\beta} + 1 + \exp(c_b))]\exp(-\beta(t - t')).$$

Let $z_i(t) = \Delta_i(t) \exp(-\beta(t-t'))MC$, $t \in [t', T_*[$. We have $|\Delta_i(t)| \le 1$ on $[t', T_*[$. Substituting into the last equation, we have

$$\frac{d}{dt}\Delta_{i}(t) < [b(t) + \beta]\Delta_{i}(t) + D[\exp(2c_{b} + c_{a})\frac{2c_{a}}{\beta} + 1 + \exp(c_{b}))],$$

$$\frac{d}{dt} - \Delta_{i}(t) < [b(t) + \beta](-\Delta_{i}(t)) + D[\exp(2c_{b} + c_{a})\frac{2c_{a}}{\beta} + 1 + \exp(c_{b}))].$$

By definition of T_* there exists $i \in \{1, \ldots, N\}$ such that $|\Delta_i(T_*)| = 1$. We will use Lemma 5 to obtain a contradiction.

Lemma 5 implies that for $||\zeta|| = D < D_* := D_0$, there exists a strictly positive 1-periodic function of the equation

$$\frac{d}{dt}\Delta(t) = [b(t) + \beta]\Delta(t) + D[\exp(2c_b + c_a)\frac{2c_a}{\beta} + 1 + \exp(c_b))],$$

such that $\max_{t \in [0,1]} \Delta(t) < 1$. Let M > 1 be such that $\frac{1}{M} < \Delta(t')$; thus, $\Delta_i(t') \leq \frac{||Z||}{MC} \leq \frac{1}{M} < \Delta(t')$. There exists $\epsilon' > t'$ such that $|\Delta_i(t)| < \Delta(t)$ on $[t', \epsilon']$; let $T_0 = \sup\{t' \leq s \leq T_0 : |\Delta_i(s)| < \Delta(s)\}$. If $t' < T_0 < T_*$, then $|\Delta_i(T_0)| = \Delta(T_0)$. Without loss of generality, assume that $\Delta_i(T_0) = \Delta(T_0)$,

then we obtain

$$\frac{d}{dt} |\Delta_i(T_0)| < [b(t) + \beta] |\Delta_i(T_0)| + D[\exp(2c_b + c_a)\frac{2c_a}{\beta} + 1 + \exp(c_b))]$$

= $[b(t) + \beta] \Delta(T_0) + D[\exp(2c_b + c_a)\frac{2c_a}{\beta} + 1 + \exp(c_b))]$
= $\frac{d}{dt} \Delta(T_0).$

Contradiction. Thus, $T_0 > T_*$, which implies $\Delta_i(T_*) < 1$, contradicting the definition of T_* . Therefore, for all $t \in [t', T]$, we have

$$||Z^*(t)|| < L \exp(-\beta(t-t'))[||Z(t')|| + ||Y|]]$$

for some constant $L = M \max(1, 2 \exp(2c_b + c_a)c_a)$.

The previous lemma does not provide exponential decay of a solution over $[t', +\infty[$. To achieve this, the strategy in the following lemma is to consider intervals [0, T] from the previous lemma, gradually increasing in size. We will approximate a solution $Z_Y(t)$ with an initial condition $Z_Y(t')$ using solutions that satisfy the previous lemma. This allows us to demonstrate the existence of a solution $Z_Y(t)$ that exponentially decays to zero over $[t', +\infty[$. We will also localize the initial condition $Z_Y(t')$.

Lemma 7. Consider the system (3) with b(t) and \mathcal{A} satisfying the stability assumption (H_{stab}) . Let $\beta \in (0, \alpha)$, then there exist K > 0 and $D_* > 0$ such that for any $D < D_*$ and any $Y \in \mathbb{R}^N$, if there exists a sequence of solutions $Z_m(t) = (Z_m^*(t)^T, z_{N+1,m}(t))^T$ of (3) with initial condition $Z_m(t') = z_{t',m}W$ ($z_{t',m} \in \mathbb{R}$, $W = (\mathbb{1}^T, 0)^T$) and a sequence (t_m)_m that tends to infinity such that $z_{N+1,m}(t_m) = 0$, then there exists a solution $Z_Y(t) = (Z_Y^*(t)^T, z_{N+1,Y}(t))^T$ of system (3) with initial condition $Z_Y(t') = Z_Y \in \mathbb{R}^N$ such that

$$||Z_Y(t)|| < K \exp(-\beta(t-t'))||Y||, \quad \forall t \ge t'.$$

Moreover, there exists a subsequence $(Z_{m_k}(t'))_k$ of $(Z_m(t'))_m$ such that $Z_Y(t') = \lim_{k \to +\infty} Z_{m_k}(t').$

Proof. Let $Z_m(t)$ and $(t_m)_m$ satisfy the assumptions of this lemma. Lemma 6 implies that

$$||Z_m(t)|| < L \exp(-\beta(t-t'))[||Y|| + ||Z_m(t')||], \quad \forall t \in [t', t_m].$$

The idea is to show that there exists C > 0 such that $||Z_m(t')|| < C||Y||$ and then extract a convergent subsequence of $Z_m(t)$. Let ζ be a continuous matrix with norm $||\zeta|| = D < D_*$. By integrating (3) for all $t \in [t', t_m]$ and all $i = 1, \ldots, N$, we have

$$z_{i,m}(t) = e(t, t')z_{i,m}(t') + F_{i,m}(t),$$

 $\mathrm{so},$

$$\begin{aligned} |F_{i,m}(t)| \\ &= |e(t,t') \int_{t'}^{t} < \zeta_{i}(s), z_{N+1,m}(s) \mathbb{1} + Z_{m}^{*}(s) + e(s,t')Y > e(t',s)ds| \\ &< D(2L + \exp(c_{b}))[||Y|| + ||Z_{m}(t')||]e(t,t') \int_{t'}^{t} \exp((\alpha - \beta)(s - t'))ds \\ &< D(2L + \exp(c_{b}))\exp(c_{b}) \frac{||Y|| + ||Z_{m}(t')||}{\alpha - \beta} \exp(-\beta(t - t')). \end{aligned}$$

Since $z_{N+1,m}(t_m) = 0$, and by integrating (3), we obtain

$$\begin{split} z_{N+1,m}(t_m) &= 0 \\ \iff z_{i,m}(t') \int_{t'}^{t_m} < A_*(s), \mathbb{1} > \exp(-\int_{t'}^s < A_*(x), \mathbb{1} > dx) ds \\ &+ \int_{t'}^{t_m} \sum_{i=1}^N [a_j(s)F_{i,m}(s)] P(t',s) ds \\ &= -\int_{t'}^{t_m} < A_*(s), e(s,t') Y > P(t',s) ds. \end{split}$$

We deduce that

$$\begin{aligned} |z_{i,m}(t')| \Big| 1 - \exp(-\int_{t'}^{t_m} < A_*(x), \mathbb{1} > dx) \Big| \\ - c_a D(2L + \exp(c_b)) \exp(c_b) \frac{||Y|| + |z_{t',m}|}{\alpha - \beta} \int_{t'}^{t_m} \exp(-\beta(s - t')) P(t', s) ds \\ < |\int_{t'}^{t_m} < A_*(s), e(s, t') Y > P(t', s) ds|. \end{aligned}$$

For $D \approx 0$ and $m \to +\infty$, we will have

$$|z_{i,m}(t')| < \frac{\exp(2c_b + c_a) \left[D \frac{c_a(2L + \exp(c_b))}{\beta(\alpha - \beta)} + \frac{c_a}{\alpha} \right]}{1 - c_a D(2L + c_b) \exp(2c_b + c_a) \frac{1}{\beta(\alpha - \beta)}} ||Y||,$$

which implies that $(||Z_m(t)||)_m$ is uniformly bounded on each interval $[t', t_m]$. Furthermore,

$$||Z_m(t)|| < L \exp(-\beta(t-t'))[||Y|| + ||Z_m(t')||] < K \exp(-\beta(t-t'))||Y||,$$

where

$$K = L \left[1 + \frac{\exp(2c_b + c_a) \left[D \frac{c_a(2L + \exp(c_b))}{\beta(\alpha - \beta)} + \frac{c_a}{\alpha} \right]}{1 - c_a D(2L + c_b) \exp(2c_b + c_a) \frac{1}{\beta(\alpha - \beta)}} \right].$$

Therefore, we can extract a convergent subsequence that converges to a solution $Z_Y(t)$ of (3) and satisfies

$$||Z_m(t)|| < K \exp(-\beta(t-t'))||Y||.$$

2.2 Ingredients for the linear form $\mathcal{L}_{t'}$

In this section, we will show the existence of a family of solutions to the system (3) that satisfies the assumptions of Lemma 7 in the previous section. To do this, we only need to determine the initial conditions $Z_m(t') = z_{t',m}W$. Note that it is sufficient to determine the sequence of real numbers $(z_{t',m})_m$.

In this section, we consider b(t) and \mathcal{A} satisfying only the stability assumption (H_{stab}) without distinction in the matrix ζ . This allows us, in particular, to deduce the linear result II^{l} .

To obtain more information about the solutions $Z_m(t)$ of Lemma 7, we will express them in terms of the fundamental matrix. Let $S(t;t') = \{s_{i,j}(t;t')\}_{\substack{1 \le i \le N+1 \\ 1 \le j \le N}}$ be the fundamental matrix of the homogeneous linear system associated with the system (3) as follows:

$$\begin{cases} \dot{X}^{*}(t) = b(t)X^{*}(t) + \zeta(t)[x_{N+1}(t)\mathbb{1} + X^{*}(t)], \\ X^{*}(t) = (x_{1}(t), \dots, x_{N}(t))^{T}, \\ \dot{X}_{N+1}(t) = [b(t) + \langle A_{*}(t), \mathbb{1} \rangle]x_{N+1}(t) + \langle A_{*}, X^{*}(t) \rangle, \end{cases}$$
(8)

where $A_*(s)$ is given by Equation (2). The solution $Z(t) = (Z^*(t)^T, z_{N+1}(t))^T$ of (3) with initial condition $Z(t') \in \mathbb{R}^N$

can be written in terms of the fundamental matrix as follows:

$$Z^*(t) = S^*(t;t')Z(t') + S_1(t;t')Y,$$
(9)

$$z_{N+1}(t) = P(t, t') z_{N+1}(t')$$
(10)

$$+ +P(t,t')\int_{t'}^{t} < A_{*}(s), S^{*}(s;t')Z(t') > P(t',s)ds + P(t,t')\int_{t'}^{t} < A_{*}(s), S_{1}(s;t')Y + e(s,t')Y > P(t',s)ds,$$

where $S^*(t;t') = \{s_{i,j}(t;t')\}_{\substack{1 \le i \le N \\ 1 \le j \le N}}$ and $S_1(t;t')$ is an operator that does not depend on Z(t') and Y such that S(t',t') = 0.

As mentioned earlier, we aim to show the existence of solutions to the system (3) that satisfy the assumptions of Lemma 7. In Lemma 7, we have $Z_m(t') = z_{t',m}W$, so for a sequence $(T_m)_m$ tending to infinity such that $Z_{m,N+1}(T_m) = 0$ and using the notation from the previous equation (10), the sequence of real numbers $(z_{t',m})_m$ must be defined as

$$z_{t',m} = -\frac{\int_{t'}^{T_m} \langle A_*(s), S_1(s;t')Y + e(s,t')Y \rangle P(t',s)ds}{\int_{t'}^{T_m} \langle A_*(s), S^*(s;t')W \rangle P(t',s)ds}.$$

In Lemma 8, we will show that this sequence $(z_{t',m})_m$ is well-defined, i.e., the denominator of the quotient on the right-hand side of the equation is nonzero.

Lemma 8. Let $S(t;t') = \{s_{i,j}(t;t')\}_{\substack{1 \le i \le N+1 \\ 1 \le j \le N}}$ be the fundamental matrix of the homogeneous linear system (8) associated with the system (3). Let $S^*(t;t') = \{s_{i,j}(t;t')\}_{\substack{1 \le i \le N \\ 1 \le j \le N}}$ Suppose that b(t) and \mathcal{A} satisfy the stability assumption (H_{stab}) . Define

$$H(t,t') := \int_{t'}^{t} \langle A_{*}(s), S^{*}(s;t')W \rangle P(t',s)ds, \qquad (11)$$
$$W := (\mathbb{1}^{T}, 0)^{T}, \ t \ge t'.$$

Then, for any $\beta \in (0, \alpha)$, there exists $D_* > 0$ such that for any continuous matrix ζ with norm $||\zeta|| < D_*$, there exists $T_W > 0$ such that for all $t \ge T_W$, we have $H(t, t') \neq 0$.

Proof. From equation (10), we deduce that $Z(t) = (S^*(t;t')W - z_{N+1}(t), z_{N+1}(t))$ is a solution of the linear homogeneous system (8) with the initial condition W, where $z_{N+1}(t)$

satisfies

$$z_{N+1}(t) = P(t, t')H(t, t'), \quad t \ge t'.$$

So, $Z(t) = (S^*(t;t')W - z_{N+1}(t), z_{N+1}(t))$ is, in particular, a solution of the nonhomogeneous linear equation (3) with Y = 0 and satisfies the assumptions of Lemma 7. Since Y = 0, we have $||Z(t)|| \equiv 0$, which contradicts the initial condition Z(t') = W.

Therefore, in the following proposition, we show that the system (3) admits a sequence of solutions that satisfy the assumptions of Lemma 7.

Proposition 9. Consider the system (3) with b(t) and \mathcal{A} satisfying the stability hypothesis (H_{stab}) . Let $\beta \in (0, \alpha)$. Then there exist $D_* > 0$ and K > 0, such that for any continuous matrix ζ with $||\zeta|| < D_*$ and any $Y \in \mathbb{R}^N$, there exists a solution $Z_Y(t) = (Z_Y^*(t)^T, z_{N+1,Y}(t))^T$ of (3) such that

$$|Z_Y(t)|| < K \exp(-\beta(t-t'))||Y||, \quad \forall t \ge t',$$

and $Z_Y(t') = \psi_{t'}(Y)W$, where

$$\psi_{t'}(Y) = \lim_{t_m \to +\infty} \frac{-1}{H(t_m, t')} \int_{t'}^{t_m} \langle A_*(s), S_1(s; t')Y + e(s, t')Y \rangle P(t', s)ds,$$

and where $(t_m)_m$ is a sequence of real numbers that tends to infinity, and $H(t_m, t')$ is given by Lemma 8.

Proof. According to Lemma 8, for any sequence $(t_m)_m$ such that $t_m > T_W$, we have $H(t_m, t') \neq 0$. From (10), the solution $Z_m(t) = (Z_m^*(t)^T, z_{N+1,m}(t))^T$ of (3) with the initial condition $z_{m,t'}W$ such that

$$z_{m,t'} = -\frac{1}{H(t_m, t')} \int_{t'}^{t_m} \langle A_*(s), S_1(s; t')Y + e(s, t')Y \rangle P(t', s)ds,$$

satisfies:

$$z_{N+1,m}(t_m) = 0 = P(t',t)z_{m,t'}H(t_m,t') + = P(t',t)\int_{t'}^{t_m} \langle A_*(s), S_1(s;t')Y + e(s,t')Y \rangle P(t',s)ds.$$

According to Lemma 7, there exists a solution $Z_Y(t)$ such that $||Z_Y(t)|| < L \exp(-\beta(t-t'))||Y||$ for all $t \ge t'$, with $Z_Y(t') = \psi_{t'}(Y)W$, where

$$\psi_{t'}(Y) = -\lim_{t_{m_k} \to \infty} \frac{\int_{t'}^{t_{m_k}} \langle A_*(s), S_1(s;t')Y + e(s,t')Y \rangle P(t',s)ds}{H(t_{m_k},t')}.$$

2.3 Decomposition of the fundamental matrix

Finally, we will prove results I^l and II^l in the context of linear systems. To ensure consistency in the proof, we will first demonstrate the general case, which is the second linear result II^l that does not require the matrix $\zeta(t)$ to be normalizing.

Proof of the second linear result Π^{l} : General Case.

• Let's show that $||R(t;t')[Y + \psi_{t'}(Y)\mathbb{1}]||$ decreases exponentially:

For any $\beta > 0$ and $||\zeta|| = D < D_*$, where D_* is defined by Lemma 7, let $\psi_{t'}(Y)$ be given by the previous proposition 9. Take $Y \in \mathbb{R}^N$, we add and subtract the same term in $\phi^t(Y)$ as follows:

$$\phi^{t}(Y) = -\psi_{t'}(Y)R(t;t')\mathbb{1} + R(t;t')[Y + \psi_{t'}(Y)\mathbb{1}].$$
 (12)

Let $R(t;t')[Y + \psi_{t'}(Y)\mathbb{1}] = z_{N+1}(t)\mathbb{1} + Z^*(t) + e(t;t')Y$ with $z_{N+1}(t)$ being a solution with the initial condition $z_{N+1}(t') = 0$ of the equation

$$\dot{z}_{N+1}(t) = [b(t) + \langle A_*(t), \mathbb{1} \rangle] z_{N+1}(t) + \langle A_*(t), R(t; t') [Y + \psi_{t'}(Y) \mathbb{1}] - z_{N+1}(t) \mathbb{1} \rangle.$$

Thus, $Z(t) = (Z^*(t), z_{N+1}(t))$ is a solution of the nonhomogeneous linear equation (3). From equation (12), we deduce that it has the initial condition $Z(t') = W\psi_{t'}(Y)$. Proposition 9 implies that

$$||Z(t)|| < K \exp(-\beta(t-t'))||Y||, \quad \forall t \ge t'.$$

Hence, $||R(t;t')[Y + \psi_{t'}(Y)\mathbb{1}]|| < [K + \exp(c_b)]\exp(-\beta(t-t'))||Y||$ for all $t \ge t'$.

• Let's show that when the system has a solution that does not decay to zero exponentially, then $\psi_t(Y) = \psi_s(R(s;t)Y)$:

Suppose that the system (1) has a solution that does not exponentially decay to zero. Let V(t) be this solution. We have:

$$\begin{aligned} R(t;s)R(s;t')Y &= \mathcal{L}_{t'}(Y)V(t) - R(t;t')[Y - \mathcal{L}_{t'}(Y)V(t')] \\ &= \mathcal{L}_s(R(s;t')Y)V(t) \\ &- R(t;s)[R(s;t')Y - \mathcal{L}_s(R(s;t')Y)V(s)]. \end{aligned}$$

Implies

$$\begin{aligned} [\mathcal{L}_{t'}(Y) &- \mathcal{L}_s(R(s;t')Y)]V(t) \\ &= R(t;s)[R(s;t')\mathcal{L}_{t'}(Y)V(t') + \mathcal{L}_s(R(s;t')Y)V(s)]. \end{aligned}$$

Since the right-hand side of the last equation satisfies

$$||R(t;s)[R(s;t')\mathcal{L}_{t'}(Y)V(t') + \mathcal{L}_{s}(R(s;t')Y)V(s)]|| < K[\exp(-\beta(t-t')) + \exp(-\beta(t-s))]||Y||.$$

While the left-hand side is a linear form multiplied by the function V(t), which does not decay exponentially to zero, we must have

$$\mathcal{L}_{t'}(Y) - \mathcal{L}_s(R(s;t')Y) = 0, \quad \forall t' \ge 0, \forall s \ge t'.$$

Now, we will prove the first linear result I^l . We consider the particular case where $\zeta(t)$ is a normalizing matrix. By Definition 2, the system (1) has a solution $V(t) = (v_1(t), \ldots, v_N(t))^T$ such that

$$\inf_{t\in\mathbb{R}}||V(t)||>0, \quad \text{and} \quad \sup_{t\in\mathbb{R}}||V(t)||<+\infty.$$

We denote in the following

$$\alpha_{-} = \inf_{t \in \mathbb{R}} ||V(t)|| > 0, \quad \text{and} \quad \alpha_{+} = \sup_{t \in \mathbb{R}} ||V(t)|| < +\infty.$$

We define in the proof below the linear form $\mathcal{L}_{t'}$ as

$$\mathcal{L}_{t'}(Y) = \frac{\psi_{t'}(Y)}{\psi_{t'}(V(t'))},$$

where $\psi_{t'}$ is defined by the previous proposition 9. We note that by uniqueness, $\mathcal{L}_{t'}$ is defined as

$$\mathcal{L}_{t'}(Y) = \lim_{t \to \infty} \frac{\int_{t'}^t \langle A_*(s), S_1(s)Y + e(s,t')Y \rangle P(t',s)ds}{\int_{t'}^t \langle A_*(s), S_1(s)V(t') + e(s,t')V(t') \rangle P(t',s)ds}.$$

Proof of linear result I^l .

Exponential stability for the mean field system

• Let's prove $\mathcal{L}_t(R(t;t')V(t')) = 1$: By the definition of $\mathcal{L}_{t'}$, we have $\mathcal{L}_t(V(t)) = 1$ for all $t \in \mathbb{R}$. Therefore,

$$\mathcal{L}_t(R(t;t')V(t')) = 1 = \mathcal{L}_{t'}(V(t')), \quad \forall t' \in \mathbb{R} \quad \forall t \ge t'.$$
(13)

 Construction of the linear form t: We add and subtract the same term in V(t), so we have

$$R(t;t')V(t') = V(t) = -\psi_{t'}(V(t'))R(t;t')\mathbb{1} + R(t;t')[V(t') + \psi_{t'}(V(t'))\mathbb{1}].$$
(14)

According to proposition 9, we have $||\psi_{t'}(V(t'))|| < K||V(t')|| < K\alpha_+$ for all $t' \ge 0$. Furthermore,

$$\begin{aligned} |\psi_{t'}(V(t'))|||R(t;t')\mathbb{1}|| &= ||V(t) - R(t;t')[V(t') + \psi_{t'}(V(t'))\mathbb{1}]|| \\ &> ||V(t)|| - ||R(t;t')[V(t') + \psi_{t'}(V(t'))\mathbb{1}]|| \\ &> \alpha_{-} - K \exp(-\beta(t-t'))||V(t')|| \\ &> \alpha_{-} - \alpha_{+} K \exp(-\beta(t-t')), \quad \forall t' \in \mathbb{R},. \end{aligned}$$

We integrate over a compact of length δ fixed such that $1 \ll \delta \ll +\infty$; let $t = t' + \delta$, we have from (1): $||R(t;t')\mathbb{1}|| \ll \exp((c_b + c_a + D_*)\delta)$

$$|\psi_{t'}(V(t'))| > \frac{\alpha_- - \alpha_+ K \exp(-\beta\delta)}{\exp((2\alpha + c_b + c_a + D)\delta)} > 0, \ \forall t' \in \mathbb{R}, \ \forall t \ge t'.$$
(15)

Let's define

$$\mathcal{L}_{t'}(Y) = \frac{\psi_{t'}(Y)}{\psi_{t'}(V(t'))}.$$

- From (15) and according to proposition 9, there exists $K_1 > 0$ such that $\mathcal{L}_{t'}(Y) < K_1 ||Y||$.
- Exponential decay: According to equation (13) for all $Y \in \mathbb{R}$

$$\begin{split} \phi^{t}(Y) &= \mathcal{L}_{t'}(Y)V(t) - R(t;t')[Y - \mathcal{L}_{t'}(Y)V(t')] \\ &= \mathcal{L}_{t'}(Y)V(t) \\ &+ R(t;t')[Y - \psi_{t'}(Y)\mathbb{1} - (\mathcal{L}_{t'}(Y)V(t') - \frac{\psi_{t'}(Y)}{\psi_{t'}(V(t'))}\psi_{t'}(V(t'))\mathbb{1})] \\ &= \mathcal{L}_{t'}(Y)V(t) \\ &+ R(t;t')[Y - \psi_{t'}(Y)\mathbb{1} - (\mathcal{L}_{t'}(Y)V(t') - \psi_{t'}(\mathcal{L}_{t'}(Y)V(t'))\mathbb{1})]. \end{split}$$

According to the previously demonstrated linear result II^l , there exists $K_* > 0$ such that we have

$$||R(t;t')[Y - \psi_{t'}(Y)\mathbb{1}]|| < K_* \exp(-\beta(t-t'))||Y||, \quad \forall t \ge t' \\ ||R(t;t')[\mathcal{L}_{t'}(Y)V(t') - \psi_{t'}(\mathcal{L}_{t'}(Y)V(t'))\mathbb{1}] < K_* \exp(-\beta(t-t'))||Y||,$$

• Finally, let's show that $\mathcal{L}_{t'}(Y) = \mathcal{L}_s(R(s;t')Y)$: We have

$$R(t;s)R(s;t')Y = \mathcal{L}_{t'}(Y)V(t) - R(t;t')[Y - \mathcal{L}_{t'}(Y)V(t')]$$

= $\mathcal{L}_s(R(s;t')Y)V(t)$
- $R(t;s)[R(s;t')Y - \mathcal{L}_s(R(s;t')Y)V(s)].$

Therefore,

$$\begin{aligned} [\mathcal{L}_{t'}(Y) - \mathcal{L}_s(R(s;t')Y)]V(t) \\ &= R(t;s)[R(s;t')\mathcal{L}_{t'}(Y)V(t') + \mathcal{L}_s(R(s;t')Y)V(s)]. \end{aligned}$$

Since $\min_{t \in \mathbb{R}} ||V(t)|| = \alpha_{-} > 0$ and as the right-hand side of the last equation verifies

$$||R(t;s)[R(s;t')\mathcal{L}_{t'}(Y)V(t') + \mathcal{L}_{s}(R(s;t')Y)V(s)]|| < K[\exp(-\beta(t-t')) + \exp(-\beta(t-s))]||Y||.$$

Thus,

$$\mathcal{L}_{t'}(Y) - \mathcal{L}_s(R(s;t')Y) = 0, \quad \forall t' \ge 0, \forall s \ge t'.$$

We have studied a class of linear systems perturbed by a time-dependent matrix ζ . We have shown that the space \mathbb{R}^N decomposes into a direct sum of subspaces $\mathbb{R}\mathbb{1} \oplus \mathcal{W}$ where \mathcal{W} is a manifold of dimension N-1 and exponentially stable. The vector subspace $\mathbb{R}\mathbb{1}$ can itself be an exponentially stable manifold. In the case where the perturbing matrix ζ is normalizing, we obtained information regarding the norm of the linear form. In the next section, we will use these results to demonstrate the stability of the nonlinear mean-field systems studied.

3 Proof of results I and II: Stability of mean-field systems

In this section, we prove the two results I and II. We will study the stability of coupled systems given by equations (P) (in particular (NP)) and exhibiting a synchronized state. We linearize the system around a synchronized orbit and then apply the stability results obtained in Section 2. Recall that the two systems (P) and (NP) are given by the following two equations:

$$\dot{x}_i = F(X, x_i), \quad 1 \le i \le N, \quad t \ge t_0, \tag{NP}$$

$$\dot{x}_i = F(X, x_i) + H_i(X, x_i), \quad 1 \le i \le N, \quad t \ge t_0,$$
 (P)

where $N \geq 2$ and $t_0 \in \mathbb{R}$ is the initial time. $F : \mathbb{R}^N \times \mathbb{R} \to \mathbb{R}$ and $H_i : \mathbb{R}^N \to \mathbb{R}$ are C^1 functions, and we denote $H = (H_1, \ldots, H_N)$. Recall that we denoted Φ^t the flow of the system (P) (including (NP)).

We have seen [9, 11] that when the norm $||H||_B$ is sufficiently small and under the assumptions (H) and (H_{*}), the global phase dispersion of a solution of the system (P) (and (NP)) with an initial condition in the synchronization region remains uniformly bounded by some constant D > 0.

3.1 Linearization of System (P)

In order to simplify the notation, define the function $F_i : \mathbb{R}^N \times \mathbb{R} \to \mathbb{R}$ as

$$F_i(Y,z) := F(Y,z) + H_i(Y,z), \quad \forall Y \in \mathbb{R}^N, \ \forall z \in \mathbb{R}, \quad \forall 1 \le 1 \le N,$$

and denote the differential of the vector function (F_1, \ldots, F_N) as $d\mathcal{F}(t)$.

Without loss of generality, for $Z \in \mathbb{R}^N$, we'll write $Z := \Phi^t(Z)$, so the elements of the matrix $dF(\Phi^t(Z)) := \{g_{i,j}\}_{i,j}$ are given by

$$\begin{cases} g_{ii}(t) = \partial_{N+1} F_i(Z, z_i) + \partial_i F_i(Z, z_i), \ 1 \le i \le N, \\ g_{ij}(t) = \partial_j F_i(Z, z_i), \ 1 \le i \ne j \le N. \end{cases}$$
(16)

The strategy in the following is to apply the linear stability results obtained in Section 2. To do this, we'll choose an appropriate linearization of the system (P). More specifically, for any $Z \in \mathbb{R}^N$, we consider the following linear system:

$$\frac{d}{dt}Y(t) = d\mathcal{F}(\Phi^t(Z))Y(t), \quad t \ge t_0, \quad Y(t) = (y_1(t), \dots, y_N(t))^T.$$
(17)

Also, recall that for $Z \in \mathbb{R}^N$, we denote $\mu_Z := \mu_Z(t)$ as the solution of the system (SNP) associated with $\Phi^t(Z)$ and with the initial condition $\mu_Z(t_0) \in \mathbb{R}$ (see Definition 10).

In the following lemma, we'll see that the system (17) can be written in the form of the linear systems (1). We denote I_N as the identity matrix of order N.

Definition 10. For $X \in \mathbb{R}^N$, we define the System (SNP) associated with $\Phi^t(X)$ as the unperturbed system given by

$$\dot{\mu}_X = F(\Phi^t(X), \mu_X), \quad t \in I_X, \tag{SNP}$$

where $I_X = [t_0, T_X)$ is the maximal interval of the solution $X(t) := \Phi^t(X)$ of the system (P) with initial condition $\phi^{t_0}(X) = X$ fore some $t_0 \in I_X$. We refer to $\mu_X(t)$ as the solution of the system (SNP) associated with $\Phi^t(X)$ with initial condition $\mu_X(t_0) \in \mathbb{R}$.

Lemma 11. Consider the coupled nonlinear system (P). Suppose that F satisfies hypotheses (H) and (H_{*}). Then, there exists $\epsilon_* > 0$ such that for all $\epsilon \in (0, \epsilon_*]$, there exists r > 0 such that for any function H satisfying $\max\{||H||_B, ||dH||_B\} < r$, there exists an open set C_r that is positively Φ^t -invariant, and for all $Z \in C_r$, there exists a diffeomorphism $t \to \mu = \mu_Z(t)$ such that the linearized system (17) is equivalent to the following linear system:

$$\frac{d}{d\mu}Y^{*}(\mu) = [b(\mu)I_{N} + \mathcal{A}(\mu) + \zeta_{Z}(\mu)]Y^{*}(\mu), \qquad (18)$$
$$\mu \ge \mu_{Z}(t_{0}), \quad Y^{*}(\mu) = Y(\tau_{Z}(\mu)),$$

where $\tau_Z : \mathbb{R} \to \mathbb{R}$ is the inverse function of $\mu(t)$, $\zeta_Z(\mu)$ is a normalizing matrix satisfying $||\zeta_Z|| < \epsilon$, $\mathcal{A}(\mu) = \{a_{i,j}(\mu) = a_j(\mu)\}_{1 \le i,j \le N}$ is a rank-1 matrix, and $b(\mu)$ is a function defined as follows:

$$a_j(\mu) = \frac{\partial_j F(\mu \mathbb{1}, \mu)}{F(\mu \mathbb{1}, \mu)}, \quad 1 \le j \le N \quad and \quad b(\mu) = \frac{\partial_{N+1} F(\mu \mathbb{1}, \mu)}{F(\mu \mathbb{1}, \mu)},$$

satisfying

$$\int_0^1 b(s) + \sum_{j=1}^N a_j(s) ds = 0, \quad and \quad \int_0^1 b(s) ds < 0,$$

meaning that the stability hypothesis (H_{stab}) is satisfied.

Proof. We consider the system (P). Suppose that F satisfies hypotheses (H) and (H_*) . Let $D \approx 0$, then according to the Theorem (I), there exists r > 0 such that for a C^1 matrix H satisfying $||H||_B < r$, there exists an open set C_r that is positively Φ^t -invariant, such that for all $t \geq t_0$ and all $Z \in C_r$, we have

$$\max_{j} |\Phi_{j}^{t}(Z) - \mu_{Z}(t)| < D \quad \text{and} \quad \max_{i,j} |\Phi_{i}^{t}(Z) - \Phi_{j}^{t}(Z)| < 2D;$$

where $\mu_Z(t)$ is a solution of the system (SNP) associated with $\Phi^t(Z)$ and with the initial condition $\mu_Z(t_0) \in \mathbb{R}$. As long as $Z \in C_r$, note that the flow $\Phi^t(Z)$ and the solution $\mu_Z(t)$ are defined for all $t \ge t_0$.

Consider the linearized system (17) at the point $\Phi^t(Z)$. For any $1 \leq i, j \leq N$, define $U_Z(t) = \{u_{i,j}^Z(t)\}_{1 \leq i,j \leq N}$ as the matrix of order N given by

$$\begin{cases} u_{ii}^{Z}(t) = \left[\partial_{N+1}H_{i}(Z, z_{i}) + [\partial_{N+1}F(Z, z_{i}) - \partial_{N+1}F(\mu_{Z}(t)\mathbb{1}, \mu_{Z}(t))]\right] \\ + \partial_{i}H_{i}(Z, z_{i}) + [\partial_{i}F(Z, z_{i}) - \partial_{i}F(\mu_{Z}(t)\mathbb{1}, \mu_{Z}(t))] \\ u_{i,j}^{Z}(t) = \partial_{j}H_{i}(Z, z_{i}) + [\partial_{j}F(Z, z_{i}) - \partial_{j}F(\mu_{Z}(t)\mathbb{1}, \mu_{Z}(t))], \end{cases}$$

so that

$$\dot{Y}(t) = [d\mathcal{F}(\mu_Z(t)\mathbb{1}) + U_Z(t)]Y(t).$$

Or equivalently,

$$\dot{y}_{i}(t) = \partial_{N+1} F(\mu_{Z}(t)\mathbb{1}, \mu_{Z}(t))y_{i}(t)$$

$$+ \sum_{j=1}^{N} [\partial_{j} F(\mu_{Z}(t)\mathbb{1}, \mu_{Z}(t)) + u_{ij}^{Z}(t)]y_{j}(t).$$
(19)

Let L > 0 and $\alpha > 0$ be defined as follows:

$$L := \sum_{0 \le i \le 2} ||d^i F||_B, \text{ and } \alpha := \min_{s \in [0,1]} F(s1,s).$$
(20)

Assume that $\max\{||H||_B, ||dH||_B\} < r$, then for all $1 \le i, j \le N$, we have

$$||u_{i,i}^Z|| < 2[r + LD], \text{ and } ||u_{i,j}^Z|| < r + LD.$$
 (21)

For $D \approx 0$, we have $\dot{\mu}_Z > \alpha - LD > 0$. In particular, $\mu_Z(t)$ is a diffeomorphism, and there exists L > 0 such that

$$\frac{F(\mu_Z \mathbb{1}, \mu_Z)}{\dot{\mu}_Z} = 1 + \theta(t),$$
$$|\theta(t)| = \left|\frac{F(\mu_Z \mathbb{1}, \mu_Z) - F(\Phi^t(Z), \mu_Z)}{\dot{\mu}_Z}\right| < \frac{LD}{\alpha - LD}.$$

Let $\mu \to \mu = \mu_Z(t)$ be the change of variable and define $Y^*(\mu) = Y(\tau_Z(\mu))$, where $\tau_Z(\mu) := \mu_Z^{-1}(\mu)$ is the inverse function of $\mu(t)$. We can use the following equation in (19):

$$\frac{d}{dt}Y(t) = \frac{d}{ds}Y^*(s)\frac{d}{dt}\mu_Z(t) = \frac{d}{ds}Y^*(s)\frac{F(\mu_Z(t)\mathbb{1},\mu_Z(t))}{\mathbb{1}+\theta(t)}.$$

Taking into account equations (21) and (19), there exists a matrix $\zeta_Z(\mu) = \{\zeta_{ij}^Z(\mu)\}_{1 \le i,j \le N}$ that depends on $\mu \in \mathbb{R}$, defined as

$$\zeta_{ij}^Z(\mu) = d\mathcal{F}(\Phi^t(Z)) \frac{\theta(t)}{F(\mu \mathbb{1}, \mu)} + u_{i,j}^Z(t) \frac{1}{F(\mu \mathbb{1}, \mu)},$$

and satisfying

$$||\zeta_Z|| < (L+r)\frac{||\theta||}{\alpha} + (r+LD)\frac{1}{\alpha} < \epsilon := (L+r)\frac{LD}{\alpha-LD} + (r+LD)\frac{1}{\alpha},$$

such that the system (17) is equivalent to

$$\frac{d}{d\mu}y_{i}^{*}(\mu) = \frac{\partial_{N+1}F(\mu\mathbb{1},\mu)}{F(\mu\mathbb{1},\mu)}y_{i}^{*}(\mu) + \sum_{j=1}^{N}\left[\frac{\partial_{j}F(\mu\mathbb{1},\mu)}{F(\mu\mathbb{1},\mu)} + \zeta_{ij}^{Z}(\mu)\right]y_{j}^{*}(\mu) + \sum_{j=1}^{N}\left[\frac{\partial_{j}F(\mu)}{F(\mu\mathbb{1},\mu)} + \zeta_{ij}^{Z}(\mu)\right]y_{j}^{*}(\mu) + \sum_{j=1}^{N}\left[\frac{\partial_{j}F(\mu)}{F(\mu)} + \zeta_{ij}^{*}(\mu)\right]y_{j}^{*}(\mu) + \sum_{j=1}^{N}\left[\frac{\partial_{j}F(\mu)}{F(\mu)} + \zeta_{ij}^{*}(\mu)\right]y_{j}^{*}(\mu) + \sum_{j=1}^{N}\left[\frac{\partial_{j}F(\mu)}{F(\mu)} + \zeta_{ij}^{*}(\mu)\right]y_{j}^{*}(\mu) + \sum_{j=1}^{N}\left[\frac{\partial_{j}F(\mu)}{F(\mu)} + \zeta_{ij}^{*}(\mu)\right]y_{j}^{*}(\mu) +$$

with $\mu \ge \mu_Z = \mu_Z(t_0)$. It can be observed that

$$V_Z^*(\mu) := \left(F_1(\Phi^{\mu_Z^{-1}(\mu)}(Z)), \dots, F_N(\Phi^{\mu_Z^{-1}(\mu)}(Z))\right) = \frac{d}{dt} \Phi^{\mu_Z^{-1}(\mu)}(Z),$$

is a solution of (18) and satisfies

$$\min_{\mu \ge \mu_Z} V_Z^*(\mu) \ge \alpha - LD - r > 0 \quad \text{and} \quad ||V_Z^*|| \le \max_{\mu \in [0,1]} F(\mu \mathbb{1}, \mu) + LD + r.$$

Therefore, the matrix ζ_Z is a normalizing matrix. Furthermore, the two hypotheses (H) and (H_*) imply that

$$\int_0^1 \frac{\partial_{N+1} F(\mu \mathbb{1}, \mu)}{F(\mu \mathbb{1}, \mu)} d\mu < 0 \text{ and } \int_0^1 \frac{\partial_{N+1} F(\mu \mathbb{1}, \mu)}{F(\mu \mathbb{1}, \mu)} + \frac{\partial_j F(\mu \mathbb{1}, \mu)}{F(\mu \mathbb{1}, \mu)} d\mu = 0.$$

This implies that the stability hypothesis (H_{stab}) is satisfied.

 \Box

3.2 Proof of Main Result (I) and (II): Stability

The proof of Main Result I and II relies on a perturbative method as follows:

Proof. Consider the coupled non-linear system (P), and suppose that F satisfies hypotheses (H) and (H_{*}). According to Lemma 11, there exists $\epsilon_* > 0$ such that for all $\epsilon \in (0, \epsilon_*]$, there exists r > 0 such that for any function H satisfying max{ $||H||_B, ||dH||_B$ } < r for $Z \in C_r$, the system (17) is equivalent to the system (18) with b(t) and \mathcal{A} satisfying the stability hypothesis (H_{stab}), and ζ being a normalizing matrix satisfying $||\zeta_Z|| < \epsilon$.

So, let $Z \in C_r$, and let $R_Z(\mu; \mu_Z)$ be the fundamental matrix of the system (18), where μ is the change of variable $t \to \mu_Z(t)$ with $\mu_Z(t)$ being a solution of the system (SNP) associated with $\Phi^t(Z)$ and with initial condition $\mu_Z := \mu_Z(t_0) \in \mathbb{R}$ as defined in definition 10. According to the linear result I^l , we deduce that there exist $\beta > 0$ and K > 0, and there exists $r_* > 0$ such that for a function H satisfying max $\{||H||_B, ||dH||_B\} < r_*$ and for all $Z \in C_{r_*}$, there exists a linear form $\mathcal{L}_{\mu_Z} : \mathbb{R}^N \to \mathbb{R}$ such that for all $Y \in \mathbb{R}^N$ and for all $\mu \ge \mu_Z$, we have

• $\mathcal{L}_{\mu_Z}(R_Z(\mu;\mu_Z)V_Z^*(\mu_Z)) = 1$ and $\mathcal{L}_{\mu_Z}(Y) < K||Y||,$

•
$$\mathcal{L}_{\mu_Z}(Y) = \mathcal{L}_{\mu_Z}(R_Z(\mu;\mu_Z)Y),$$

•
$$||R_Z(\mu;\mu_Z)[Y - \mathcal{L}_{\mu_Z}(Y)V_Z^*(\mu_Z)]|| < K||Y||\exp(-\beta(\mu - \mu_Z)).$$

Furthermore, the solution $Y^*(\mu)$ with initial condition $Y^*(\mu_Z)$ can be written as follows:

$$R_Z(\mu;\mu_Z)Y^*(\mu) = \mathcal{L}_{\mu_Z}(Y^*(\mu_Z))V_Z^*(\mu) + R_Z(\mu;\mu_Z)[Y^*(\mu_Z) - \mathcal{L}_{\mu_Z}(Y^*(\mu_Z))V_Z^*(\mu_Z)].$$

Let $S_Z(t;t_0)$ be the fundamental matrix of the system (17). Since the solution Y(t) of the system (17) with initial condition Y(t') = Y satisfies $Y(t) = Y(\tau_Z(\mu)) = Y^*(\mu)$, it follows that $Y^*(\mu_Z) = Y$ where $\mu \to \tau_Z(\mu)$ is the inverse of $t \to \mu_Z(t)$, then by the change of variable $\mu := \mu_Z(t) \to t$, we obtain that Y(t) can be written as follows:

$$Y(t) = S_Z(t;t_0)Y = Y^*(\mu) = R_Z(\mu_Z(t);\mu_Z(t_0))Y$$

= $\mathcal{L}_{\mu_Z}(Y)V_Z(t)$
+ $R_Z(\mu_Z(t);\mu_Z(t_0))[Y - \mathcal{L}_{\mu_Z}(Y)V_Z(t)],$ (22)

for all $t \ge t_0$, where $V_Z(t) = V_Z^*(\mu_Z(t)) = \frac{d}{dt}\Phi^t(Z)$. On the other hand, the system (P) can also be written in the form:

$$\frac{d}{dt}\Phi^t(Z) = F(\Phi^t(Z)) \implies \frac{d}{dt}d\Phi^t(Z) = dF(\Phi^t(Z))d\Phi^t(Z)$$

Therefore, $d\Phi^t(Z) = S_Z(t; t_0)$. Item 1. Stability:

Let $X, Y \in C_{r_*}$ such that the segment z(s) = (1 - s)X + sY satisfies $z(s) \in C_{r_*}$ for all $s \in [0, 1]$. We have:

$$\Phi^{t}(Y) - \Phi^{t}(X) = \int_{0}^{1} \frac{d\Phi^{t}(z(s))}{ds} ds = \int_{0}^{1} d\Phi^{t}(z(s)) ds(Y - X)$$
$$= \int_{0}^{1} S_{z(s)}(t; t_{0}) ds(Y - X).$$

According to the decomposition (22) and knowing that $||V_{z(s)}|| < ||F||_B + ||H||_B$, we obtain:

$$\begin{split} ||\Phi^{t}(Y) - \Phi^{t}(X)|| &\leq \left| \left| \int_{0}^{1} \mathcal{L}_{\mu_{z(s)}}(Y - X) V_{z(s)}(t) ds \right| \right| \\ &+ \left| \left| \int_{0}^{1} R_{z(s)}(\mu_{z(s)}(t); \mu_{z(s)}(t_{0})) \left[(Y - X) - \mathcal{L}_{\mu_{z(s)}}(Y - X) V_{z(s)} \right] ds \right| \right| \\ &\leq K[||F||_{B} + ||H_{B}||] ||Y - X|| + K \exp\left(-\beta(\mu_{z(s)}(t) - \mu_{z(s)}(t_{0})) \right) ||Y - X|| \\ &\leq K[||F||_{B} + ||H_{B}|| + 1] ||Y - X||, \quad \forall t \geq t_{0}. \end{split}$$

Item 2. Exponentially Stable Submanifold:

Let $X \in C_{r_*}$. In the following, we show that there exists an exponentially stable submanifold at X. The idea is to find trajectories of the form z(s)connecting z(0) := X and $z(1) := Y \in C_r$ such that for all $s \in [0, 1]$, the quantity $\frac{dz(s)}{ds}$ lies in the kernel of the linear form $\mathcal{L}_{\mu_{z(s)}}$. This allows us to use equation (22) and cancel the term that does not exponentially decay to zero as follows:

$$\begin{split} ||\Phi^{t}(Y) - \Phi^{t}(X)| &= ||\int_{0}^{1} d\Phi^{t}(z(s))ds \frac{dz(s)}{s}|| \\ &\leq ||\int_{0}^{1} \mathcal{L}_{\mu_{z(s)}}(\frac{dz(s)}{s})V_{z(s)}(t)ds|| \\ &+ ||\int_{0}^{1} R_{z(s)}(\mu_{z(s)}(t);\mu_{z(s)})\Big((\frac{dz(s)}{s}) - \mathcal{L}_{\mu_{z(s)}}(\frac{dz(s)}{s})V_{z(s)}\Big)ds|| \\ &= \int_{0}^{1} R_{z(s)}(\mu_{z(s)}(t);\mu_{z(s)})\Big((\frac{dz(s)}{s}) - \mathcal{L}_{\mu_{z(s)}}(\frac{dz(s)}{s})V_{z(s)}\Big)ds \\ &\leq \exp(-\frac{\beta}{\alpha - LD}(t - t_{0}))\max_{s \in [0,1]} ||\frac{dz(s)}{ds}||, \end{split}$$
(23)

where $\alpha - LD > 0$ since $\dot{\mu}_Z > \alpha - LD$ as given in the proof of the previous lemma 11. To ensure that $\frac{dz(s)}{ds}$ lies in the kernel of the linear form $\mathcal{L}_{\mu_{z(s)}}$, it is sufficient that $z(s) := z(\xi, s)$ satisfies the following equation:

$$\frac{d}{ds}z(\xi,s) = \xi - \mathcal{L}_{\mu_{z(\xi,s)}}(\xi)V_{z(\xi,s)}(t_0),$$
(24)

with initial condition $Z(\xi, 0) = X$ where $\xi \in \mathbb{R}^N$. For $||\xi|| \approx 0$, the solution $z(\xi, s)$ exists for all $s \in [0, 1]$. In the rest of this proof, we consider vectors ξ with norms close to zero such that $z(\xi, s) \in C_{r_*}$ for all $s \in [0, 1]$. As $\mathcal{L}_{\mu_Z}(V_{z(\xi,s)}(t_0)) = 1$, we eventually obtain $\mathcal{L}_{\mu_{z(\xi,s)}}(\xi)(\frac{d}{ds}z(\xi, s)) = 0$, which implies equation (23) and concludes the proof. We consider vectors $\xi \in \mathbb{R}^{N-1}$ in the hyperplane defined by $\mathcal{L}_{\mu_X}(\xi) = 0$. We now show that the map $\xi \to \frac{d}{d\xi}z(\xi, 1) \approx Id$ with $Id : \mathbb{R}^{N-1} \to \mathbb{R}^N$ is the identity matrix, i.e., a C^1 -diffeomorphism, which defines the exponentially stable submanifold. We have:

$$\frac{d}{\xi}\frac{d}{s}z(\xi,s) = Id_N$$

$$- \frac{d}{d\xi} \Big[\mathcal{L}_{\mu_{z(\xi,s)}}(\xi) \Big] V_{z(\xi,s)}(t_0) - \mathcal{L}_{\mu_{z(\xi,s)}}(\xi) \frac{d}{d\xi} \Big[V_{z(\xi,s)}(t_0) \Big].$$
(25)

The differentiability of the flow with respect to each variable implies that the linear form \mathcal{L}_{μ_Z} is differentiable with respect to Z. Since \mathcal{L}_{μ_Z} is linear with respect to ξ , by Taylor's theorem, there exists a matrix Π of size $N - 1 \times N$ and $s' \in [0, s]$ such that:

$$\mathcal{L}_{\mu_{z(\xi,s)}}(\xi) = \mathcal{L}_{\mu_X}(\xi) + \Pi \xi(z(\xi,s) - X)$$

= $\Pi \xi(z(\xi,s) - Z(\xi,0)) = \Pi \xi \frac{d}{ds} z(\xi,s')s.$

Therefore, $\mathcal{L}_{\mu_{z(\xi,s)}}(\xi)$ is of second order with respect to ξ and has the form:

$$\mathcal{L}_{z(\xi,s)}(\xi) \approx \xi^2$$

For $\xi \approx 0$, there exists M > 0 such that $||\frac{d}{d\xi}\mathcal{L}_{z(\xi,s)}(\xi)|| < M||\xi||$. Considering that:

$$\frac{d}{d\xi} \Big[V_{z(\xi,s)}(t_0) \Big] = \frac{d}{\xi} F(z(\xi,s)) = dF(\Phi^t(z(\xi,s))) \frac{d}{\xi} z(\xi,s),$$

and as $||dF(\Phi^t(z(\xi, s)))|| < L$ (as given in equation (20)), we can obtain $||\frac{d}{\xi}z(\xi, s)|| < M'$. Finally, there exists a constant M'' > 0 such that equation (25) satisfies:

$$||\frac{d}{\xi}\frac{d}{s}z(\xi,s) - Id_N|| < M''\xi,$$

for $\xi \approx 0$. For such ξ , the function:

$$z(\xi,1) = X + \int_0^1 \frac{d}{\xi} \frac{d}{s} z(\xi,s) ds,$$

is a C^1 -diffeomorphism, which defines an exponentially stable submanifold \mathcal{W}_{stab} of dimension N-1 such that the points in \mathcal{W}_{stab} satisfy equation (23), i.e., for all $Y \in \mathcal{W}_{stab}$:

$$||\Phi^t(Y) - \Phi^t(X)|| < K \exp(-\frac{\beta}{\alpha - LD}(t - t_0))$$

for all $t \ge t_0$, with K > 0 being an upper bound of $K(\xi) := \max_{s \in [0,1]} ||\frac{dz(s,\xi)}{ds}||$ for $\xi \approx 0$, and $Y = z(\xi, 1)$.

In conclusion, this section has highlighted the relationship between the synchronization hypothesis and the stability hypothesis. The coupled system (P) (or (NP)) satisfying the synchronization hypothesis can be linearized around synchronized orbits and satisfy the stability hypothesis, which allows us to deduce stability and exponential stability of the coupled system. In the case of the system (NP) and when the perturbation of the system (P) is 1-periodic, we deduce that the periodic orbit mentioned in result II is exponentially stable. In other words, we have ultimately deduced the existence of a stable limit cycle.

References

- J.T. Ariaratnam and S.H. Strogatz, Phase Diagram for the Winfree Model of Coupled Nonlinear Oscillators, Phys. Rev. Lett, 2001, v 86, pp. 4278
- [2], Ko, Dongnam and Ha, Seung-Yeal and Lee, Euntaek and Shim, Woojoo, Collective behaviors of stochastic agent-based models and applications to finance and optimization, Mathematical Models and Methods in Applied Sciences, v. 33, n. 07, pp. 1373-1436,(2023),
- [3] Ko, Dongnam and Ha, Seung-Yeal and Yoon, Jaeyoung, Emerging Asymptotic Patterns in a Winfree Ensemble with Higher-Order Couplings, 2023, Springer-Verlag, Berlin, Heidelberg, v. 185, nu. 1, Acta Appl. Math. (2023)
- [4] Kang, Myeongju and Rehmeier, Marco, Emergence of phase-locked states for a deterministic and stochastic Winfree model with inertia, Communications in Mathematical Sciences, v. 21, n. 7, (2023)
- [5] Seung-Yeal Ha and Doheon Kim and Bora Moon, Interplay of random inputs and adaptive couplings in the Winfree model, Communications on Pure and Applied Analysis, 2021, v 20, n 11, pp.3975-4006
- [6] S.Y. Ha and J. Park and S. W. Ryoo, Emergence of phase-locked states for the Winfree model in a large coupling regime, Discrete and Continuous Dynamical Systems, Series A, 35, 2015, v 8, pp. 3417–3436
- [7] Y. Kuramoto, International Symposium on Mathematical Problems in Theoretical Physics, Lecture Notes in Physics, Springer, New York, 1975, v 39, pp. 420–20
- [8] W. Oukil, Ordinary differential equations defined by a trigonometric polynomial field: behaviour of the solutions, Dynamical Systems,v 38, nu 2, pp. 217-235, Taylor & Francis, (2023)
- [9] Oukil, W, Synchronization in abstract mean field models, eid = arXiv:1703.07692
- [10] W. Oukil and Ph. Thieullen and A. Kessi, Invariant cone and synchronization state stability of the mean field models, Dynamical Systems, v 34, nu 3, pp. 422-433, Taylor & Francis, (2019)

- [11], W. Oukil and A. Kessi and Ph. Thieullen, Synchronization hypothesis in the Winfree model, Dynamical Systems, v 32, nu 3, pp. 326-339, 2017, publisher = Taylor& Francis,
- [12] Sakaguchi, Hidetsugu and Shinomoto, Shigeru and Kuramoto, Yoshiki, Local and Grobal Self-Entrainments in Oscillator Lattices", Progress of Theoretical Physics, v. 77, n. 5, pp. 1005-1010, (1987)
- [13] A. T. Winfree, Biological rhythms and the behavior of populations of coupled oscillators, J. Theor. Bio, 1967, v 16, pp. 15–42